

Research Article

Quantitative Global Estimates for Generalized Double Szász-Mirakjan Operators

Mehmet Ali Özarslan and Hüseyin Aktuğlu

Eastern Mediterranean University, Gazimagusa, Cyprus, Mersin 10, Turkey

Correspondence should be addressed to Mehmet Ali Özarslan; mehmetali.ozarslan@emu.edu.tr

Received 15 December 2012; Accepted 8 May 2013

Academic Editor: Jingxin Zhang

Copyright © 2013 M. Ali Özarslan and H. Aktuğlu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce the generalized double Szász-Mirakjan operators in this paper. We obtain several quantitative estimates for these operators. These estimates help us to determine some function classes \mathcal{S} (including some Lipschitz-type spaces) which provide uniform convergence on the whole domain $[0, \infty) \times [0, \infty)$.

1. Introduction

The well-known Szász-Mirakjan operators are defined on the space \mathcal{A}_1 as follows:

$$S_n(f; x) = e^{-nx} \sum_{k=0}^{\infty} f\left(\frac{k}{n}\right) \frac{(nx)^k}{k!}, \quad (1)$$

where \mathcal{A}_1 is the set of all real functions on $[0, \infty)$ such that the right-hand side in (1) make sense for all $n > 0$ and $x \in [0, \infty)$. By modifying the Szász-Mirakjan operators as

$$D_n(f; x) = e^{-nu_n(x)} \sum_{k=0}^{\infty} f\left(\frac{k}{n}\right) \frac{(nu_n(x))^k}{k!}, \quad (2)$$

where $\{u_n(x)\}$ is a sequence of real-valued, continuous functions defined on $[0, \infty)$ with $0 \leq u_n(x) < \infty$, it has been shown in [1] that if one let

$$u_n^*(x) := \frac{-1 + \sqrt{4n^2x^2 + 1}}{2n}, \quad n \in \mathbb{N}, \quad (3)$$

then the operators defined by

$$D_n^*(f; x) := S_n(f; u_n^*(x)) \quad (4)$$

preserve the test function $e_2(x) = x^2$ and provide a better error estimation than the operators $S_n(f; x)$ for all

$f \in C_B([0, \infty))$ and for each $x \in [0, \infty)$. Note that $C_B([0, \infty))$ denotes the space of all bounded and continuous functions on $[0, \infty)$. On the other hand, by letting

$$v_n(x) := x - \frac{1}{2n}; \quad n \in \mathbb{N}, \quad (5)$$

it has been shown in [2] that the operators defined by

$$V_n^*(f; x) := S_n(f; v_n(x)) \quad (6)$$

do not preserve the test functions $e_1(x) = x$ and $e_2(x) = x^2$ but provide the best error estimation among all the Szász-Mirakjan operators for all $f \in C_B([0, \infty))$ and for each $x \in [1/2, \infty)$. For the other linear positive operator families which preserve $e_2(x) = x^2$, we refer [3–9]. On the other hand, in [10, 11] the authors considered some operators preserving $e_1(x) = x$.

Favard was the first to introduce the double Szász-Mirakjan operators [12]:

$$S_n(f; x, y) = e^{-n(x+y)} \sum_{k,l=0}^{\infty} f\left(\frac{k}{n}, \frac{l}{n}\right) \frac{(nx)^k}{k!} \frac{(ny)^l}{l!}, \quad f \in \mathcal{A}_2, \quad (7)$$

where \mathcal{A}_2 is the set of all real functions on $[0, \infty) \times [0, \infty)$ such that the right-hand side in (7) has a meaning for all $n > 0$ and $x, y \in [0, \infty)$. Recently, Dirik and Demirci have

introduced and investigated different variants of the general double Szász-Mirakjan operators:

$$\begin{aligned} D_n(f; x, y) : S_n(f; u_n(x), v_n(y)) \\ = e^{-n(u_n(x)+v_n(y))} \\ \times \sum_{k,l=0}^{\infty} f\left(\frac{k}{n}, \frac{l}{n}\right) \times \frac{(nu_n(x))^k (nv_n(y))^l}{k! l!}, \\ f \in \mathcal{A}_2. \end{aligned} \quad (8)$$

In [13], they considered the case of operators

$$\begin{aligned} u_n^{(1)}(x) &:= \frac{-1 + \sqrt{4n^2 x^2 + 1}}{2n}, \\ v_n^{(1)}(y) &:= \frac{-1 + \sqrt{4n^2 y^2 + 1}}{2n}, \\ n &\in \mathbb{N}, \end{aligned} \quad (9)$$

which preserve the test function $e_{2,0}(x, y) + e_{0,2}(x, y) := x^2 + y^2$ and provide a better error estimation than the operators $S_n(f; x, y)$ for all $f \in C_B([0, \infty) \times [0, \infty))$ and for each $x, y \in [0, \infty)$. On the other hand, in [14], they considered the case

$$\begin{aligned} u_n^{(2)}(x, \alpha) &:= \frac{-(n\alpha + 1) + \sqrt{4n^2(x^2 + \alpha x) + (n\alpha + 1)^2}}{2n}, \\ v_n^{(2)}(y, \beta) &:= \frac{-(n\beta + 1) + \sqrt{4n^2(y^2 + \beta y) + (n\beta + 1)^2}}{2n}, \\ n &\in \mathbb{N}, \alpha, \beta \in \mathbb{R}. \end{aligned} \quad (10)$$

Note that for this case, the operators $D_n(f; x, y)$ do not preserve any test function (i.e., $e_{0,0}(x, y) = 1$, $e_{1,0}(x, y) = x$, $e_{0,1}(x, y) = y$, and $e_{2,0}(x, y) + e_{0,2}(x, y) = x^2 + y^2$) but provide a better error estimation than the operators $S_n(f; x, y)$ for all $f \in C_B([0, \infty) \times [0, \infty))$ and $x, y \in [0, 1]$.

Finally, we should note that, following the similar arguments as used in [2], the best error estimation among all the general double Szász-Mirakjan operators can be obtained from the case:

$$u_n^{(3)}(x) := x - \frac{1}{2n}, \quad v_n^{(3)}(y) := y - \frac{1}{2n}, \quad n \in \mathbb{N}, \quad (11)$$

for all $f \in C_B([0, \infty) \times [0, \infty))$ and $x, y \in [1/2, \infty)$.

For the operators $D_n(f; x, y)$ the following Lemma is straightforward.

Lemma 1. Let $\mathbf{x} = (x, y)$, $\mathbf{t} = (t, s)$, $e_{i,j}(\mathbf{x}) = x^i y^j$, $i, j = 0, 1, 2$, and $\psi_{\mathbf{x}}^2(\mathbf{t}) = \|\mathbf{t} - \mathbf{x}\|^2$. Then, for each $x, y \geq 0$ and $n > 1$, one has

- (a) $D_n(e_{0,0}; x, y) = 1$,
- (b) $D_n(e_{1,0}; x, y) = u_n(x)$, $D_n(e_{0,1}; x, y) = v_n(y)$,

$$(c) D_n(e_{2,0} + e_{0,2}; x, y) = u_n^2(x) + v_n^2(y) + ((u_n(x) + v_n(y))/n),$$

$$(d) D_n(\psi_{\mathbf{x}}^2(\mathbf{t}); x, y) = (u_n(x) - x)^2 + (v_n(y) - y)^2 + ((u_n(x) + v_n(y))/n).$$

2. Global Results

In this section we first introduce the following Lipschitz-type space:

$$\begin{aligned} \text{Lip}_M^*(\alpha) &:= \left\{ f \in C([0, \infty) \times [0, \infty)) : \right. \\ &\quad |f(\mathbf{t}) - f(\mathbf{x})| \leq M \frac{\|\mathbf{t} - \mathbf{x}\|^\alpha}{(\|\mathbf{t}\| + x + y)^{\alpha/2}}; \\ &\quad \left. t, s; x, y \in (0, \infty) \right\}, \end{aligned} \quad (12)$$

where $\mathbf{t} = (t, s)$, $\mathbf{x} = (x, y)$, M is any positive constant, and $0 < \alpha \leq 1$.

We should note that this space is the bivariate extension of Lipschitz-type space considered earlier by Szasz [15]. For the space $\text{Lip}_M^*(\alpha)$ with $0 < \alpha \leq 1$, we have the following approximation result.

Theorem 2. For any $f \in \text{Lip}_M^*(\alpha)$, $\alpha \in (0, 1]$ and for each $x, y \in (0, \infty)$, $n \in \mathbb{N}$, one has

$$\begin{aligned} &|D_n(f; x, y) - f(x, y)| \\ &\leq \frac{M}{(x + y)^{\alpha/2}} \left[(u_n(x) - x)^2 + (v_n(y) - y)^2 \right. \\ &\quad \left. + \frac{u_n(x) + v_n(y)}{n} \right]^{\alpha/2}. \end{aligned} \quad (13)$$

Proof. Take $\alpha = 1$. Then, for $f \in \text{Lip}_M^*(1)$ and for each $x, y \in (0, \infty)$, we get

$$\begin{aligned} |D_n(f; x, y) - f(x, y)| &\leq D_n(|f(\mathbf{t}, s) - f(\mathbf{x}, y)|; x, y) \\ &\leq MD_n\left(\frac{\|\mathbf{t} - \mathbf{x}\|}{(\|\mathbf{t}\| + x + y)^{1/2}}; x, y\right) \\ &\leq \frac{M}{(x + y)^{1/2}} D_n(\|\mathbf{t} - \mathbf{x}\|; x, y). \end{aligned} \quad (14)$$

Using the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned}
 & |D_n(f; x, y) - f(x, y)| \\
 & \leq \frac{M}{(x+y)^{1/2}} \sqrt{D_n(\psi_x^2(\mathbf{t}); x, y)} \\
 & = \frac{M}{(x+y)^{1/2}} \\
 & \quad \times \sqrt{(u_n(x) - x)^2 + (v_n(y) - y)^2 + \frac{u_n(x) + v_n(y)}{n}}.
 \end{aligned} \tag{15}$$

Secondly let $0 < \alpha < 1$. Then, for $f \in \text{Lip}_M^*(\alpha)$ and for each $x, y \in (0, \infty)$, we have

$$\begin{aligned}
 |D_n(f; x, y) - f(x, y)| & \leq D_n(|f(t, s) - f(x, y)|; x, y) \\
 & \leq MD_n\left(\frac{\|\mathbf{t} - \mathbf{x}\|^\alpha}{(\|\mathbf{t}\| + x + y)^{\alpha/2}}; x, y\right) \\
 & \leq \frac{M}{(x+y)^{\alpha/2}} D_n(\|\mathbf{t} - \mathbf{x}\|^\alpha; x, y).
 \end{aligned} \tag{16}$$

Applying the Hölder inequality with $p = 2/\alpha$ and $q = 2/(2 - \alpha)$, we have, for any $f \in \text{Lip}_M^*(\alpha)$,

$$\begin{aligned}
 |D_n(f; x, y) - f(x, y)| & \leq \frac{M}{(x+y)^{\alpha/2}} [D_n(\psi_x^2(\mathbf{t}); x, y)]^{\alpha/2} \\
 & = \frac{M}{(x+y)^{\alpha/2}} \\
 & \quad \times \left[(u_n(x) - x)^2 + (v_n(y) - y)^2 \right. \\
 & \quad \left. + \frac{u_n(x) + v_n(y)}{n} \right]^{\alpha/2},
 \end{aligned} \tag{17}$$

Hence, the result. \square

The following lemma will be used in the rest of the paper.

Lemma 3. One has, for each $x, y > 0$,

$$\begin{aligned}
 & D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right) \\
 & \leq \frac{1}{\sqrt{x}} \sqrt{(u_n(x) - x)^2 + \frac{u_n(x)}{n}} \\
 & \quad + \frac{1}{\sqrt{y}} \sqrt{(v_n(y) - y)^2 + \frac{v_n(y)}{n}}.
 \end{aligned} \tag{18}$$

Proof. Using the fact that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ ($a, b \geq 0$), we get

$$\begin{aligned}
 & D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right) \\
 & = e^{-n(u_n(x) + v_n(y))} \sum_{k,l=0}^{\infty} \sqrt{\left(\sqrt{\frac{k}{n}} - \sqrt{x}\right)^2 + \left(\sqrt{\frac{l}{n}} - \sqrt{y}\right)^2} \\
 & \quad \times \frac{(nu_n(x))^k}{k!} \frac{(nv_n(y))^l}{l!} \\
 & \leq e^{-nu_n(x)} \sum_{k=0}^{\infty} \left|\sqrt{\frac{k}{n}} - \sqrt{x}\right| \frac{(nu_n(x))^k}{k!} \\
 & \quad + e^{-nv_n(y)} \sum_{l=0}^{\infty} \left|\sqrt{\frac{l}{n}} - \sqrt{y}\right| \frac{(nv_n(y))^l}{l!} \\
 & = e^{-nu_n(x)} \sum_{k=0}^{\infty} \frac{|k/n - x|}{\sqrt{k/n} + \sqrt{x}} \frac{(nu_n(x))^k}{k!} \\
 & \quad + e^{-nv_n(y)} \sum_{l=0}^{\infty} \frac{|l/n - y|}{\sqrt{l/n} + \sqrt{y}} \frac{(nv_n(y))^l}{l!} \\
 & \leq \frac{e^{-nu_n(x)}}{\sqrt{x}} \sum_{k=0}^{\infty} \left|\frac{k}{n} - x\right| \frac{(nu_n(x))^k}{k!} \\
 & \quad + \frac{e^{-nv_n(y)}}{\sqrt{y}} \sum_{l=0}^{\infty} \left|\frac{l}{n} - y\right| \frac{(nv_n(y))^l}{l!}.
 \end{aligned} \tag{19}$$

Finally, applying the Cauchy-Schwarz inequality, we write

$$\begin{aligned}
 & D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right) \\
 & \leq \frac{1}{\sqrt{x}} \sqrt{e^{-nu_n(x)} \sum_{k=0}^{\infty} \left(\frac{k}{n} - x\right)^2 \frac{(nu_n(x))^k}{k!}} \\
 & \quad + \frac{1}{\sqrt{y}} \sqrt{e^{-nv_n(y)} \sum_{l=0}^{\infty} \left(\frac{l}{n} - y\right)^2 \frac{(nv_n(y))^l}{l!}} \\
 & = \frac{1}{\sqrt{x}} \sqrt{(u_n(x) - x)^2 + \frac{u_n(x)}{n}} \\
 & \quad + \frac{1}{\sqrt{y}} \sqrt{(v_n(y) - y)^2 + \frac{v_n(y)}{n}}.
 \end{aligned} \tag{20}$$

Using Lemma 1, we get the result. \square

Recall that, for all $f \in C_B([0, \infty) \times [0, \infty))$, the modulus of f denoted by $\omega(f; \delta)$ is defined as

$$\omega(f; \delta) := \sup \left\{ |f(t, s) - f(x, y)| : \sqrt{(t-x)^2 + (s-y)^2} < \delta, (t, s), (x, y) \in [0, \infty) \times [0, \infty) \right\}. \quad (21)$$

Theorem 4. Let $f^*(x, y) = f(x^2, y^2)$. Then one has, for each $x, y > 0$,

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega(f^*; \delta_n(x, y)), \quad (22)$$

where

$$\delta_n(x, y) := \frac{1}{\sqrt{x}} \sqrt{(u_n(x) - x)^2 + \frac{u_n(x)}{n}} + \frac{1}{\sqrt{y}} \sqrt{(v_n(y) - y)^2 + \frac{v_n(y)}{n}}. \quad (23)$$

Proof. We directly have

$$\begin{aligned} & |D_n(f; x, y) - f(x, y)| \\ & \leq D_n(|f(t, s) - f(x, y)|; x, y) \\ & = D_n(|f^*(\sqrt{t}, \sqrt{s}) - f^*(\sqrt{x}, \sqrt{y})|; x, y) \\ & \leq D_n\left(\omega\left(f^*; \sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}\right); x, y\right) \\ & = e^{-n(u_n(x) + v_n(y))} \\ & \quad \times \sum_{k, l=0}^{\infty} \omega\left(f^*; \sqrt{\left(\sqrt{\frac{k}{n}} - \sqrt{x}\right)^2 + \left(\sqrt{\frac{l}{n}} - \sqrt{y}\right)^2}; x, y\right) \\ & \quad \times \frac{(nu_n(x))^k (nv_n(y))^l}{k! l!}. \end{aligned} \quad (24)$$

Therefore,

$$\begin{aligned} & |D_n(f; x, y) - f(x, y)| \\ & = e^{-n(u_n(x) + v_n(y))} \sum_{k, l=0}^{\infty} \frac{(nu_n(x))^k (nv_n(y))^l}{k! l!} \end{aligned}$$

$$\begin{aligned} & \times \omega\left(f^*; \frac{\sqrt{(\sqrt{k/n} - \sqrt{x})^2 + (\sqrt{l/n} - \sqrt{y})^2}}{D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right)} \right. \\ & \quad \times D_n\left(\left((\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2\right)^{1/2}; x, y\right); x, y\right). \end{aligned} \quad (25)$$

Because of the fact that

$$\omega(f; \lambda\delta) \leq (1 + \lambda)\omega(f; \delta), \quad (26)$$

we have

$$\begin{aligned} & |D_n(f; x, y) - f(x, y)| \\ & \leq \omega\left(f^*; D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right)\right) \\ & \quad \times e^{-n(u_n(x) + v_n(y))} \\ & \quad \times \sum_{k, l=0}^{\infty} \left[1 + \frac{\sqrt{(\sqrt{k/n} - \sqrt{x})^2 + (\sqrt{l/n} - \sqrt{y})^2}}{D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right)} \right] \\ & \quad \times \frac{(nu_n(x))^k (nv_n(y))^l}{k! l!}, \end{aligned} \quad (27)$$

and hence

$$\begin{aligned} & |D_n(f; x, y) - f(x, y)| \\ & \leq 2\omega\left(f^*; D_n\left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y\right)\right). \end{aligned} \quad (28)$$

Finally, using Lemma 3, the proof is completed. \square

Theorem 5. Let $f^*(x, y) = f(x^2, y^2)$. Let

$$\begin{aligned} f^* \in Lip_M(\alpha) & := \{f^* \in C_B([0, \infty) \times [0, \infty)) : \\ & |f^*(\mathbf{t}) - f^*(\mathbf{x})| \leq M\|\mathbf{t} - \mathbf{x}\|^\alpha; \\ & t, s; x, y \in (0, \infty)\}, \end{aligned} \quad (29)$$

where $\mathbf{t} = (t, s)$, $\mathbf{x} = (x, y)$, M is any positive constant, and $0 < \alpha \leq 1$. Then

$$|D_n(f; x, y) - f(x, y)| \leq M\delta_n^\alpha(x, y), \quad (30)$$

where $\delta_n(x, y)$ is the same as in Theorem 4.

Proof. We directly have

$$\begin{aligned}
 & |D_n(f; x, y) - f(x, y)| \\
 & \leq D_n(|f(t, s) - f(x, y)|; x, y) \\
 & = D_n(|f^*(\sqrt{t}, \sqrt{s}) - f^*(\sqrt{x}, \sqrt{y})|; x, y) \\
 & \leq MD_n\left(\left((\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2\right)^{\alpha/2}; x, y\right) \\
 & = Me^{-n(u_n(x) + v_n(y))} \\
 & \quad \times \sum_{k, l=0}^{\infty} \left(\left(\sqrt{\frac{k}{n}} - \sqrt{x} \right)^2 + \left(\sqrt{\frac{l}{n}} - \sqrt{y} \right)^2 \right)^{\alpha/2} \\
 & \quad \times \frac{(nu_n(x))^k (nv_n(y))^l}{k! l!}.
 \end{aligned} \tag{31}$$

Applying the Hölder inequality with $p = 1/\alpha$ and $q = 1/(1 - \alpha)$, we have

$$\begin{aligned}
 & |D_n(f; x, y) - f(x, y)| \\
 & \leq M \left[D_n \left(\sqrt{(\sqrt{t} - \sqrt{x})^2 + (\sqrt{s} - \sqrt{y})^2}; x, y \right) \right]^{\alpha}.
 \end{aligned} \tag{32}$$

Using Lemma 3, we get the result. \square

3. Concluding Remarks

In this section we show that taking $u_n(x) = x$ and $v_n(y) = y$ or $u_n(x) = u_n^{(i)}(x)$ and $v_n(y) = v_n^{(i)}(y)$, $i = 1, 3$, in Theorems 2, 4, and 5 gives global results. Also we present the results obtained by Theorems 2, 4, and 5 for $u_n(x) = u_n^{(2)}(x)$ and $v_n(y) = v_n^{(2)}(y)$.

Corollary 6. For any $f \in Lip_M^*(\alpha)$, $\alpha \in (0, 1]$ and for all $x, y \in (0, \infty)$, $n \in \mathbb{N}$, one has

$$|D_n(f; x, y) - f(x, y)| \leq \frac{M}{n^{\alpha/2}}, \tag{33}$$

uniformly as $n \rightarrow \infty$, for the following pairs of $u_n(x)$ and $v_n(x)$:

- (i) $u_n(x) = x$ and $v_n(y) = y$,
- (ii) $u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n$,
- (iii) $u_n(x) = x - 1/2n$ and $v_n(y) = y - 1/2n$.

Proof. (i) Taking $u_n(x) = x$ and $v_n(y) = y$ in (13), we directly have

$$|D_n(f; x, y) - f(x, y)| \leq \frac{M}{n^{\alpha/2}}. \tag{34}$$

(ii) Taking $u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n$ in (13) gives

$$\begin{aligned}
 & |D_n(f; x, y) - f(x, y)| \\
 & = \frac{M}{(x + y)^{\alpha/2}} \left[\frac{1}{n} \left(x + y - x\sqrt{4n^2x^2 + 1} \right. \right. \\
 & \quad \left. \left. - y\sqrt{4n^2y^2 + 1} + 2nx^2 + 2ny^2 \right) \right]^{\alpha/2} \\
 & \leq \frac{M}{(x + y)^{\alpha/2}} \left[\frac{1}{n} \left(x + y - x\sqrt{4n^2x^2} \right. \right. \\
 & \quad \left. \left. - y\sqrt{4n^2y^2} + 2nx^2 + 2ny^2 \right) \right]^{\alpha/2} \\
 & = \frac{M}{n^{\alpha/2}}.
 \end{aligned} \tag{35}$$

(iii) Taking $u_n(x) = x - 1/2n$ and $v_n(y) = y - 1/2n$ in (13), we have

$$\begin{aligned}
 & |D_n(f; x, y) - f(x, y)| \\
 & = \frac{M}{(x + y)^{\alpha/2}} \left[\left(-\frac{1}{2n} \right)^2 + \left(-\frac{1}{2n} \right)^2 + \frac{x + y - 1/n}{n} \right]^{\alpha/2} \\
 & \leq \frac{M}{(x + y)^{\alpha/2}} \left[\frac{1}{2n^2} (2nx + 2ny - 1) \right]^{\alpha/2} \\
 & = \frac{M}{(x + y)^{\alpha/2}} \left[\frac{1}{n} (x + y) \right]^{\alpha/2} \\
 & = \frac{M}{n^{\alpha/2}}.
 \end{aligned} \tag{36}$$

\square

Corollary 7. Let $f^*(x, y) = f(x^2, y^2)$. Then one has

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega \left(f^*; \frac{2}{\sqrt{n}} \right), \tag{37}$$

uniformly as $n \rightarrow \infty$, for the following pairs of $u_n(x)$ and $v_n(x)$:

- (i) $u_n(x) = x$ and $v_n(y) = y$,
- (ii) $u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n$,
- (iii) $u_n(x) = x - 1/2n$ and $v_n(y) = y - 1/2n$.

Proof. (i) Taking $u_n(x) = x$ and $v_n(y) = y$ in (23), we directly have

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega \left(f^*; \frac{2}{\sqrt{n}} \right). \tag{38}$$

(ii) Taking $u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n$ in (23) gives

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega(f^*; \delta_n(x, y)), \quad (39)$$

where

$$\begin{aligned} \delta_n(x, y) &:= \frac{1}{\sqrt{x}\sqrt{y}} \left(\sqrt{y} \sqrt{\frac{1}{n} \left(x - x\sqrt{4n^2x^2 + 1} + 2nx^2 \right)} \right. \\ &\quad \left. + \sqrt{x} \sqrt{\frac{1}{n} \left(y - y\sqrt{4n^2y^2 + 1} + 2ny^2 \right)} \right) \\ &\leq \frac{1}{\sqrt{x}\sqrt{y}} \left(\sqrt{y} \sqrt{\frac{1}{n}x} + \sqrt{x} \sqrt{\frac{1}{n}y} \right) \\ &= \frac{2}{\sqrt{n}}. \end{aligned} \quad (40)$$

(iii) Taking $u_n(x) = x - 1/2n$ and $v_n(y) = y - 1/2n$ in (23), we have

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega(f^*; \delta_n(x, y)), \quad (41)$$

where

$$\begin{aligned} \delta_n(x, y) &:= \frac{1}{\sqrt{x}} \sqrt{\left(\frac{1}{2n}\right)^2 + \frac{x}{n} - \frac{1}{2n^2}} \\ &\quad + \frac{1}{\sqrt{y}} \sqrt{\left(\frac{1}{2n}\right)^2 + \frac{y}{n} - \frac{1}{2n^2}} \\ &= \frac{1}{2\sqrt{x}\sqrt{y}} \left(\sqrt{x} \sqrt{\frac{1}{n^2}(4ny - 1)} \right. \\ &\quad \left. + \sqrt{y} \sqrt{\frac{1}{n^2}(4nx - 1)} \right) \\ &\leq \frac{1}{\sqrt{x}\sqrt{y}} \left(\sqrt{x} \sqrt{\frac{1}{n}y} + \sqrt{y} \sqrt{\frac{1}{n}x} \right) \\ &= \frac{2}{\sqrt{n}}. \end{aligned} \quad (42)$$

□

Corollary 8. Let $f^*(x, y) = f(x^2, y^2)$, and let

$$\begin{aligned} f^* \in Lip_M(\alpha) &:= \{f^* \in C_B([0, \infty) \times [0, \infty)) : \\ |f^*(\mathbf{t}) - f^*(\mathbf{x})| &\leq M\|\mathbf{t} - \mathbf{x}\|^\alpha; \quad (43) \\ t, s; x, y &\in (0, \infty)\}, \end{aligned}$$

where $\mathbf{t} = (t, s)$, $\mathbf{x} = (x, y)$, M is any positive constant, and $0 < \alpha \leq 1$. Then

$$|D_n(f; x, y) - f(x, y)| \leq M\left(\frac{4}{n}\right)^{\alpha/2}, \quad (44)$$

uniformly as $n \rightarrow \infty$, for the following pairs of $u_n(x)$ and $v_n(y)$:

$$(i) \quad u_n(x) = x \text{ and } v_n(y) = y,$$

$$(ii) \quad u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n \text{ and } v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n,$$

$$(iii) \quad u_n(x) = x - 1/2n \text{ and } v_n(y) = y - 1/2n.$$

Proof. (i) Taking $u_n(x) = x$ and $v_n(y) = y$ in (23), we directly have

$$|D_n(f; x, y) - f(x, y)| \leq M\left(\frac{4}{n}\right)^{\alpha/2}. \quad (45)$$

(ii) Taking $u_n(x) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n(y) = (-1 + \sqrt{4n^2y^2 + 1})/2n$ in (23) gives

$$|D_n(f; x, y) - f(x, y)| \leq M\delta_n^{\alpha/2}(x, y), \quad (46)$$

where

$$\begin{aligned} \delta_n(x, y) &:= \delta_n(x, y) \\ &:= \frac{1}{\sqrt{x}\sqrt{y}} \\ &\quad \times \left(\sqrt{y} \sqrt{\frac{1}{n} \left(x - x\sqrt{4n^2x^2 + 1} + 2nx^2 \right)} \right. \\ &\quad \left. + \sqrt{x} \sqrt{\frac{1}{n} \left(y - y\sqrt{4n^2y^2 + 1} + 2ny^2 \right)} \right) \\ &\leq \frac{1}{\sqrt{x}\sqrt{y}} \left(\sqrt{y} \sqrt{\frac{1}{n}x} + \sqrt{x} \sqrt{\frac{1}{n}y} \right) \\ &= \frac{2}{\sqrt{n}}. \end{aligned} \quad (47)$$

(iii) Taking $u_n(x) = x - 1/2n$ and $v_n(y) = y - 1/2n$ in (23), we have

$$|D_n(f; x, y) - f(x, y)| \leq M\delta_n^{\alpha/2}(x, y), \quad (48)$$

where

$$\begin{aligned}
 \delta_n(x, y) &:= \frac{1}{\sqrt{x}} \sqrt{\left(\frac{1}{2n}\right)^2 + \frac{x}{n} - \frac{1}{2n^2}} \\
 &\quad + \frac{1}{\sqrt{y}} \sqrt{\left(\frac{1}{2n}\right)^2 + \frac{y}{n} - \frac{1}{2n^2}} \\
 &= \frac{1}{2\sqrt{x}\sqrt{y}} \left(\sqrt{x} \sqrt{\frac{1}{n^2}(4ny-1)} \right. \\
 &\quad \left. + \sqrt{y} \sqrt{\frac{1}{n^2}(4nx-1)} \right) \\
 &\leq \frac{1}{\sqrt{x}\sqrt{y}} \left(\sqrt{x} \sqrt{\frac{1}{n}y} + \sqrt{y} \sqrt{\frac{1}{n}x} \right) \\
 &= \frac{2}{\sqrt{n}}.
 \end{aligned} \tag{49}$$

Remark 9. Corollaries 7 and 8 conclude that f is a real continuous and bounded function on $[0, \infty) \times [0, \infty)$ and if $f^*(x, y) = f(x^2, y^2)$ is uniformly continuous on $[0, \infty) \times [0, \infty)$, then $D_n(f)$ converges uniformly to f as $n \rightarrow \infty$. Note that the one variable version of Corollary 7 was given in [16].

Corollary 10. Take

$$\begin{aligned}
 u_n(x) &= u_n^{(2)}(x, \gamma) \\
 &= \frac{-(n\gamma + 1) + \sqrt{4n^2(x^2 + \gamma x) + (n\gamma + 1)^2}}{2n}, \\
 v_n(y) &= v_n^{(2)}(y, \beta) \\
 &= \frac{-(n\beta + 1) + \sqrt{4n^2(y^2 + \beta y) + (n\beta + 1)^2}}{2n},
 \end{aligned} \tag{50}$$

where $\alpha, \beta \in \mathbb{R}$. Then

(i) for any $f \in \text{Lip}_M^*(\alpha)$, $\alpha \in (0, 1]$ and for each $x, y \in (0, \infty)$, $n \in \mathbb{N}$, one has

$$\begin{aligned}
 |D_n(f; x, y) - f(x, y)| \\
 \leq \frac{M}{[2n(x+y)]^{\alpha/2}} [\delta(x, \gamma) + \delta(y, \beta)]^{\alpha/2},
 \end{aligned} \tag{51}$$

where

$$\begin{aligned}
 \delta(x, \gamma) &= \left[(2x + \gamma) \right. \\
 &\quad \left. \times \left(n\gamma - \sqrt{n^2(2x + \gamma)^2 + 2n\gamma + 1 + 2nx + 1} \right) \right],
 \end{aligned} \tag{52}$$

(ii) let $f^*(x, y) = f(x^2, y^2)$. Then one has for each $x, y > 0$

$$|D_n(f; x, y) - f(x, y)| \leq 2\omega(f^*; \delta(x, \gamma) + \delta(y, \beta)), \tag{53}$$

where

$$\begin{aligned}
 \delta(x, \gamma) &= \frac{1}{\sqrt{2x}} \\
 &\quad \times \left(\frac{1}{n} (2x + \gamma) \left(n\gamma - \sqrt{n^2(2x + \gamma)^2 + 2n\gamma + 1} \right. \right. \\
 &\quad \left. \left. + 2nx + 1 \right) \right)^{1/2},
 \end{aligned} \tag{54}$$

(iii) let $f^*(x, y) = f(x^2, y^2)$, and let

$$f^* \in \text{Lip}_M(\alpha) := \{f^* \in C_B([0, \infty) \times [0, \infty)) :$$

$$|f^*(\mathbf{t}) - f^*(\mathbf{x})| \leq M \|\mathbf{t} - \mathbf{x}\|^\alpha; \tag{55}$$

$$t, s; x, y \in (0, \infty) \},$$

where $\mathbf{t} = (t, s)$, $\mathbf{x} = (x, y)$, M is any positive constant, and $0 < \alpha \leq 1$. Then

$$|D_n(f; x, y) - f(x, y)| \leq M [\delta(x, \gamma) + \delta(y, \beta)]^{\alpha/2}, \tag{56}$$

where $\delta(x, \gamma)$ is the same given in Corollary 10(ii).

It should be mentioned that, for $\alpha = 0$ and $\beta = 0$, $u_n^{(2)}(x, 0) = (-1 + \sqrt{4n^2x^2 + 1})/2n$ and $v_n^{(2)}(y, 0) = (-1 + \sqrt{4n^2y^2 + 1})/2n$. Therefore, Corollary 10(i), Corollary 10(ii), and Corollary 10(iii) reduce to Corollary 6(ii), Corollary 7(ii), and Corollary 8(ii), respectively.

References

- [1] O. Duman and M. A. Özarslan, "Szász-Mirakjan type operators providing a better error estimation," *Applied Mathematics Letters*, vol. 20, no. 12, pp. 1184–1188, 2007.
- [2] M. A. Özarslan and O. Duman, "A new approach in obtaining a better estimation in approximation by positive linear operators," *Communications de la Faculté des Sciences de l'Université d'Ankara A1*, vol. 58, no. 1, pp. 17–22, 2009.
- [3] O. Agratini, "Linear operators that preserve some test functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 2006, Article ID 94136, 11 pages, 2006.
- [4] O. Duman, M. A. Özarslan, and H. Aktuğlu, "Better error estimation for Szász-Mirakjan-beta operators," *Journal of Computational Analysis and Applications*, vol. 10, no. 1, pp. 53–59, 2008.
- [5] H. Gonska, P. Pişul, and I. Raşa, "General King-type operators," *Results in Mathematics*, vol. 53, no. 3–4, pp. 279–286, 2009.

- [6] J. P. King, "Positive linear operators which preserve x^2 ," *Acta Mathematica Hungarica*, vol. 99, no. 3, pp. 203–208, 2003.
- [7] N. I. Mahmudov, "Korovkin-type theorems and applications," *Central European Journal of Mathematics*, vol. 7, no. 2, pp. 348–356, 2009.
- [8] M. A. Özarslan and O. Duman, "Local approximation results for Szász-Mirakjan type operators," *Archiv der Mathematik*, vol. 90, no. 2, pp. 144–149, 2008.
- [9] L. Rempulska and K. Tomczak, "Approximation by certain linear operators preserving x^2 ," *Turkish Journal of Mathematics*, vol. 33, no. 3, pp. 273–281, 2009.
- [10] O. Duman, M. A. Özarslan, and B. D. Vecchia, "Modified Szász-Mirakjan-Kantorovich operators preserving linear functions," *Turkish Journal of Mathematics*, vol. 33, no. 2, pp. 151–158, 2009.
- [11] M. Örkücü and O. Doğru, " q -Szász-Mirakjan-Kantorovich type operators preserving some test functions," *Applied Mathematics Letters*, vol. 24, no. 9, pp. 1588–1593, 2011.
- [12] J. Favard, "Sur les multiplicateurs d'interpolation," *Journal de Mathématiques Pures et Appliquées. Neuvième Série*, vol. 23, pp. 219–247, 1944.
- [13] F. Dirik and K. Demirci, "Modified double Szász-Mirakjan operators preserving $x^2 + y^2$," *Mathematical Communications*, vol. 15, no. 1, pp. 177–188, 2010.
- [14] F. Dirik, Demirci, and K. Szasz, "Mirakjan type operators of two variables providing a better estimation on $[0; 1] \times [0; 1]$," *Matematicki Vesnik*, vol. 63, no. 1, pp. 59–66, 2011.
- [15] O. Szasz, "Generalization of S. Bernstein's polynomials to the infinite interval," *Journal of Research of the National Bureau of Standards*, vol. 45, pp. 239–245, 1950.
- [16] J. de la Cal and J. Cárcamo, "On uniform approximation by some classical Bernstein-type operators," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 625–638, 2003.