

PROJECTABLE ALMOST COMPLEX CONTACT STRUCTURES

BY D. E. BLAIR, S. ISHIHARA AND G. D. LUDDEN

A complex manifold of complex dimension $2m+1$ is said to be a *complex contact manifold* if it admits an open covering $\{u_\alpha\}$ such that on each u_α there is a holomorphic 1-form ω_α with $\omega_\alpha \wedge (d\omega_\alpha)^m \neq 0$ on $u_\alpha \cap u_\beta \neq \emptyset$, $\omega_\beta = f\omega_\alpha$ for some non-vanishing holomorphic function f . In general such a structure is not given by a global 1-form ω ; in fact this is the case for a compact complex manifold if and only if its first Chern class vanishes [6]. However, a complex contact manifold is the base space of a principal fibre bundle with 1-dimensional fibres and real contact structure. Homogeneous complex contact manifolds were studied by Boothby in [3].

It is also shown in [6] that the structural group of the tangent bundle of a Hermitian contact manifold M is reducible to $(Sp(m) \cdot Sp(1)) \times U(1)$ where $Sp(m) \cdot Sp(1) = Sp(m) \times Sp(1) / \{\pm I\}$ and hence equivalently M admits the following local structure tensors. Let F denote the almost complex structure and g the Hermitian metric on M . Then each coordinate neighborhood admits tensor fields G, H of type $(1, 1)$ and vector fields U, V with covariant forms u and v such that (G, U, V, g) and (H, U, V, g) are metric f -structures with complemented frames (see e.g. [1]), $FU = V$ and $GH = -HG = F + v \otimes U - u \otimes V$. In the overlap of coordinate neighborhoods we have

$$\begin{aligned} G' &= aG + bH, & u' &= au + bv, \\ H' &= -bG + aH, & v' &= -bu + av \end{aligned} \tag{0.1}$$

with $a^2 + b^2 = 1$. Such a structure is called an *almost complex contact structure* [5] and our first project here will be to give an equivalent definition in terms of global tensor fields.

A standard example of a complex contact manifold is the odd-dimensional complex projective space PC^{2m+1} . It is also well known that PC^{2m+1} is a fibre space over the quaternionic projective space PH^m with fibres $S^2 \approx PC^1$. In sections 3 and 4 we generalize this situation to a projectable almost complex contact structure on a Kählerian manifold.

§ 1. Almost Complex Contact Structures

In terms of the above local tensor fields G, H, U, V we can define global

Received June 14, 1976

tensor fields Σ of type (1, 3) and S of type (1, 1). For local vector fields X, Y, Z set

$$\Sigma_{XY}Z = g(GX, Y)GZ + g(HX, Y)HZ \quad (1.1)$$

and

$$SX = u(X)U + v(X)V. \quad (1.2)$$

It is then easy to check using equations (0.1) that Σ and S are globally defined. Note also that S is a projection tensor field of rank 2, i.e. $S^2 = S$. For a unit vector $A \in T_p M$ with $SA = 0$, let

$$\sigma_A = \{B \in T_p M \mid g(A, B) = 0, \|B\| = 1, \underline{\Sigma}(A, B, A, B) = 1\},$$

where $\underline{\Sigma}(X, Y, Z, W) = g(\Sigma_{XY}Z, W)$ and $[\sigma_A]$ the subspace of $T_p M$ generated by σ_A .

The following properties of Σ and S are now easily deduced. 1)-8) are straightforward computations using equations (1.1) and (1.2) and elementary properties of metric f -structures. For 9) given A set $B = GA$ and it is easy to see that $B \in \sigma_A$.

- 1) $SF = FS$
- 2) $\Sigma_{XY} = -\Sigma_{YX}$
- 3) $\Sigma_{XY}^2 = \underline{\Sigma}(X, Y, X, Y)(-I + S)$
- 4) $\Sigma_{XY}S = S\Sigma_{XY} = 0$
- 5) $\Sigma_{XY}F = -F\Sigma_{XY}$
- 6) $\Sigma_{XFY}F = \Sigma_{XY}$
- 7) $\underline{\Sigma}(X, Y, Z, W) = \underline{\Sigma}(Z, W, X, Y)$
- 8) $\Sigma_{X \Sigma_{YZ} X} W = g(X, (I - S)X)\Sigma_{YZ}W$
- 9) $\sigma_A \neq \emptyset$ for any unit vector A with $SA = 0$ and at any point p of M .

Conversely we will show that an almost Hermitian manifold M with structure tensors (F, g) admitting global tensor fields Σ and S satisfying 1)-9) is an almost complex contact manifold. We first give several lemmas.

LEMMA 1.1. For $B \in \sigma_A$, $\Sigma_{AB}A = B$, $SB = 0$ and σ_A is invariant under F .

Proof. Since $\underline{\Sigma}(A, B, A, B) = g(\Sigma_{AB}A, B) = 1$ to show that $\Sigma_{AB}A = B$ it suffices to show that $\Sigma_{AB}A$ is a unit vector.

$$g(\Sigma_{AB}A, \Sigma_{AB}A) = -g(\Sigma_{AB}^2 A, A) = -g(-A + SA, A) = 1$$

by 2), 7) and 3), since A is a unit vector and $SA = 0$. Now $SB = S\Sigma_{AB}A = 0$ by 4). Finally for the invariance by F ,

$$\Sigma_{AFB}A = -\Sigma_{AFB}F^2A = F\Sigma_{AB}A = FB,$$

from which $\underline{\Sigma}(A, FB, A, FB) = 1$ and $g(FB, A) = g(\Sigma_{AFB}A, A) = 0$.

LEMMA 1.2. For any unit vector $B \in \sigma_A$ set $C = FB \in \sigma_A$, then

$$\Sigma_{AB}\Sigma_{AC}=F-SF.$$

Proof. If we take an arbitrary vector $D \in T_p M$ then, using (3) and (6), we have

$$\begin{aligned}\Sigma_{AB}\Sigma_{AC}D &= \Sigma_{AB}\Sigma_{AFB}D = -\Sigma_{AB}\Sigma_{AFB}F^2D \\ &= -\Sigma_{AB}\Sigma_{AB}FD = (I-S)FD.\end{aligned}$$

LEMMA 1.3. For any orthonormal pair $\{B, C\} \in \sigma_A$,

$$\Sigma_{AB}\Sigma_{AC} = -\Sigma_{AC}\Sigma_{AB}.$$

Proof. First, using (3), we have

$$\begin{aligned}\Sigma_{AB+C}^2 &= \Sigma(A, B+C, A, B+C)(-I+S) \\ &= 2(-I+S).\end{aligned}$$

On the other hand, we obtain

$$\begin{aligned}\Sigma_{AB+C}^2 &= (\Sigma_{AB} + \Sigma_{AC})^2 \\ &= 2(-I+S) + (\Sigma_{AB}\Sigma_{AC} + \Sigma_{AC}\Sigma_{AB}).\end{aligned}$$

Thus we have $\Sigma_{AB}\Sigma_{AC} + \Sigma_{AC}\Sigma_{AB} = 0$.

LEMMA 1.4. $\dim[\sigma_A] = 2$.

Proof. Take B and C as in Lemma 1.1 and assume that there is a unit vector $D \in [\sigma_A]$ such that D is orthogonal to B and C . They by Lemmas 1.2 and 1.3 we have

$$\Sigma_{AB}\Sigma_{AC}\Sigma_{AD} = \Sigma_{AD}\Sigma_{AB}\Sigma_{AC},$$

and so

$$(F-SF)\Sigma_{AD} = \Sigma_{AD}(F-SF).$$

Thus, using (1) and (4), we obtain

$$F\Sigma_{AD} = \Sigma_{AD}F,$$

which contradicts (5). Therefore, $[\sigma_A]$ is necessarily of dimension 2.

LEMMA 1.5. For any vectors $B, C \in T_p M$, satisfying $\underline{\Sigma}(B, C, B, C) = 1$, $\Sigma_{BC}A \in \sigma_A$.

Proof. Using (8), we have

$$\Sigma_{A\Sigma_{BC}}A = \Sigma_{BC}A,$$

from which it follows that $\Sigma_{BC}A \in \sigma_A$.

LEMMA 1.6. Take a unit vector $A \in T_p M$ with $SA=0$ and a unit vector $B \in \sigma_A$. Put $C=FB \in \sigma_A$. The $\Sigma_{AB}D$ and $\Sigma_{AC}D$ are orthonormal, where D is an arbitrary unit vector at p such that $SD=0$.

Proof. $g(\Sigma_{AB}D, \Sigma_{AB}D) = -g(\Sigma_{AB}^2 D, D) = g(D - SD, D) = 1$ and similarly $\Sigma_{AC}D$ is also a unit vector. Finally

$$\begin{aligned} g(\Sigma_{AB}D, \Sigma_{AC}D) &= -g(\Sigma_{AB}D, \Sigma_{AFB}F^2D) = -g(\Sigma_{AB}D, \Sigma_{AB}FD) \\ &= g(\Sigma_{AB}D, F\Sigma_{AB}D) = 0. \end{aligned}$$

Summing up Lemmas 1.4, 1.5 and 1.6, we have

PROPOSITION 1. Take a unit vector $A \in T_p M$ such that $SA=0$ and a unit vector $B \in \sigma_A$. Put $C=FB \in \sigma_A$. Then, for any unit vector $D \in T_p M$ with $SD=0$, $\Sigma_{AB}D$ and $\Sigma_{AC}D$ form an orthonormal basis of $[\sigma_D]$.

LEMMA 1.7. Take A, B and C as in Proposition 1. Then, for any $D, E \in T_p M$,

$$\Sigma_{DE} = \underline{\Sigma}(A, B, D, E)\Sigma_{AB} + \underline{\Sigma}(A, C, D, E)\Sigma_{AC}.$$

Proof. When D (or E) satisfies $SD=D$ (or $SE=E$), then both sides of the equation above vanish because of (4). So, D and E may be assumed to satisfy $SD=SE=0$ and also that D and E are unit. First, we consider the case in which E is orthogonal to σ_D . Linearizing (3) we have $\Sigma_{XY}\Sigma_{XZ} + \Sigma_{XZ}\Sigma_{XY} = 2\underline{\Sigma}(X, Y, X, Z)(-I+S)$. Thus if $Y \in \sigma_D$, Σ_{DE} anti-commutes with Σ_{DY} and Σ_{DFY} and hence Σ_{DE} commutes with $\Sigma_{DY}\Sigma_{DFX}$ which by Lemma 1.2 is equal to $F-SF$. Therefore using (1) and (4)

$$F\Sigma_{DE} = (F-SF)\Sigma_{DE} = \Sigma_{DE}(F-SF) = \Sigma_{DE}F,$$

from which by (5) and the non-singularity of F we have $\Sigma_{DE}=0$ and again both sides of the above equation vanish.

Finally we consider the case where $E \in \sigma_D$. For simplicity set $a = g(\Sigma_{AB}D, E)$ and $b = g(\Sigma_{AC}D, E)$. Then as $\{\Sigma_{AB}D, \Sigma_{AC}D\}$ is an orthonormal basis of $[\sigma_D]$,

$$E = a\Sigma_{AB}D + b\Sigma_{AC}D.$$

Using (8) we have

$$\begin{aligned} \Sigma_{DE}A &= a\Sigma_{D\Sigma_{AB}D}A + b\Sigma_{D\Sigma_{AC}D}A \\ &= a\Sigma_{AB}A + b\Sigma_{AC}A \\ &= aB + bC. \end{aligned}$$

Using (8) again

$$\Sigma_{DE} = \Sigma_{A\Sigma_{DE}A} = a\Sigma_{AB} + b\Sigma_{AC},$$

which is the desired formula.

Take a suitable coordinate neighborhood u of an arbitrary point p of M and a unit vector field A in u . Then there is in u a unit vector field B belonging to σ_A at each point of u . On putting $C=FB \in \sigma_A$ we define locally in u two tensor fields G and H of type $(1, 1)$ respectively by

$$G = \Sigma_{AB}, \quad H = \Sigma_{AC}.$$

Then setting $F^H = F - FS$ and using (3) and (4) and Lemma 1.3, we have

$$\begin{aligned} (F^H)^2 &= G^2 = H^2 = -I + S, \\ GH &= -HG = F^H, \quad HF^H = -F^H H = G, \quad F^H G = -GF^H = H, \\ F^H S &= SF^H = GS = SG = HS = SH = 0. \end{aligned} \quad (1.3)$$

Next, (1), (2), and (7) imply

$$\begin{aligned} g(F^H X, Y) &= -g(F^H Y, X), \\ g(GX, Y) &= -g(GY, X), \quad g(HX, Y) = -g(HY, X), \end{aligned}$$

for all X and Y . By Lemma 1.7, a local expression for Σ_{XY} in u is the following

$$\Sigma_{XY} = g(GX, Y)G + g(HX, Y)H. \quad (1.4)$$

We now take another coordinate neighborhood $u'(u \cap u' \neq \emptyset)$ and define G' and H' as in u , say $G' = \Sigma_{A'B'}$ and $H' = \Sigma_{A'C'}$. By the formula of Lemma 1.7

$$\begin{aligned} \Sigma_{A'B'} &= \underline{\Sigma}(A, B, A', B')\Sigma_{AB} + \underline{\Sigma}(A, C, A', C')\Sigma_{AC}, \\ \Sigma_{A'C'} &= \underline{\Sigma}(A, B, A', C')\Sigma_{AB} + \underline{\Sigma}(A, C, A', C')\Sigma_{AC}. \end{aligned}$$

Setting $a = \underline{\Sigma}(A, B, A', B')$ and $b = \underline{\Sigma}(A, C, A', C')$ we have that

$$1 = g(\Sigma_{A'B'}A', B') = \underline{\Sigma}(A, B, A', B')^2 + \underline{\Sigma}(A, C, A', C')^2 = a^2 + b^2$$

and

$$\begin{aligned} \underline{\Sigma}(A, C, A', C') &= -g(\Sigma_{AFB}F^2A', FB') = -g(\Sigma_{AB}FA', FB') \\ &= g(F\Sigma_{AB}A', FB') = g(\Sigma_{AB}A', B') = a, \\ \underline{\Sigma}(A, B, A', C') &= -g(\Sigma_{AFB}FA', F^2C') = -g(F\Sigma_{AC}A', FB') \\ &= -g(\Sigma_{AC}A', B') = -b, \end{aligned}$$

so that $G' = aG + bH$ and $H' = -bG + aH$.

THEOREM 1. *Let (M, G, F) be an almost Hermitian manifold. Then M is an almost complex contact manifold if and only if M admits a global tensor field Σ of type $(1, 3)$ and a projection tensor field S of rank 2 satisfying 1)-9).*

§ 2. Horizontal and Vertical Tensors

Given a vector field X on an almost Hermitian manifold (M, g, F) with almost complex contact structure (g, F, Σ, S) , $X^V = SX$ and $X^H = X - X^V$ will be called the *vertical* and the *horizontal parts* of X , respectively. For a 1-form ω , $\omega^V = \omega \circ S$ and $\omega^H = \omega - \omega^V$ will be called the *vertical* and the *horizontal parts* of ω , respectively. We now define, for a function f , $f^H = f^V = f$. Then we easily have

$$(2.1) \quad \begin{aligned} (fX + hY)^H &= f^H X^H + h^H Y^H, & (fX + hY)^V &= f^V X^V + h^V Y^V, \\ (f\omega + h\pi)^H &= f^H \omega^H + h^H \pi^H, & (f\omega + h\pi)^V &= f^V \omega^V + h^V \pi^V, \end{aligned}$$

where f, h are arbitrary functions and ω, π are arbitrary 1-forms.

We now define the *horizontal part* T^H of an arbitrary tensor field T . Assume that the operation of taking the horizontal part satisfies

$$(2.2) \quad (P+Q)^H = P^H + Q^H, \quad (P \otimes U)^H = P^H \otimes U^H,$$

where P and Q are arbitrary tensor fields of the same type and U another arbitrary tensor field, then by using (2.1) we can inductively define the horizontal part T^H of an arbitrary tensor field T on M .

§ 3. Almost Complex Contact Structures which are Projectable

The Riemannian connection is denoted by ∇ in a Kählerian manifold M with almost complex contact structure (g, F, Σ, S) . We define a tensor field P of type (1, 2) by

$$(3.1) \quad P_X Y = ((\nabla_Y S)X)^H.$$

Note that

$$(3.2) \quad S P_X = 0.$$

Next, differentiating covariantly $S^2 = S$ we have

$$(3.3) \quad P_{S X} = P_X$$

and differentiating covariantly (1)

$$(3.4) \quad P_{F X} = F P_X.$$

LEMMA 3.1. *When $P=0$, a Kählerian manifold M of complex dimension $2m+1$ with almost complex contact structure (g, F, Σ, S) is locally a product of two Kählerian manifolds of complex dimensions $2m$ and 1 respectively.*

Proof. If $P=0$, (3.1) implies

$$(\nabla_Y (SX))^H = ((\nabla_Y S)X)^H = 0,$$

which means that the distribution determined by S and its complement are parallel. This with $SF=FS$ proves the lemma.

We now consider the following conditions :

- (P1) for any vector $A \in T_p M$, there are two vectors $B, C \in T_p M$ such that $P_A = \Sigma_{BC}$, $\underline{\Sigma}(B, C, B, C) = ag(SA, SA)$ with constant a ;
(P2) $(\nabla_{SX} \underline{\Sigma})^H = 0$.

When an almost complex contact structure (g, F, Σ, S) satisfies the conditions (P1) and (P2), it is said to be *projectable*.

In this section, the almost complex contact structure (g, F, Σ, S) is assumed to be projectable. Then 3)-4) and (P1) imply

$$(3.5) \quad P_x^2 = ag(SX, SX)(-I + S)$$

for some a and

$$(3.6) \quad P_x S = 0$$

is equivalent to

$$(3.7) \quad S(\nabla_{SY} S) = \nabla_{SY} S.$$

Thus we now have from (1) and (3.7)

PROPOSITION 2. *In a Kählerian manifold M with almost complex contact structure (g, F, Σ, S) which satisfies (P1), the distribution determined by S is integrable and each of its integral submanifolds is totally geodesic and holomorphic.*

Since (P1) is satisfied, restricting ourselves to a coordinate neighborhood u in which (1.4) is established, we find

$$(3.8) \quad P_x = c(u(X)G + v(X)H)$$

with local 1-forms u and v defined in u , where the associated vector fields U of u and V of v satisfy $\|U\|^2 = \|V\|^2 = 1$, $g(U, V) = 0$, i. e.,

$$(3.9) \quad S = u \otimes U + v \otimes V.$$

(3.8) implies that

$$(3.10) \quad (\nabla(SX))^H = c(u(x)G + v(x)H).$$

The fundamental 2-form Φ of the Kählerian manifold (M, g, F) is defined by $\Phi(X, Y) = g(FX, Y)$. We now define in M a tensor field \underline{A} of type $(0, 4)$ by

$$(3.11) \quad \underline{A} = \Phi^H \otimes \Phi^H + \underline{\Sigma},$$

which is horizontal, that is, $\underline{A}^H = \underline{A}$. Then, using (1.3) and (3.8), we can verify that in u

$$P_U \cdot \underline{A} = 0, \quad P_V \cdot \underline{A} = 0,$$

where $P_X \cdot$ denotes the action of P_X as a derivation. Thus, using (3.9), we obtain

$$(3.13) \quad P_X \cdot \underline{A} = 0.$$

Since $\nabla F = 0$, we find

$$(3.14) \quad (\nabla_{S_X} \underline{A})^H = 0$$

as a consequence of (P2). As is well known, the Lie derivative $\mathcal{L}_{S_X} \underline{A}$ is given by

$$\mathcal{L}_{S_X} \underline{A} = \nabla_{S_X} \underline{A} + P_X \cdot \underline{A}$$

(See e. g. Yano [8]). Thus we have

$$(3.15) \quad (\mathcal{L}_{X^V} \underline{A})^H = 0.$$

LEMMA 3.2. *If an almost complex contact structure (g, F, Σ, S) is projectable, then*

$$(\mathcal{L}_{X^V} \underline{A}^H)^H = 0.$$

On the other hand, by Proposition 2, each integral submanifold of the distribution determined by S is totally geodesic. Thus we have (see Ishihara and Konishi [5])

LEMMA 3.3. *If an almost complex contact structure (g, F, Σ, S) is projectable, then*

$$(\mathcal{L}_{X^V} g^H)^H = 0.$$

We now put

$$A = \Phi \otimes F + \Sigma.$$

Then Lemmas 3.2 and 3.3 imply

LEMMA 3.4. *If an almost complex contact structure (g, F, Σ, S) is projectable, then*

$$(\mathcal{L}_{X^V} A^H)^H = 0.$$

§ 4. Submersion of a Kählerian Manifold with Almost Complex Contact Structure

Let (M, G, F) be a Kählerian manifold of complex dimension $2m+1$ with almost complex contact structure (g, F, Σ, S) , which is projectable, and \tilde{M} a manifold of real dimension $4m$. Suppose that there is a differential mapping $\pi: M \rightarrow \tilde{M}$ which is of rank $4m$ everywhere and satisfies $\pi(M) = \tilde{M}$ and that for each point p of \tilde{M} , $\pi^{-1}(p)$ is a connected integral submanifold of the distribution

determined by S . In such a case, the Kählerian manifold M with almost complex contact structure is said to have a *fibred Riemannian structure* $\pi: M \rightarrow \tilde{M}$ and \tilde{M} is called the *base space*. When M is compact and the distribution \mathcal{D} determined by S is regular, M has a fibred Riemannian structure if \tilde{M} is defined as the set of all maximal integral submanifolds of \mathcal{D} , $\pi: M \rightarrow \tilde{M}$ being defined by $\pi(p) = \mathcal{D}p$, $p \in M$, where $\mathcal{D}p$ is the maximal integral submanifold passing through p , and \tilde{M} is naturally topologized.

Consider a Kählerian manifold M with almost complex contact structure (g, F, Σ, S) , which is projectable, and with fibred Riemannian structure $\pi: M \rightarrow \tilde{M}$. Then, taking account of arguments developed in [5], we see by Lemma 3.4 that the tensor field A is projectable in M and its projection is a tensor field \tilde{A} of type (1, 3) in the base space \tilde{M} . The metric tensor g in M is, by Lemma 3.3, projectable and its projection \tilde{g} defines a Riemannian structure on \tilde{M} . Thus, (2)-(9) implies that (\tilde{g}, \tilde{A}) is an almost quaternionic structure in the base space \tilde{M} (see Blair and Showers [2]). Thus, summing up, we have

THEOREM 2. *Suppose that (M, g, F) is a Kählerian manifold with almost complex contact structure (g, F, Σ, S) , which is projectable. Assume moreover that (M, g, F) has a fibred Riemannian structure $\pi: M \rightarrow \tilde{M}$. Then (\tilde{g}, \tilde{A}) is an almost quaternionic structure in the base space \tilde{M} , where \tilde{g} and \tilde{A} are the projections of g and A , respectively.*

If in a Kählerian manifold M satisfying the conditions given in Theorem 2

$$(\nabla A^H)^H = 0$$

holds, then the projection \tilde{A} of A in \tilde{M} is covariantly constant. Thus in such a case (\tilde{g}, \tilde{A}) is a quaternionic Kählerian structure (see Ishihara [4]). Thus we have

THEOREM 3. *If, in a Kählerian manifold M satisfying the conditions given in Theorem 2, $(\nabla A^H)^H = 0$, then (\tilde{g}, \tilde{A}) is a quaternionic Kählerian structure in the base space \tilde{M} .*

Taking account of Lemma 3.1, we easily have

PROPOSITION 3. *If a Kählerian manifold M of complex dimension $2m+1$ with almost complex contact structure (g, F, Σ, S) , which is projectable, satisfies the condition $P=0$, then M is locally a product of Kählerian manifolds (M_1, g_1, F_1) of complex dimension $2m$ and (M_2, g_2, F_2) of complex dimension 1, where M_1 admits quaternion structure (g_1, A_1) .*

PROPOSITION 4. *If, in a Kählerian manifold M satisfying the conditions given in Proposition 3 $(\nabla A^H)^H = 0$ then M_1 admits a quaternionic Kählerian structure (g_1, A_1) with vanishing Ricci tensor (see Ishihara [4]).*

BIBLIOGRAPHY

- [1] D. E. BLAIR, Geometry of Manifolds with Structural Group $u(n) \times O(S)$, J. Diff. Geom. 4 (1970) 155-167.
- [2] D. E. BLAIR AND D. K. SHOWERS, A Note on Quaternionic Geometry, to appear.
- [3] W. M. BOOTHBY, Homogeneous Complex Contact Manifolds, Proc. Symposia in Pure Math. of A. M. S., Vol. III, Diff. Geom., 1961, 144-154.
- [4] S. ISHIHARA, Quaternion Kählerian manifolds, J. Diff. Geom. 9 (1974) 483-500.
- [5] S. ISHIHARA AND M. KONISHI, Complex Almost Contact Structures and Fiberings, to appear.
- [6] S. KOBAYASHI, Remarks On Complex Contact Manifolds, Proc. of A. M. S. 10 (1959), 164-167.
- [7] S. SASAKI, Almost Contact Manifolds I, II, III, Lecture Notes, Tôhoku University, 1965.
- [8] K. YANO, Theory of Lie Derivatives, North Holland, Amsterdam, 1957.

MICHIGAN STATE UNIVERSITY
EAST LANSING
DEPARTMENT OF MATHEMATICS
TOKYO INSTITUTE OF TECHNOLOGY
MICHIGAN STATE UNIVERSITY
EAST LANSING