

ON A METRIC INDUCED BY ANALYTIC CAPACITY

BY NOBUYUKI SUITA

Dedicated to Professor Yukinari Tôki on his 60th birthday

1. In our previous paper [6] we gave a conjecture that the metrics $c_{\beta}(z)|dz|$ and $\sqrt{\pi\tilde{K}(z, z)|dz|}$ have negative curvatures ≤ -4 ; here $c_{\beta}(z)$ and $\tilde{K}(z, z)$ are the capacity and the Bergman kernel of exact analytic differentials on an open (non-trivial) Riemann surface. In the present paper we shall show that the curvature of the metric $c_B(z)|dz| \leq -4$ for plane regions $\Omega \notin O_{AB}$, where $c_B(z)$ denotes the analytic capacity of Ω at z . In order to verify $c_B(z) \in C^2$ we prove that $c_B(z)$ is real analytic. This enables us to answer a question of Havinson [4], namely “Does the sequence of extremal functions ψ_n in the dual problem of Schwarz’s lemma in Ω_n converge as $\{\Omega_n\}$ exhausts Ω ?”.

2. Let Ω be a plane region $\notin O_{AB}$. The analytic capacity $c_B(\zeta)$ is given by $\sup |f'(\zeta)|$ in the family of analytic functions satisfying $f(\zeta)=0$ and $|f(z)| \leq 1$. Let $\{\Omega_n\}$ be a canonical exhaustion of Ω such that the boundary of Ω_n consists of a finite number of analytic curves. Let $c_n(\zeta)$ be the analytic capacity of Ω_n . Then $\{c_n(\zeta)\}$ is decreasing and tends to $c_B(\zeta)$. There exist extremal functions f_n such that $f'_n(\zeta)=c_n(\zeta)$ and $f'_0(\zeta)=c_B(\zeta)$. It is known that those extremal functions are unique [4]. The function $\log c_B(\zeta)$ is subharmonic [2].

In every Ω_n there exists the Szegő kernel $k_n(z, \zeta)$ and its adjoint kernel $l_n(z, \zeta)$ [3]. $k_n(z, \zeta)$ is hermitian and analytic with respect to z and $\bar{\zeta}$. Further the following facts are known [3]:

$$(1) \quad f_n(z) = \frac{k_n(z, \zeta)}{l_n(z, \zeta)} \quad \text{with} \quad c_n(\zeta) = 2\pi k_n(\zeta, \zeta)$$

and

$$|k_n(z, \zeta)|^2 \leq |k_n(z, z)| |k_n(\zeta, \zeta)|.$$

Thus $k_n(z, \zeta)$ is uniformly bounded on every compact subset and hence forms a normal family of analytic functions of two variables $z, \bar{\zeta}$. We will show that $\{k_n(z, \zeta)\}$ converges to a function $k(z, \zeta)$ uniformly on every compact subset of Ω .

Suppose that there exist two limit functions $k(z, \zeta)$ and $k^*(z, \zeta)$. We may assume $0 \in \Omega$. The difference $k(z, \zeta) - k^*(z, \zeta)$ has an expansion in a polydisc $\{|z| < r\} \times \{|\zeta| < r\}$

Received May 19, 1972.

$$k(z, \zeta) - k^*(z, \zeta) = \sum_{\nu, \mu=0}^{\infty} a_{\nu\mu} z^{\nu} \bar{\zeta}^{\mu} \quad \text{with} \quad a_{\nu\mu} = \bar{a}_{\mu\nu}.$$

Since $k(z, z) = k^*(z, z) = c_B(z)/2\pi$, by setting $z = |z|e^{i\theta}$ we have $a_{00} = 0$ and by induction

$$\sum_{\nu+\mu=n} a_{\nu\mu} e^{i(\nu-\mu)\theta} = 0.$$

Hence $a_{\nu\mu} = 0$ and $k(z, \zeta)$ coincides with $k^*(z, \zeta)$. $\{f_n(z)\}$ clearly converges to $f_0(z)$ uniformly on every compact subset of Ω . Thus the sequence $\{l_n(z, \zeta)\}$ converges to a meromorphic function $l(z, \zeta)$ uniformly on every compact subset of $\Omega - \zeta$. This yields

THEOREM 1. *The sequence $\{k_n(z, \zeta)\}$ converges uniformly on every compact subset of Ω . The analytic capacity $c_B(z)$ is real analytic. Further the sequence $\{l_n(z, \zeta)\}$ converges uniformly on every compact subset of $\Omega - \zeta$.*

3. The last statement gives an affirmative answer to a question of Havinson [4]. For a moment suppose $\infty \in \Omega$. Take an exhaustion $\{\Omega_n\}$ of Ω with $\infty \in \Omega_1$. Let $\tilde{E}_1(\Omega_n)$ be the family of functions ϕ analytic in Ω_n satisfying $\phi(\infty) = 0$ and

$$\int_{\partial\Omega_n} |\phi| ds \leq 1,$$

where ds is the length element on $\partial\Omega_n$.

As the duality relation in Schwarz's lemma, Havinson [4] showed

$$c_n(\infty) = \min_{\phi \in \tilde{E}_1(\Omega_n)} \int_{\partial\Omega_n} |1 + \phi| ds$$

and that the extremal function ϕ_n exists uniquely in $\tilde{E}_1(\Omega_n)$ and satisfies

$$-if_n(z)(1 + \phi_n(z))dz = |1 + \phi_n(z)|ds$$

where f_n is the extremal function in no. 2 with $\zeta = \infty$. He proved that every subsequence of $\{\phi_n\}$ contains a convergent subsequence and conjectured that $\{\phi_n\}$ itself is convergent as $n \rightarrow \infty$.

On the other hand the square of $l_n(z, \infty)$ is expressed as

$$l_n(z, \infty)^2 = \frac{1}{4\pi^2} \left(1 + \frac{b_1}{z} + \dots \right)$$

near the point at infinity and has no zeros in Ω_n . The fundamental relation $l_n(z, \infty)ds = i\overline{k_n(z, \infty)}d\bar{z}$ along $\partial\Omega_n$ [3] and (1) show $-il_n(z, \infty)^2 f_n(z)dz > 0$ along $\partial\Omega_n$ which implies that $(1 + \phi_n(z))/l_n(z, \infty)^2$ is real and that $\phi_n(z) = 4\pi^2 l_n(z, \infty)^2 - 1$. The validity of his conjecture is deduced from Theorem 1.

4. We now turn to the estimation of the curvature of the metric $c_B(z)|dz|$. Let $\lambda|dz|$ and $\mu|dz|$ with $\lambda, \mu \geq 0$ be two metrics on a plane region Ω . Following

Ahlfors [1], we call $\mu|dz|$ a *supporting metric* of $\lambda|dz|$ at the point ζ if 1) $\lambda = \mu$ at ζ and 2) $\lambda - \mu \geq 0$ in a neighborhood of ζ . Then we prepare

LEMMA. *If $\mu|dz|$ is a supporting metric of $\lambda|dz|$ and if both λ and μ are of class C^2 , then the curvature of $\mu|dz|$ dominates that of $\lambda|dz|$.*

Proof. The curvature $\kappa(\lambda; z)$ of $\lambda|dz|$ at z is given by

$$\kappa(\lambda; z) = \frac{-\Delta \log \lambda}{\lambda^2}.$$

Since $\log(\lambda/\mu)$ assume a local minimum at $z = \zeta$, we have $\Delta \log(\lambda/\mu) \geq 0$, which implies the assertion.

We state

THEOREM 2. *The curvature of the metric $c_B(z)|dz| \leq -4$.*

Proof. Let $f_0(z)$ be the extremal function with $c_B(\zeta) = f_0'(\zeta)$. Set $F(z, \eta) = (f_0(z) - f_0(\eta)) / (1 - \bar{f}_0(\eta)f_0(z))$. We have $|F(z, \eta)| \leq 1$ and

$$\frac{|f_0'(\eta)|}{1 - |f_0(\eta)|^2} \leq c_B(\eta).$$

The metric $F'(z, z)|dz|$ has the curvature -4 at every point z except for the zeros z_0 of f_0' . Thus $c_B(z)|dz|$ has a supporting metric $|F'(z, z)|dz|$ at every point $\zeta \in \Omega$. We have $\kappa(c_B; z) \leq -4$.

There remains a problem to decide the equality statement in Theorem 2. It is plausible that if $\kappa(c_B; z) = -4$ at one point z , Ω is conformally equivalent to the unit disc less a (possible) closed set expressed as a countable union of compact $N_{\mathfrak{B}}$ sets.

5. A recent development of the theory of conformal metrics was given by Heins [5]. He defined an S-K metric by a metric $\lambda|dz|$ such that

1) λ is nonnegative and upper semi-continuous, and

$$2) \quad \lim_{r \rightarrow 0} \frac{1}{r^2} \left(\frac{1}{2\pi} \int_0^{2\pi} \log \lambda(\zeta + re^{i\theta}) d\theta - \log \lambda(\zeta) \right) \geq \lambda(\zeta)^2.$$

As he pointed out, the operator in the left hand side in 2) is one fourth of the generalized lower Laplacian and if $\log \lambda \in C^2$, the condition 2) reduces to

$$\Delta \log \lambda(\zeta) \geq 4\lambda(\zeta)^2.$$

A sufficient condition for a metric to be S-K is the existence of a supporting metric with curvature ≤ -4 at every point [5]. From this Theorem 2 again follows.

This remark works for the metric $c_{SB}(z)|dz|$ associated with bounded univalent functions. Let \mathcal{F} be the family of univalent functions satisfying $|g| \leq 1$ on Ω . $c_{SB}(\zeta)$ is defined by

$$c_{SB}(\zeta) = \sup_{g \in \mathcal{F}} |g'(\zeta)|.$$

If $\mathcal{F} \neq \phi$, there exists an extremal function satisfying $c_{SB}(\zeta) = g'_0(\zeta)$ with $J_0(\zeta) = 0$ in \mathcal{F} . Thus we can construct a supporting metric at every point $\zeta \in \Omega$ as before. However $c_{SB}(z)$ is not necessarily differentiable (an example is $c_{SB}(z)$ for an annulus $1 < |z| < r$ at the points on the circle $|z| = \sqrt{r}$). Hence we must stop at the result that $c_{SB}(z)|dz|$ is an S-K metric on Ω .

REFERENCES

- [1] AHLFORS, L. V., An extension of Schwarz's lemma. *Trans Amer. Math. Soc.* **43** (1938), 359-364.
- [2] AHLFORS, L. V., AND A. BEURLING, Conformal invariants and function-theoretic null-sets. *Acta Math.* **83** (1950), 101-129.
- [3] BERGMAN, S., The kernel function and conformal mapping. *Math. Surveys* **5**, Amer. Math. Soc. Providence, R. I. (1950).
- [4] HAVINSON, S. Ja., Analytic capacity of sets, joint nontriviality of various classes of analytic functions and the Schwarz lemma in arbitrary domains. *Amer. Math. Soc. Transl. (2)* **43** (1964), 215-266.
- [5] HEINS, M., On a class of conformal metrics. *Nagoya Math. J.* **21** (1962), 1-60.
- [6] SUITA, N., On capacities and kernels of Riemann surfaces. *Arch. Rat. Mech. Anal.* **46** (1972), 212-217.

DEPARTMENT OF MATHEMATICS,
TOKYO INSTITUTE OF TECHNOLOGY.