

A NOTE ON THE SET OF PERIODS FOR KLEIN BOTTLE MAPS

JAUME LLIBRE

By using the Nielsen fixed point theory we characterize the set of periods attained by continuous self-maps on the Klein bottle belonging to a given homotopy class.

1. Introduction. In dynamical systems, it is often the case that topological information can be used to study qualitative and quantitative properties (like the set of periods) of the system. This note deals with the problem of determining the set of periods (of the periodic points) of a continuous self-map of the Klein bottle. Our interest in this problem comes from the fact that the unique manifolds in dimensions 1 and 2 with zero Euler characteristic are the circle, torus and Klein bottle, and for the first two the structure of the set of periods of their continuous self-maps has been determined.

To fix terminology, suppose f is a continuous self-map on the manifold M . A *fixed point* of f is a point x in M such that $f(x) = x$. We shall call x a *periodic point of period n* if x is a fixed point of f^n but is not fixed by any f^k , for $1 \leq k < n$. We denote by $\text{Per}(f)$ the set of natural numbers corresponding to periods of periodic orbits of f .

Even for continuous self-maps f on the circle the relation between the degree of f and the set $\text{Per}(f)$ is interesting and nontrivial (see [5], [3] and for more details [2]). Let \mathbb{N} denote the set of natural numbers. Suppose f has degree d .

- (1) For $d \notin \{-2, -1, 0, 1\}$, $\text{Per}(f) = \mathbb{N}$.
- (2) For $d = -2$, $\text{Per}(f)$ is either \mathbb{N} or $\mathbb{N} \setminus \{2\}$.
- (3) For $d = -1, 0$, $\text{Per}(f) \supset \{1\}$.
- (4) For $d = 1$, the set $\text{Per}(f)$ can be empty.

Recently, in [1] these results have been extended to continuous self-maps on the 2-dimensional torus, and many of them to the m -dimensional torus with $m > 2$.

The goal of this note is to provide a similar description of the set of periods for continuous self-maps on the Klein bottle, or simply *Klein bottle maps*.

Let K be the Klein bottle and let f be a Klein bottle map. If p is a base point for K , then f is homotopic to a continuous map h , $f \sim h$, such that $h(p) = p$ and a presentation of the group $\Pi(K, p)$ is given by two generators a and b and the relation $abab^{-1} = 1$. Halpern in [6] shows that $h_p(a) = a^u$ and $h_p(b) = a^w b^v$ for some integers u , v and w , where h_p is the induced endomorphism on $\Pi(K, p)$. In what follows u and v will be called the *integers associated to f* .

Now our main result can be stated as follows. Here, $2\mathbb{N}$ denotes the set of all even natural numbers.

THEOREM. *Let f be a Klein bottle map and let u and v its associated integers.*

(1) *Suppose $|u| > 1$. Then*

$$\begin{aligned} \text{Per}(f) &= \mathbb{N} && \text{if } |v| \neq 1, \text{ and} \\ \text{Per}(f) &\supset \mathbb{N} \setminus 2\mathbb{N} && \text{if } v = -1. \end{aligned}$$

Furthermore, the last result cannot be improved without additional hypothesis on f .

(2) *Suppose $|u| \leq 1$. Then*

$$\begin{aligned} \text{Per}(f) &= \mathbb{N} && \text{if } v \notin \{-2, -1, 0, 1\}, \\ \text{Per}(f) &\supset \mathbb{N} \setminus \{2\} && \text{if } v = -2, \\ \text{Per}(f) &\supset \{1\} && \text{if } v \in \{-1, 0\}. \end{aligned}$$

Furthermore, these last two results cannot be improved without additional hypotheses on f .

(3) *If $v = 1$ then the set $\text{Per}(f)$ can be empty.*

This theorem will be proved in the next section. Its proof is based in the results of Halpern [6] on the Nielsen numbers of Klein bottle maps. Unfortunately this nice paper of Halpern remains unpublished.

As we shall see the Nielsen numbers of the Klein bottle map f only depend on its associated integers. Then, since the Nielsen numbers are homotopy invariants, the characterization of the set of periods given in the theorem works for all Klein bottle maps homotopic to f .

I thank Christopher McCord for his comments on the Nielsen fixed point theory.

2. Periods and Nielsen numbers. We start by recalling the definition of Nielsen number, for more details see, for instance, [4], [7] or [8]. Suppose that $f: E \rightarrow E$ is a continuous map of a compact ENR. The Nielsen number $N(f)$ is defined as follows. First an equivalence relation \sim is defined on the set F of fixed points of f . Two fixed points x, y are equivalent, $x \sim y$, provided there is a path γ in E from x to y such that $f \circ \gamma$ and γ are homotopic and the homotopy fixes endpoints. The set of equivalence classes F/\sim is known to be finite and each equivalence class is compact. Each of these equivalence classes will be called a fixed point class. Using a fixed point index, we may assign an index to each fixed point class. The *Nielsen number* is defined as the number of fixed point classes with nonzero index.

Notice that from the definition of Nielsen number, f has at least $N(f)$ fixed points. If $g \sim f$ then g has at least $N(f)$ fixed points, and $N(g) = N(f)$.

In what follows f will be a Klein bottle map, and u and v its associated integers. Halpern in [6] proves that

$$(a) \quad N(f^n) = \begin{cases} |u^n(v^n - 1)| & \text{if } |u| > 1, \\ |v^n - 1| & \text{if } |u| \leq 1, \end{cases}$$

for all $n \geq 1$.

Define A_n for $n \geq 1$ inductively as follows. For $u \neq 1$, set

$$A_1 = N(f)$$

and

$$A_n = N(f^n) - \sum_{k < n, k|n} A_k \quad \text{for } n > 1,$$

where $k|n$ denotes k divides n .

For $u = 1$, set

$$A_1 = N(f)$$

and

$$A_n = N(f^n) - \sum_{k < n, k|n, \frac{n}{k} \text{ odd}} A_k \quad \text{for } n > 1.$$

For a Klein bottle map g and $n \geq 1$, let $P_n(g)$ be the set of periodic points of g of period n . The cardinality of a set S is denoted by $\text{Card}(S)$. Halpern also shows that

$$(b) \quad \min\{\text{Card}(P_n(g)): g \sim f\} = A_n \quad \text{for all } n \geq 1.$$

In particular, from (b) it follows that $A_n \geq 0$ for all $n \geq 1$.

Furthermore, except for two special cases $u \geq 1$ and $v = -1$, or $u = 1$ and $|v| > 1$, Halpern shows that there exists a Klein bottle map $g \sim f$ such that

$$(c) \quad \text{Card}(P_n(g)) = A_n \quad \text{for all } n \geq 1.$$

Finally, he shows for each $m \geq 1$ that there exists a Klein bottle map $g \sim f$ such that

$$(d) \quad \text{Card}(P_n(g)) = A_n \quad \text{for } 1 \leq n \leq m.$$

The next proposition plays a main role in the proof of Theorem.

PROPOSITION. *The following statements hold.*

(1) *Suppose $|u| > 1$. If $|v| \neq 1$ then*

$$(e) \quad \sum_{k < n, k|n} N(f^k) < N(f^n)$$

for all $n > 1$. If $v = -1$ then (e) holds for all odd $n > 1$.

(2) *Suppose $|u| \leq 1$. If $v \notin \{-2, -1, 0, 1\}$ then (e) holds for all $n > 1$. If $v = -2$ then (e) holds for all $n > 2$.*

Proof. Suppose $|u| > 1$. We separate the proof of (1) into four cases.

Case 1. $v = 0$. From (a) we obtain (e) as follows:

$$\sum_{k=1}^{n-1} N(f^k) = \sum_{k=1}^{n-1} |u|^k = \frac{|u|^n - |u|}{|u| - 1} < |u|^n = N(f^n).$$

Case 2. $v = -1$. Then $N(f^n)$ is 0 for n even, and $2|u|^n$ for n odd. Hence, for n odd the proof of (e) follows as in Case 1.

Case 3. $v > 1$. For $n > 1$ we have

$$\begin{aligned} \sum_{k=1}^{n-1} N(f^k) &= \sum_{k=1}^{n-1} (|u|^k v^k - |u|^k) = \frac{|u|^n v^n - |u|v}{|u|v - 1} - \frac{|u|^n - |u|}{|u| - 1} \\ &< \frac{|u|^n v^n - |u|v - |u|^n + |u|}{|u|v - 1} < \frac{|u|^n v^n - |u|^n}{|u|v - 1} \\ &< |u|^n v^n - |u|^n = N(f^n). \end{aligned}$$

Case 4. $v < -1$. For $n > 1$ we have

$$\begin{aligned} \sum_{k=1}^{n-1} N(f^k) &= \sum_{k=1}^{n-1} |u|^k |v^k - 1| = \sum_{k=1}^{n-1} |uv|^k + \sum_{k=1}^{n-1} (-1)^{k+1} |u|^k \\ &= \frac{|uv|^n - |uv|}{|uv| - 1} + \frac{|u| + (-1)^n |u|^n}{1 + |u|}. \end{aligned}$$

Therefore, if $n > 1$ is odd, clearly we have

$$\sum_{k=1}^{n-1} N(f^k) < |uv|^n + |u|^n = N(f^n);$$

and if $n > 2$ is even, we have

$$\begin{aligned} \sum_{k=1}^{n-2} N(f^k) &= \frac{|uv|^{n-1} - |uv|}{|uv| - 1} + \frac{|u| + (-1)^{n-1} |u|^{n-1}}{1 + |u|} \\ &< |uv|^{n-1} + |u|^{n-1} = |u|^{n-1} (|v|^{n-1} + 1) \\ &< |u|^n (|v|^n - 1) = N(f^n). \end{aligned}$$

So (e) holds for all $n > 2$. Suppose $n = 2$. Since

$$N(f) = |u|(|v| + 1) < |u|^2(|v|^2 - 1) = N(f^2),$$

(e) also holds for $n = 2$. This completes the proof of (1).

Suppose $|u| \leq 1$. Then, from (a), $N(f^n) = |v^n - 1|$. We separate the proof of (2) into two cases.

Case 1: $v \geq 2$. Then for $n > 1$ we have

$$\sum_{k=1}^{n-1} |v^k - 1| = \frac{v^n - v}{v - 1} - (n - 1) < v^n - 1 < |v^n - 1|,$$

and (e) follows.

Case 2: $v \leq -2$. We have

$$|v^k - 1| = \begin{cases} |v|^k - 1 & \text{for } k \text{ even,} \\ |v|^k + 1 & \text{for } k \text{ odd.} \end{cases}$$

Therefore for n odd we have

$$\sum_{k=1}^{n-1} |v^k - 1| = \frac{|v|^n - |v|}{|v| - 1} < |v|^n + 1 = |v^n - 1|,$$

and (e) follows. If n is even then for $v \leq -3$ we have

$$\sum_{k=1}^{n-1} |v^k - 1| = \frac{|v|^n - |v|}{|v| - 1} + 1 < |v|^n - 1 = |v^n - 1|,$$

and (e) follows.

If n is even and $v = -2$ we get

$$\sum_{k=1}^{n-1} |v^k - 1| = \frac{2^n - 2}{2 - 1} = 2^n - 1 = |v^n - 1|.$$

However, if $n > 2$ then there exists $k \in \{1, 2, \dots, n-1\}$ which does not divide n . For this k we have $|v^k - 1| > 0$, so (e) also holds. \square

The proof of statement (2) of Proposition 1 is essentially the proof for continuous self-maps on the circle, because for such maps the Nielsen number is $|d^n - 1|$ where d denotes the degree of the maps, see for instance [2].

Proof of the Theorem. From the definition of the numbers A_k , clearly we have $A_k \leq N(f^k)$ for all $k \geq 1$. Therefore, if (e) is true for some $n > 1$, we have, for this n , that

$$A_n = N(f^n) - \sum_{k < n, k|n} A_k \geq N(f^n) - \sum_{k < n, k|n} N(f^k) > 0$$

when $u \neq 1$, and

$$\begin{aligned} A_n &= N(f^n) - \sum_{k < n, k|n, \frac{n}{k} \text{ odd}} A_k \geq N(f^n) - \sum_{k < n, k|n, \frac{n}{k} \text{ odd}} N(f^k) \\ &\geq N(f^n) - \sum_{k < n, k|n} N(f^k) > 0 \end{aligned}$$

when $u = 1$. From (b) it follows that $\text{Per}(f) \supset \{n \in N : A_n > 0\}$. Hence, by the proposition we obtain the first part of statements (1) and (2) of the theorem.

From (c) and (d) the second part of statements (1) and (2), and statement (3) follow immediately. \square

REFERENCES

- [1] L. Alsedà, S. Baldwin, J. Llibre, R. Swanson and W. Szlenk, *Minimal sets of periods for torus maps via Nielsen numbers*, submitted, 1991.
- [2] L. Alsedà, J. Llibre and M. Misiurewicz, *Combinatorial dynamics and entropy in dimension one*, submitted, 1990.

- [3] L. Block, J. Guckenheimer, M. Misiurewicz and L. S. Young, *Periodic Points and Topological Entropy of One Dimensional Maps*, Springer Lecture Notes in Math., **819** (1980), 18–34.
- [4] R. Brown, *The Lefschetz Fixed Point Theorem*, Scott, Foresman and Company, 1971.
- [5] L. S. Efremova, *Periodic orbits and a degree of a continuous map of a circle* (in Russian), *Diff. and Integer Equations (Gor'kii)*, **2** (1978), 109–115.
- [6] B. Halpern, *Periodic points on the Klein bottle*, preprint, 1979.
- [7] B. Jiang, *Lectures on Nielsen fixed point theory*, *Contemporary Math.*, **14** (1983).
- [8] T. H. Kiang, *The Theory of Fixed Point Classes*, Springer, 1989.

Received May 13, 1991. Partially supported by a DGICYT grant number PB90-0695.

NORTHWESTERN UNIVERSITY
EVANSTON, IL 60208-2730

UNIVERSITAT AUTÒNOMA DE BARCELONA
08193 BELLATERRA
BARCELONA, SPAIN

