

# Computation of Five- and Six-Dimensional Bieberbach Groups

Carlos Cid and Tilman Schulz

## CONTENTS

- 1. Introduction
- 2. The Classification Problem
- 3. Deciding Torsion-freeness
- 4. Five-Dimensional Bieberbach Groups
- 5. Six-Dimensional Bieberbach Groups

Electronic Availability

References

---

This paper deals with the computation and classification of 5- and 6-dimensional torsion-free crystallographic groups, known as *Bieberbach groups*. We describe the basis of an algorithm that decides torsion-freeness of a crystallographic group as well as the triviality of its centre. The computations were done using the computer package CARAT, which handles enumeration, construction, recognition and comparison problems for crystallographic groups up to dimension 6.

The complete list of isomorphism types of Bieberbach groups up to dimension 6 can be found online.

---

## 1. INTRODUCTION

We say that a group  $G$  is an  $n$ -dimensional crystallographic group  $G$  if  $G$  contains a normal, torsion-free, maximal abelian subgroup  $V$ , of rank  $n$  and finite index. Thus an  $n$ -dimensional crystallographic group satisfies the short exact sequence

$$0 \longrightarrow V \longrightarrow G \longrightarrow P \longrightarrow 1,$$

where

$$P \leq \mathrm{GL}(n, \mathbb{Z}) \cong \mathrm{Aut}(V)$$

is a finite group acting faithfully on  $V$ . The groups  $P$  and  $V$  are called the *point-group (or holonomy group)* and *translation subgroup* of  $G$ , respectively. Crystallographic groups arise as discrete, irreducible subgroups of the group of isometries of the  $n$ -dimensional Euclidean space; see [Charlap 1986], for example. We say that a crystallographic group  $G$  is a *Bieberbach group* if it is torsion-free. Bieberbach groups also appear as fundamental groups of compact, connected, flat Riemannian manifolds (flat manifolds for short). Then  $X$  is a flat manifold of dimension  $n$  if and only if its fundamental group  $G$  is a  $n$ -dimensional Bieberbach group. Furthermore,  $G$  determines  $X$  up to affine equivalence; see [Charlap 1986], for example.

A referee pointed out that the classification described here implies estimates for the volume of certain 6 and 7 dimensional hyperbolic manifolds; cf. [Szczeptański 1996].

## 2. THE CLASSIFICATION PROBLEM

The problem of classifying isomorphism types of  $n$ -dimensional crystallographic groups goes back over a hundred years. Bieberbach's Third Theorem [Bieberbach 1912] states that for every  $n \in \mathbb{N}$ , the set of representatives of isomorphism classes of crystallographic (and Bieberbach) groups is finite. The classification for dimensions 1, 2 and 3 was found at the end of the nineteenth century, using mainly geometric techniques. H. Zassenhaus [1948] presented an algorithm, based on group-theoretical concepts, that allows the calculation of the set of representatives of isomorphism classes of  $n$ -dimensional crystallographic groups for any  $n$ .

If  $G$  is a  $n$ -dimensional crystallographic group, it follows from Bieberbach's Theorems [1911; 1912] that the point-group of  $G$  is isomorphic to a finite subgroup of  $\mathrm{GL}(n, \mathbb{Z})$ . Let  $\varphi_n$  be the set of representatives of conjugacy classes of finite subgroups of  $\mathrm{GL}(n, \mathbb{Z})$ . It follows from the Jordan–Zassenhaus Theorem that the set  $\varphi_n$  is finite for every  $n$ . For instance, for  $n = 1, 2, 3, 4, 5, 6$ , the set  $\varphi_n$  has 2, 13, 73, 710, 6079 and 85311 elements, respectively [Plesken and Schulz 2000].

Once given the set  $\varphi_n$  of representatives of conjugacy classes of finite subgroups of  $\mathrm{GL}(n, \mathbb{Z})$  and the generators of the normalizer in  $\mathrm{GL}(n, \mathbb{Z})$  of each element of  $\varphi_n$ , Zassenhaus' algorithm allows one to calculate the set of representatives of isomorphism classes of  $n$ -dimensional crystallographic groups. In [Brown et al. 1978] one can find the list of crystallographic groups for dimensions  $\leq 4$ . The numbers for dimensions 5 and 6 (see Table 1) were reported in [Plesken and Schulz 2000].

Note that  $n = 4$  seems to be the last dimension where a *complete list* of isomorphism classes of  $n$ -dimensional crystallographic groups still makes sense, since the increase in the number of isomorphism classes for higher dimensions does not allow a complete list in readable form. An alternative kind of classification for crystallographic group up to dimension 6 is suggested in [Plesken and Schulz 2000].

For every finite group  $P$ , Auslander and Kuranishi [1957] have shown that there is a Bieberbach group having point-group isomorphic to  $P$ . Following [Hiller and Sah 1986], we will call a finite group  $P$  *primitive* if it can be realized as point-group of a Bieberbach group  $G$  with finite commutator quotient. In contrast to the result of Auslander and Kuranishi, not every finite group is primitive. Hiller and Sah [1986] proved that a finite group  $P$  is primitive if and only if no cyclic Sylow  $p$ -subgroup of  $P$  has a normal complement. Let  $X$  be a flat manifold that has  $G$  as its fundamental group. Then it is well known that the first Betti number of the manifold  $X$  is zero if and only if the commutator quotient of  $G$  is finite, which is also equivalent to the triviality of the centre of  $G$  [Hiller and Sah 1986]. Due to the *Calabi construction* [Calabi 1957],  $n$ -dimensional Bieberbach groups with trivial centre have a relevant role in the classification of  $n$ -dimensional Bieberbach groups in general.

## 3. DECIDING TORSION-FREENESS

Even though the problem of classification of isomorphism types of crystallographic groups dates back to the nineteenth century, the idea of studying and classifying the torsion-free ones in particular came much later, with the study of flat manifolds. In [Brown et al. 1978] one can find the list of Bieberbach groups up to dimension 4; there are 1, 2, 10 and 74 Bieberbach groups, respectively, in these low dimensions. Below we describe the basis of an algorithm that decides torsion-freeness of a crystallographic group and verifies whether it has trivial centre.

The description of a crystallographic group  $G$  is based on the following property: If  $x_1, x_2, \dots, x_k$  are the generators of the point-group  $P \leq \mathrm{GL}(n, \mathbb{Z})$  of  $G$ , then a generating set of  $G$  can be given in the form

$$\left\{ \begin{pmatrix} x_1 & v_1 \\ 0 & 1 \end{pmatrix}, \dots, \begin{pmatrix} x_k & v_k \\ 0 & 1 \end{pmatrix} \right\} \cup \left\{ \begin{pmatrix} \mathrm{Id} & v \\ 0 & 1 \end{pmatrix} : v \in \mathbb{Z}^n \right\},$$

where  $v_1, \dots, v_k \in \mathbb{Q}^n$  and  $\mathrm{Id}$  is the  $n \times n$  identity matrix. We set

$$g_i = \begin{pmatrix} x_i & v_i \\ 0 & 1 \end{pmatrix}.$$

The following lemma gives the condition that a crystallographic group  $G$  must satisfy in order to be torsion-free.

**Lemma 3.1.** *Let  $V$  be a torsion-free  $\mathbb{Z}P$ -module, and  $G$  the extension of  $P$  by  $V$  corresponding to  $\alpha \in H^2(P, V)$ . Then  $G$  is torsion-free if and only if  $\text{res}_H^P(\alpha) \neq 0$  for every cyclic subgroup  $H$  of prime order of  $P$ .*

Let  $\varphi : G \rightarrow P \cong G/V$  be the natural homomorphism. Let  $g \in G$  be an element of finite order. Without loss of generality, we can assume that  $g^p = 1$  for some prime  $p$ . And since  $V$  is torsion-free,  $\varphi(g)$  is an element of order  $p$  in  $P$ , and  $\text{res}_H^P(\alpha) = 0$ , where  $H = \langle \varphi(g) \rangle$ . Conversely, assume that  $\text{res}_H^P(\alpha) = 0$ . Then  $\varphi^{-1}(H)$  splits and therefore  $G$  has an element of finite order.

**Remark 3.2.** Let  $V, G, P$  and  $\alpha$  be as given in Lemma 3.1, and define  $t(n, x) := 1 + \dots + x^{n-1}$ . Let  $g \in G$  given by  $g = \begin{pmatrix} h & \nu \\ 0 & 1 \end{pmatrix}$ , where  $\varphi(g) = h \in P$  has order  $m$ . Then  $\text{res}_{\langle h \rangle}^P(\alpha) = 0$  if and only if  $t(m, h)\nu \in t(m, h)V$ .

A trivial calculation shows that one needs only to do such tests for representatives of conjugacy classes of cyclic subgroups of prime order of  $P$ .

In short, the main points of the algorithm are :

- (1) Given the generators  $g_1, g_2, \dots, g_k$  of  $G$ , obtain coset representatives of  $V$  in  $G$  corresponding to the point-group  $P$  (via  $\varphi$ );
- (2) In view of Lemma 3.1, restrict the list from (1) to those elements giving rise to representatives of conjugacy classes of elements of prime order in  $P$ ;
- (3) For every element  $g \in G$  from the list (2), test if  $t(p, \varphi(g))v_g \in t(p, \varphi(g))V$ , where  $v_g$  is the translation component of  $g$  and  $p$  is a prime such that  $\varphi(g)^p = e$ . If so, the solution of this  $\mathbb{Z}$ -linear system will give an element of order  $p$  in  $G$ . Otherwise,  $G$  is torsion-free.

The centre of a space group in general is the lattice fixed by  $P$ , and it is easy to determine by solving  $(\varphi(g_i) - 1)v = 0$ , for  $i = 1, \dots, k$ . Note that this calculation has to be done only once for all groups in a specific  $\mathbb{Q}$ -class.

The algorithm has been implemented in  $C$  and uses as input the results presented in [Plesken and

Schulz 2000]. The computer programs and data which were used to obtain these results are part of the package CARAT, which is also available at the address given in the last section. The package, the algorithms, a lot of the underlying theory are described in [Opgenorth et al. 1998; Plesken and Schulz 2000].

As a result, we have the following count of  $n$ -dimensional crystallographic groups and Bieberbach groups, for  $n \leq 6$ .

dim.	cryst.	Bieb.	dim.	cryst.	Bieb.
1	2	1	4	4783	74
2	17	2	5	222,018	1060
3	219	10	6	28,927,922	38746

**TABLE 1.** Number of crystallographic and Bieberbach groups in each dimension up to 6.

#### 4. FIVE-DIMENSIONAL BIEBERBACH GROUPS

Through our calculations, we have obtained **1060** Bieberbach groups in dimension 5, of which 101 have trivial centre. On working first with those 5-dimensional Bieberbach groups with trivial centre, we classified the possible isomorphism types of point-groups.

Let  $(C_n)^k$  denote the direct product of  $k$  copies of the cyclic group of order  $n$ ,  $D_n$  the dihedral group of order  $n$ ,  $S_n$  and  $A_n$  the symmetric and alternating groups, and  $\Gamma$  the group of order 16 and nilpotence class 2, isomorphic to  $(C_2 \times C_2) \rtimes C_4$ .

**Theorem 4.1.** *A finite group  $P$  can be realized as point-group of a five-dimensional Bieberbach group with trivial centre if and only if it is isomorphic to  $(C_2)^k$ , for  $2 \leq k \leq 4$ ,  $C_2 \times C_4$ ,  $D_{12}$ ,  $D_8$ ,  $D_8 \times C_2$  or  $\Gamma$ .*

This list shows that [Szczepański 1990, Theorem 1], where the group  $C_4 \times C_2 \times C_2$  is included in the list of finite groups, is not correct. The given example of a Bieberbach group with such point-group and trivial centre is in fact not torsion-free (for instance, the element  $c_2b_2a_2$  has order 4).

By considering all 1060 groups, the theorem below, that lists the finite groups that can be realized as the point-group of a 5-dimensional Bieberbach group, agrees with [Szczepański 1996, Theorem 1].

family symbol	isomorphism type of point-group	number of classes		
		$\mathbb{Q}$	$\mathbb{Z}$	affine
1, 1, 1; 1; 1	$C_2 \times C_2$	1	1	1
1, 1; 1, 1; 1	$C_2 \times C_2$	1	2	2
1, 1; 1; 1; 1	$C_2 \times C_2 \times C_2$	2	3	17
1; 1; 1; 1; 1	$C_2 \times C_2 \times C_2, C_2 \times C_2 \times C_2 \times C_2$	3	8	44
2-1, 2-1; 1	$D_8$	1	1	1
2-1; 1, 1; 1	$D_8$	2	3	3
2-1; 1; 1; 1	$C_2 \times C_4, C_2 \times D_8, D_8$	6	13	25
2-1; 2-1; 1	$\Gamma$	2	2	4
2-2; 1; 1; 1	$C_2 \times S_3$	2	4	4
total		<b>20</b>	<b>37</b>	<b>101</b>

TABLE 2. Five-dimensional Bieberbach groups with trivial centre (first Betti number zero).

**Theorem 4.2.** *A finite group  $P \neq \{e\}$  can be realized as point-group of a 5-dimensional Bieberbach group if and only if it is isomorphic to  $C_n$ , for  $n = 2, \dots, 6, 8, 10, 12, (C_2)^k$ , for  $2 \leq k \leq 4, C_2 \times C_4, C_4 \times C_2 \times C_2, C_3 \times C_3, C_6 \times C_2, C_6 \times C_3, C_6 \times C_2 \times C_2, C_{12} \times C_2, S_3, D_8, D_{12}, D_8 \times C_2, S_3 \times C_3, D_{12} \times C_2, A_4, A_4 \times C_2, A_4 \times C_2 \times C_2, S_4$  or  $\Gamma$ .*

Tables 2 and 3 classify the 5-dimensional Bieberbach groups, using the notation of [Opgenorth et al. 1998; Plesken and Schulz 2000].

### 5. SIX-DIMENSIONAL BIEBERBACH GROUPS

Tables 4 and 5 classify 6-dimensional Bieberbach groups. As to the isomorphism types of point groups, the notation  $[n, k]$  refers to the classification of solvable groups with small order given in GAP [Schönert et al. 1994] and  $n$  stands for the order of the group in question.

Note that in the notation  $C_3 \times C_4, C_3 \times C_8$  and  $C_3^2 \times C_2$  the top  $C_2$  acts by inverting, in the latter case on both components.

family symbol	isomorphism type of point-group	number of classes		
		$\mathbb{Q}$	$\mathbb{Z}$	affine
1, 1, 1, 1, 1	$C_1$	1	1	1
1, 1, 1, 1; 1	$C_2$	2	3	3
1, 1, 1; 1, 1	$C_2$	2	5	5
1, 1, 1; 1; 1	$C_2^2$	2	8	21
1, 1; 1, 1; 1	$C_2^2$	2	9	31
1, 1; 1; 1; 1	$C_2^2, C_2^3$	4	43	236
1; 1; 1; 1; 1	$C_2^3, C_2^4$	3	22	290
2-1', 2-1'; 1	$C_4$	1	1	1
2-1; 1, 1, 1	$C_4$	1	2	2
2-1; 1; 1; 1	$C_2 \times C_4, C_4, D_8$	8	36	68
2-1; 1; 1; 1	$C_2^2 \times C_4, C_2 \times C_4, C_2 \times D_8, D_8$	8	62	179
2-1; 2-1; 1	$C_2 \times C_4, \Gamma$	2	5	12
2-2', 2-2'; 1	$C_2 \times C_3, C_3$	2	2	2
2-2; 1, 1, 1	$C_2 \times C_3, C_3$	2	3	3
2-2; 1, 1; 1	$C_2^2 \times C_3, C_2 \times C_3, C_2 \times S_3, S_3$	10	27	31
2-2; 1; 1; 1	$C_2^3 \times C_3, C_2^2 \times C_3, C_2^2 \times S_3, C_2 \times S_3$	7	12	42
2-2; 2-1; 1	$C_2 \times C_3 \times C_4, C_3 \times C_4$	3	3	3
2-2; 2-2; 1	$C_2 \times C_3, C_2 \times C_3^2, C_3^2, C_3 \times S_3$	4	8	10
3; 1, 1	$A_4, C_2 \times A_4$	2	4	4
3; 1; 1	$C_2 \times A_4, C_2^2 \times A_4, S_4$	5	9	11
4-1'; 1	$C_8$	1	1	1
4-2'; 1	$C_3 \times C_4$	1	1	1
4-3'; 1	$C_2 \times C_5, C_5$	2	2	2
total		<b>75</b>	<b>269</b>	<b>959</b>

TABLE 3. Five-dimensional Bieberbach groups with nontrivial centre.

family symbol	isomorphism type of point-group	number of classes		
		$\mathbb{Q}$	$\mathbb{Z}$	affine
1, 1, 1, 1; 1; 1	$C_2^2$	1	1	1
1, 1, 1; 1, 1; 1	$C_2^2$	1	2	2
1, 1, 1; 1; 1; 1	$C_2^3$	2	3	18
1, 1; 1, 1; 1, 1	$C_2^2$	1	4	4
1, 1; 1, 1; 1; 1	$C_2^3$	2	7	62
1, 1; 1; 1; 1; 1	$C_2^i, 3 \leq i \leq 4$	5	36	791
1; 1; 1; 1; 1; 1	$C_2^i, 3 \leq i \leq 5$	6	49	2727
2-1, 2-1; 1, 1	$D_8$	1	1	1
2-1, 2-1; 1; 1	$C_2 \times D_8, D_8$	2	7	11
2-1; 1, 1, 1; 1	$D_8$	2	3	3
2-1; 1, 1; 1, 1	$D_8$	1	2	3
2-1; 1, 1; 1; 1	$C_2 \times C_4, C_2 \times D_8, D_8$	15	67	173
2-1; 1; 1; 1; 1	$C_2^2 \times C_4, C_2^2 \times D_8, C_2 \times D_8$	16	113	883
2-1; 2-1; 1, 1	$C_2 \times D_8, \Gamma$	3	7	9
2-1; 2-1; 1; 1	$C_2 \times C_4, C_2 \times D_8, C_2 \times \Gamma, \Gamma, [16, 10], [32, 33], [32, 36]$	20	74	197
2-1; 2-1; 2-1	$\Gamma, [32, 33]$	2	4	5
2-2; 1, 1; 1; 1	$C_2^2 \times S_3, C_2 \times S_3$	6	13	13
2-2; 1; 1; 1; 1	$C_2^2 \times S_3$	6	13	71
2-2; 2-1; 1; 1	$D_{24}, [24, 11]$	4	10	10
3; 1; 1; 1	$C_2 \times S_4$	3	3	4
3; 2-2; 1	$C_2 \times S_4, S_4$	2	9	9
4-1'; 1; 1	$D_{16}$	1	1	1
4-1; 1; 1	$[16, 13], [16, 8], [32, 47]$	3	3	3
4-1; 2-1	$[32, 46], [64, 250]$	3	3	3
total		<b>108</b>	<b>435</b>	<b>5004</b>

TABLE 4. Six-dimensional Bieberbach groups with trivial centre.

**ELECTRONIC AVAILABILITY**

The complete list of isomorphism types of Bieberbach groups up to dimension 6 can be found at <http://wwwb.math.rwth-aachen.de/carat/>.

**REFERENCES**

[Auslander and Kuranishi 1957] L. Auslander and M. Kuranishi, “On the holonomy group of locally Euclidean spaces”, *Ann. of Math. (2)* **65** (1957), 411–415.

[Bieberbach 1911] L. Bieberbach, “Über die Bewegungsgruppen der Euklidischen Räume I”, *Mathematische Annalen* **70** (1911), 297–336.

[Bieberbach 1912] L. Bieberbach, “Über die Bewegungsgruppen der Euklidischen Räume II”, *Mathematische Annalen* **72** (1912), 400–412.

[Brown et al. 1978] H. Brown, R. Bülow, J. Neubüser, H. Wondratschek, and H. Zassenhaus, *Crystallographic groups of four-dimensional space*, Wiley Monographs in Crystallography, Wiley-Interscience, New York, 1978.

[Calabi 1957] E. Calabi, “Closed locally euclidean four-dimensional manifolds”, *Bull. Amer. Math. Soc.* **63** (1957), 135.

[Charlap 1986] L. Charlap, *Bieberbach groups and flat manifolds*, Springer, New York, 1986.

[Hiller and Sah 1986] H. Hiller and C.-H. Sah, “Holonomy of flat manifolds with  $b_1 = 0$ ”, *Quart. J. Math. Oxford Ser. (2)* **37**:146 (1986), 177–187.

[Opgenorth et al. 1998] J. Opgenorth, W. Plesken, and T. Schulz, “Crystallographic algorithms and tables”, *Acta Cryst. Sect. A* **54**:5 (1998), 517–531.

[Plesken and Schulz 2000] W. Plesken and T. Schulz, “Counting crystallographic groups in low dimensions”, *Experiment. Math.* **9**:3 (2000), 407–411.

[Schönert et al. 1994] M. Schönert et al., *GAP: Groups, algorithms, and programming*, 4th ed., Lehrstuhl D für Mathematik, RWTH Aachen, 1994. See <http://www-gap.dcs.st-and.ac.uk/~gap>.

[Szczepeński 1990] A. Szczepeński, “Five-dimensional Bieberbach groups with trivial centre”, *Manuscripta Math.* **68**:2 (1990), 191–208.

family symbol	isomorphism type of point-group	number of classes		
		$\mathbb{Q}$	$\mathbb{Z}$	affine
1, 1, 1, 1, 1, 1	$C_1$	1	1	1
1, 1, 1, 1, 1; 1	$C_2$	2	3	3
1, 1, 1, 1; 1, 1	$C_2$	2	5	5
1, 1, 1, 1; 1; 1	$C_2^2$	2	8	22
1, 1, 1; 1, 1, 1	$C_2$	1	3	3
1, 1, 1; 1, 1; 1	$C_2^2$	3	21	87
1, 1, 1; 1; 1; 1	$C_2^i, 2 \leq i \leq 3$	4	55	440
1, 1; 1, 1; 1, 1	$C_2^2$	1	9	29
1, 1; 1, 1; 1; 1	$C_2^i, 2 \leq i \leq 3$	4	98	1078
1, 1; 1; 1; 1; 1	$C_2^i, 3 \leq i \leq 4$	7	214	8403
1; 1; 1; 1; 1; 1	$C_2^i, 3 \leq i \leq 5$	5	158	13839
2-1', 2-1'; 1, 1	$C_4$	1	2	2
2-1', 2-1'; 1; 1	$C_2 \times C_4, C_4$	2	6	8
2-1, 2-1; 1; 1	$D_8$	2	14	25
2-1; 1, 1, 1, 1	$C_4$	1	2	2
2-1; 1, 1, 1; 1	$C_2 \times C_4, C_4, D_8$	8	37	89
2-1; 1, 1; 1, 1	$C_2 \times C_4, C_4, D_8$	4	38	88
2-1; 1, 1; 1; 1	$C_2^2 \times C_4, C_2 \times C_4, C_2 \times D_8, D_8$	21	392	2187
2-1; 1; 1; 1; 1	$C_2^3 \times C_4, C_2^2 \times C_4, C_2^2 \times D_8, C_2 \times C_4, C_2 \times D_8, D_8$	18	525	4972
2-1; 2-1; 1, 1	$C_2 \times C_4, C_4^2, [16, 10], \Gamma$	4	24	71
2-1; 2-1; 1; 1	$C_2^2 \times C_4, C_2 \times C_4, C_2 \times D_8, C_2 \times \Gamma, C_4^2, C_4 \times D_8, [16, 10], \Gamma, [32, 33], [32, 36]$	20	173	757
2-2', 2-2'; 1, 1	$C_2 \times C_3, C_3$	2	3	3
2-2', 2-2'; 1; 1	$C_2^2 \times C_3, C_2 \times C_3$	3	4	4
2-2, 2-2; 1; 1	$C_2 \times S_3, S_3$	2	7	7
2-2; 1, 1, 1, 1	$C_2 \times C_3, C_3$	2	3	3
2-2; 1, 1, 1; 1	$C_2^2 \times C_3, C_2 \times C_3, C_2 \times S_3, S_3$	10	27	33
2-2; 1, 1; 1, 1	$C_2^2 \times C_3, C_2 \times C_3, C_2 \times S_3, S_3$	5	20	26
2-2; 1, 1; 1; 1	$C_2^3 \times C_3, C_2^2 \times C_3, C_2^2 \times S_3, C_2 \times S_3$	18	118	391
2-2; 1; 1; 1; 1	$C_2^i \times C_3, 2 \leq i \leq 4, C_2^i \times S_3, 1 \leq i \leq 3$	15	107	652
2-2; 2-1; 1, 1	$C_2 \times C_3 \times C_4, C_3 \times C_4$	3	8	10
2-2; 2-1; 1; 1	$C_3 \times C_4, C_2 \times (C_3 \times C_4), C_2^i \times C_3 \times C_4, 0 \leq i \leq 2, C_2 \times C_3 \times D_8, C_2 \times C_4 \times S_3, C_2 \times D_{24}, C_3 \times D_8, C_4 \times S_3, D_{24}, [24, 11]$	26	98	140
2-2; 2-2; 1, 1	$C_2^i \times C_3^j, 1 \leq i, j \leq 2, C_2 \times C_3 \times S_3, C_3^2, C_3 \times S_3$	8	21	32
2-2; 2-2; 1; 1	$C_2^2 \times C_2, C_2 \times (C_3^2 \times C_2), C_2^i \times C_3^j, 1 \leq i, j \leq 2, C_2 \times C_3 \times S_3, C_2 \times S_3, C_3 \times S_3, S_3 \times S_3$	20	65	89
3; 1, 1, 1	$A_4, C_2 \times A_4$	2	4	4
3; 1, 1; 1	$C_2 \times A_4, C_2^2 \times A_4, C_2 \times S_4, S_4$	11	37	47
3; 1; 1; 1	$C_2^2 \times A_4, C_2^3 \times A_4, C_2 \times S_4$	8	18	57
3; 2-1; 1	$C_2 \times C_4 \times A_4, C_4 \times A_4$	3	5	7
3; 2-2; 1	$A_4, C_2 \times A_4, C_2^2 \times A_4, C_2 \times C_3 \times A_4, C_3 \times A_4, C_3 \times S_4$	9	30	55
4-1'; 1, 1	$C_8$	1	2	2
4-1'; 1; 1	$C_2 \times C_8, C_8, D_{16}$	3	8	11
4-1; 1; 1	$(C_3 \times C_8), C_3 \times Q_8, Q_8, [16, 11], [16, 13], [16, 8], [32, 31]$	8	15	23
4-2'; 1, 1	$C_3 \times C_4$	1	1	1
4-2'; 1; 1	$C_3 \times C_4, D_{24}$	2	3	3
4-2; 1; 1	$C_3 \times C_4, C_2 \times C_3 \times S_3, C_3 \times (C_3 \times C_4), C_3 \times D_8, C_3 \times S_3$	5	10	12
4-3'; 1, 1	$C_2 \times C_5, C_5$	2	3	3
4-3'; 1; 1	$C_2^2 \times C_5, C_2 \times C_5, C_2 \times D_{10}, D_{10}$	5	8	8
5-1; 1	$C_2 \times [80, 52], [80, 52]$	2	3	8
total		<b>289</b>	<b>2416</b>	<b>33742</b>

TABLE 5. Six-dimensional Bieberbach groups with nontrivial centre.

[Szczepański 1996] A. Szczepański, “Holonomy groups of five-dimensional Bieberbach groups”, *Manuscripta Math.* **90**:3 (1996), 383–389.

[Zassenhaus 1948] H. Zassenhaus, “Über einen Algorithmus, zur Bestimmung der Raumgruppen”, *Commentarii Mathematici Helvetici* **21** (1948), 117–141.

Carlos Cid, Lehrstuhl B für Mathematik, RWTH-Aachen, Templergraben 64, 52064 Aachen, Germany  
(cfcid@momo.math.rwth-aachen.de)

Tilman Schulz, Lehrstuhl B für Mathematik, RWTH-Aachen, Templergraben 64, 52064 Aachen, Germany  
(tilman@momo.math.rwth-aachen.de)

Received May 25, 2000; accepted in revised form September 12, 2000