

Indices (Index of Notation, General Index), 717-730

DOI: [10.3792/euclid/9781429799928-13](https://doi.org/10.3792/euclid/9781429799928-13)


from

Advanced Algebra *Digital Second Edition*

Anthony W. Knapp

Full Book DOI: [10.3792/euclid/9781429799928](https://doi.org/10.3792/euclid/9781429799928)

ISBN: 978-1-4297-9992-8



Anthony W. Knapp
81 Upper Sheep Pasture Road
East Setauket, N.Y. 11733–1729, U.S.A.
Email to: aknapp@math.stonybrook.edu
Homepage: www.math.stonybrook.edu/~aknapp

Title: Advanced Algebra
Cover: Content of the Snake Lemma; see page 185.

Mathematics Subject Classification (2010): 11–01, 13–01, 14–01, 16–01, 18G99, 55U99, 11R04, 11S15, 12F99, 14A05, 14H05, 12Y05, 14A10, 14Q99.

First Edition, ISBN-13 978-0-8176-4522-9

©2007 Anthony W. Knapp
Published by Birkhäuser Boston

Digital Second Edition, not to be sold, no ISBN

©2016 Anthony W. Knapp
Published by the Author

All rights reserved. This file is a digital second edition of the above named book. The text, images, and other data contained in this file, which is in portable document format (PDF), are proprietary to the author, and the author retains all rights, including copyright, in them. The use in this file of trade names, trademarks, service marks, and similar items, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.

All rights to print media for the first edition of this book have been licensed to Birkhäuser Boston, c/o Springer Science+Business Media Inc., 233 Spring Street, New York, NY 10013, USA, and this organization and its successor licensees may have certain rights concerning print media for the digital second edition. The author has retained all rights worldwide concerning digital media for both the first edition and the digital second edition.

The file is made available for limited noncommercial use for purposes of education, scholarship, and research, and for these purposes only, or for fair use as understood in the United States copyright law. Users may freely download this file for their own use and may store it, post it online, and transmit it digitally for purposes of education, scholarship, and research. They may not convert it from PDF to any other format (e.g., EPUB), they may not edit it, and they may not do reverse engineering with it. In transmitting the file to others or posting it online, users must charge no fee, nor may they include the file in any collection of files for which a fee is charged. Any exception to these rules requires written permission from the author.

Except as provided by fair use provisions of the United States copyright law, no extracts or quotations from this file may be used that do not consist of whole pages unless permission has been granted by the author (and by Birkhäuser Boston if appropriate).

The permission granted for use of the whole file and the prohibition against charging fees extend to any partial file that contains only whole pages from this file, except that the copyright notice on this page must be included in any partial file that does not consist exclusively of the front cover page. Such a partial file shall not be included in any derivative work unless permission has been granted by the author (and by Birkhäuser Boston if appropriate).

Inquiries concerning print copies of either edition should be directed to Springer Science+Business Media Inc.

INDEX OF NOTATION

This list indexes recurring symbols introduced in Chapters I through X (pages 1–648). For recurring symbols introduced in *Basic Algebra*, see the list of Notation and Terminology on pages xxiii–xxvi. Some of the latter notation has been repeated here for the reader’s convenience.

In the list below, each piece of notation is regarded as having a key symbol. The first group consists of those items for which the key symbol is a fixed Latin letter, and the items are arranged roughly alphabetically by that key symbol. The next group consists of those items for which the key symbol is a Greek letter. The final group consists of those items for which the key symbol is a variable or a nonletter, and these are arranged by type. To locate an item below, first proceed on the assumption that the key symbol is a Latin or Greek letter; if the item does not appear to be in the list, then treat it as if its key symbol is a variable or a letter.

- | | |
|---|---|
| \mathbb{A} , \mathbb{A}_K , 389, 559
\mathbb{A}^n , \mathbb{A}_K^n , 455, 559
$\mathcal{A}(K, \text{Gal}(K/F), a)$, 137
\mathcal{A}_F , 542, 543
\mathcal{A}_F^* , 542, 543
$\tilde{\mathcal{A}}(V)$, 579
$\tilde{\mathcal{A}}$, 570
$\tilde{\mathcal{A}}_d$, 570
$\tilde{\mathcal{A}}(V)$, 584
$\tilde{\mathcal{A}}(V)_d$, 585
${}^a\varphi$, 639
$\mathcal{B}(F)$, 126
$\mathcal{B}(K/F)$, 127
\mathcal{C} , 330
$\mathcal{C}(\mathfrak{a})$, 620
\mathcal{C}_R , 169
$\mathcal{C}(V(\mathfrak{a}))$, 633
$\mathcal{C}_{\mathbb{F}}$, 532, 549
$\mathcal{C}_{\mathbb{F},0}$, 534
E^c , complement, xxiii
coimage f , 240 | coker f , 175
$\mathcal{D}(\xi)$, 279
$\mathcal{D}(K/F)$, 372
$D_{\mathbb{F}}$, 532, 549
$D_{\mathbb{F},0}$, 534
$D_{\mathbb{K}}$, 267
$D(\Gamma)$, 267
$\text{Diff}(\mathbb{F})$, 547
$\text{Div}(\omega)$, 548
d_{-1} , 194
d_n , 153, 154
$X = \{(X_n, d_n)\}_{n=-\infty}^{\infty}$, 174
$\dim R$, 424
$\text{Ext}_R^n(A, B)$, 223
e_i, f_i, g , 275, 354
$\langle e_{j_1}, \dots, e_{j_k} \rangle$, 619
$\text{ext}_R^n(A, B)$, 223
$\mathbb{F}_q[[X]]$, 347
$\mathbb{F}_q((X))$, 347
$F_{\mathfrak{p}}$, 346
Fr_q , 437 |
|---|---|

- f_v , 533
 G_P , 368
 $\text{Gal}(F_2/F_1)$, 434
 \leq_{GLEX} , \leq_{GREVLEX} , 494
 g , 538
 g_x , 538
 $\mathcal{H}(s, \mathfrak{a})$, 633
 $\mathcal{H}_a(s, \mathfrak{a})$, 621, 626
 $H(s, \mathfrak{a})$, 633
 $H_a(s, \mathfrak{a})$, 625, 628
 H_j , 620
 $H_n(X)$, 153, 172
 $H^n(X)$, 153, 174
 $H_*(X)$, 172
 $H^*(X)$, 174
 $H_n(G, M)$, 209
 $H^n(G, M)$, 147
 $\text{Hom}_R(A, B)$, 169
 $h(D)$, 7, 14
 $h_{\mathbb{K}}$, 299
 \mathbb{I}, \mathbb{I}_K , 390
 \mathbb{I}^1 , 390
 \mathcal{I} , 330, 393, 576
 $\tilde{\mathcal{I}}$, 576
 $I = (r_1, r_2)$, 38
 $I = \langle r_1, r_1 \rangle$, 38
 $I(E)$, 560
 $I(P)$, 571
 $I(P, F \cap G)$, 474
 $I(P, L \cap F)$, 467
 $\text{image } f$, 240
 $J(\xi)$, 272
 $\overline{K(S)}$, 409
 $\overline{K(E)}$, 412
 \mathbb{k} , 528, 559
 $\mathbb{k}(V)$, 580, 585
 \mathbb{k}' , 531
 $\mathcal{L}(A)$, 544
 $L(A)$, 535
 $L(s, \chi)$, 63
 $\text{LCM}(X^\alpha, X^\beta)$, 501
 Log , 289
 $\text{LM}(f)$, $\text{LC}(f)$, $\text{LT}(f)$, 496
 $\text{LT}(I)$, 497
 \leq_{LEX} , 493
 $\ell(A)$, 536
 \lim_{\leftarrow} , 439
 \mathcal{M} , 493, 620
 M_P , 600
 M_x , 431
 \mathfrak{m}_P , 600
 \mathfrak{m}_x , 431
 $m_P(F)$, 474
 $N(I)$, 39, 273
 $N_{A/F}(\cdot)$, 165
 $N_{K/F}(\cdot)$, norm, xxvi
 $\text{Nrd}_{A/F}(\cdot)$, 165
 $\mathcal{O}(U)$, 580, 582, 587, 641
 $\mathcal{O}_P(U)$, 582, 587
 $\mathcal{O}_P(V)$, 580, 585
 R^o , opposite ring, xxiv
 $\text{ord}_v(A)$, 532
 \mathbb{P}^2 , 456
 \mathbb{P}^n , 457, 570
 \mathbb{P}_K^n , 457
 \mathcal{P} , 330, 393
 $P_{\mathbb{F}}$, 532, 549
 P_v , 322, 533
 \mathbb{Q}_p , 316, 318
 $\mathcal{R}(f, g)$, 451
 $R(f, g)$, 451
 $R(f_1, F)$, 514
 R_p , 346
 R_v , 322, 533
 R_x , 431
 Residue , 542
 $\text{Residue}_{p(v)}$, 541
 r_1, r_2 , 348, 383
 $\text{rad } A$, 78
 $S(f_1, f_2)$, 502

S_∞ , 391
 $S^{-1}R$, localization, xxvi
 $\text{Spec } A$, 639
 $(\text{Spec } A, \mathcal{O})$, 641
 $\text{Tor}_n^R(A, B)$, 224
 $\text{Tr}_{A/F}(\cdot)$, 165
 $\text{Tr}_{K/F}(\cdot)$, trace, xxvi
 $\text{Trd}_{A/F}(\cdot)$, 165
 A^t , transpose, xxiii
 $\text{tor}_n^R(A, B)$, 224
 $\text{tr. deg } R$, 424
 $V(C)$, $V_K(C)$, 455–456
 $V(I)$, 429
 $V(S)$, 559, 571
 $V(f_1, \dots, f_k)$, 559
 $V_{\mathbb{F}}$, 532
 $v_P(\cdot)$, 321
 v_∞ , 328
 $X(S)$, 388
 X^α , 494, 620
 $x_j(P)$, 559
 $\mathbb{Z}(\Gamma)$, 268
 \mathbb{Z}_p , 318
 $\widehat{\mathbb{Z}}$, 437
 $\mathbb{Z}G$, integral group ring, xxv

Greek

α_i , 369
 β_0 , 574
 β_0^t , 574
 β_i , 369, 575
 $\delta(A)$, 543
 δ_{ij} , Kronecker δ , xxiii
 ε , 149, 195
 η_i , 369
 ι , 390, 391
 σ , 617, 646
 $\sigma_1, \dots, \sigma_n$, 288, 383
 χ_0 , 62
 ω , 185, 547

Functors given by subscripts and superscripts

R^\times , units, xxiv
 R_P , localization, xxvi
 X^+ , 194
 K_{alg} , algebraic closure, 434
 K_{sep} , separable algebraic closure, 434
 M^G , invariants, 208
 M_G , coinvariants, 209
 \widehat{M} , dual fractional ideal, 372
 M_p , 376
 L^Φ , 460

Specific functions

$\alpha = (\alpha_1, \dots, \alpha_n)$, multi-index, 494
 $|\alpha|$, 620
 $\left(\frac{a}{p}\right)$, Legendre symbol, 8
 $\left(\frac{m}{n}\right)$, Jacobi symbol, 68
 $[K : F]$, degree, xxvi
 $|\cdot|_p$, 316
 $|\cdot|$, absolute value, 331
 $\|\cdot\|$, norm, 356
 $(x)_0$, $(x)_\infty$, 532

Isolated symbols

\sim , Brauer equivalent, 124
 \simeq , homotopic, 154
 ∂_n , 153, 172
 ∂_{-1} , 194
 \prod' , restricted direct product, 388

Operations on sets and classes

RG , group algebra, xxv
 \sqrt{T} , radical, 405
 $K[X_1, \dots, X_{n+1}]_d$, 458
 $A \xrightarrow{u} B$, morphism, 235

Miscellaneous

(x) , principal divisor, 532
 $(x_i)_{i \in I}$, 388

$I = (r_1, r_2)$, generated ideal, 38	$\varphi = \{(E, \varphi_E)\}$, rational map, 595
$I = \langle r_1, r_2 \rangle$, 38	$X = \{(X_n, \partial_n)\}_{n=-\infty}^{\infty}$, 171
$[x, y, w]$, point in \mathbb{P}^2 , 459	$(F, \cdot _F)$, valued field, 342
$[x_0, \dots, x_n]$, point in \mathbb{P}^n , 570	$\{\mathcal{O}(U), \rho_{VU}\}$, presheaf, 640

INDEX

- Abel, 521
- abelian category, 238
- abelian group
 - divisible, 196
 - torsion, 169
- abelian Lie algebra, 78
- absolute discriminant, 35, **267**
- absolute norm of ideal, 39, **273**
- absolute value, 289, 331
 - archimedean, 289
 - discrete, 338
 - nontrivial, 332
 - normalized, 383, 384, 385, 386
 - of idele, 390
 - trivial, 331
- acyclic resolution, 219
- additive category, 233
- additive functor, **170**, 178
- adele, 389
- adjoint, 252
- affine algebraic set, 559
 - dimension of, 566
 - irreducible, 563
- affine coordinate ring, 579
- affine curve, irreducible, 529
- affine Hilbert function, 621, 626
- affine Hilbert polynomial, 625, 628
- affine hypersurface, irreducible, **430**, 562
- affine local coordinates, 461
- affine n -space, **455**, 559
- affine plane curve, 455
 - irreducible, 430, 524, 562
- affine plane line, 455
- affine scheme, 642
- affine variety, 429, 562
- algebra, xxv
 - abelian Lie, 78
 - central, 111
 - central simple, 111
 - crossed-product, 137
 - cyclic, 122, 162, 163
 - generalized quaternion, 121
 - Lie, 77
 - semisimple associative, 80
 - semisimple Lie, 79
 - simple associative, 80
 - simple Lie, 79
 - solvable Lie, 78
 - tensor product for, 104
 - Weyl, 85
- algebra polynomial, 164
- algebraic closure, separable, 434
- algebraic set
 - affine, 559
 - irreducible affine, 563
 - projective, 571
- algebraically independent, 409
- aligned primitive forms, 25
- archimedean, 331, 333, 346
 - absolute value, 289
 - place, 383
 - valuation, 289
- Artin product formula, **387**, 390, 395
- Artin reciprocity, 265
- Artin's Theorem, 89
- Artinian ring, 87
- associated prime ideal, 446
- associated translation, 622
- associated vector subspace, 622
- associative algebra
 - semisimple, 80
 - simple, 80
- augmentation map, 149
- Baer, 168
- base field, 327
- base space, 640
- Bayer–Stillman ordering, 494

- Bezout, 449
 Bezout's Theorem, 447, 453, 465, 471, 487, **488**
 bidegree, 617
 bifunctor, 223
 bihomogeneous polynomial, 617
 binary quadratic form, 3, **12**
 similar, 74
 birational, 595
 map, 595
 birationally equivalent, 595
 Blichfeldt, 293
 boundary, 172
 map, 172
 operator, 172
 bounded sequence, 317
 bracket, 78
 Brauer equivalent, 124
 Brauer group, 126
 relative, 127
 Brauer's Lemma, 91
 Buchberger, 450
 Buchberger's algorithm, 506

 canonical class, 551
 canonical divisor, 551
 Cartan, E., 79
 Cartan, H., 168
 category
 abelian, 238
 additive, 233
 good, 169
 Cauchy sequence, 317
 Cayley, 77
 central algebra, 111
 central simple algebra, 111
 centralizer, 114
 chain complex, 171
 double, 257
 in abelian category, 240
 tensor product for, 258
 chain map, 154, **155**, 173
 character
 Dirichlet, 62
 genus, 74
 multiplicative, 61
 principal Dirichlet, 62
 Chase, 141

 Chevalley, 165, 168
 Chinese Remainder Theorem, **xxv**, 30, 69, 106, 314, 341, 367, 480, 483
 class field, Hilbert, 265
 class field theory, 265
 class group
 form 28
 ideal, 42, 265, **299**, 330, 393
 class number, **299**, 393
 Dirichlet, 7, 14
 co-invariant, 209
 co-invariants functor, 209
 coboundary, 174
 map, 174
 operator, 174
 cochain complex, 173
 cochain map, **154**, 174
 cocycle, 174
 codomain of morphism, 232
 cohomology, 153, 174
 sheaf, 168, 171, 218, **643**
 coimage in abelian category, 240
 cokernel, 175
 cokernel of morphism, 236
 universal mapping property of, 236
 common discriminant divisor, 272
 common index divisor, **272**, 287, 310, 371
 commutator ideal, 78
 complete presheaf, 641
 complete valued field, 343
 equal-characteristic case, 398
 unequal-characteristic case, 398
 completion, 342
 universal mapping property of, 343
 complex, 171
 chain, 171
 cochain, 173
 double, 257
 flat, 259
 in abelian category, 240
 place, 383
 composition formula, 24
 condition (C1), **165**, 518
 cone, **572**, 633
 conic, 458
 conjugate, 266, 288, 383
 connecting homomorphism, 185, 187
 connecting morphism in abelian category, 248

- convergent infinite product, 51
- convergent sequence, 317
- coordinate, 455, 559
 - affine local, 461
- coordinate hyperplane, 620
- coordinate ring
 - affine, 579
 - homogeneous, 584
- coordinate subspace, 619
- coproduct, xxv
- correspondence, one-one, xxiii
- countable, xxiii
- Cramer, 448
- Cramer's paradox, 449
- Cramer's rule, 448
- crossed-product algebra, 137
- cubic, 458
 - extension, pure, 280
 - number field, **279**, 302
 - twisted, 562
- cubical singular chain, 172
- cubical singular homology, 172
- cup product, 256
- curve, affine plane, 455
- curve, elliptic, 648
- curve, irreducible, 604
 - affine, 529
 - affine plane, 430, 524, 562
- curve projective plane, 458
- cycle, 172
- cyclic algebra, 122, 162, 163
- cyclotomic field, 309

- decomposition group, 368
- Dedekind, 77
- Dedekind Discriminant Theorem, **275**, 371, 379, 381
- Dedekind domain, xxvi, 266
 - extension of, xxvi, 327, 417
- Dedekind example, **287**, 302, 310
- Dedekind's Theorem on Differents, 376
- defined at a point, 580, 585
- degenerate, 172
- degree, 153
 - of divisor, 533
 - of inseparability, 415
 - residue class, 275, 354, 533
 - total, 457

- transcendence, 413
- derived functor, 204
 - formation of, 205
 - long exact sequence for, 211, 214
- Dickson, 122
- different, 279
 - relative, 279, 372
- differential, 543, 547
- differential form, 541
- dimension
 - geometric, 565
 - Krull, 403, **424**, 426, 528, 529, 564, 566, 605, 619, 630, 639
 - of affine algebraic set, 566
 - of affine variety, 563
 - of zero locus, 423
- Diophantus, 1
- direct product, restricted, 388
- direct sum in additive category, 233
- directed set, 438
- Dirichlet, 2, 24, 77
- Dirichlet box principle, 297
- Dirichlet character modulo m , 62
- Dirichlet class number, 7, 14
- Dirichlet L function, 63
- Dirichlet pigeonhole principle, 297
- Dirichlet series, 56
- Dirichlet Unit Theorem, **290**, 292, 384, 390, 395
- Dirichlet's Theorem, 7, **50**
- discrete, 290
- discrete absolute value, 338
- discrete valuation, 322
 - defined over \mathbb{k} , 529
- discriminant, 12
 - absolute, 35, **267**
 - field, 35, 264, **267**
 - fundamental, 33
 - of commutative semisimple algebra, 382
 - of ordered basis, 267
 - relative, 275, 381
- discriminant divisor, 272
- divisible abelian group, 196
- divisible module, 251
- division algorithm, generalized, 499
- divisor, 532
- divisor class, **532**, 549
- divisor, principal, 532
- domain of morphism, 232

- dominant rational map, 595
- Double Centralizer Theorem, 115
- double chain complex, 257
- dual of fractional ideal, 372

- Eckmann, 168
- Eilenberg, 168
- Eisenstein, 12
- Eisenstein polynomial, 402
- elimination ideal, 512
- Elimination Theorem, 512
- elimination type ordering, 494, 512
- elliptic curve, 648
- enough injectives, 202
- enough projectives, 202
- epi, 233
- epimorphism, 233
- equal-characteristic case, 398
- equivalence class of forms
 - ordinary, 13
 - proper, 13
- equivalence of
 - absolute values, 333
 - completions, 383
 - forms, 13, 32
 - forms, improper, 13
 - forms, proper, 13, 32
 - ideals, 40, 298
 - ideals, narrow, 40
 - ideals, strict, 40, 298
 - morphisms, 242
- Euler, 1, 3, 9, 50
- Euler product, 50, 54, **60**
 - first-degree, 60
- Euler's Theorem, 516, 646
- exact complex, 175
- exact functor, 179
 - left, 182
 - right, 183
- exact on injectives, 222
- exact on projectives, 222
- exact sequence, 175
 - in abelian category, 240
 - long, 187, 188
 - short, 175
 - split, 200
- Exchange Lemma, 412
- Ext functor, 223

- extension
 - normal, 435
 - of Dedekind domain, xxvi, 327, 417
 - of integrally closed domain, 610
 - of valued field, 358
 - purely transcendental, 409
- Extension Theorem, 512

- factor set, 133
 - trivial, 135
- Fermat, 1, 3, 9
- field discriminant, 35, 264, **267**
- field of formal Laurent series, 347
- field of fractions, xxv
- field polynomial, 266
- fine sheaf, 218
- finiteness of class number, 390
- first-degree Euler product, 60
- flabby sheaf, 218
- flat complex, 259
- flat module, 257
- form
 - binary quadratic, 3, 12
 - class group, 28
 - negative definite, 14
 - positive definite, 14
 - primitive aligned, 25
 - reduced primitive, 18, 21
- Fourier inversion formula for finite abelian
 - groups, 61
- fractional ideal, 321
 - principal, 321
 - relative dual of, 372
- free resolution, 152, 195
- Freudenthal, 168
- Frobenius element, 437
- Frobenius's Theorem about division algebras
 - over the reals, **118**, 160
- function field, **419**, 528, 580, 582, 585, 587
 - in one variable, 326, 382, 528, 529
 - in r variables, 419
- functor
 - additive, **170**, 178
 - co-invariants, 209
 - derived, 204
 - exact, 179
 - Ext, 223
 - global-sections, 218

- homology-of-groups, 209
- invariants, 208
- left exact, 182
- right exact, 183
- Tor, 224
- functorial, 177
- functoriality of long exact sequence of
 - derived functors, 215, 218
- functoriality with long exact sequence, 191
- functoriality with snake diagram, 190
- fundamental discriminant, 33
- fundamental parallelotope, 293
- Fundamental Theorem of Galois Theory, 443
- fundamental unit, 36, 288

- Galois, 77
- Galois group, 434
- gap sequence, 557
- Gauss, 1, 3, 9, 24, 77
- Gauss's group, 5, **28**
- Gelfand, 348
- generalized division algorithm, 499
- generalized quaternion algebra, 121
- generalized resultant, 514
- genus, 32, 539, 556, 557
 - principal, 33
- genus character, 74
- genus group, **33**, 70, 73
- geometric dimension, 565
- germ, 584
- global field, 382
- global-sections functor, 218
- good category, 169
- graded lexicographic ordering, 493
- graded monomial ordering, 627
- graded reverse lexicographic ordering, 494
- Gröbner, 450
- Gröbner basis, 450, **497**, 564
 - minimal, 508
 - reduced, 509
- Grothendieck, 638

- Haar measure, 385
- Halphen, 450
- Hamilton, 77
- Hensel, 279
- Hensel's Lemma, **349**, 351, 353, 399
- Herstein, 130

- Hilbert, 404
- Hilbert Basis Theorem, **xxvi**, 491, 560
- Hilbert class field, 265
- Hilbert function, 633
 - affine, 621, 626
- Hilbert polynomial, 633
 - affine, 625, 628
- Hilbert's Theorem 90, 71, **145**
- homogeneous coordinate ring, 584
- homogeneous ideal, 458, 570
- homogeneous member of homogeneous
 - coordinate ring, 585
- homogeneous Nullstellensatz, **572**, 586, 635
- homogeneous polynomial, 457
- homology, 153, 172
 - cubical singular, 172
 - simplicial, 172
- homology-of-groups functor, 209
- homomorphism, 78
 - connecting, 185, 187
 - inflation, 254
 - of valued field, 342
 - restriction, 254
- homotopic, 154, 173, 174, 193, 198
- homotopy, 173, 174, 193, 198
- Hopf, H., 167
- Hopkins, 92
- Hurewicz, 167
- hyperplane coordinate, 620
- hypersurface, irreducible affine, **430**, 562
- hypersurface, irreducible projective, 573

- ideal
 - fractional, 321
 - in Lie algebra, 78
 - principal fractional, 321
 - valuation, 322
- ideal class group, 42, 265, **299**, 330
- idele, 390
- idele class group, 393
- idempotent, 91, 369
- idempotent primitive, 369
- image in abelian category, 240
- Implicit Function Theorem, 428, 600
- improper equivalence of forms, 13
- independent algebraically, 409
- index, 272
 - ramification, 275, 354

- inertia group, 370
- inertia subfield, 368
- inflation homomorphism, 254
- inflation-restriction sequence, 254
- injective, 195
 - in abelian category, 241
- injective module, 195
- injective resolution, 199, 205
- inseparable element, 414
- integral closure, xxvi, 610
- integral domain, xxv
- integral element, xxvi
- integrally closed, xxvi
- intersection multiplicity, 467, **474**
- intersection number, 467
- invariant, 208
- invariants functor, 208
- inverse limit, 439
 - standard, 439
- inverse system, 438
- irreducible
 - affine algebraic set, 563
 - affine curve, 529
 - affine hypersurface, **430**, 562
 - affine plane curve, 430, 524, 562
 - closed set, 564, 573
 - curve, 604
 - element, xxv
 - ideal, 446
 - projective hypersurface, 573
- irredundant, 446
- isomorphic idempotents, 97
- isomorphism, 78
 - of valued field, 342
 - of varieties, 591
- Jacobi, 521
- Jacobi identity, 77
- Jacobi symbol, 68
- Jacobson radical, 89
- kernel of morphism, 235
 - universal mapping property of, 235
- Koszul, 168
- Kronecker, 77
- Krull dimension, 403, **424**, 426, 528, 529, 564, 566, 605, 619, 630, 639
- Kummer, 77
- Kummer's criterion, 275
- Künneth Theorem, 258–259
- Lagrange, 1, 4
- Langlands reciprocity, 265
- largest domain, 583, 595
- Lasker–Noether Decomposition Theorem, **446**, 639
- lattice, 290
- Law of Quadratic Reciprocity, 3, **8**
- least common multiple, 501
- left adjoint, 252
- left Artinian ring, 87
- left exact functor, 182
- left Noetherian ring, 87
- left semisimple ring, 81
- Legendre, 1, 4
- Legendre symbol, 8
- Leibniz, 7
- Leray, 168
- Levi, E. E., 79
- lexicographic ordering, 493
- Lie algebra, 77
 - abelian, 78
 - semisimple, 79
 - simple, 79
 - solvable, 78
- Lie subalgebra, 78
- line
 - affine plane, 455
 - at infinity, 459
 - projective, 458
- Liouville, 521
- local expression, 462
- local field, 383
- local morphism, 642
- local ring, xxvi
 - at a point, 580, 582, 585, 587
- local/global approach, 371
- localization, xxvi
- locus of common zeros, 429, 559, 571
- long exact sequence, 187, 188
 - functoriality with, 191
 - of derived functors, 211, 214
 - functoriality of, 215, 218
- Mac Lane, 168, 420
- Macaulay, 627

- maps of a good category, 169
- matrix units, 101
- member in abelian category, 242
- minimal Gröbner basis, 50
- Minkowski, 301, 302
- Minkowski Lattice-Point Theorem, **293**, 384
- modules of a good category, 169
- monic, 232
- mono, 232
- monomial, 457
 - reduced, 646
- monomial ideal, 619
- monomial ordering, 493
 - graded, 627
- monomorphism, 232
- morphism, 169
 - local, 642
 - of affine scheme, 642
 - of ringed space, 642
 - of varieties, 591
- multiplicative, 60
- multiplicative character, 61
 - strictly, 60
- multiplicity of a tangent line, 478

- Nakayama's Lemma, **xxv**, 120, 605, 606
- narrow equivalence of ideals, 40
- natural, 177
- negative, xxiii
- negative definite form, 14
- negatively oriented, 40
- neighbor, 21
 - on the left, 21
 - on the right, 21
- nil left ideal, 89
- nilpotent element, 89
- nilpotent left ideal, 80, 90
- Noether Normalization Lemma, 612
- Noether–Jacobson Theorem, 130
- Noetherian, xxvi
- Noetherian ring, 87
- Noetherian topological space, 564
- nonarchimedean, 331, 335, 338
- nonarchimedean place, 383
- nonsingular curve, 604
- nonsingular point, 429, 600, 601
- nontrivial absolute value, 332
- norm, 165, 356
 - of ideal, 39
 - absolute, 273
- normal extension, 435
- normalized absolute value, 383, 384, 385, 486
- Nullstellensatz, 403, **404**, 428, 455, 480, 487, 510, 516, 518, 524, 526, 529, 559, 561, 563, 572, 579, 580, 581
 - homogeneous, **572**, 586, 635
- number field, xxvi
 - cubic, **279**, 302
 - cyclotomic, 309
 - quadratic, **35**, 69, 263, 269

- Oka, 168
- one-one correspondence, xxiii
- order, 532
- order of vanishing, 474
- ordering
 - Bayer–Stillman type, 494
 - from tuple of weight vectors, 494
 - graded lexicographic, 493
 - graded monomial, 627
 - graded reverse lexicographic, 494
 - k -elimination type, 494, 512
 - lexicographic, 493
 - monomial, 493
 - total, 493
- ordinary equivalence class of forms, 13
- oriented, 40
- orthogonal idempotents, 99, 369
- Ostrowski, 348
- Ostrowski's Theorem, 336

- p -adic absolute value, 316
- p -adic integer, 279, 318
- p -adic integer, 346
- p -adic metric, 316
- p -adic number, 279, 316, 318
- p -adic number, 346
- parallelootope, 548
 - fundamental, 293
- Peirce decomposition, 95
- perfect field, **418**, 554
- place, 383
- plane, projective, 456
- plane curve
 - affine, 455
 - irreducible affine, 430, 524, 562

- projective, 458
- plane line, affine, 455
- Plücker, 450
- point, 455, 456, 459, 559
- points at infinity, 459
- pole part, 537
- pole set, 581, 585
- positive, xxiii
- positive definite form, 14
- positively oriented, 40
- presheaf, 640
 - complete, 641
- primary ideal, 445
- prime element, xxv
- prime ideal, xxv
 - associated, 446
- primitive, 12
- primitive form
 - aligned, 25
 - reduced, 18, 21
- primitive idempotent, 369
- primitively represent, 14
- principal Dirichlet character, 62
- principal divisor, 532
- principal fractional ideal, 321
- principal genus, 33
- problem
 - ideal-equality, 510
 - ideal-membership, 507
 - proper-ideal, 507
- product, xxv
- profinite group, 441
- projective, 192
 - algebraic set, 571
 - closure, 575
 - hypersurface, irreducible, 573
 - in abelian category, 241
 - limit, 439
 - line, 458
 - module, 192
 - n -space, 457
 - plane, 456
 - plane curve, 458
 - resolution, 195, 205
 - transformation, 460
 - variety, 572
- proper equivalence class of forms, 13
 - forms over \mathbb{Q} , 32
 - forms over \mathbb{Z} , 13
- pullback, 242
- pure cubic extension, 280
 - type of, 281
- purely inseparable element, 415
- purely inseparable extension, 416
- purely transcendental extension, 409
- pushout, 202, 243
- quadratic form, binary, 3, 12
- quadratic form, similar, 74
- quadratic number field, **35**, 69, 263, 269
- quadratic reciprocity, 3, **8**, 68
- quartic, 458
- quasi-affine variety, 568
- quasiprojective variety, 573
- quaternion algebra, 121
- radical
 - associative algebra, 80
 - ideal, 405
 - Jacobson, 89
 - of Lie algebra, 78
 - Wedderburn–Artin, 89, **91**
- ramification index, 275, 354
- ramified, 367
- ramify, 264, 275, 308
- rational function, 580, 585
- rational map, 595
 - dominant, 595
- rational point, 455, 456, 457, 459
- real place, 383
- reciprocity
 - Artin, 265
 - Langlands, 265
 - quadratic, 3, **8**, 68
- reduced Gröbner basis, 509
- reduced monomial, 646
- reduced norm, 165
- reduced polynomial, 165
- reduced primitive form, 18, 21
- reduced trace, 165
- reducible ideal, 446
- regular at a point, 580, 582, 585
- regular function at a point, 587
- regular function on an open set, 580, 582, 587, 641

- regular point, 429
- relative Brauer group, 127
- relative different, 279, 372
- relative discriminant, 275, 381
- relative dual of fractional ideal, 372
- represent, 14
 - primitively, 14
- residue class degree, 275, 354, 533
- residue class field, 322
- Residue Theorem, 543
- resolution, 194
 - acyclic, 219
 - free, 152, 195
 - injective, 199, 205
 - projective, 195, 205
 - standard, 149
- restricted direct product, 388
- restriction homomorphism, 254
- resultant, 449, **451**
 - generalized, 514
- Riemann, 521
- Riemann hypothesis, 530
- Riemann sphere, 328
- Riemann surface, 522
- Riemann zeta function, 52, 58
- Riemann–Roch Theorem, 520, 522, 523, 530, 540, 543, **551**, 552, 648
- Riemann’s inequality, 538
- right adjoint, 252
- right Artinian ring, 87
- right exact functor, 183
- right Noetherian ring, 87
- right semisimple ring, 81
- ring of formal power series, 347
- ringed space, 642

- S -polynomial, 502
- scheme, 642
 - affine, 642
 - defined over a ring, 643
- Schmidt, 422
- Schreier, 168
- Schur’s Lemma, 83
- section, 641
- Segre embedding, **617**, 646
- Segre variety, 617
- semisimple
 - associative algebra, 80
 - Lie algebra, 78
 - module, xxiv
 - ring, 81, 84
- separable
 - algebraic closure, 434
 - element, 414
 - extension, 415
 - polynomial, 414
 - semisimple algebra over a field, 109
- separably generated extension, 419
- separating transcendence basis, 419
- sheaf, 168, **640**
 - fine, 218
 - flabby, 218
 - cohomology, 168, 171, 218, **643**
 - structure, 641
- short exact sequence, 175
 - in abelian category, 241
- similar binary quadratic form, 74
- simple
 - associative algebra, 80
 - Lie algebra, 79
 - module, xxiv, 80
 - ring, 85
- simplicial homology, 172
- singular cube, 172
- singular homology, 172
- singular point, 429, 600, 601
- Skolem–Noether Theorem, 113
- snake diagram, **185**, 261
 - functoriality with, 190
- Snake Lemma, **185**, 248
- solution of problem
 - ideal-equality, 510
 - ideal-membership, 507
 - proper-ideal, 507
- solvable Lie algebra, 78
- spectral sequence, 171
- spectrum, 639
- split algebra, 127
- split exact sequence, 200
- splitting field, 127
- stalk, 640
- standard inverse limit, 439
- standard resolution, 149
- standard subset, 622
- Stickelberger’s condition, 309
- Stone, 638

- strict equivalence of ideals, 40, 298
- strictly multiplicative, 60
- strong approximation property, 374
- Strong Approximation Theorem, **373**, 390, 391
- structure sheaf, 641
- subalgebra, Lie, 78
- summation by parts, 56

- tangent lines, 478
- tensor product of
 - algebras, 104
 - chain complexes, 258
 - fields, 104
- Theorem 90, Hilbert's, 71, **145**
- Tor functor, 224
- Tornheim, 349
- torsion abelian group, 169
- torsion submodule, 257
- total degree, 457
- total ordering, 493
- totally ramified, 367
- trace, 165
- transcendence basis, **409**, 424
 - existence, 411
 - separating, 419
- transcendence degree, 413
- transcendence set, 409
- translate of form, 26
- triangular ring, 88
- trivial absolute value, 331
- trivial factor set, 135
- twisted cubic, 562
- type of pure cubic extension, 281

- ultrametric inequality, 316, 331
- unequal-characteristic case, 398
- uniformizer, 323
- uniformizing element, 323
- unit, **xxiv**, 36, 288
 - fundamental, 36, 288
- unital, **xxiv**
- Universal Coefficient Theorem, 261
- universal mapping property of
 - cokernel, 236
 - completion of valued field, 343
 - kernel, 235
- unramified, 367
- valuation, 322, 331
 - archimedean, 289
 - discrete, 322, 529
- valuation ideal, 322
- valuation ring, 322
- valued field, 342
 - complete, 343
 - extension of, 358
 - homomorphism of, 342
 - isomorphism of, 342
- variety, 590
 - affine, 429, 562
 - as a scheme, 643
 - projective, 572
 - quasi-affine, 568
 - quasiprojective, 573
 - Segre, 617

- Weak Approximation Theorem, **340**, 374
- Wedderburn, 79, 86, 164
- Wedderburn–Artin radical, 89, **91**
- Wedderburn's Main Theorem, 94
- Wedderburn's Theorem about finite division rings, **117**, 160
- Wedderburn's Theorem about semisimple rings, 83
- Weierstrass, 521
- Weierstrass gap, 557
- Weierstrass point, 557
- Weierstrass valuation, 557
- weight vectors, 494
- Weil, 530, 541, 543
- Weyl algebra, 85

- Zariski closure, 561, 578
- Zariski topology, 560, 571
- Zariski's Theorem, 403, **431**, 525, 558, 600, 601, 605, 606
- zero locus, 455, 559, 571
- zero member, 245
- zero morphism, 233
- zero object, 233
- zeta function, 530
 - Riemann, 52, 58