

Schubert calculus and puzzles

Allen Knutson

Abstract.

These are notes for four lectures given at the Osaka summer school on Schubert calculus in 2012, presenting the geometry from the unpublished [arXiv:1008.4302](https://arxiv.org/abs/1008.4302) giving an extension of the puzzle rule for Schubert calculus to equivariant K -theory, while eliding some of the combinatorial detail. In particular, §3 includes background material on equivariant cohomology and K -theory.

Since that school, I have extended the results to arbitrary interval positroid varieties (not just those arising in Vakil’s geometric Littlewood-Richardson rule), in the preprint [Kn2].

CONTENTS

1. Schubert varieties and interval positroid varieties	185
2. Vakil’s Littlewood-Richardson rule, from shifting	193
3. Equivariant and K - extensions	200
4. Puzzles for the Belkale-Kumar Schubert calculus of other partial flag manifolds	207

§1. Schubert varieties and interval positroid varieties

In my first lecture I’ll present a family of varieties interpolating between Schubert and Richardson, called “interval positroid varieties”.

1.1. Schubert varieties

Definitions:

- $M_{k \times n} := k \times n$ matrices over \mathbb{C} .
- the **Stiefel manifold** $M_{k \times n}^{\text{rank } k}$ is the open subset in which the rows are linearly independent.

- the **Grassmannian** $Gr_k(\mathbb{C}^n) \cong GL(k) \backslash M_{k \times n}^{\text{rank } k}$ is the space of k -planes in \mathbb{C}^n .

Two matrices in $M_{k \times n}^{\text{rank } k}$ give the same k -plane if they're related by row operations. To kill that ambiguity, put things in **reduced row-echelon form**. From there we can associate a discrete invariant, a **bit string** like 0010101101 with 0s in the k pivot positions and 1s in the remaining $n - k$ (**careful**: not the reverse!). Let $\binom{n}{k}$ denote the *set* of such bit strings.

Examples:

- (1) Most matrices in $M_{k \times n}^{\text{rank } k}$ give 000...11111.
- (2) If all the columns are zero except the last k , the bit string is 111...00000.
- (3) If the k th column is in the span of the first $k - 1$, but there are no other dependencies, the bit string is 00...0101...1111.

Our reference for Grassmannian Schubert varieties, and their properties such as the following, is [Fu97].

Proposition 1.1. (1) *There is a decomposition of the Grassmannian indexed by $\lambda \in \binom{n}{k}$, into complex cells X_λ° . The codimension $\ell(\lambda)$ of the λ cell is the number of **inversions** in λ , where a 1 occurs somewhere left of a 0.*

- (2) *These cells give bases for homology and cohomology.*
- (3) *To figure out which cell a matrix is in, look at rank $[1, i] :=$ the rank of the first i columns, for each $i \in [1, n]$.*
- (4) *The closure of a cell (or, its preimage in the Stiefel manifold) satisfies a bunch of determinantal conditions.*
- (5) *Fix λ . Let $P_\lambda \subset M_{k \times n}$ be the vector space of matrices where the i th row has 0s left of the i th 0 in λ . Then $P_\lambda \cap M_{k \times n}^{\text{rank } k} \rightarrow Gr_k(\mathbb{C}^n)$ has image X_λ° .*

The closures $\{\overline{X_\lambda^\circ}\}$ of these **Bruhat cells** in the Grassmannian are called **Schubert varieties** and denoted X_λ , $\lambda \in \binom{n}{k}$. Hodge proved that the determinantal equations (vanishing of the **Plücker coordinates** on the affine cone over the Grassmannian) give prime ideals, i.e. define these as schemes.

- Exercise 1.2.** (1) *Fix $\lambda \in \binom{n}{k}$. Show that the rank conditions on those initial intervals $[1, i]$ for which 10 occur in positions $i, i + 1$ of λ imply all of λ 's other rank conditions.*
- (2) *Let $B \leq GL(n)$ denote the upper triangular matrices. Show that the Bruhat cells are exactly the B -orbits (acting on $M_{k \times n}$ on the right).*

- (3) Show the same remains true if we replace B by $N = B'$, the upper triangular matrices with 1s on the diagonal.

1.2. Schubert calculus

Let $[X_\lambda]$ always denote the element of cohomology. Since we have a basis, we know $[X_\lambda][X_\mu] = \sum_\nu c_{\lambda\mu}^\nu [X_\nu]$. The study of these **Schubert structure constants** (and generalizations thereof) is **Schubert calculus**.

Theorem 1.3 (e.g. [Kl73]). X_λ and $w_0 \cdot X_\mu$ intersect transversely, where $w_0 \in S_n \leq GL(n)$ reverses $M_{k \times n}$ left/right (really, by a right multiplication).

Let $X^\mu := w_0 \cdot X_{w_0 \cdot \mu}$ be the **opposite Schubert variety**, where $w_0 \cdot \mu$ again means reverse left/right. Since w_0 lies in the connected group $GL(n)$, we have $[X_\mu] = [X^{w_0 \cdot \mu}]$, so $[X_\lambda][X_\mu] = [X_\lambda][X^{w_0 \cdot \mu}] = [X_\lambda \cap X^{w_0 \cdot \mu}]$ (the last by Kleiman transversality).

First example: let $\lambda = \mu = 010$, each defining the space of matrices $\{[0 * *]\}$, so those points in the projective plane $Gr_1(\mathbb{C}^3)$ that lie on the yz line. Then $X_\lambda = X_\mu$ so they don't intersect transversely. But $X_\lambda \cap w_0 \cdot X_\mu = \{[0 * 0]\}$, the y point. Which is homologous to the Schubert point $\{[0 0 *]\}$.

An intersection $X_\lambda^\mu := X_\lambda \cap X^\mu$ of a Schubert and opposite Schubert variety is called a **Richardson variety**¹. Schubert calculus then becomes the question of computing the cohomology classes Poincaré dual to the X_λ^μ , in the Schubert basis. To determine the smallest Richardson variety containing a matrix, look at $\{\text{rank } [1, i]\}$ and $\{\text{rank } [j, n]\}$ for $i, j \leq n$.

Proposition 1.4. Define the **Bruhat order** $\lambda \leq \mu$ on $\binom{n}{k}$ by requiring the i th 1 in λ to occur left of the i th 1 in μ , $i = 1, \dots, n - k$.

- (1) If $\lambda \leq \mu$, then $\dim X_\lambda^\mu$ is nonempty of dimension $\ell(\mu) - \ell(\lambda)$.
- (2) X_λ^λ is a point.
- (3) If $\lambda \not\leq \mu$, then $X_\lambda^\mu = \emptyset$.
- (4) Let $P_\lambda^\mu \subset M_{k \times n}$ denote the vector space of matrices in which the i th row is supported between the i th 0 in λ and the i th 0 in

¹Warning: Richardson also studied nice nilpotent orbits in Lie algebras, and non-cognoscenti often guess incorrectly that “Richardson variety” refers to the closure of a Richardson orbit. It doesn't seem to have been used that way in the literature, or at least I prefer to believe that.

μ . Then $P_\lambda^\mu \cap M_{k \times n}^{\text{rank } k} \rightarrow Gr_k(\mathbb{C}^n)$ has image X_λ^μ .

E.g. $\lambda = 1100110111, \mu = 1101110101,$

$$P_\lambda^\mu = \begin{pmatrix} 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & * & * & * & * & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & * & * & * \end{pmatrix}$$

Corollary 1.5. $\int_{Gr_k(\mathbb{C}^n)} [X_\lambda][X^\mu] = \delta_{\lambda\mu}$. In particular, if we define the more symmetric **Schubert intersection numbers**

$$c_{\lambda\mu\nu} := \int_{Gr_k(\mathbb{C}^n)} [X_\lambda][X_\mu][X^\nu],$$

then $c_{\lambda\mu}^\nu = c_{\lambda \ \mu \ w_0 \cdot \nu}$.

1.3. First positivity result

Proposition 1.6. (1) (Borel) Let B act on a nonempty projective scheme (or more generally, a complete one). Then there is a B -fixed point.

(2) Any complete scheme over \mathbb{C} has a natural fundamental class in homology: the \mathbb{N} -combination of the classes of its top-dimensional (geometric) components, where the coefficients are the lengths of the local rings at the generic points.

(3) (Grothendieck, Mumford) Let Y be a projective scheme (in a moment, the Grassmannian) and X a subscheme. There is a functorially associated “Hilbert scheme” parametrizing a family of subschemes of Y , all of whom have the same homology class (and more specifically, the same “K-class”, discussed later), and this moduli space is projective. (Our reference for Hilbert schemes is [EH00].)

(4) Hence every subscheme X of the Grassmannian is homologous to a scheme union X' of Schubert varieties.

(5) Hence each $c_{\lambda\mu}^\nu \geq 0$.

Proof. (1) Filter B by normal subgroups, so that the subquotients are 1-dimensional (most groups can’t be filtered like this). Prove the theorem for those two groups, $(\mathbb{C}, +)$ and $(\mathbb{C}^\times, \cdot)$. Then use induction.

(2) As stated, this is a definition, reducing to the case of X a variety (reduced and irreducible). Then we punt, appealing either to the statement that complex varieties are triangulable (1930s), or have resolutions of their singularities (1960s).

(3) Beyond our scope.

- (4) By the functoriality, since B acts on $Gr_k(\mathbb{C}^n)$ it acts on X 's Hilbert scheme. Let X_0 be a B -fixed point on there. Then X_0 's support must be a union of B -orbits, which are the Schubert varieties.
- (5) Apply the previous to X_μ^ν , obtaining some B -invariant degeneration X_0 with components X^λ and multiplicities $d_{\lambda\mu}^\nu$, i.e. $d_{\lambda\mu}^\nu \in \mathbb{N}$ is the length of the local ring at the generic point of $X^\lambda \subseteq X_0$. Then $[X_\mu^\nu] = [X_0] = \sum_\lambda d_{\lambda\mu}^\nu [X^\lambda]$. Multiply by $[X_\lambda]$, integrate, and we get $\int_{Gr_k(\mathbb{C}^n)} [X_\mu^\nu][X^\lambda] = d_{\lambda\mu}^\nu$, so $c_{\lambda\mu}^\nu = d_{\lambda\mu}^\nu$ and hence is nonnegative.

Q.E.D.

Examples:

- (1) Let X be a conic in the xyz -plane. Then the Hilbert scheme is the \mathbb{P}^5 of all conics. The B -invariant one is the double line $x^2 = 0$.
- (2) Let X be a disjoint union of two lines in \mathbb{P}^3 . First let one fall across the other, giving a union of two lines with an extra point embedded at the intersection (reduced everywhere else). Then let those lines fall atop one another, giving a double (Schubert) line with an extra embedded (Schubert) point, which doesn't contribute to the homology class (but will contribute to the K -class).

This hints at some other related subschemes in \mathbb{P}^3 : plane conics union a disjoint point. Indeed, this Hilbert scheme has two components, one consisting of pairs of lines and the other of conics plus points. (It's connected, which is true for all Hilbert schemes of projective space, a theorem of Hartshorne.)


- (3) A Schubert calculus example: $X_{0101}^{1010} \subset Gr_2(\mathbb{C}^4) \cong Gr_1(\mathbb{C}\mathbb{P}^3)$, which interpreted projectively is the $\mathbb{P}^1 \times \mathbb{P}^1$ worth of lines that touch the xy line and zw line. Move the latter continuously to xz , and we get lines that touch xy and xz . That's the (reduced) scheme $X_{1001} \cup X_{0110}$, with cohomology class $1[X_{1001}] + 1[X_{0110}]$.

So any subscheme of a Grassmannian (or more generally, a space with an action of a unipotent group N with finitely many orbits) leads naturally to a "Schubert calculus" problem with nonnegative integer answers.


The real game, then: how to calculate these natural numbers in a *manifestly positive* way, i.e. by counting some combinatorial objects?

There are many known ways (and sometimes-tricky bijections between them); the one we'll give is to use *puzzles* [KnTao03, KnTaoWood04].

The following three shapes, with edges labeled 0 or 1, are the **puzzle pieces**. They may be rotated but not reflected.



A **puzzle** is a size n triangle made of puzzle pieces, glued so as to have the edge labels match. The two possible puzzles with NW,NE sides labeled 0101.



Theorem 1.7. [KnTao03, KnTaoWood04] $c'_{\lambda\mu} = \#\{\text{puzzles with } \lambda, \mu, \nu \text{ on their NW, NE, S sides respectively, all read left-to-right}\}$. Equivalently, $c_{\lambda\mu\nu} = \#\{\text{puzzles with } \lambda, \mu, \nu \text{ on their NW, NE, S sides respectively, all read clockwise}\}$.

So these two puzzles compute the example (3) above.

In the remainder we'll indicate a geometric proof of this theorem, following [Kn1], and discuss harder problems requiring more puzzle pieces.

Note that in addition to the obvious Z_3 rotational symmetry of puzzles, which matches half of the S_3 symmetry of Schubert intersection numbers, one can flip a puzzle over while exchanging 0s and 1s, which matches the symmetry coming from Grassmannian duality. However, it is hard to see directly that the puzzle product is commutative. (The nicest self-contained combinatorial proof of this is in [P08].)

1.4. Interval rank varieties

Before giving an answer to this, we generalize the problem just a little, beyond Richardson varieties to *interval rank varieties*.

Given a matrix $M \in M_{k \times n}$, let's compute all $\{\text{rank } [i, j]\}_{i \leq j}$. These numbers are weakly increasing in j and $-i$, by 0 or 1 only, with the additional restriction that one doesn't see the pattern

$$\begin{matrix} r & r+1 \\ r & r \end{matrix}$$

anywhere in the matrix (or really, triangle) r .

Define a partial permutation matrix π to be one with *at most* one 1 in any row/column. To avoid confusion we refer to and draw the 1s as **dots** \bullet .

Proposition 1.8. *Such rank matrices r correspond 1:1 to upper triangular partial permutation matrices π under the correspondence*

$$r_{ij} = |[i, j]| - \#\{\text{dots in } \pi \text{ weakly SW of box } [i, j]\}, \quad \forall 1 \leq i \leq j \leq n.$$

If M has rank k , then π will have $n - k$ dots.

As before, many of these rank conditions imply others. Define the **diagram**² of π by crossing out strictly South and West (but not South-west) of each dot, and any rows or columns without dots, and taking the remaining boxes. (So each box with a dot is in the diagram.) Define the **essential set**³ as the Northeast corners of the diagram boxes.

For example,

$$\pi = \begin{pmatrix} \cdot & \cdot & \cdot & \bullet & \cdot & \cdot \\ & \cdot & \bullet & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & \bullet \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix} \text{ has diagram } \begin{pmatrix} + & + & - & \boxed{e} & | & \boxed{e} \\ & + & \boxed{e} & | & | & \boxed{e} \\ & & + & + & + & - \\ & & & + & + & \boxed{e} \\ & & & & + & + \\ & & & & & + \end{pmatrix}$$

with $-$, $|$, $+$ to indicate the crossing-out, and e s to indicate the essential boxes.

²This is closely related to the Rothe diagram used e.g. in Fulton's essential set definition [Fu92], but not the same, both for being flipped East/West and for the "strictly".

³It's actually possible to cut this down further, but we're using this name in analogy to [Fu92].

It is easy to show that given two adjacent rank conditions e.g. from (i, j) and $(i + 1, j)$, one implies the other, so very few are not implied by one of their neighbors, and this was worked out in [Fu92], where the essential set was introduced (slightly differently from here).

Proposition 1.9. *The rank conditions $\text{rank } [i, j] \leq r_{ij}$ in the essential set imply all the other rank conditions.*

(More specifically, any other rank condition is implied by some single essential condition; one doesn't have to cleverly combine them.)

- Exercise 1.10.**
- (1) Show that π has a unique essential rank condition, at (i, j) , iff its dots are the diagonal of some square with NE corner (i, j) .
 - (2) Find a π for $n = 4$ with three “essential” rank conditions, one of which is actually implied by the other two taken together.
 - (3) Show that the following are equivalent:
 - All the essential rank conditions are in the first row.
 - The dots are NW/SE, and in the first $n - k$ rows.
 - Π_r is a Schubert variety.

Theorem 1.11. *Let π be an upper triangular partial permutation with $n - k$ dots. Then the scheme Π_π defined by the rank conditions in proposition 1.8 is reduced and irreducible. Moreover, any intersection of such schemes is reduced.*

I hoped we would have time to prove this later, but we didn't. The quickest proof I know is in [Kn, §7.3], combined with [Kn1, theorem 1.8].

If we simply imposed random rank conditions, with some matrix r , what different schemes could we get? Different r s can give the same scheme: anywhere $r_{ij} > r_{i,j+1}$, we can cut down r_{ij} to $r_{i,j+1}$ without changing the scheme, and anywhere $r_{ij} + 1 < r_{i,j+1}$, we can cut down $r_{i,j+1}$ to $r_{ij} + 1$ without changing the scheme, and similarly for j instead of i . So it's enough to consider matrices r that only increase by 0, 1 as one goes North or East.

That leaves the forbidden pattern from before.

Proposition 1.12. *Let*

$$\begin{matrix} m & m+1 \\ m & m \end{matrix}$$

occur in the middle of a rank matrix, and Π_r be the associated scheme. Then $\Pi_r = \Pi_{r_1} \cup_{\Pi_{r_{12}}} \Pi_{r_2}$, where

$$r_1 = \begin{matrix} m & m \\ m & m' \end{matrix}, \quad r_2 = \begin{matrix} m & m+1 \\ m-1 & m' \end{matrix}, \quad r_{12} = \begin{matrix} m & m \\ m-1 & m \end{matrix}$$

where the t is in position (i, j) . More specifically, define $F^\circ \subseteq \mathbb{P}^1 \times Gr_k(\mathbb{C}^n)$ as

$$F^\circ := \bigcup_{t \in \mathbb{A}^1} \{t\} \times \begin{pmatrix} 1 & & & & & \\ & 1 & & & & t \\ & & 1 & & & \\ & & & \ddots & & \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix} \cdot X,$$

then F as its closure (adding the $t = \infty$ fiber), and $sh_{i \rightarrow j} X := F \cap (\{\infty\} \times Gr_k(\mathbb{C}^n))$.

Examples:

- (1) Let X be a single point, the coordinate k -plane \mathbb{C}^S that uses the k coordinates $S \subseteq \{1, \dots, n\}$. Then $sh_{i \rightarrow j} \{\mathbb{C}^S\} = \{\mathbb{C}^{sh_{i \rightarrow j} S}\}$, a first link of the two notions.
- (2) Let X be the divisor $p_S = 0$, given by the vanishing of the Plücker coordinate. Then $sh_{i \rightarrow j} X = \{p_{sh_{j \rightarrow i} S} = 0\}$, backwards. One should think of X as corresponding to $\binom{n}{k} \setminus \{S\}$, the set of coordinate subspaces lying in X .
- (3) Generalizing both examples, let $C \subseteq 2^{\binom{n}{k}}$ be a collection of k -element subsets, and define $W_C \subseteq Gr_k(\mathbb{C}^n)$ as the vanishing set $\{p_S = 0, S \notin C\}$. Then $sh_{i \rightarrow j} W_C = W_{sh_{i \rightarrow j} C}$. (W is for Neil White, who first considered these schemes in [W75].)
- (4) Let $X = \mathbb{C}^{01} \amalg \mathbb{C}^{10} \subseteq Gr_1(\mathbb{C}^2)$, defined by the equation $p_{01}p_{10} = 0$. Then moving it by t as in the definition, it becomes $(p_{01} + tp_{10})p_{10} = 0$, so as $t \rightarrow \infty$ we get the double point $(p_{10})^2 = 0$.

By construction, F is a flat family over \mathbb{P}^1 , with the effect that all of its fibers have the same Hilbert polynomial (as subvarieties of the Grassmannian, hence of projective space under the Plücker embedding). One way to determine F_∞ is to give an upper bound $(F_\infty)^+$ on it by determining (what are a priori) some of the equations that hold on it, and then to show that that upper bound has the same Hilbert polynomial as $F_0 = X$.

Proposition 2.2. *Let $X = Y \cap \{p_S = 0\}$, where Y is a $sh_{i \rightarrow j}$ -invariant variety, and $Y \supseteq \{\mathbb{C}^S, \mathbb{C}^{sh_{j \rightarrow i} S}\}$ (remember, backwards). Then $sh_{i \rightarrow j} X = Y \cap \{p_{sh_{j \rightarrow i} S} = 0\}$.*

Proof. By the condition, both $p_S = 0$ and $p_{sh_{j \rightarrow i} S} = 0$ define nonzero elements of Y 's coordinate ring. Since Y is a variety, they

both define non-zerodivisors. Modding out a non-zerodivisor of degree 1 replaces the Hilbert polynomial $h(d)$ by its difference $h(d) - h(d - 1)$, so the two have the same Hilbert polynomial.

Following the equations gives the containment \subseteq , so by the equality of Hilbert polynomials they are equal. Q.E.D.

Our key result is similar-sounding, in that it is about having only one non-shift-invariant rank condition, but that condition may involve several equations. I thank Nicolas Ford for catching this mistake from a previous version.

Proposition 2.3. *Let r be a rank matrix defining some Π_r , and let $a < b$. Assume that for*

- *one of the essential rank conditions is on $[a + 1, b]$, and*
- *for the others $[i, j]$, $\text{sh}_{b \rightarrow a}[i, j] = [i, j]$.*

Then $\text{sh}_{a \rightarrow b}\Pi_r$ is contained in a union of various $\Pi_{r'}$, whom can be determined through repeated use of proposition 1.12.

We can always ensure these hypotheses hold, assuming Π_r is not a Schubert variety; let $[a + 1, b]$ be the essential box with maximum a , then maximum b . (The not-Schubert condition says that $a + 1 \geq 2$.) Consequently, we have a combinatorial algorithm with which to degenerate any interval rank variety in the Grassmannian towards a union of Schubert varieties, and can thereby determine its homology class!

To spell this out further, we introduce also the combinatorial and geometric sweeping operations. The **combinatorial sweep** $\Psi_{i \rightarrow j}C$ of a collection C is just $C \cup \text{sh}_{i \rightarrow j}C$. The **geometric sweep** $\Psi_{i \rightarrow j}X$ is the image of the projection to $Gr_k(\mathbb{C}^n)$ of the family F from the beginning of 2.2, so $\Psi_{i \rightarrow j}X \supseteq X \cup \text{sh}_{i \rightarrow j}X$.

Proposition 2.4. • *If X is irreducible, then so too are F and $\Psi_{i \rightarrow j}X$.*

- *If any two of $X, \text{sh}_{i \rightarrow j}X, \Psi_{i \rightarrow j}X$ are equal, so is the third.*
- *If not, then $\dim \Psi_{i \rightarrow j}X = \dim X + 1$.*

Theorem 2.5. *Assume the setup of proposition 2.3. There is necessarily a dot in column b , which we'll call the wandering dot.*

Then $\Psi_{a \rightarrow b}\Pi_\pi = \Pi_\sigma$, where σ is constructed from π by moving the wandering dot up from whatever row k to row a , and the dot (if any) in row a down to that row k .

The π' are constructed from this σ . If there is no dot in column $b - 1$, one π' comes from moving the wandering dot in σ left one square to column $b - 1$.

For the other π' , consider the dots in σ that are minimally NW of the wandering dot (nobody else is in between). Each π' comes from moving the wandering dot West, while one of them moves East, ending in the same two columns. (If this causes the wandering dot to end up in the lower triangle, discard this misbegotten π' .)

For example, let

$$\pi = \begin{pmatrix} \cdot & \cdot & \cdot & \bullet & \cdot & \cdot \\ & \cdot & \bullet & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & \bullet \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix} \quad \text{with diagram} \quad \begin{pmatrix} + & + & - & \boxed{e} & | & \boxed{e} \\ & + & \boxed{e} & | & | & \boxed{e} \\ & & + & + & + & - \\ & & & + & + & \boxed{e} \\ & & & & + & + \\ & & & & & + \end{pmatrix}.$$

where the es are essential boxes. Let $(a, b) = (3, 6)$, above the SErnmost. Then

$$\sigma = \begin{pmatrix} \cdot & \cdot & \cdot & \bullet & \cdot & \cdot \\ & \cdot & \bullet & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & W & \cdot \\ & & & \cdot & \uparrow & \cdot \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix}, \quad \pi' = \begin{pmatrix} \cdot & \cdot & \cdot & \bullet & \cdot & \cdot \\ & \cdot & \bullet & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & W & \leftarrow \\ & & & \cdot & \cdot & \cdot \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix},$$

$$\begin{pmatrix} \cdot & \cdot & \cdot & \bullet & \cdot & \cdot \\ & \cdot & \rightarrow & \cdot & \cdot & \bullet \\ & & W & \cdot & \cdot & \leftarrow \\ & & & \cdot & \cdot & \cdot \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix}, \quad \begin{pmatrix} \cdot & \cdot & \cdot & \rightarrow & \cdot & \bullet \\ & \cdot & \bullet & \cdot & \cdot & \cdot \\ & & \cdot & W & \cdot & \leftarrow \\ & & & \cdot & \cdot & \cdot \\ & & & & \cdot & \cdot \\ & & & & & \cdot \end{pmatrix},$$

where the W indicates the wandering dot, and the arrows are only to show where dots have most recently moved from.

We won't hack through the combinatorics to prove this from proposition 2.3. A different approach is taken in [Kn1], where this is related to Lascoux's "transition formula" for Schubert polynomials.

Exercise 2.6. (1) Check the theorem in this example, using the algorithm from proposition 2.3.

(2) Do the same for all $n = 3$ examples.

(3) Do the same for all $n = 4, k = 2$ examples.

2.3. Vakil’s degeneration order

Geometric shifts were introduced⁴ in [Va06], where he considers the following **degeneration order** of shifts:

$$\begin{aligned}
 & n - 1 \rightarrow n, \\
 & n - 2 \rightarrow n, \quad n - 2 \rightarrow n - 1, \\
 & n - 3 \rightarrow n, \quad n - 3 \rightarrow n - 1, \quad n - 3 \rightarrow n - 2, \\
 & \quad \quad \quad \vdots \\
 & 1 \rightarrow n, \quad 1 \rightarrow n - 1, \quad \dots \quad 1 \rightarrow 4, \quad 1 \rightarrow 3, \quad 1 \rightarrow 2.
 \end{aligned}$$

Proposition 2.7 (essentially in [Va06]). *Let π be an upper triangular partial permutation matrix, and $a < b$. Assume*

- *the dots in rows $[1, a]$ are NW/SE,*
- *the dots in rows $[a + 1, n]$ are NW/SE, and*
- *the j dots in rows $[a + 1, n]$ are in the first such rows, $[a + 1, a + j]$.*

If $(a + 1, b)$ is not an essential box of π , then $\text{sh}_{a \rightarrow b} \Pi_\pi = \Pi_\pi$, so the unique π' is just π .

Otherwise $\text{sh}_{a \rightarrow b} \Pi_\pi \neq \Pi_\pi$, and we compute its components $\Pi_{\pi'}$ with theorem 2.5. There are at most two π' .

Let $(a, b)'$ be the next shift after (a, b) , i.e. $(a, b)' = (a, b - 1)$ unless $a = b - 1$, in which case $(a, b)' = (a - 1, n)$. Then each π' satisfies the conditions of the first paragraph, for $(a, b)'$.

Vakil only shows that the components of the shift have length 1, i.e. that the shift is generically reduced, whereas the above says that it’s actually reduced. We’ll need this actual reducedness later in order to compute in K -theory.

Example. Start with

$$\begin{aligned}
 \pi &= \begin{pmatrix} \cdot & \bullet & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \bullet & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}, & \text{diagram} & \begin{pmatrix} - & e & | & e \\ & + & + & - \\ & & + & e \\ & & & + \end{pmatrix}, & (a, b) &= (2, 4), \\
 \sigma &= \begin{pmatrix} \cdot & \bullet & \cdot & \cdot \\ \cdot & \cdot & \cdot & W \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}.
 \end{aligned}$$

⁴Though not under that name – he didn’t connect them with Erdős-Ko-Rado shifting theory.

Then the two π' are

$$\left(\begin{array}{cccc} \cdot & \bullet & \cdot & \cdot \\ & W & \leftarrow & \\ & & \cdot & \\ & & & \cdot \end{array} \right) \text{ which is Schubert, and } \left(\begin{array}{cccc} \cdot & \rightarrow & \cdot & \bullet \\ & W & \cdot & \leftarrow \\ & & \cdot & \\ & & & \cdot \end{array} \right)$$

with diagram $\begin{pmatrix} + & - & + & e \\ & e & + & + \\ & & + & + \\ & & & + \end{pmatrix}$.

For the latter one take $(a, b) = (1, 2)$, and get

$$\sigma = \begin{pmatrix} \cdot & W & \cdot & \cdot \\ & \uparrow & \cdot & \bullet \\ & & \cdot & \cdot \\ & & & \cdot \end{pmatrix}, \quad \pi' = \begin{pmatrix} W & \leftarrow & \cdot & \cdot \\ & \cdot & \cdot & \bullet \\ & & \cdot & \cdot \\ & & & \cdot \end{pmatrix}.$$

The final result is thus

$$\left[\Pi \left(\begin{pmatrix} \cdot & \bullet & \cdot & \cdot \\ & \cdot & \cdot & \cdot \\ & & \cdot & \bullet \\ & & & \cdot \end{pmatrix} \right) \right] = \left[\Pi \left(\begin{pmatrix} \cdot & \bullet & \cdot & \cdot \\ & \cdot & \cdot & \cdot \\ & & \cdot & \cdot \\ & & & \cdot \end{pmatrix} \right) \right] + \left[\Pi \left(\begin{pmatrix} \bullet & \cdot & \cdot & \cdot \\ & \cdot & \cdot & \cdot \\ & & \cdot & \bullet \\ & & & \cdot \end{pmatrix} \right) \right]$$

or

$$[X_{0101}^{1010}] = [X_{1001}] + [X_{0110}].$$

2.4. Partial puzzles

Imagine a would-be puzzle with μ on NE and ν on South, both read left-right, but no other edge labels. Shear this to fit into the upper triangle where π s live, so the ν ends up on the diagonal and the μ on the upper right. For the rest of this section we'll draw puzzles in this way.

Now we can think about Vakil's order as specifying a sequence of squares in the matrix upper triangle, or rhombi in the puzzle triangle. Note that once one gets to the (a, a) triangles, they are uniquely fillable, so it's harmless to add them as (a, a) at the end of each line of the degeneration order.

Define an (a, b) -**partial puzzle** to be a filling of the puzzle triangle with puzzle pieces, but only through positions (a, b) as indexed in the

above paragraph. Its **puzzle path** is the labels along its top, whose shape

- starts at $(0, 0)$ in matrix coordinates (row,column),
- goes SE to (a, a) ,
- goes E to (a, b) ,
- has a **kink** up to $(a - 1, b)$,
- goes E to $(a - 1, n)$,
- finally up to $(0, n)$.

Note that in this correspondence, the edges of the matrix boxes may cut puzzle rhombi in half. As such, we have to ditch our rhombus puzzle piece in favor of a triangle with labels $1, 0, R$ clockwise, and declare that a puzzle can't have any R s on its boundary.⁵ The main idea: we'll associate an interval rank variety to each puzzle path, such that the components of $\text{sh}_{a \rightarrow b}$ correspond to the ways to fill in one more square in the matrix. The partial permutation matrix should have the properties in proposition 2.7, in particular being NW/SE in each half of the triangle (above/below the path).

The construction, in the upper half: first draw little **rays** inside the puzzle

- left from each \uparrow
- up from each \uparrow and \leftarrow
- left from the kink if it's \leftarrow , in which case also make the next \leftarrow to its left get an upward pointing ray.

We require there be the same number of up and left rays for the puzzle path to be **viable**. If there are, put the dots on the intersections of these rays, in the only NW/SE way possible.

In the lower half, draw rays down from each \ominus . If the kink is \uparrow , and there is a \ominus to its right, the \ominus gets a downward ray, and these two rays meet at a dot. Put dots on all remaining downward rays NW/SE, as N as possible (à la proposition 2.7). Lots of pictures appear in [Kn1].

Write this association puzzle path \mapsto partial permutation matrix as $\gamma \mapsto \pi(\gamma)$.

Theorem 2.8. *Let γ be a puzzle path.*

If the labels on the kink and the edge just leftward are not $\leftarrow \uparrow$, then

- (1) *there is a unique way to put in two puzzle pieces,*
- (2) *the new path γ' has the same $\pi()$, and*

⁵Exercise: prove that if a sheared “puzzle” without this new condition has the same number of 1 s on the N and E sides, and no R s there, then it has no R s on the SW side either.

(3) $\Pi_{\pi(\gamma)}$ is $\text{sh}_{a \rightarrow b}$ -invariant.

If those labels are indeed $\oplus \mathbb{1}$, then there are either one or two ways to fill two triangles and remain viable. The corresponding $\{\gamma'\}$ correspond to the components of $\text{sh}_{a \rightarrow b} \Pi_{\pi(\gamma)} \neq \Pi_{\pi(\gamma)}$.

In this way, filling in the puzzle corresponds to keeping track of the components encountered during Vakil's degeneration.

The proof of this is a several-page case check [Kn1], quite straightforward yet somehow miraculous.⁶ The hard work was really in theorem 2.5.

§3. Equivariant and K -extensions

3.1. K -homology

To begin with, let A be a commutative ring, and consider its finitely generated modules, under direct sum. The K -homology group $K_{\bullet}(A)$ is freely generated by the set of isomorphism classes $[M]$ of finitely generated A -modules M , “modulo exact sequences”, meaning we impose $[M_2] = [M_1] + [M_3]$ for any exact sequence $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$. In particular, $[M_1 \oplus M_3] = [M_1] + [M_3]$. If $A \rightarrow B$ is a ring homomorphism making B a finitely generated A -module, we get a map $K_{\bullet}(B) \rightarrow K_{\bullet}(A)$, contravariantly.

One can easily soup this up by assuming A is graded and only using graded modules, or a group G acts on A and its modules, etc., in which case we'd write $K_{\bullet}^G(A)$.

We can reinterpret this ring-theoretic construction in terms of affine schemes, where the statement becomes “Given an affine scheme X , define its K -homology using coherent sheaves, and this gives a *covariant* functor for finite maps (proper with finite fibers).” At that point one can leave out the word “affine” and obtain a K -homology theory for all schemes and finite maps. (Using higher sheaf cohomology, one can extend this to proper maps.)

The most important example to understand will be the following. Let $A = \mathbb{C}[x, y]$, considered as a bigraded ring. Then we have an exact sequence of bigraded modules and maps

$$0 \rightarrow A/\langle xy \rangle \rightarrow (A/\langle x \rangle) \oplus (A/\langle y \rangle) \rightarrow A/\langle x, y \rangle \rightarrow 0.$$

Hence $[A/\langle xy \rangle] = [A/\langle x \rangle] + [A/\langle y \rangle] - [A/\langle x, y \rangle]$.

⁶The combinatorics in [Kn2], of “IP pipe dreams”, is better adapted than puzzles are to the geometry of shifting, and the proof looks less miraculous. After which the connecting of IP pipe dreams to puzzles is quite short.

3.1.1. *Comparison with ordinary homology.* Given a (closed) subscheme $X \subseteq Y$, we can associate a K -class $[X] \in K_\bullet(Y)$, really shorthand for $[\mathcal{O}_X]$ where \mathcal{O}_X is the structure sheaf of X . In the example above, we have the equation

$$[\text{union of axes}] = [x\text{-axis}] + [y\text{-axis}] - [\text{origin}]$$

in $K_\bullet^{T^2}(\mathbb{C}^2)$. This differs from what we would expect in ordinary homology, where we wouldn't have the last term.

One can make this precise; filter $K_\bullet(Y)$ according to the dimension of the support of the sheaf. Then there is a map from the associated graded ring to $H_*(Y)$ (or even better, to the Chow group, which then maps to $H_*(Y)$).

In the example from before, we computed the homology class of X_{0101}^{1010} by degenerating it to $X_{1001} \cup X_{0110}$, glued along X_{1010} . Therefore we get the equation on K -classes,

$$[X_{0101}^{1010}] = [X_{1001}] + [X_{0110}] - [X_{1010}].$$

3.2. K -cohomology

Not only can we \oplus sheaves, we can \otimes them, suggesting we make $K_\bullet(Y)$ into a ring. But it turns out that \otimes is not well-defined on K -equivalence classes, i.e. tensoring with M is not exact.

Of course, tensoring with a free module is just repeated direct sum, so that'd be okay. More generally tensoring with projective modules is fine; in scheme-theoretic language, use (finite-dimensional) vector bundles instead of all coherent sheaves. So define the **K -cohomology** $K^\bullet(Y)$ in the same way as $K_\bullet(Y)$, but only using exact sequences of vector bundles over Y . All of the following are easy except half of (5), for which it seems difficult to find a reference.

Proposition 3.1. (1) K^\bullet is a contravariant functor from schemes to rings.

- (2) $K^\bullet(Y)$ acts on $K_\bullet(Y)$, a “cap product” in K -theory.
- (3) $K_\bullet(Y)$ comes with a “fundamental class” $[Y]$.
- (4) There is a “Poincaré map” $K^\bullet(Y) \rightarrow K_\bullet(Y)$, taking a vector bundle $[\mathcal{V}]$ to its sheaf of sections, i.e. to $[\mathcal{V}] \cap [Y]$.
- (5) If Y is smooth and proper, this is an isomorphism (“Poincaré duality”).
- (6) The Schubert varieties on the Grassmannian (or any other G/P) give bases for K -homology and K -cohomology.

(The onto-ness of the Poincaré map is not hard to see; it is based on Hilbert's theorem that modules over polynomial rings have free resolutions of finite length.)

So now we have ***K*-theoretic Schubert calculus** to compute: what is the cup product $[X_\lambda][X_\mu]$, where these denote the elements of *K*-cohomology constructed using Poincaré duality?


For the geometry, we need to soup up Vakil’s proposition 2.7 using proposition 1.12: when $\text{sh}_{i \rightarrow j} \Pi_\pi$ has two components, their intersection is another interval rank variety, whose class must therefore be subtracted, as in the example $[X_{0101}^{1010}] = [X_{1001}] + [X_{0110}] - [X_{1010}]$.

3.2.1. *Second positivity result* The minuses may seem to kill the positivity statement $c_{\lambda\mu}^\nu \geq 0$ from before, but this can be fixed:

Theorem 3.2. [Buc02, Bri02] *In $K^\bullet(Gr_k(\mathbb{C}^n))$, $(-1)^{|\nu|-|\lambda|-|\mu|} c_{\lambda\mu}^\nu \geq 0$.*

The history is a little weird – Anders Buch gave an explicit combinatorial formula for $(-1)^{|\nu|-|\lambda|-|\mu|} c_{\lambda\mu}^\nu$ (in terms of tableaux, later bijected to *K*-puzzles, only much later explained geometrically), *after* which Michel Brion gave an abstract geometric proof that holds for general flag manifolds. Usually the geometry comes first (as, in this very story told, it has for all *G/P* other than Grassmannians).

The cleanest way to puzzlify Buch’s theorem is with the ***K*-piece**, which is twice the size of other pieces ($4x$ the area), and cannot be rotated. Try it out in the $[X_{0101}]^2$ case.



Theorem 3.3. [Buc02] *In $K^\bullet(Gr_k(\mathbb{C}^n))$, $(-1)^{|\nu|-|\lambda|-|\mu|} c_{\lambda\mu}^\nu$ is the number of puzzles using the usual three pieces and now the *K*-piece, with λ, μ, ν on the NW, NE and S sides respectively, each left-to-right.*

K-puzzles have a very weird Z_3 rotational symmetry. Why weird? The obvious analogue of corollary 1.5 in *K*-theory is

$$\int_{Gr_k(\mathbb{C}^n)} [X_\lambda][X^\mu] = \begin{cases} 1 & \text{if } \lambda \leq \mu \\ 0 & \text{if not} \end{cases}$$

where \int denotes the pushforward to a point in *K*-homology. But there is a less obvious analogue [Buc02, §8],

$$\int_{Gr_k(\mathbb{C}^n)} [X_\lambda][X^\mu](1 - [X_{box}]) = \delta_{\lambda\mu}$$

with which one can interpret the number of K -puzzles with λ, μ, ν clockwise as the S_3 -symmetric number

$$K \int_{Gr_k(\mathbb{C}^n)} [X_\lambda][X_\mu][X_\nu](1 - [X_{box}]).$$

3.3. Equivariant K -theory

As we said above, the definition of K -theory of a scheme X extends in a trivial way to G -equivariant K -theory (homology or cohomology). One extreme case is X a point, in which case a G -equivariant sheaf on X is just a G -representation, and $K^\bullet(X)$ is the representation ring $\text{Rep}(G)$.

The natural group to use in Schubert calculus is B . However, since the only irreps V of B are 1-dimensional (apply Borel’s theorem to $\mathbb{P}V$), every rep is K -equivalent to a sum of 1-d reps, which are thus really reps of $B/[B, B] \cong T$. The effect is that $K_B^\bullet(pt) \cong K_T^\bullet(pt)$ canonically, and it’s traditional to use T -equivariant instead of B -equivariant K -theory.

Proposition 3.4. *The K -theoretic Schubert classes $[\mathcal{O}_{X_\lambda}]$ are a basis of $K_T^\bullet(Gr_k(\mathbb{C}^n))$, as a module over $K_T^\bullet(pt) \cong \text{Rep}(T)$, a Laurent polynomial ring.*

I like to call the elements of $K_T^\bullet(pt)$ (and later, of $H_T^*(pt)$) “equivariant numbers”.

To keep track of the relation of the additively written group $T^* := \text{Hom}(T, \mathbb{C}^\times)$ and its multiplicative role inside $K_*^T(pt)$, it helps to denote the generators of $K_*^T(pt)$ by e^λ , not λ . In the application to Schubert calculus on Grassmannians, we’ll write $y_i \in T^*$ for the representation taking $\text{diag}(t_1, \dots, t_n) \mapsto t_i$.

Example: $Gr_1(\mathbb{C}^2) = \mathbb{P}^1$. The T -fixed points are $s = [* 0]$ and $n = [0 *]$, but only the latter point is Schubert. To compute the K_T -class $[s]$ as a combination of $[n]$ and $[\mathbb{P}^1]$, start with the surjections $\mathcal{O}_{\mathbb{P}^1} \rightarrow \mathcal{O}_{\{s\}}$. The kernel of this is *nonequivariantly* $\mathcal{O}(-1) \rightarrow \mathcal{O}_{\mathbb{P}^1}$, the dual of the section σ of $\mathcal{O}(1)$ vanishing at s (once, and nowhere else, hence the 1). Equivariantly, we have to twist by the weight of σ , which we can determine by its restriction to n , giving the equivariant exact sequence

$$0 \rightarrow \mathcal{O}(-1) \otimes \mathbb{C}_{-y_1} \rightarrow \mathcal{O}_{\mathbb{P}^1} \rightarrow \mathcal{O}_s \rightarrow 0$$

Hence

$$[\mathcal{O}(-1) \otimes \mathbb{C}_{-y_1}] + [\mathcal{O}_{\{s\}}] = [\mathcal{O}_{\mathbb{P}^1}]$$

or

$$[\mathcal{O}(-1)] = ([\mathbb{P}^1] - [s]) \exp(y_1) \text{ and similarly } [\mathcal{O}(-1)] = ([\mathbb{P}^1] - [n]) \exp(y_2).$$

hence

$$[s] = (1 - \exp y_2 - y_1)[\mathbb{P}^1] + \exp y_2 - y_1[n].$$

If $S \rightarrow T$ is a homomorphism of groups, there is an obvious map $K_T^\bullet(X) \rightarrow K_S^\bullet(Y)$. In the case that S is the trivial group, this factors through setting each $e^\lambda = 1$, or “ $\lambda \mapsto 0$ ”.

Theorem 3.5 (see [HL]). *If a torus T acts on a smooth projective variety M , then the map*

$$K_T^\bullet(M) \Big/ \langle \{e^\lambda - 1\}, \lambda \in T^* \rangle \rightarrow K^\bullet(M)$$

is an isomorphism.

*While less relevant to us, it's also true that the natural **localization map***

$$K_T^\bullet(M) \rightarrow K_T^\bullet(M^T) \cong K_T^\bullet(M) \otimes K_T^\bullet(pt)$$

is an inclusion, as was used frequently in T. Ikeda's and T. Lam's lectures, where it was written $\alpha \mapsto (\alpha|_f)_{f \in M^T}$ for M^T isolated, following [KnTao03].

On the Grassmannian and other flag manifolds G/P , an abstract positivity result was proven in [AGriMil], and is a little complicated to state. With more puzzle pieces, one can compute the K_T -class of X_λ^μ in their positive sense.

However, that is not Schubert calculus, because $[X^\mu] \neq [X_{w_0 \cdot \mu}]$ as *equivariant* classes. Before, we could turn $X_{w_0 \cdot \mu}$ into X^μ by multiplying it with w_0 , which lives in the connected group $GL(n)$, so this preserved the homology and K -homology classes.

In any case the puzzle rule for the K_T -class of $[X_\lambda^\mu]$ becomes intricate enough that we won't bother detailing it here. One very unfortunate thing is that the matching rules are *no longer completely local*; one must sometimes look ahead along a row to see if a puzzle piece is allowed. *Update*: the nonlocality is more natural-looking in the “IP pipe dream” formulation in [Kn2].

3.4. Equivariant cohomology

First we describe some properties, then some interpretation, and finally puzzles. There are many references, one being [Ki84].

- (1) Equivariant cohomology H_G^* is a functor from $\{G\text{-spaces and } G\text{-equivariant maps}\}$ to (supercommutative) rings, with a natural transformation to ordinary H^* .

- (2) Since every G -space has a canonical G -equivariant map to the (G -invariant) point, and triangles connecting these commute, we could instead say H_G^* takes values in $H_G^*(pt)$ -algebras.
- (3) If M is compact with a G -invariant even-dimensional cell decomposition, then $H_G^*(M)$ is a free module over $H_G^*(pt)$ with a basis indexed by the cells.
- (4) If G is a torus T , then $H_T^*(pt)$ is the symmetric algebra in the weight lattice T^* of T (where weights are given degree 2, making it commutative).
- (5) If M is a complex projective manifold, then over \mathbb{Q} the natural map

$$H_T^*(M) / \langle \lambda \in T^* \rangle \rightarrow H^*(M)$$

is an isomorphism, and the localization map $H_T^*(M) \rightarrow H_T^*(M^T) \cong H^*(M^T) \otimes H_T^*(pt)$ is injective.

Why aren't we proceeding from an actual definition of equivariant cohomology? The odd thing is that, in contrast to the easy equivariant extension of K -theory, actually defining equivariant cohomology requires infinite-dimensional spaces. (There is a de Rham theory, called the Cartan model, that does not.) As such it is generally easier to proceed from its universal properties rather than worrying about the details of a concrete definition.

We now interpret the ring structure. What is *ordinary* cohomology ring structure? The product $[X] \cup [Y]$ for $X, Y \subseteq Z$ measures the difficulty in disentangling X from Y (say, for X, Y, Z compact oriented manifolds). A topologist would say we should perturb Y to miss X as much as possible, i.e. become transverse, and then $[X] \cup [Y] = [X \cap Y]$.

Now assume a group acts on Z , and X, Y are invariant. Then we may not be able to perturb Y *while keeping it invariant*. In keeping with not actually defining equivariant cohomology, we won't derive the following result from first principles. (But just as homology is related to the associated graded of K -theory, one should see this as the leading-order terms as $t \rightarrow 1$ of the formula after proposition 3.4.)

Proposition 3.6. *Let T act on 1-dimensional spaces \mathbb{C}, V with weights $0, \lambda \in T^*$, so and n, s denote the T -fixed points $[* 0], [0 *]$. Then as classes in $H_T^2(\mathbb{P}(\mathbb{C} \oplus V))$,*

$$[s] = \lambda [\mathbb{P}(\mathbb{C} \oplus V)] + [n].$$

Here $[n]$ and $[s]$ are degree 2 because the points are real codimension 2 inside the line $\mathbb{P}(\mathbb{C} \oplus V)$, and the third term is degree 2 because we put T^* into $H_T^2(pt)$. This formula accords with the fact that one can

T -equivariantly deform n to s iff λ happens to be 0. Or, that passing from equivariant to ordinary corresponds to imposing $\lambda = 0$.

Corollary 3.7. *If we pick the shift $\text{sh}_{a \rightarrow b}$ of an interval rank variety Π_π as in proposition 2.3, then*


$$[\Pi_\pi] = [\text{sh}_{a \rightarrow b} \Pi_\pi] + (y_a - y_b)[\Psi_{a \rightarrow b} \Pi_\pi].$$

Proof. Recall the total space F of the family degenerating Π_π to its shift. This has a projection to \mathbb{P}^1 , and when we pull back the H_T^* equation from 3.6, we get

$$[\{0\} \times \Pi_\pi] = [\{\infty\} \times \text{sh}_{a \rightarrow b} \Pi_\pi] + (y_a - y_b)[F].$$

One must then check that the projection $F \rightarrow \Psi_{a \rightarrow b} \Pi_\pi$ has degree 1, which is in [Kn1] and I will skip here. Pushing the equation above forward along that projection, we get the desired equation. Q.E.D.

The effect is, we should get an extra correction term with a factor of the equivariant number $y_a - y_b$ each time $\text{sh}_{a \rightarrow b} \Pi_\pi \neq \Pi_\pi$. We know when the latter happens, from theorem 2.8: the labels on the kink and edge immediately left must be \uparrow . The **equivariant puzzle piece** we create to fit in there looks like



and may not be rotated. To determine the factor $y_a - y_b$ it contributes, we drop lines SW and SE from it, coming out of the puzzle at the South edges a, b . Then the $H_T^*(pt)$ structure constant is

$$c_{\lambda\mu}^\nu = \sum_P \prod_{\text{eqvt rhombi } \rho \text{ in } P} (y_{a(\rho)} - y_{b(\rho)}).$$

The proof of this formula in [KnTao03] worked backwards from the “most equivariant” case $c_{\lambda\lambda}^\lambda$, and gaily divided by factors $y_a - y_b$ throughout in its derivation. Such a derivation becomes impossible if one specializes to ordinary cohomology in advance, taking each $y_i \mapsto 0$.

§4. Puzzles for the Belkale-Kumar Schubert calculus of other partial flag manifolds

There is a degeneration of the cohomology ring of G/P in [BeKu06] that is more useful for their applications to inequalities in linear algebra problems. (Each nonvanishing structure constant in ordinary Schubert calculus implies a certain inequality [BS], but this list of inequalities is very redundant. To have a less redundant list, it's actually nice to replace most of the structure constants by zero.)

Unlike ordinary cohomology, this Belkale-Kumar product is not functorial under projections $G/P \rightarrow G/Q$, so each flag manifold must be handled separately. Let us consider only GL_n/P , where the reductive part of P is $\prod_{i=1}^m GL(n_i)$. Then the Schubert classes can be indexed not by bit strings λ, μ, ν as on the Grassmannian, but by words π, ρ, σ in $1 \dots m$ where the letter i is used n_i times.

- Theorem 4.1.** (1) [BeKu06] *If the B-K structure constant $d_{\pi\rho}^\sigma$ is nonzero, then it matches the actual structure constant $c_{\pi\rho}^\sigma$.*
- (2) [KnP] *$d_{\pi\rho}^\sigma$ is the number of puzzles with boundary labels π, ρ, σ , made of (i, i, i) -triangles and (i, j, i, j) -rhombi where $i > j$.*
- (3) [KnP] *It factors as $\prod_{i < j \leq m} c_{\pi_{ij}\rho_{ij}}^{\sigma_{ij}}$, where π_{ij} is π with all other letters removed (likewise ρ, σ).*

Around 2000 (so, long before [BeKu06]) I circulated among a small number of people, a conjectural puzzle rule for actual Schubert calculus on GL_n/P . I'm still very pleased with this beautiful conjecture and quite annoyed that it's wrong – already for 3-step flag manifolds in 5-space, it defines a noncommutative ring. *Ç'est la vie*. However, for quantum Schubert calculus purposes Anders Buch was interested in 2-step flag manifolds [BuKrTam03], where this conjecture seems to be correct! One can think of it with all triangular pieces

$$(0, 0, 0), (1, 1, 1), (2, 2, 2), (1, 0, 10), (2, 0, 20), (2, 1, 21), (2, (10), 2(10)), \\ ((21), 0, (21)0)$$

with edge-labels $0, 1, 2, 10, 20, 21, 2(10), (21)0$. As before only single numbers can appear on the boundary of a puzzle. Examples appear in [BuKrTam03]. *Update:* Indeed, the 2-step conjecture is now a theorem, proven in [BuKrPTam03].

References

- [AGriMil] DAVID ANDERSON, STEPHEN GRIFFETH, EZRA MILLER: Positivity and Kleiman transversality in equivariant K -theory of homogeneous spaces. <http://arxiv.org/abs/0808.2785>
- [BeKu06] PRAKASH BELKALE, SHRAWAN KUMAR: Eigenvalue problem and a new product in cohomology of flag varieties, *Inventiones Math.* 166, 185–228 (2006). <http://arxiv.org/abs/math.AG/0407034>
- [BS] ARKADY BERENSTEIN, REYER SJAMAAR: Coadjoint orbits, moment polytopes, and the Hilbert–Mumford criterion, *J. Amer. Math. Soc.* 13 (2000), 433–466. <http://de.arxiv.org/abs/math.SG/9810125>
- [Bri02] MICHEL BRION: Positivity in the Grothendieck group of complex flag varieties, *J. Algebra* (special volume in honor of Claudio Procesi) 258 (2002), 137–159. <http://arxiv.org/abs/math/0105254>
- [Buc02] ANDERS BUCH: A Littlewood–Richardson rule for the K -theory of Grassmannians, *Acta Math.* 189 (2002), 37–78. <http://arxiv.org/abs/math.AG/0004137>
- [BuKrTam03] ———, ANDREW KRESCH, HARRY TAMVAKIS: Gromov–Witten invariants on Grassmannians, *J. Amer. Math. Soc.* 16 (2003), 901–915. <http://arxiv.org/abs/math.AG/0306388>
- [BuKrPTam03] ———, ———, KEVIN PURBHOO, HARRY TAMVAKIS: The puzzle conjecture for the cohomology of two-step flag manifolds, preprint 2014. <http://arxiv.org/abs/1401.1725>
- [EH00] DAVID EISENBUD AND JOE HARRIS: *The geometry of schemes*. New York: Springer, 2000.
- [EKR61] PÁL ERDŐS, CHAO KO, RICHARD RADO: Intersection theorems for systems of finite sets, *Quarterly Journal of Mathematics, Oxford Series, series 2* (1961) 12: 313–320. <http://dx.doi.org/10.1093/qmath/12.1.313>
- [Fu92] WILLIAM FULTON: Flags, Schubert polynomials, degeneracy loci, and determinantal formulas, *Duke Math. J.* 65 (1992), no. 3, 381–420.
- [Fu97] ———: *Young tableaux, with applications to representation theory and geometry*. Cambridge University Press, 1997.
- [HL] MEGUMI HARADA, GREG LANDWEBER: Surjectivity for Hamiltonian G -spaces in K -theory. <http://de.arxiv.org/abs/math.SG/0503609>
- [Ki84] FRANCES CLARE KIRWAN: *Cohomology of quotients in symplectic and algebraic geometry*. Vol. 31. Princeton University Press, 1984.
- [Kl73] STEVEN KLEIMAN: The transversality of a general translate, *Compositio Mathematica* (1974), Volume: 28, Issue: 3, page 287–297.
- [KnTao03] ALLEN KNUTSON, TERENCE TAO: Puzzles and (equivariant) cohomology of Grassmannians, *Duke Math. J.* 119 (2003), no. 2, 221–260. <http://arxiv.org/abs/math.AT/0112150>
- [KnTaoWood04] ———, ———, CHRIS WOODWARD: The honeycomb model of $GL(n)$ tensor products II: Puzzles determine facets of the Littlewood–Richardson cone, *J. Amer. Math. Soc.* 17 (2004), no. 1, 19–48. <http://arxiv.org/abs/math.CO/0107011>

- [Kn1] ———: Puzzles, positroid varieties, and equivariant K -theory of Grassmannians.
<http://arxiv.org/abs/1008.4302>
- [KnP] ———, KEVIN PURBHOO: Product and puzzle formulæ for GL_n Belkale-Kumar coefficients. *Electronic Journal of Combinatorics*, 18(1) (2011) no. 1, p76. <http://arxiv.org/abs/1008.4979>
- [Kn] ———: Frobenius splitting, point-counting, and degeneration.
<http://arxiv.org/abs/0911.4941>
- [Kn2] ———: Schubert calculus and shifting of interval positroid varieties.
<http://arxiv.org/abs/1408.1261>
- [P08] KEVIN PURBHOO: Puzzles, tableaux, and mosaics. *J. Algebraic Combin.* 28 (2008), no. 4, 461–480.
- [Va06] RAVI VAKIL: A geometric Littlewood-Richardson rule, *Annals of Math.* 164 (2006), 371–422.
<http://annals.math.princeton.edu/annals/2006/164-2/p01.xhtml>
- [W75] NEIL WHITE: The bracket ring of a combinatorial geometry, I. *Transactions of the AMS* 202 (1975) 79–95.

Cornell University, Department of Mathematics, Ithaca, NY, USA
E-mail address: allenk@math.cornell.edu