Advanced Studies in Pure Mathematics 32, 2001 Groups and Combinatorics — in memory of Michio Suzuki pp. 289–290

On the Lattice of all Subgroups of a Finite Noncyclic Simple Group

Walter Feit and M.A. Shahabi

In some of his earliest work Suzuki studied the lattice L(G) of all subgroups of a group G. Amongst other results he showed that if Gis a finite noncyclic simple group and $L(H) \approx L(G \times G)$ then $H \approx$ $G \times G$ [2, Theorem 19, p.53]. (Here of course \approx denotes either a lattice isomorphism or a group isomorphism as appropriate.) In particular this implies that if G and H are finite simple groups with $L(G \times G) \approx$ $L(H \times H)$ then $G \approx H$. (The case that G is cyclic of prime order is clear.) If G is cyclic of prime order then L(G) has just 2 elements so that $L(G) \approx L(H)$ implies only that H is cyclic of some prime order and so need not be isomorphic to G. However this leaves open the natural question of whether a finite noncyclic simple group is characterized by its lattice of subgroups. The purpose of this note is to show that by using the classification of the finite simple groups and further results from [2] that this is the case. More precisely:

Theorem. Let G and H be finite noncyclic simple groups. Then $G \approx H$ if and only if $L(G) \approx L(H)$.

It is clear that if $G \approx H$ then $L(G) \approx L(H)$. The proof of the converse needs some deep results. If G is a finite group and p is a prime let $n_p(G)$ denote the number of elements of order p in G. Then G contains exactly $n_p(G)/(p-1)$ subgroups of order p. The following will be needed:

- (I) [2, Theorem 15, p.51] If G is a finite noncyclic simple group and H is a finite group with $L(G) \approx L(H)$ then H is a finite noncyclic simple group of the same order as G.
- (II) [2, Theorem 8, p.45] Let G be a finite noncyclic simple group and let p be a prime. If φ is a lattice isomorphism of L(G)

Received December 9, 1998.

Revised April 26, 2000.