

Table of contents

Chap. 1 Existence and main properties of \mathbf{W}

1.0 Introduction

1.1 Existence of \mathbf{W} and first properties

1.1.1 A few more notations

1.1.2 A Feynman-Kac penalisation result

1.1.3 Definition of \mathbf{W}

1.1.4 Study of the canonical process under $W_\infty^{(\lambda\delta_0)}$

1.1.5 Some remarkable properties of \mathbf{W}

1.1.6 Another approach to Theorem 1.1.6

1.1.7 Relation between \mathbf{W} and other penalisations (than the Feynman-Kac ones)

1.2 W -Brownian martingales associated to \mathbf{W}

1.2.1 Definition of the martingales $(M_t(F), t \geq 0)$

1.2.2 Examples of martingales $(M_t(F), t \geq 0)$

1.2.3 A decomposition Theorem for positive Brownian supermartingales

1.2.4 A decomposition result for the martingale $(M_t(F), t \geq 0)$

1.2.5 A penalisation Theorem, for functionals in class \mathcal{C}

1.2.6 Some other results about the martingales $(M_t(F), t \geq 0)$

1.3 Invariant measures related to \mathbf{W}_x and $\mathbf{\Lambda}_x$

1.3.1 The process $(\mathcal{X}_t, t \geq 0)$

1.3.2 The measure $\mathbf{\Lambda}_x$

1.3.3 Invariant measures for the process $((X_t, L_t^\bullet), t \geq 0)$

1.3.4 Invariant measures for the process $(L_t^{X_t-\bullet}, t \geq 0)$

Chap. 2 Existence and properties of $\mathbf{W}^{(2)}$

2.1 Existence of $\mathbf{W}^{(2)}$

2.1.1 Notation and Feynman-Kac penalisations in two dimensions

2.1.2 Existence of the measure $\mathbf{W}^{(2)}$

2.2 Properties of $\mathbf{W}^{(2)}$

2.2.1 Some notation

2.2.2 Description of the canonical process under $W_\infty^{(2,q_0)}$

2.2.3 Another description of the measure $\mathbf{W}^{(2)}$

2.3 Study of the winding process under $\mathbf{W}^{(2)}$

2.3.1 Spitzer's Theorem

2.3.2 An analogue of Spitzer's Theorem

2.4 $W^{(2)}$ -martingales associated to $\mathbf{W}^{(2)}$

2.4.1 Definition of $(M_t^{(2)}(F), t \geq 0)$

2.4.2 A decomposition Theorem for positive $W^{(2)}$ -supermartingales

2.4.3 A decomposition Theorem of the martingales $(M_t^{(2)}(F), t \geq 0)$

Chap. 3 The analogue of the measure \mathbf{W} for a class of linear diffusions

3.1 Main hypotheses and notations

3.1.1 Our framework is that of Salminen-Vallois-Yor

3.1.2 The semi-group of $(X_t, t \geq 0)$

3.1.3 The local time process

3.1.4 The process X conditioned not to vanish

3.1.5 A useful Proposition

3.2 The σ -finite measure \mathbf{W}^*

3.2.1 Definition of \mathbf{W}^*

3.2.2 Some properties of \mathbf{W}^*

3.2.3 Relation between the measure \mathbf{W}^* and penalisations

3.3 The example of Bessel processes with dimension d ($0 < d < 2$)

3.3.1 Transcription of our notation in the context of Bessel processes

3.3.2 The measure $\mathbf{W}^{(-\alpha)}$

3.3.3 Relations between $\mathbf{W}^{(-\alpha)}$ ($d = 2(1 - \alpha)$) and Feynman-Kac penalisations

3.4 Another description of $\mathbf{W}^{(-\alpha)}$ (and of \mathbf{W}_g^*)

3.5 Penalisations of α -stable symmetric Lévy process ($1 < \alpha \leq 2$)

3.5.1 Notation and classical results

3.5.2 Definition of the σ -finite measure \mathbf{P}

3.5.3 The martingales $(M_t(F), t \geq 0)$ associated with \mathbf{P}

3.5.4 Relations between \mathbf{P} and penalisations

Chap. 4 An analogue of the measure \mathbf{W} for discrete Markov chains

4.0 Introduction

4.1 Construction of the σ -finite measures $(\mathbb{Q}_x, x \in E)$

4.1.1 Notation and hypothesis

4.1.2 A family of new measures

4.1.3 Definition of the measures $(\mathbb{Q}_x, x \in E)$

4.2 Some properties of $(\mathbb{Q}_x, x \in E)$

4.2.1 Martingales associated with $(\mathbb{Q}_x, x \in E)$

4.2.2 Properties of the canonical process under $(\mathbb{Q}_x, x \in E)$

4.2.3 Dependence of \mathbb{Q}_x on x_0

4.2.4 Dependence of \mathbb{Q}_x on ϕ

4.3 Some examples

4.3.1 The standard random walk

4.3.2 The "bang-bang random walk"

4.3.3 The random walk on a tree

4.3.4 Some more general conditions for the existence of ϕ

4.3.5 The standard random walk on \mathbb{Z}^2