

Research Article

Homoclinic Orbits for a Class of Nonperiodic Hamiltonian Systems

Wenping Qin, Jian Zhang, and Fukun Zhao

Department of Mathematics, Yunnan Normal University, Yunnan, Kunming 650092, China

Correspondence should be addressed to Fukun Zhao, fukunzhao@163.com

Received 6 December 2011; Accepted 16 January 2012

Academic Editor: Svatoslav Staněk

Copyright © 2012 Wenping Qin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the following nonperiodic Hamiltonian system $\dot{z} = \mathcal{J}H_z(t, z)$, where $H \in C^1(\mathbb{R} \times \mathbb{R}^{2N}, \mathbb{R})$ is the form $H(t, z) = (1/2)B(t)z \cdot z + R(t, z)$. We introduce a new assumption on $B(t)$ and prove that the corresponding Hamiltonian operator has only point spectrum. Moreover, by applying a generalized linking theorem for strongly indefinite functionals, we establish the existence of homoclinic orbits for asymptotically quadratic nonlinearity as well as the existence of infinitely many homoclinic orbits for superquadratic nonlinearity.

1. Introduction and Main Results

In this paper, we are interested in the existence of homoclinic orbits of the Hamiltonian system

$$\dot{z} = \mathcal{J}H_z(t, z), \quad (\text{HS})$$

where $z = (p, q) \in \mathbb{R}^N \times \mathbb{R}^N = \mathbb{R}^{2N}$, $\mathcal{J} = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}$, and $H \in C^1(\mathbb{R} \times \mathbb{R}^{2N}, \mathbb{R})$ is the form

$$H(t, z) = \frac{1}{2}B(t)z \cdot z + R(t, z) \quad (1.1)$$

with $B(t) \in C(\mathbb{R}, \mathbb{R}^{4N^2})$ being a $2N \times 2N$ symmetric matrix valued function, and $R \in C^1(\mathbb{R} \times \mathbb{R}^{2N}, \mathbb{R})$. Here, by a homoclinic orbit of (HS), we mean a solution of the equation satisfying $z(t) \neq 0$ and $z(t) \rightarrow 0$ as $|t| \rightarrow \infty$.

Establishing the existence of homoclinic orbits for system like (HS) is one of the most important problems in the theory of Hamiltonian systems. In very recent years, many

authors devoted to the existences of homoclinic orbits for Hamiltonian systems via critical point theory. For example, see [1–5] for the second-order systems and [6–18] for the first-order systems. Coti-Zelati et al. first considered the system (HS) in [6], and they obtained a homoclinic orbit for strictly convex Hamiltonian system. The existence of infinitely many homoclinic orbits was established in [15]. Subsequently, Hofer and Wycsocki removed the convexity assumptions in [13]. Later, suppose that $R(t, z)$ and $B(t)$ depend periodically on t , the existence of homoclinic orbit for (HS) was considered in [7, 8, 12, 16, 18].

Without assumption of periodicity, the problem is quite different in nature. In [10], Ding and Li first obtained one homoclinic orbit for the nonperiodic system (HS), see also [9, 11] and the references therein for recent works on this direction.

Motivated by [9, 11], in this paper, we introduce a new nonperiodic assumption on $B(t)$ as the following.

(B₀) $B(t) \in C(\mathbb{R}, \mathbb{R}^{2N \times 2N})$, there exists $r_0 > 0$ such that, for any $h > 0$,

$$|\{t \in \mathbb{R} : |t - t_1| \leq r_0, \mathcal{J}_0 B(t) < h\}| \rightarrow 0, \quad \text{as } |t_1| \rightarrow \infty, \quad (1.2)$$

where \mathcal{J}_0 will be defined in Section 2; we regard a real function $U(x)$ as a symmetric matrix $U(x)I_{2N \times 2N}$, and, for two given matrix valued functions $L_1(t)$ and $L_2(t)$, we say that $L_1(t) \leq L_2(t)$ if and only if

$$\max_{\xi \in \mathbb{R}^{2N}, |\xi|=1} (L_1(t) - L_2(t))\xi \cdot \xi \leq 0, \quad (1.3)$$

and $L_1(t) > L_2(t)$ if and only if $L_1(t) \leq L_2(t)$ does not hold. Obviously, (B₀) holds if $B(t)$ satisfies.

(B₁) For any $b > 0$, the set $\Lambda^b := \{t \in \mathbb{R} : \mathcal{J}_0 B(t) < b\}$ is nonempty and has finite Lebesgue measure.

One of the main ingredients of our work is two steps, the first is to show that the spectrum of Hamiltonian operator

$$A := -\left(\mathcal{J} \frac{d}{dt} + B\right) \quad (1.4)$$

consists of a sequence of eigenvalues with finite multiplicity which is unbounded from above and below (see Lemma 2.2), and hence the energy functional corresponding to (HS) is strongly indefinite; the second is to show that the working space E has compact embedding property (see Lemma 2.3).

First, we handle the asymptotically quadratic case. In what follows, $\widehat{R}(t, z) := (1/2)R_z(t, z)z - R(t, z)$. Suppose that

(R₁) $R \in C^1(\mathbb{R} \times \mathbb{R}^{2N}, [0, \infty))$ and $R_z(t, z) = o(|z|)$ as $|z| \rightarrow 0$ uniformly in t ,

(R₂) there exists a bounded function $R_\infty \in C(\mathbb{R}, \mathbb{R})$ such that $|R_z(t, z) - R_\infty(t)z|/|z| \rightarrow 0$ as $|z| \rightarrow \infty$ uniformly in t , and $b := \inf_{t \in \mathbb{R}} R_\infty(t) > \inf[(0, \infty) \cap \sigma(A)]$, where $\sigma(A)$ denotes the spectrum of A ,

(R₃) $\widehat{R}(t, z) > 0$ if $z \neq 0$, $\widehat{R}(t, z) \rightarrow \infty$ as $|z| \rightarrow \infty$.

We recall that a solution z_1 of (HS) is called a least energy solution if it possesses minimum energy among all solutions, that is,

$$\Phi(z_1) = \theta := \inf\{\Phi(z) \mid z \text{ is a nonzero solution of (HS)}\}, \tag{1.5}$$

where Φ is the energy functional

$$\Phi(z) := - \int_{\mathbb{R}} \left(\frac{1}{2} \mathcal{J}\dot{z} \cdot z + H(t, z) \right) dt. \tag{1.6}$$

Let k be the number of eigenvalues of the operator A lying in $(0, b)$, our main results are the following.

Theorem 1.1. *Suppose (B_0) , (R_1) – (R_3) be satisfied. Then, (HS) has a least energy solution. Moreover, if $R(t, z)$ is even in z , then (HS) has at least k pairs of solutions.*

Next, we consider the superquadratic case. Assume

$$(R_4) \ R(t, z) \cdot |z|^{-2} \rightarrow \infty \text{ as } |z| \rightarrow \infty \text{ uniformly in } t,$$

$$(R_5) \ \widehat{R}(t, z) > 0 \text{ if } z \neq 0, \text{ and there are } \mu > 1 \text{ and } r, c_1 > 0, \text{ such that, for } |z| \geq r, |R_z(t, z)|^\mu \leq c_1 \widehat{R}(t, z) |z|^\mu.$$

Theorem 1.2. *Suppose (B_0) , (R_1) , and (R_4) – (R_5) be satisfied. Then, (HS) has a least energy solution. Moreover, if $R(t, z)$ is even in u . Then, (HS) has infinitely many solutions.*

Remark 1.3. The assumptions (R_1) , (R_4) – (R_5) imply that $\widehat{R}(t, z) \rightarrow \infty$ uniformly in t as $|z| \rightarrow \infty$.

2. Variational Framework and Linking Structure

In order to establish a variational setting for the system (HS), in this section, we first study the spectrum of the Hamiltonian operator.

Observe that, since we have assumed (B_0) about $B(t)$, $A = -(\mathcal{J}(d/dt) + B(t))$ is a self-dajoint operator on $L^2(\mathbb{R}, \mathbb{R}^{2N})$ with $\mathfrak{D}(A) \subset H^1(\mathbb{R}, \mathbb{R}^{2N})$. Let $|\cdot|_q$ denote the usual L^q -norm, $(\cdot, \cdot)_2$ denote the usual L^2 inner product, c, c_i, d , or d_i stand for different positive constants. Let $\sigma(A), \sigma_d(A)$, and $\sigma_e(A)$ be the spectrum of A , the discrete spectrum of A , and the essential spectrum of A , respectively. Observe that $\mathfrak{D}(A)$ is a Hilbert space with the graph inner product

$$(z_1, z_2)_A := (Az_1, Az_2)_2 + (z_1, z_2)_2, \tag{2.1}$$

and the induced norm $\|z\|_A := (z, z)_A^{1/2}$.

Set $\mathcal{J}_0 := \begin{pmatrix} 0 & I_N \\ I_N & 0 \end{pmatrix}$ and $A_0 := \mathcal{J}(d/dt) + \mathcal{J}_0$ (self-adjoint with $\mathfrak{D}(A_0) = H^1(\mathbb{R}, \mathbb{R}^{2N})$); thus, $A_0^2 = -d^2/dt^2 + 1$. Let $|A_0|$ denote the absolute value of A_0 , and we have

$$\|A_0 z\|_2^2 = |A_0 z|_2^2 = (A_0 z, A_0 z)_2 = (A_0^2 z, z) = \left(\left(-\frac{d^2}{dt^2} + 1 \right) z, z \right)_2 = |\nabla z|_2^2 + |z|_2^2, \quad (2.2)$$

which implies that

$$\|z\|_{H^1} = \|A_0 z\|_2, \quad (2.3)$$

for all $z \in \mathfrak{D}(A_0^2) = H^2(\mathbb{R}, \mathbb{R}^{2N})$, hence for all $z \in H^1(\mathbb{R}, \mathbb{R}^{2N})$ because of the density of H^2 in H^1 .

Lemma 2.1. *For any $z \in \mathfrak{D}(A) \subset H^1(\mathbb{R}, \mathbb{R}^{2N})$, there exists $d > 0$ such that*

$$\|z\|_{H^1} = \|A_0 z\|_2 \leq d \|z\|_A. \quad (2.4)$$

Proof. Let A_1 be the restriction of A_0 to $\mathfrak{D}(A)$, A_1 is a linear operator from $\mathfrak{D}(A)$ to L^2 . We claim that A_1 is closed. Indeed, let $z_n \xrightarrow{\|\cdot\|_A} z$ and $A_1 z_n \xrightarrow{\|\cdot\|_2} w$. Then, $z \in \mathfrak{D}(A)$, and since A_0 is closed, $A_1 z_n = A_0 z_n \xrightarrow{\|\cdot\|_2} A_0 z = A_1 z$, hence the claim holds. Now the Closed Graph Theorem implies that $A_1 : \mathfrak{D}(A) \rightarrow L^2$ is a bounded linear operator, so $\|A_0 z\|_2 = \|A_1 z\|_2 \leq d \|z\|_A$ for all $z \in \mathfrak{D}(A)$. This together with (2.3) implies (2.4). \square

Lemma 2.2. *Suppose (B_0) holds. Then $\sigma(A) = \sigma_d(A)$.*

Proof. Similar to the idea of [9, 19], for any $h > 0$, if (B_0) holds, set

$$(\mathcal{J}_0 B(t) - h)^+ := \begin{cases} \mathcal{J}_0 B(t) - h, & \text{if } \mathcal{J}_0 B(t) - h \geq 0, \\ 0, & \text{if } \mathcal{J}_0 B(t) - h < 0, \end{cases} \quad (2.5)$$

and $(\mathcal{J}_0 B(t) - h)^- = (\mathcal{J}_0 B(t) - h) - (\mathcal{J}_0 B(t) - h)^+$. Thus, $A = A_2 + \mathcal{J}_0(\mathcal{J}_0 B(t) - h)^-$, where

$$A_2 = -\mathcal{J} \frac{d}{dt} + \mathcal{J}_0(\mathcal{J}_0 B(t) - h)^+ + h \mathcal{J}_0. \quad (2.6)$$

Since $\mathcal{J}_0^2 = I$ and $\mathcal{J}_0\mathcal{J} = -\mathcal{J}\mathcal{J}_0$, we have, for $z \in \mathfrak{D}(A)$,

$$\begin{aligned}
 (A_2z, A_2z)_2 &= \left| \left(-\mathcal{J} \frac{d}{dt} + \mathcal{J}_0(\mathcal{J}_0B(t) - h)^+ \right) z + h\mathcal{J}_0z \right|_2^2 \\
 &= \left| \left(-\mathcal{J} \frac{d}{dt} + \mathcal{J}_0(\mathcal{J}_0B(t) - h)^+ \right) z \right|_2^2 + h^2|z|_2^2 + (-\mathcal{J}\dot{z}, h\mathcal{J}_0z)_2 + (h\mathcal{J}_0z, -\mathcal{J}\dot{z})_2 \\
 &\quad + ((\mathcal{J}_0(\mathcal{J}_0B(t) - h)^+)z, h\mathcal{J}_0z)_2 + ((h\mathcal{J}_0z, \mathcal{J}_0(\mathcal{J}_0B(t) - h)^+)z)_2 \\
 &= \left| \left(-\mathcal{J} \frac{d}{dt} + \mathcal{J}_0(\mathcal{J}_0B(t) - h)^+ \right) z \right|_2^2 + h^2|z|_2^2 + 2h((\mathcal{J}_0B(t) - h)^+z, z)_2 \\
 &\geq h^2|z|_2^2.
 \end{aligned} \tag{2.7}$$

Thus, $\sigma(A_2) \subset \mathbb{R} \setminus (-h, h)$.

We claim that $\sigma_e(A) \cap (h, h) = \emptyset$. Assume by contradiction that there is $v \in \sigma_e(A)$ with $|v| < h$. Let $\{z_n\} \subset \mathfrak{D}(A)$ with $|z_n|_2 = 1$, $z_n \rightarrow 0$ in L^2 and $|(A - v)z_n|_2 \rightarrow 0$. Moreover, by (B₀), one can check that the multiplication operator $z \rightarrow \mathcal{J}_0(\mathcal{J}_0B(t) - h)^-z$ is compact. In fact, let $\{z_n\}$ be bounded in $\mathfrak{D}(A)$, without loss of generality, we may assume $z_n \rightarrow 0$ in $\mathfrak{D}(A)$. Next we show that $|\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-z_n|_2 \rightarrow 0$ in L^2 . For every $R > 0$, define $B_R(0) = \{t \in \mathbb{R} : |t| < R\}$ and $B_R^c = \mathbb{R} \setminus B_R(0)$. Let $\{t_i\}$ be a sequence of points in B_R^c satisfying $B_R^c \subset \cup_{i=1}^\infty B(t_i, r_0)$ and such that each point t is contained in at most 2 such balls $B(t_i, r_0)$. Let $B := \{t \in B_R^c : \mathcal{J}_0B(t) < h\}$, choose $s \in (1, 3)$ and $s' = s/(s - 1)$. We get

$$\begin{aligned}
 \int_{B_R^c} |\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-z_n|^2 &\leq \sum_{i=1}^\infty \int_{B(t_i, r_0) \cap B} |\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-z_n|^2 \\
 &\leq \sum_{i=1}^\infty \left(\int_{B(t_i, r_0) \cap B} |z_n|^{2s} \right)^{1/s} \left(\int_{B(t_i, r_0) \cap B} |\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-|^{2s'} \right)^{1/s'} \\
 &\leq C_R^{2s'} \sum_{i=1}^\infty |B(t_i, r_0) \cap B|^{1/s'} \left(\int_{B(t_i, r_0) \cap B} |z_n|^{2s} \right)^{1/s} \\
 &\leq 2C_R^{2s'} \varepsilon_R \|z_n\|_A^2,
 \end{aligned} \tag{2.8}$$

where $\varepsilon_R = \sup_i |B(t_i, r_0) \cap B|^{1/s'}$, $C_R = \sup_i \|\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-\|_M$ (for real matrix valued function $Q(x) = (q_{jk}(x))_{2N \times 2N}$, $\|Q(x)\|_M = \max_{1 \leq j \leq 2N} \sum_{k=1}^{2N} |q_{jk}(x)|$).

Here, we used (2.4) of Lemma 2.1 and that H^1 embeds into L^{2s} . Assumption (B₀) implies that $\varepsilon_R \rightarrow 0$ as $R \rightarrow \infty$; thus,

$$\int_{B_R^c} |\mathcal{J}_0(\mathcal{J}_0B(t) - h)^-z_n|^2 \rightarrow 0. \tag{2.9}$$

On the other hand,

$$\int_{B_R(0)} |\mathcal{J}_0(\mathcal{J}_0 B(t) - h)^- z_n|^2 \leq \left(\int_{B_R(0)} |z_n|^{2s} \right)^{1/s} \left(\int_{B_R(0)} |\mathcal{J}_0(\mathcal{J}_0 B(t) - h)^-|^{2s'} \right)^{1/s'} \rightarrow 0, \quad (2.10)$$

since $H^1 \hookrightarrow L^2_{\text{loc}}$ is compact. Thus, $|\mathcal{J}_0(\mathcal{J}_0 B(t) - h)^- z_n|_2 \rightarrow 0$, we have

$$\begin{aligned} o(1) &= |(A - \nu)z_n|_2 = |A_2 z_n - \nu z_n + \mathcal{J}_0(\mathcal{J}_0 B(t) - h)^- z_n|_2 \\ &\geq |A_2 z_n|_2 - |\nu| - o(1) \\ &\geq h - |\nu| - o(1), \end{aligned} \quad (2.11)$$

which implies that $0 < h - |\nu| \leq 0$, a contradiction. So $\sigma_e(A) \cap (-h, h) = \emptyset$. Since $h > 0$ is arbitrary, it follows that $\sigma(A) = \sigma_d(A)$. \square

From Lemma 2.2, we know that the operator A has a sequence of eigenvalues

$$\cdots \lambda_{-k} \leq \cdots \leq \lambda_{-1} \leq 0 < \lambda_1 \leq \cdots \leq \lambda_k \cdots \quad (2.12)$$

with $\lambda_{\pm k} \rightarrow \pm\infty$ as $k \rightarrow \infty$, and corresponding eigenfunctions $\{e_{\pm k}\}_{k \in \mathbb{N}}$ form an orthogonal basis in L^2 . Observe that we have an orthogonal decomposition

$$L^2 = L^- \oplus L^0 \oplus L^+, \quad z = z^- + z^0 + z^+, \quad (2.13)$$

such that A is negative definite on L^- and positive definite on L^+ and $L^0 = \ker A$. Let $P^0 : L^2 \rightarrow L^0$ be the projection. Set $E := \mathfrak{D}(|A|^{1/2})$ be the domain of the self-adjoint operator $|A|^{1/2}$ which is a Hilbert space equipped with the inner product

$$(z, w) = \left(|A|^{1/2} z, |A|^{1/2} w \right)_2 + \left(P^0 z, P^0 w \right)_2 \quad (2.14)$$

and norm $\|z\| = (z, z)^{1/2}$. Let $E^\pm := \overline{\text{span}\{e_{\pm k}\}_{k \in \mathbb{N}}}$, $E^0 = \ker A$. Then, $E = E^- \oplus E^0 \oplus E^+$ is an orthogonal decomposition of E .

Similar to [19], on $\mathfrak{D}(A)$, we introduce an inner product

$$\langle z, w \rangle_A = (Az, Aw)_2 + \left(P^0 z, P^0 w \right)_2 = (|A|z, |A|w)_2 + \left(P^0 z, w \right)_2 \quad (2.15)$$

whose induced norm will be denoted by $|z|_A$. Since 0 is at most an isolated eigenvalue of finite multiplicity, it is clear that $|\cdot|_A$ and $\|\cdot\|_A$ are equivalent on $\mathfrak{D}(A)$: $d_1 |z|_A \leq \|u\|_A \leq d_2 |z|_A$, for all $z \in \mathfrak{D}(A)$. Define

$$\widehat{A} := |A| + P^0. \quad (2.16)$$

Then, $\mathfrak{D}(A) = \mathfrak{D}(\hat{A})$. Noting that $P^0|A| = |A|P^0 = 0$, for $z, w \in \mathfrak{D}(\hat{A})$, we have

$$\begin{aligned} (\hat{A}z, \hat{A}w)_2 &= (|A|z, |A|w)_2 + (P^0z, |A|w)_2 + (|A|z, P^0w)_2 + (P^0z, P^0w)_2 \\ &= (|A|z, |A|w)_2 + (P^0z, P^0w)_2 = \langle z, w \rangle_A, \end{aligned} \quad (2.17)$$

hence

$$|z|_A = |\hat{A}z|_2, \quad \forall z \in \mathfrak{D}(A). \quad (2.18)$$

Observe that, for all $z \in \mathfrak{D}(A)$ and $w \in \mathfrak{D}(|A|^{1/2})$,

$$\begin{aligned} (\hat{A}^{1/2}z, \hat{A}^{1/2}w)_2 &= (\hat{A}z, w)_2 = \left((|A| + P^0)z, w \right)_2 = (|A|z, w)_2 + (P^0z, w)_2 \\ &= (|A|^{1/2}z, |A|^{1/2}w)_2 + (P^0z, P^0w)_2 = (z, w). \end{aligned} \quad (2.19)$$

Consequently, since $\mathfrak{D}(A) = \mathfrak{D}(\hat{A})$ is a core of $\hat{A}^{1/2}$, we have

$$(z, w) = (\hat{A}^{1/2}z, \hat{A}^{1/2}w)_2 \quad \forall z, w \in \mathfrak{D}(|A|^{1/2}), \quad (2.20)$$

which implies in particular that

$$\|z\| = \left| \hat{A}^{1/2}z \right|_2 \quad \forall z \in E. \quad (2.21)$$

By complex interpolation theory, we have $H^{1/2} = [L^2, H^1]_{1/2}$ (see Theorem 2.4.1 [20]). Since $\mathfrak{D}(|A_0|^0) = L^2$ and $\|z\|_{H^1} = \|A_0|z|_2$, one has

$$H^{1/2} = \left[\mathfrak{D}(|A_0|^0), \mathfrak{D}(|A_0|) \right]_{1/2} \quad (2.22)$$

with equivalent norms. It then follows from Theorem 1.18.10 of [20] that

$$H^{1/2} = \left[\mathfrak{D}(|A_0|^0), \mathfrak{D}(|A_0|) \right]_{1/2} = \mathfrak{D}(|A_0|^{1/2}), \quad (2.23)$$

hence $\|z\|_{H^{1/2}}$ and $\|A_0|^{1/2}z|_2$ are equivalent norms on $H^{1/2}$,

$$d_3 \|z\|_{H^{1/2}} \leq \left| A_0|^{1/2}z \right|_2 \leq d_4 \|z\|_{H^{1/2}} \quad \forall z \in H^{1/2}(\mathbb{R}, \mathbb{R}^{2N}). \quad (2.24)$$

Lemma 2.3. *E embeds continuously into $H^{1/2}(\mathbb{R}, \mathbb{R}^{2N})$. Moreover, E embeds compactly into $L^p(\mathbb{R}, \mathbb{R}^{2N})$ for all $p \in [2, \infty)$.*

Proof. By (2.4) and (2.18), there exists $d_5 > 0$ such that

$$\|A_0|z|_2 \leq d_5 \left| \widehat{A}z \right|_2 = \left| \left(d_5 \widehat{A} \right) z \right|_2 \quad \forall z \in \mathfrak{D}(A). \quad (2.25)$$

Thus, $(|A_0|z, z)_2 \leq (d_5 \widehat{A}z, z)_2$ for all $u \in \mathfrak{D}(A)$ (see Proposition III 8.11 of [21]). This implies

$$\left| |A_0|^{1/2} z \right|_2^2 = (|A_0|z, z)_2 \leq (d_5 \widehat{A}z, z)_2 = d_5 \left| \widehat{A}^{1/2} z \right|_2^2 \quad (2.26)$$

for all $z \in \mathfrak{D}(A)$ (see Proposition III 8.12 of [21]). Since $\mathfrak{D}(A)$ is core of $\widehat{A}^{1/2}$, we obtain that $\left| |A_0|^{1/2} z \right|_2^2 \leq d_5 \left| \widehat{A}^{1/2} z \right|_2^2$ for all $z \in E$. This combination with (2.21) shows that

$$\left| |A_0|^{1/2} z \right|_2^2 \leq d_5 \|z\|^2 \quad \forall z \in E, \quad (2.27)$$

which, together with (2.24), implies that

$$\|z\|_{H^{1/2}} \leq d_6 \|z\| \quad \forall z \in E. \quad (2.28)$$

This proves that the embedding $E \hookrightarrow H^{1/2}(\mathbb{R}, \mathbb{R}^{2N})$ is continuous. In order to show that the embedding $E \hookrightarrow L^p(\mathbb{R}, \mathbb{R}^{2N})$ is compact for all $p \in [2, \infty)$, it suffices to prove that $E \hookrightarrow L^2$ is compact. Set $L_k := \text{span}\{e_{-k}, \dots, e_{-1}, e_1, \dots, e_k\}$. Let $P_k : E \rightarrow L_k$ denote the orthogonal projector. Consider a weakly converging sequence $z_n \rightharpoonup z$ in E . Denote $w_n = z_n - z$ and $K := \sup_n \|w_n\|^2$. Given $\varepsilon > 0$, we choose $k \in \mathbb{N}$ so that $K/\nu_k < \varepsilon/2$, where $\nu_k := |\lambda_{-k}| + \lambda_k$. Since $P_k w_n \rightarrow 0$ as $n \rightarrow \infty$, then there exists $n_0 \in \mathbb{N}$ such that $\|P_k w_n\|^2 < \varepsilon/2$ for all $n \geq n_0$. Let $\{E(\tau)\}_{\tau \in \mathbb{R}}$ be the spectral family of A . It follows from

$$\begin{aligned} \|w_n\|^2 &\geq \|(I - P_k)w_n\|^2 = \left(|A|^{1/2}(I - P_k)w_n, |A|^{1/2}(I - P_k)w_n \right)_2 \\ &= \int_{-\infty}^{\lambda_{-k}} |\tau| d|E(\tau)(I - P_k)w_n|_2^2 + \int_{\lambda_k}^{\infty} \tau d|E(\tau)(I - P_k)w_n|_2^2 \\ &\geq (|\lambda_{-k}| + \lambda_k) |(I - P_k)w_n|_2^2 \end{aligned} \quad (2.29)$$

that

$$|(I - P_k)w_n|_2^2 \leq \frac{\|w_n\|^2}{|\lambda_{-k}| + \lambda_k} = \frac{\|w_n\|^2}{\nu_k} < \frac{\varepsilon}{2}. \quad (2.30)$$

Then,

$$|w_n|_2^2 = |P_k w_n|_2^2 + |(I - P_k)w_n|_2^2 < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \quad (2.31)$$

for all $n \geq n_0$. This proves that $z_n \rightarrow z$ in L^2 . \square

Set $\Psi(z) = \int_{\mathbb{R}} R(t, z(t)) dt$, by assumptions and Lemma 2.3, $\Psi(z) \in C^1(E, \mathbb{R})$ and $\Psi'(z)w = \int_{\mathbb{R}} R_z(t, z(t))w(t) dt$, for all $z, w \in E$. Now, let us consider the function

$$\Phi(z) = \frac{1}{2} \left(\|z^+\|^2 - \|z^-\|^2 \right) - \Psi(z) \quad (2.32)$$

for $z = z^+ + z^0 + z^- \in E$, then $\Phi \in C^1(E, \mathbb{R})$. Moreover, for $\Psi \in C_0^\infty(\mathbb{R})$,

$$\Phi'(z)\Psi = \int_{\mathbb{R}} (-\mathcal{J}\dot{z}(t) + B(t)z(t) - R_z(t, z(t)), \Psi(t)) dt. \quad (2.33)$$

It follows that critical points of $\Phi(z)$ are solutions of (HS). Moreover, if z is a solution of (HS), by Lemma 2.3, $R_z(t, z) \in L^s(\mathbb{R}, \mathbb{R}^{2N})$ for any $s \in [2, \infty)$. Thus, $R_z(t, z) \in L^2$. A standard argument shows that z is also a homoclinic orbit of (HS) (see [12]). So if $z \neq 0$ is a solution of (HS), then z is a homoclinic orbit of (HS).

Now, we discuss the linking structure of Φ .

Lemma 2.4. *Suppose (R_1) , (R_2) are satisfied. Then, there is a $\rho > 0$ such that $\kappa := \inf \Phi(\partial B_\rho \cap E^+) > 0$.*

Proof. Observe that, given $\varepsilon > 0$, there is $C_\varepsilon > 0$ such that

$$|R_z(t, z)| \leq \varepsilon|z| + C_\varepsilon|z|^{p-1}, \quad (2.34)$$

$$|R(t, z)| \leq \varepsilon|z|^2 + C_\varepsilon|z|^p, \quad (2.35)$$

for all (t, z) , where $p \in [2, \infty)$. For $z \in E^+$, by Lemma 2.3 and (2.35), we have

$$\begin{aligned} \Phi(z) &= \frac{1}{2} \|z\|^2 - \int_{\mathbb{R}} R(t, z) \\ &\geq \frac{1}{2} \|z\|^2 - C \left(\varepsilon \|z\|^2 + C_\varepsilon \|z\|^p \right). \end{aligned} \quad (2.36)$$

Choosing an appropriate ε , we see that the desired conclusion holds for some $\rho > 0$. \square

For the asymptotically quadratic case, let $b := \inf R_\infty(t)$, and we arrange all the eigenvalues (counted with multiplicity) of A in $(0, b)$ by $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k < b$ and let e_j denote the corresponding eigenfunctions $Ae_j = \lambda_j e_j$ for $j = 1, 2, \dots, k$. Set $Y_0 := \text{span}\{e_1, e_2, \dots, e_k\}$. Note that

$$\lambda_1 |z|_2^2 \leq \|z\|^2 \leq \lambda_k |z|_2^2 \quad \forall z \in Y_0. \quad (2.37)$$

For any subspace W of Y_0 , set $E_W = E^- \oplus E^0 \oplus W$.

Lemma 2.5. *Let (R_1) – (R_3) be satisfied and $\rho > 0$ given by Lemma 2.4. Then, for any subspace W of Y_0 , $\sup \Phi(E_W) < \infty$, and there is a $R_W > 0$ such that $\sup \Phi(E_W \setminus \bar{B}_{R_W}) < \inf \Phi(B_\rho \cap E^+)$.*

Proof. It is sufficient to prove that $\Phi(z) \rightarrow -\infty$ in E_W as $\|z\| \rightarrow \infty$. If not, then there are $M > 0$ and $\{z_j\} \subset E_W$ with $\|z_j\| \rightarrow \infty$ such that $\Phi(z_j) \geq -M$ for all j . Denote $v_j := z_j/\|z_j\|$, passing to a subsequence if necessary, $v_j \rightharpoonup v$, $v_j^- \rightharpoonup v^-$, $v_j^0 \rightarrow v^0$, and $v_j^+ \rightarrow v^+$. Now, we have

$$\frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - \int_{\mathbb{R}} \frac{R(t, z_j)}{\|z_j\|^2} = \frac{\Phi(z_j)}{\|z_j\|^2} \geq \frac{-M}{\|z_j\|^2}, \quad (2.38)$$

since $R(t, z) \geq 0$. If $v^+ = 0$ and $v^0 = 0$, it follows from (2.38) that

$$\frac{1}{2} \|v_j^-\|^2 + \int_{\mathbb{R}} \frac{R(t, z_j)}{\|z_j\|^2} \leq \frac{1}{2} \|v_j^+\|^2 + \frac{M}{\|z_j\|^2} \rightarrow 0 \quad (2.39)$$

as $j \rightarrow \infty$. Thus, we have $\|v_j^-\| \rightarrow 0$ and $\int_{\mathbb{R}} (R(t, z_j)/\|z_j\|^2) \rightarrow 0$, this is a contradiction with $\|v_j\| = 1$. Therefore, there are three possibilities: (i) $v^+ \neq 0$ and $v^0 = 0$, (ii) $v^+ = 0$ and $v^0 \neq 0$, (iii) $v^+ \neq 0$ and $v^0 \neq 0$.

If (i) holds, by (2.37), one has

$$\begin{aligned} \|v^+\|^2 - \|v^-\|^2 - \int_{\mathbb{R}} R_{\infty}(t)v^2 &\leq \|v^+\|^2 - \|v^-\|^2 - b|v|_2^2 \\ &\leq -(b - \lambda_k)|v^+|_2^2 - \|v^-\|^2 - b|v^-|_2^{2-b|v^0|_2^2} - b|v^-|_2^2 < 0. \end{aligned} \quad (2.40)$$

If (ii) or (iii) holds, similar to (i), one has

$$\|v^+\|^2 - \|v^-\|^2 - \int_{\mathbb{R}} R_{\infty}(t)v^2 < 0. \quad (2.41)$$

Then, there exists $a > 0$ such that

$$\|v^+\|^2 - \|v^-\|^2 - \int_{-a}^a R_{\infty}(t)v^2 < 0. \quad (2.42)$$

Letting $\tilde{R}(t, z) := R(t, z) - (1/2)R_{\infty}(t)z^2$, then $|\tilde{R}(t, z)| \leq cz^2$ for some $c > 0$ and $\tilde{R}(t, z)/z^2 \rightarrow 0$ as $|z| \rightarrow \infty$ uniformly in t . Hence, by Lebesgues dominated convergence theorem, we have

$$\lim_{j \rightarrow \infty} \int_{-a}^a \frac{\tilde{R}(t, z_j)}{\|z_j\|^2} = \lim_{j \rightarrow \infty} \int_{-a}^a \frac{\tilde{R}(t, z_j)}{|z_j|^2} |v_j|^2 = 0. \quad (2.43)$$

Thus, (2.38)–(2.43) imply that

$$\begin{aligned} 0 &\leq \lim_{j \rightarrow \infty} \left(\frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - \int_{-a}^a \frac{R(t, z_j)}{\|z_j\|^2} \right) \\ &\leq \frac{1}{2} \left(\|v^+\|^2 - \|v^-\|^2 - \int_{-a}^a R_\infty(x) v^2 \right) < 0. \end{aligned} \tag{2.44}$$

Now, the desired conclusion is obtained from this contradiction. \square

For the superquadratic case, we define $Y_n := \text{span}\{e_1, \dots, e_n\}$ and $E_n = E^- \oplus E^0 \oplus Y_n$, $n \in \mathbb{N}$. Then, Y_n is a finite dimensional subspace of E^+ and

$$\lambda_1 |z|_2^2 \leq \|z\|^2 \leq \lambda_n |z|_2^2 \quad \forall z \in Y_n. \tag{2.45}$$

By (R_4) , there exists $R > 0$ and $\alpha > 0$ such that

$$R(t, z) \geq \alpha |z|^2 \geq \lambda_n |z|_2^2 \quad \forall |z| \geq R. \tag{2.46}$$

Lemma 2.6. *Let (R_1) , (R_4) – (R_5) be satisfied and $\rho > 0$ given by Lemma 2.4. Then, there is a sequence $\{R_n\}$ with $R_n > \rho$ such that $\sup \Phi(E_n) < \infty$ and $\sup \Phi(E_n \setminus \bar{B}_n) < \inf \Phi(B_\rho \cap E^+)$, where $\bar{B}_n = \{z \in E_n : \|z\| \leq R_n\}$.*

Proof. Similar to proof of Lemma 2.5. If (i) holds, by (2.45) and (2.46), it is easy to prove that

$$\begin{aligned} \|v^+\|^2 - \|v^-\|^2 - \alpha \int_{\mathbb{R}} |v|^2 &\leq \lambda_n \|v^+\|^2 - \|v^-\|^2 - \alpha |v|_2^2 \\ &= -(\alpha - \lambda_n) |v^+|_2^2 - \|v^-\|^2 - \alpha |v^-|_2^2 - \alpha |v^0|_2^2 < 0. \end{aligned} \tag{2.47}$$

If (ii) or (iii) holds, similar to (i), one has

$$\|v^+\|^2 - \|v^-\|^2 - \alpha \int_{\mathbb{R}} |v|^2 < 0. \tag{2.48}$$

Hence, there exists $a > 0$ such that

$$\|v^+\|^2 - \|v^-\|^2 - \alpha \int_{-a}^a |v|^2 < 0. \tag{2.49}$$

Note that

$$\begin{aligned}
\frac{\Phi(z_j)}{\|z_j\|^2} &\leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - \alpha \int_{-a}^a \frac{R(t, z_j)}{\|z_j\|^2} \\
&= \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - \alpha \int_{-a}^a |v_j|^2 \right) - \int_{-a}^a \frac{R(t, z_j) - (\alpha/2)|z_j|^2}{\|z_j\|^2} \\
&\leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - \alpha \int_{-a}^a |v_j|^2 \right) + \int_{(-a, a) \cap \{t \in \mathbb{R} : |z_j| \leq R\}} \frac{(\alpha/2)|z_j|^2 - R(t, z_j)}{\|z_j\|^2} \\
&\leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - \alpha \int_{-a}^a |v_j|^2 \right) + \frac{2\alpha C_R}{\|z_j\|^2},
\end{aligned} \tag{2.50}$$

where $C_R = \sup\{R(t, z) : t \in (-a, a), |z| \leq R\}$. Thus, (2.38) and (2.49) imply that

$$\begin{aligned}
0 &\leq \lim_{j \rightarrow \infty} \left(\frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - \int_{-a}^a \frac{R(t, z_j)}{\|z_j\|^2} \right) \\
&\leq \frac{1}{2} \left(\|v^+\|^2 - \|v^-\|^2 - \alpha \int_{\Omega} |v|^2 \right) < 0,
\end{aligned} \tag{2.51}$$

a contradiction. This proves the lemma. \square

As a consequence, we have the following.

Lemma 2.7. *Under the assumptions of Theorem 1.1 (Theorem 1.2), then letting $e \in Y_0(Y_n)$ with $\|e\| = 1$, there is $R_1 > \rho > 0$, such that $\Phi|_{\partial Q} \leq \kappa$, where $\kappa > 0$ be given by Lemma 2.4, $Q := \{u = u^- + u^0 + se : u^- + u^0 \in E^- \oplus E^0, s \geq 0, \|u\| \leq R_1\}$.*

3. The $(C)_c$ Condition

In this section, we discuss the properties of the $(C)_c$ sequences. Recall that a sequence $\{z_n\} \subset E$ is said to be a $(C)_c$ sequence if $\Phi(z_n) \rightarrow c$ and $(1 + \|z_n\|)\Phi'(z_n) \rightarrow 0$, Φ is said to satisfy the $(C)_c$ condition if any $(C)_c$ sequence has a convergent subsequence.

Lemma 3.1. *Under the assumptions of Theorem 1.1 or Theorem 1.2, then any $(C)_c$ -sequence is bounded.*

Proof. Let $\{z_j\} \subset E$ be such that

$$\Phi(z_j) \rightarrow c, \quad (1 + \|z_j\|)\Phi'(z_j) \rightarrow 0. \tag{3.1}$$

Then, there is constant $C > 0$ such that we have

$$\Phi(z_j) - \frac{1}{2}\Phi'(z_j)z_j = \int_{\mathbb{R}} \widehat{R}(t, z_j) \leq C. \tag{3.2}$$

Suppose to the contrary that $\{z_j\}$ is unbounded. Setting $v_j := z_j/\|z_j\|$, then $\|v_j\| = 1$, $v_j = v_j^+ + v_j^0 + v_j^- \in E^+ \oplus E^0 \oplus E^-$. After passing a subsequence, we have $v_j \rightarrow v$, $v_j^0 \rightarrow v^0$, and $\gamma = \lim_{j \rightarrow \infty} \|v_j^+ + v_j^-\|^2$ exists. Moreover, $|v_j|_s \leq C_s$ for all $s \in [2, \infty)$. For γ , we have only the following two cases: $\gamma = 0$ or $\gamma > 0$.

First, we consider $\gamma = 0$, then $\|v_j^0\| = |v_j^0|_2 \rightarrow 1 = |v^0|_2$. Denote $\tilde{z}_j = z_j^+ + z_j^-$, $\tilde{v}_j = v_j^+ + v_j^-$. Note that by (R_3) , for any $\ell > 0$, there exists a $h > 0$ such that

$$\ell \leq \widehat{R}(t, z), \quad \forall |z| \geq h. \tag{3.3}$$

For $\delta > 0$, set $\Omega_\delta := \{t \in \mathbb{R} : |v^0(t)| \geq 2\delta\}$ and $\Omega_{j\delta} := \{t \in \mathbb{R} : |\tilde{v}_j(t)| \geq \delta\}$. Since $v^0 \in C^1(\mathbb{R})$ and $|v^0|_2 = 1$, $|\Omega_\delta| > 0$ for all δ small. Moreover, we have

$$|\Omega_{j\delta}| \leq \frac{1}{\delta^2} \int_{\Omega_{j\delta}} |\tilde{v}_j|^2 \leq \frac{c}{\delta^2} \|\tilde{v}_j\|^2 = \frac{c}{\delta^2} \|v_j^+ + v_j^-\|^2 \rightarrow 0 \quad \text{as } j \rightarrow \infty. \tag{3.4}$$

Hence, $|\Omega_\delta \setminus \Omega_{j\delta}| \rightarrow |\Omega_\delta|$ as $j \rightarrow \infty$. Therefore, there exists $N > 0$ such that $|v_j(t)| \geq \delta/2$ for all $t \in \Omega_\delta \setminus \Omega_{j\delta}$ with $j \geq N$; thus, $|z_j(t)| \geq (\delta/2)\|z_j\| \geq h$ for $j \geq N$. From this and (3.3), we get

$$\int_{\mathbb{R}} \widehat{R}(t, z_j) \geq \int_{\Omega_\delta \setminus \Omega_{j\delta}} \ell = \ell |\Omega_\delta \setminus \Omega_{j\delta}|, \tag{3.5}$$

contradicting (3.2) since ℓ is arbitrary.

Hence, $\gamma > 0$, observe that

$$\Phi'(z_j)(z_j^+ - z_j^-) = \|z_j\|^2 \left(\|\tilde{v}_j\|^2 - \int_{\mathbb{R}} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \right). \tag{3.6}$$

Hence,

$$\int_{\mathbb{R}} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \rightarrow \gamma^2. \tag{3.7}$$

Set

$$g(r) := \inf \left\{ \widehat{R}(t, z) \mid t \in \mathbb{R}, z \in \mathbb{R}^{2N} \text{ with } |z| \geq r \right\} \tag{3.8}$$

for $r \geq 0$. By (R_3) , $g(r) > 0$ for all $r > 0$ and $g(r) \rightarrow \infty$ as $r \rightarrow \infty$. For $0 \leq a < b$, let

$$\begin{aligned} \Omega_{j(a,b)} &:= \{t \in \mathbb{R} \mid a \leq |z_j(t)| < b\} \\ C_a^b &:= \inf \left\{ \frac{\widehat{R}(t, z)}{|z|^2} \mid t \in \mathbb{R}, z \in \mathbb{R}^{2N} \text{ with } a \leq |z(x)| < b \right\}. \end{aligned} \tag{3.9}$$

By (3.2), it is easy to prove that

$$|\Omega_{j(b,\infty)}| \leq \frac{C}{g(b)} \rightarrow 0 \quad (3.10)$$

as $b \rightarrow \infty$ uniformly in t , and, for any fixed $0 < a < b$,

$$\int_{\Omega_{j(a,b)}} |v_j|^2 = \frac{1}{\|z_j\|^2} \int_{\Omega_{j(a,b)}} |z_j|^2 \leq \frac{C}{C_a^b \|z_j\|^2} \rightarrow 0 \quad (3.11)$$

as $j \rightarrow \infty$, and

$$\int_{\Omega_{j(b,\infty)}} |v_j|^s \leq \left(\int_{\Omega_{j(b,\infty)}} |v_j|^{2s} \right)^{1/2} |\Omega_{j(b,\infty)}|^{1/2} \rightarrow 0 \quad (3.12)$$

for any $s \in [2, \infty)$, as $b \rightarrow \infty$ uniformly in j . Let $0 < \varepsilon < \gamma^2/3$. By (R_1) , there is a $a_\varepsilon > 0$ such that

$$|R_z(t, z)| < \varepsilon|z|, \quad \forall |z| \leq a_\varepsilon. \quad (3.13)$$

Consequently,

$$\int_{\Omega_{j(0,a_\varepsilon)}} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \leq \int_{\Omega_{j(0,a_\varepsilon)}} \varepsilon |v_j^+ - v_j^-| |v_j| \leq \frac{\varepsilon}{C_2} |v_j|_2^2 < \varepsilon \quad (3.14)$$

for all j .

By (R_2) , there is some $c > 0$ such that

$$|R_z(t, z)| < c|z|, \quad \forall (t, z). \quad (3.15)$$

By (3.11) and Hölder inequality, setting $\mu = 2\nu/(\nu - 1) > 2$ and $\nu' = \mu/2 = \nu/(\nu - 1)$, we can take large b_ε such that

$$\begin{aligned} \int_{\Omega_{j(b_\varepsilon,\infty)}} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} &\leq \left(\int_{\Omega_{j(b_\varepsilon,\infty)}} \frac{|R_z(t, z_j)|^\nu}{|z_j|^\nu} \right)^{1/\nu} \left(\int_{\Omega_{j(b_\varepsilon,\infty)}} (|v_j^+ - v_j^-| |v_j|)^{\nu'} \right)^{1/\nu'} \\ &\leq c |\Omega_{j(b_\varepsilon,\infty)}|^{1/\nu} \left(\int_{\Omega_{j(b_\varepsilon,\infty)}} |v_j^+ - v_j^-|^\mu \right)^{1/\mu} \left(\int_{\Omega_{j(b_\varepsilon,\infty)}} |v_j|^\mu \right)^{1/\mu} \leq \varepsilon \end{aligned} \quad (3.16)$$

for all j . By (3.11), there is j_0 such that

$$\int_{\Omega_j(a_\varepsilon, b_\varepsilon)} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \leq c \int_{\Omega_j(a_\varepsilon, b_\varepsilon)} |v_j^+ - v_j^-||v_j| \leq c|v_j|_2 \left(\int_{\Omega_j(a_\varepsilon, b_\varepsilon)} |v_j|^2 \right)^{1/2} \leq \varepsilon \tag{3.17}$$

for all $j \geq j_0$. By (3.14)–(3.17), we have

$$\limsup_{j \rightarrow \infty} \int_{\mathbb{R}} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \leq 3\varepsilon < \gamma^2 \tag{3.18}$$

which contradicts (3.7). □

Next, we deal with the superquadratic case. Note that $2\mu' = 2\mu/(\mu - 1) < \infty$, by (3.12), we can take large $b_\varepsilon > r$ such that

$$\begin{aligned} & \int_{\Omega_j(b_\varepsilon, \infty)} \frac{R_z(t, z_j)(v_j^+ - v_j^-)|v_j|}{|z_j|} \\ & \leq \left(\int_{\Omega_j(b_\varepsilon, \infty)} \left(\frac{|R_z(t, z_j)|}{|z_j|} \right)^\mu \right)^{1/\mu} \left(\int_{\Omega_j(b_\varepsilon, \infty)} (|v_j^+ - v_j^-||v_j|)^{\mu'} \right)^{1/\mu'} \\ & \leq \left(\int_{\mathbb{R}} c_1 \widehat{R}(t, z_j) \right)^{1/\mu} \left(\int_{\mathbb{R}} |v_j^+ - v_j^-|^{2\mu'} \right)^{1/2\mu'} \left(\int_{\Omega_j(b_\varepsilon, \infty)} |v_j|^{2\mu'} \right)^{1/2\mu'} < \varepsilon \end{aligned} \tag{3.19}$$

for all j . Clearly, (3.17) remains true. Now, the combination of (3.14), (3.17) and (3.19) implies again a contradiction with (3.7).

Let $\{z_n\}$ be an arbitrary $(C)_c$ sequence. By Lemma 3.1, it is bounded; hence, we may assume without loss of generality that $z_n \rightarrow z$ in E , $z_n \rightarrow z$ in L^q for $q \geq 2$. Plainly, z is a critical point of Φ .

Lemma 3.2. *Under the assumptions of Theorem 1.1 or Theorem 1.2, then Φ satisfies $(C)_c$ condition for all $c > 0$.*

Proof. Let $\{z_n\}$ be an arbitrary $(C)_c$ sequence. By (2.35), Lemmas 2.3 and 3.1, it follows from

$$\begin{aligned} \int_{\mathbb{R}} (R_z(t, z_m) - R_z(t, z_n))(z_m^+ - z_n^+) & \leq \left(\int_{\mathbb{R}} (R_z(t, z_m) - R_z(t, z_n))^2 \right)^{1/2} \left(\int_{\mathbb{R}} |z_m^+ - z_n^+|^2 \right)^{1/2} \\ & \leq c \left(\int_{\mathbb{R}} |z_m - z_n|^2 \right)^{1/2} \rightarrow 0 \\ o(1) & = (\Phi'(z_m) - \Phi'(z_n), z_m^+ - z_n^+) \\ & = \|z_m^+ - z_n^+\|^2 + \int_{\mathbb{R}} (R_z(t, z_m) - R_z(t, z_n))(z_m^+ - z_n^+) \\ & = \|z_m^+ - z_n^+\|^2 + o(1) \end{aligned} \tag{3.20}$$

that $z\|z_m^+ - z_n^+\|^2 = o(1)$ as $m, n \rightarrow \infty$. Similarly, we can prove that $z\|z_m^- - z_n^-\|^2 = o(1)$ as $m, n \rightarrow \infty$. Since $\dim E^0 < \infty$, thus $z_n^0 \rightarrow z^0$. So $\{z_n\}$ is a Cauchy sequence in E . Now, the conclusion follows from the completeness of E . \square

4. Proof of Theorems

In this section, we give the proof for our Theorems 1.1 and 1.2. Let E be a Banach space with direct sum $E = X \oplus Y$ and corresponding projections P_X, P_Y onto X, Y . Let $\mathcal{S} \subset X^*$ be a dense subset, for each $s \in \mathcal{S}$, there is a seminorm on E defined by

$$p_s : E \rightarrow \mathbb{R}, \quad p_s(z) : |s(x)| + \|y\| \quad \text{for } z = x + y \in E. \quad (4.1)$$

We denote by $\mathcal{T}_{\mathcal{S}}$ the topology induced by seminorm family $\{p_s\}$, w^* denotes the weak*-topology on E^* . Now, some notations are needed. For a functional $\Phi \in C^1(E, \mathbb{R})$, we write $\Phi_a = \{z \in E \mid \Phi(z) \geq a\}$, $\Phi^b = \{z \in E \mid \Phi(z) \leq b\}$, and $\Phi_a^b = \Phi_a \cap \Phi^b$. Recall that Φ is said to be weakly sequentially lower semicontinuous, if for any $z_j \rightharpoonup z$ in E , one has $\Phi(z) \leq \liminf_{j \rightarrow \infty} \Phi(z_j)$, and Φ' is said to be weakly sequentially continuous if $\lim_{j \rightarrow \infty} \Phi'(z_j)w = \Phi'(z)w$ for each $w \in E$.

Suppose

- (Φ_0) for any $c \in \mathbb{R}$, superlevel Φ_c is $\mathcal{T}_{\mathcal{S}}$ -closed, and $\Phi' : (\Phi_c, \mathcal{T}_{\mathcal{S}}) \rightarrow (E^*, w^*)$ is continuous;
- (Φ_1) for any $c > 0$, there exists $\xi > 0$ such that $\|z\| < \xi \|P_Y z\|$ for all $z \in \Phi_c$;
- (Φ_2) there exists $\rho > 0$ such that $\kappa := \inf \Phi(S_\rho \cap Y) > 0$, where $S_\rho := \{z \in E : \|z\| = \rho\}$;
- (Φ_3) there exists a finite-dimensional subspace $Y_0 \subset Y$ and $R > \rho$ such that we have for $E_0 := X \oplus Y_0$ and $B_0 := \{z \in E_0 : \|z\| \leq R\}$ that $\bar{c} := \sup \Phi(E_0) < \infty$ and $\sup \Phi(E_0 \setminus B_0) < \inf \Phi(B_\rho \cap Y)$;
- (Φ_4) there exists an increasing sequence of finite-dimensional subspaces $Y_n \subset Y$ and a sequence of numbers $R_n > \rho$ such that we have for $E_n := X \oplus Y_n$ and $B_n := \{z \in E_n : \|z\| \leq R_n\}$ that $\bar{c} := \sup \Phi(E_n) < \infty$ and $\sup \Phi(E_n \setminus B_n) < \inf \Phi(B_\rho \cap Y)$;
- (Φ_5) Φ satisfies the $(C)_c$ condition for $c > 0$.

Now, we state three critical point theorems which will be used later (see [22]).

Theorem 4.1. *Let $\Phi_0 - \Phi_2$ be satisfied, and suppose there are $R > \rho > 0$ and $e \in Y$ with $\|e\| = 1$ such that $\sup \Phi(\partial Q) \leq \kappa$ where $Q := \{z = x + te : x \in X, t \geq 0, \|z\| < R\}$. If Φ satisfies the $(C)_c$ condition for all $c \leq \bar{c}$, then Φ has a critical point u with $\kappa \leq \Phi(z) \leq \bar{c}$.*

Theorem 4.2. *Assume Φ is even and $(\Phi_0), (\Phi_2) - (\Phi_3)$ be satisfied. Then, Φ has at least $m := \dim Y_0$ pairs of critical points with critical values less or equal to \bar{c} provided Φ satisfies the $(C)_c$ condition for all $c \in [\kappa, \bar{c}]$.*

Theorem 4.3. *Assume Φ is even with $\Phi(0) = 0$, and $(\Phi_0) - (\Phi_2), (\Phi_4) - (\Phi_5)$ be satisfied. Then, Φ has an unbounded sequence of critical values.*

Lemma 4.4. *Under the assumptions of Theorem 1.1 or Theorem 1.2, then, Ψ is nonnegative, weakly sequentially lower semicontinuous, and Ψ' is weakly sequentially continuous. Moreover, Ψ' is compact.*

In virtue of a standard way of [23], by Lemma 2.3, it is easy to prove this lemma, so we omit it here.

Lemma 4.5. Φ satisfies (Φ_1) .

Proof. For any $c > 0$ and $u \in \Phi_c$, using the fact that $R(t, z) \geq 0$, one has

$$0 < c \leq \frac{1}{2}(\|z^+\| - \|z^-\|) - \int_{\mathbb{R}} R(t, z) \leq \frac{1}{2}\|z^+\|. \quad (4.2)$$

This yields

$$\|z^+\| > \|z^-\|, \quad \|z^+\|^2 \geq 2c. \quad (4.3)$$

Thus it suffices to show that there exists $C > 0$ such that $\|z^0\| \leq C\|z^+\|$. If it is not true, then there is a sequence $\{z_j\} \subset \Phi_c$ such that

$$\|z_j\|^2 \geq \|z_j^0\|^2 \geq j\|z_j^+\|^2. \quad (4.4)$$

By (4.3), $\|z_j^0\| \rightarrow \infty$ as $j \rightarrow \infty$, hence $\|z_j\| \rightarrow \infty$ as $j \rightarrow \infty$. Set $v_j = z_j/\|z_j\|$, then $\|v_j\|^2 = 1$ and $\|v_j^+\|^2 \leq 1/j \rightarrow 0$, hence $\|v_j^-\| \rightarrow 0$ as $j \rightarrow \infty$. Therefore, we can assume there is a subsequence denoted still by $\{v_j\}$, $v_j \rightarrow v$, thus $v_j \rightarrow v^0$ and $\|v^0\| = 1$. Recall that $\tilde{R}(t, z) = R(t, z) - (1/2)R_\infty(t)z^2$, then $\tilde{R}(t, z)/|z|^2 \rightarrow 0$ as $|z| \rightarrow \infty$ uniformly in t . Therefore, since $|z_j(t)| \rightarrow \infty$ for $v^0(t) \neq 0$, we have

$$\begin{aligned} \int_{\mathbb{R}} \frac{\tilde{R}(t, z_j)}{\|z_j\|^2} &\leq \int_{\mathbb{R}} \frac{\tilde{R}(t, z_j)|v_j - v|^2}{|z_j|^2} + \int_{\mathbb{R}} \frac{\tilde{R}(t, z_j)|v|^2}{|z_j|^2} \\ &\leq \int_{v^0(t) \neq 0} \frac{\tilde{R}(t, z_j)|v|^2}{|z_j|^2} + c|v_j - v|_2^2 \rightarrow 0. \end{aligned} \quad (4.5)$$

By (4.4),

$$\|z_j^- + z_j^0\|^2 \geq (j-1)\|z_j^+\|^2 \geq (j-1)\left(2c + \|z_j^-\|^2 + 2 \int_{\mathbb{R}} R(t, z_j)\right) \quad (4.6)$$

or

$$\|z_j^0\|^2 \geq (j-1)2c + (j-2)\|z_j^-\|^2 + 2(j-1) \int_{\mathbb{R}} R(t, z_j), \quad (4.7)$$

which implies

$$\begin{aligned} \frac{1}{2(j-1)} &\geq \int_{\mathbb{R}} \frac{R(t, z_j)}{\|z_j\|^2} = \frac{1}{2} \int_{\mathbb{R}} R_{\infty}(t) |v_j|^2 + \int_{\mathbb{R}} \frac{\tilde{R}(t, z_j) |v_j|^2}{|z_j|^2} \\ &\geq \frac{1}{2} b |v_j|_2^2 + o(1), \end{aligned} \quad (4.8)$$

consequently, $v^0 = 0$, a contradiction.

Next, we handle the superquadratic case. Similar to the above proof, passing to a subsequence if necessary, we have $v_j \rightarrow v^0$ and $\|v^0\| = 1$. So there exists $r > 0$ such that

$$\int_{-r}^r |v^0|^2 > 0. \quad (4.9)$$

By (R_4) , we know that there exist $m > 0$ and C_m such that $R(t, z) \geq C_m |z|^2$ for $|z| \geq m$, and we have

$$\begin{aligned} \frac{c}{\|z_j\|^2} &\leq \frac{\Phi(z_j)}{\|z_j\|^2} \leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - C_m \int_{-r}^r \frac{R(t, z_j)}{\|z_j\|^2} \\ &= \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - C_m \int_{-r}^r |v_j|^2 \right) - \int_{-r}^r \frac{R(t, z_j) - (C_m/2) |z_j|^2}{\|z_j\|^2} \\ &\leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - C_m \int_{-r}^r |v_j|^2 \right) + \int_{(-r, r) \cap \{t \in \mathbb{R} : |z_j| \leq m\}} \frac{(C_m/2) |z_j|^2 - R(t, z_j)}{\|z_j\|^2} \\ &\leq \frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 - C_m \int_{-r}^r |v_j|^2 \right) + \frac{2rC_m}{\|z_j\|^2}. \end{aligned} \quad (4.10)$$

Then,

$$\begin{aligned} 0 &\leq \lim_{j \rightarrow \infty} \left(\frac{1}{2} \left(\|v_j^+\|^2 - \|v_j^-\|^2 \right) - \int_{-r}^r \frac{R(t, z_j)}{\|z_j\|^2} \right) \\ &\leq -C_m \int_{-r}^r |v^0|^2 < 0, \end{aligned} \quad (4.11)$$

a contradiction. □

Proof of Theorem 1.1. Existence of a Least Energy Solution

With $X = E^- \oplus E^0$ and $Y = E^+$, the condition (Φ_0) holds by Lemma 4.4 and (Φ_1) holds by Lemma 4.5. Lemma 2.4 implies (Φ_2) . Lemma 2.7 shows that Φ possesses the linking structure of Theorem 4.1, and Lemma 3.2 implies Φ satisfies (Φ_5) . Therefore, Φ has at least one critical point z with $\Phi(z) \geq \kappa > 0$. Let $\mathcal{K} = \{z \in E \setminus \{0\} : \Phi'(z) = 0\}$ be the set of nontrivial points of Φ .

Then, $\mathcal{K} \neq \emptyset$. We claim that $\theta = \inf\{\Phi(z) \mid z \in \mathcal{K}\}$ is achieved. Let $\{z_j\} \subset \mathcal{K}$ be a minimizing sequence for θ , then $\{z_j\}$ is bounded by Lemma 3.1. By Lemma 3.2, there exists a renamed subsequence such that $z_j \rightarrow z$, and

$$\theta = \lim_{j \rightarrow \infty} \Phi(z_j) = \Phi(z) \quad (4.12)$$

from which it follows that θ is achieved.

Multiplicity

Φ is even provided $R(t, z)$ is even in z . Lemma 2.5 says that Φ satisfies (Φ_3) with $\dim Y_0 = k$. Therefore, Φ has at least k pairs of nontrivial critical points by Theorem 4.2. \square

Proof of Theorem 1.2. Existence of a Least Energy Solution

Repeating the above proof, we know that θ is achieved by some nonzero critical point.

Existence of Infinitely Many Solutions

Φ is even provided $R(t, z)$ is even in z . Lemma 2.6 says that Φ satisfies (Φ_4) . Therefore, Φ has an unbounded critical values by Theorem 4.3, and hence the equation (HS) has infinitely many solutions. \square

Acknowledgments

This work is supported partly by NSFC (11061040), NSFY of Yunnan Province (2008CD112 and 2009CD043), and the Foundation of Education Committee of Yunnan Province, China.

References

- [1] V. Coti-Zelati and P. H. Rabinowitz, "Homoclinic orbits for second order Hamiltonian systems possessing superquadratic potentials," *Journal of the American Mathematical Society*, vol. 4, no. 4, pp. 693–727, 1991.
- [2] Y. H. Ding, "Existence and multiplicity results for homoclinic solutions to a class of Hamiltonian systems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 25, no. 11, pp. 1095–1113, 1995.
- [3] Y. H. Ding and M. Girardi, "Periodic and homoclinic solutions to a class of Hamiltonian systems with the potentials changing sign," *Dynamic Systems and Applications*, vol. 2, no. 1, pp. 131–145, 1993.
- [4] W. Omana and M. Willem, "Homoclinic orbits for a class of Hamiltonian systems," *Differential and Integral Equations*, vol. 5, no. 5, pp. 1115–1120, 1992.
- [5] P. H. Rabinowitz, "Homoclinic orbits for a class of Hamiltonian systems," *Proceedings of the Royal Society of Edinburgh*, vol. 114, no. 1-2, pp. 33–38, 1990.
- [6] V. Coti-Zelati, I. Ekeland, and . Séré, "A variational approach to homoclinic orbits in Hamiltonian systems," *Mathematische Annalen*, vol. 288, no. 1, pp. 133–160, 1990.
- [7] Y. H. Ding, "Multiple homoclinics in a Hamiltonian system with asymptotically or super linear terms," *Communications in Contemporary Mathematics*, vol. 8, no. 4, pp. 453–480, 2006.
- [8] Y. H. Ding and M. Girardi, "Infinitely many homoclinic orbits of a Hamiltonian system with symmetry," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 38, no. 3, pp. 391–415, 1999.
- [9] Y. H. Ding and L. Jeanjean, "Homoclinic orbits for a nonperiodic Hamiltonian system," *Journal of Differential Equations*, vol. 237, no. 2, pp. 473–490, 2007.

- [10] Y. H. Ding and S. J. Li, "Homoclinic orbits for first order Hamiltonian systems," *Journal of Mathematical Analysis and Applications*, vol. 189, no. 2, pp. 585–601, 1995.
- [11] Y. H. Ding and C. Lee, "Existence and exponential decay of homoclinics in a nonperiodic superquadratic Hamiltonian system," *Journal of Differential Equations*, vol. 246, no. 7, pp. 2829–2848, 2009.
- [12] Y. H. Ding and M. Willem, "Homoclinic orbits of a Hamiltonian system," *Journal of Applied Mathematics and Physics*, vol. 50, no. 5, pp. 759–778, 1999.
- [13] H. Hofer and K. Wysocki, "First order elliptic systems and the existence of homoclinic orbits in Hamiltonian systems," *Mathematische Annalen*, vol. 288, no. 3, pp. 483–503, 1990.
- [14] P. H. Rabinowitz and K. Tanaka, "Some results on connecting orbits for a class of Hamiltonian systems," *Mathematische Zeitschrift*, vol. 206, no. 3, pp. 473–499, 1991.
- [15] E. Séré, "Existence of infinitely many homoclinic orbits in Hamiltonian systems," *Mathematische Zeitschrift*, vol. 209, no. 1, pp. 27–42, 1992.
- [16] A. Szulkin and W. Zou, "Homoclinic orbits for asymptotically linear Hamiltonian systems," *Journal of Functional Analysis*, vol. 187, no. 1, pp. 25–41, 2001.
- [17] K. Tanaka, "Homoclinic orbits in a first order superquadratic Hamiltonian system: convergence of subharmonic orbits," *Journal of Differential Equations*, vol. 94, no. 2, pp. 315–339, 1991.
- [18] J. Wang, J. Xu, and F. Zhang, "Homoclinic orbits for superlinear Hamiltonian systems without Ambrosetti-Rabinowitz growth condition," *Discrete and Continuous Dynamical Systems*, vol. 27, no. 3, pp. 1241–1257, 2010.
- [19] Y. H. Ding and B. Ruf, "Solutions of a nonlinear Dirac equation with external fields," *Archive for Rational Mechanics and Analysis*, vol. 190, no. 1, pp. 57–82, 2008.
- [20] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, vol. 18, North-Holland Publishing, Amsterdam, Holland, 1978.
- [21] D. E. Edmunds and W. D. Evans, *Spectral Theory and Differential Operators*, Clarendon Press, Oxford, UK, 1987.
- [22] T. Bartsch and Y. Ding, "Deformation theorems on non-metrizable vector spaces and applications to critical point theory," *Mathematische Nachrichten*, vol. 279, no. 12, pp. 1267–1288, 2006.
- [23] M. Willem, *Minimax Theorems*, Birkhäuser, Berlin, Germany, 1996.