



# Invariant measures and a stability theorem for locally Lipschitz stochastic delay equations<sup>1</sup>

I. Stojkovic and O. van Gaans

*Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands.*

*E-mails: [stojkov@math.leidenuniv.nl](mailto:stojkov@math.leidenuniv.nl); [vangaans@math.leidenuniv.nl](mailto:vangaans@math.leidenuniv.nl)*

Received 29 June 2009; revised 8 October 2010; accepted 23 October 2010

---

**Abstract.** We consider a stochastic delay differential equation with exponentially stable drift and diffusion driven by a general Lévy process. The diffusion coefficient is assumed to be locally Lipschitz and bounded. Under a mild condition on the large jumps of the Lévy process, we show existence of an invariant measure. Main tools in our proof are a variation-of-constants formula and a stability theorem in our context, which are of independent interest.

**Résumé.** Nous considérons une équation différentielle stochastique retardée à dérive exponentiellement stable et conduite par un processus de Lévy général. Le coefficient de la diffusion est seulement supposé satisfaire une condition lipschitzienne locale et être borné. En supposant une condition additionnelle faible sur les grands sauts du processus de Lévy, nous démontrons l'existence d'une mesure invariante. Les principaux ingrédients de la preuve sont une formule pour les variations des constantes et un théorème de stabilité par rapport aux perturbations des conditions initiales, qui sont d'un intérêt indépendant.

MSC: 34K50; 60G48

*Keywords:* Delay equation; Invariant measure; Lévy process; Semimartingale; Skorohod space; Stability; Tightness; Variation-of-constants formula

---

## 1. Introduction

The main purpose of this paper is to show existence of an invariant measure for a stochastic delay differential equation of the form

$$dX(t) = \left( \int_{[-\alpha, 0]} X(t+s) \mu(ds) \right) dt + F(X)(t-) dL(t), \quad (1)$$

where  $L$  is a Lévy process,  $\alpha$  a positive real, and  $\mu$  is a signed Borel measure on  $[-\alpha, 0]$ . The diffusion coefficient  $F$  may be a function on  $\mathbb{R}$  or a functional depending on the segment  $(X(t+s): -\alpha \leq s \leq t)$  of the solution  $X$ . If the underlying deterministic equation, that is Eq. (1) with  $F = 0$ , is exponentially stable, it may be expected that the stochastic equation has an invariant measure under suitable conditions on  $F$ . Existence of invariant measures has been shown for an increasingly general class of coefficients  $F$ . In 1982 Wolfe [25] dealt with the case where  $\mu$  is a (negative) point mass at 0 and  $F$  is a constant. In 2000, Gushchin and Küchler [10] extended to the case of the general delay measure  $\mu$ . More recently, Reiß et al. [21] considered nonlinear coefficients  $F$ . They assume a global Lipschitz condition on  $F$ , boundedness of  $F$ , and continuity of  $F$  with respect to the Skorohod topology. In each of these works

---

<sup>1</sup>Supported by a 'VIDI subsidie' (639.032.510) of the Netherlands Organisation for Scientific Research (NWO).

the analysis depends on a variation-of-constants formula for Eq. (1). In the case of global Lipschitz  $F$ , such a formula has been proved in [22].

The theory of stochastic equations in a setting beyond globally Lipschitz conditions has been vastly extended during the last years in the field of stochastic partial differential equations, see, e.g., [4,5,9,12,18,19] for some recent developments. Also in the field of stochastic delay differential equations there is interest in results on equations with coefficients that are not globally Lipschitz. In particular, models in financial mathematics can naturally involve a combination of delay, processes with jumps, and locally Lipschitz coefficients [23], formula (3.6).

Our main contribution here is extending the results of [21] to equations with diffusion coefficients that are only locally Lipschitz instead of globally Lipschitz. Moreover, the continuity with respect to the Skorohod topology is relaxed to a condition that is considerably better suited for verification in examples. Included in our analysis is the proof of a variation-of-constants formula for (1) for locally Lipschitz  $F$ .

If the diffusion coefficient  $F$  is only locally Lipschitz with linear growth (see Definition 4.1 for the precise formulation), the eventual Feller property and the variation-of-constants formula, which play a key role in [21,22], do not follow from the results given there. Establishing these results is the main content of this article. We do so by approximating the locally Lipschitz diffusion coefficient by globally Lipschitz coefficients in a suitable sense. The difficulty is to verify that the solutions of the equations with the approximated coefficients converge to the solution of (1) in an appropriate sense and that the limit inherits the desired properties. It turns out that the reduction steps in the proof of the variation-of-constants formula in [22] have to be changed. The extension to locally Lipschitz coefficients has to be done before increasing the generality of the other components and these steps have to be adapted accordingly. The proof of the eventual Feller property is based on new estimates, which relax the conditions on  $F$  even in the globally Lipschitz case. Moreover, we prove a stability theorem.

Two comments on the form of (1) are in order. First, in the spirit of [20], Chapter V, we present our results for the one-dimensional equation. At the cost of more complicated notation our arguments can be extended to equations in  $\mathbb{R}^n$ . Second, (1) is formulated with a linear drift term. However, nonlinear drift terms are covered as well by our theory, due to the generality of the noise processes that we allow. By doubling the dimension and including deterministic components in the process  $L$ , locally Lipschitz nonlinearities in the drift are included.

One may wonder how rich is the scope of equations with bounded locally Lipschitz coefficients. On one hand, our generalization is interesting from a theoretical point of view. An important question is what stochastic perturbations that can be added to a stable linear delay differential equation so that the stability persists in the form of existence of an invariant measure. Our results are a natural step in expanding the generality of this theory. On the other hand, natural transformations of equations with unbounded globally Lipschitz coefficients may yield equations with bounded coefficients that are only locally Lipschitz, a situation that fits in our setting (see Example 7.4).

For other approaches to stochastic delay differential equations and invariant measures, see, e.g., [6,15–17].

The outline of our arguments is roughly as follows. If the diffusion coefficient  $F$  in (1) maps the Skorohod space of real valued càdlàg functions on  $[-\alpha, 0]$  into itself and satisfies a suitable locally Lipschitz and growth condition, then it is known that Eq. (1) has a unique solution  $X$  for any initial process on  $[-\alpha, 0]$  (see [13]). The solution itself is, however, not a Markov process. Instead, one can consider the segments  $X_t = (X(t+a))_{-\alpha \leq a \leq 0}$  of the solution process. If a solution  $X(t)$  of Eq. (1) is such that all of the segments  $X_t$  have the same distribution, then the solution itself is stationary as well. Therefore we want to apply the Krylov–Bogoliubov method to the segment process, and for that we need the state space to be separable. If the driving process  $L$  has continuous paths,  $(X_t)_t$  takes values in  $C[-\alpha, 0]$ , which is separable with the supremum norm  $\|\cdot\|_\infty$ . In general,  $L$  may have jumps and then  $(X_t)_t$  is a process with as state space the Skorohod space  $D[-\alpha, 0]$  of càdlàg functions. This space is not separable under  $\|\cdot\|_\infty$ , but it is separable when endowed with the Skorohod metric. In order to apply the Krylov–Bogoliubov method and obtain an invariant measure, we need the eventual Feller property (see (13) and (14) in Section 4) and tightness of the segment process given by (1). We follow the approach of [21], where the tightness is obtained by means of suitable estimates on the semimartingale characteristics.

In Section 2 we give a brief review of the facts about the Skorohod space and deterministic delay equations that we need in the sequel. In Section 3 we obtain the variation-of-constants formula. In Section 4 we give a precise formulation of the input of Eq. (1) and introduce the segment process. Section 5 deals with tightness of the segment process. The stability theorem is proved in Section 6 and the Markov and eventual Feller properties and the existence of an invariant measure are established in Section 7.

## 2. Preliminaries

All the processes we consider are defined on the same filtered probability space  $(\Omega, \mathcal{F}_t, \mathcal{F}, \mathbb{P})$ . Since we are going to work with a Markov process whose state space is the Skorohod space we recall some facts about it. For  $a < b$ , let  $D[a, b]$  and  $D[a, \infty)$  denote the linear spaces of all real-valued càdlàg functions defined on  $[a, b]$  and  $[a, \infty)$ , respectively. Similarly, for  $t_0 > 0$ , let  $\mathbb{D}[0, t_0]$  denote the space of adapted càdlàg processes on  $[0, t_0]$  and likewise  $\mathbb{D}[0, \infty)$ . On  $D[a, \infty)$  the Skorohod metric is given by

$$d_S(\varphi, \psi) = \inf_{\lambda \in \Lambda[a, \infty)} (\|\varphi \circ \lambda - \psi\|_\infty + \|\lambda\|),$$

where  $\Lambda[a, \infty)$  is the set of all increasing  $[a, \infty) \rightarrow [a, \infty)$  homeomorphisms and

$$\|\lambda\| = \sup_{a \leq s < t, s \neq t} \left| \log \frac{\lambda(t) - \lambda(s)}{t - s} \right|.$$

The space  $(D[a, \infty), d_S)$  is complete and separable. Similarly, there is a complete separable metric on  $D[a, b]$ , which we also denote by  $d_S$ .

We will also use a weaker metric on  $D[-\alpha, 0]$ . Consider an arbitrary  $\beta > \alpha$ . We extend a  $\varphi \in D[-\alpha, 0]$  to  $\bar{\varphi} \in D[-\beta, 0]$  by  $\bar{\varphi}(t) = \varphi(t)$  for  $t \in [-\alpha, 0]$  and  $\bar{\varphi}(t) = 0$  for  $t \in [-\beta, -\alpha)$ . Let the metric  $d_\beta$  on  $D[-\alpha, 0]$  be defined by

$$d_\beta(\varphi, \psi) := d_{D[-\beta, 0]}(\bar{\varphi}, \bar{\psi}), \quad \varphi, \psi \in D[-\alpha, 0],$$

where  $d_{D[-\beta, 0]}$  denotes the Skorohod metric on  $D[-\beta, 0]$ . It is straightforward to verify that  $d_\beta(\varphi, \psi) \leq d_S(\varphi, \psi)$  for all  $\varphi, \psi \in D[-\alpha, 0]$ , where  $d_S$  still denotes the Skorohod metric on  $D[-\alpha, 0]$ . The metric  $d_\beta$  is actually independent of  $\beta$  and one could even choose  $\beta = \infty$ .

**Lemma 2.1.** (1)  $(D[-\alpha, 0], d_\beta)$  is a Polish space.

(2)  $d_\beta$  and  $d_S$  generate the same Borel  $\sigma$ -algebra  $\mathcal{B}(D[-\alpha, 0])$ .

**Proof.** (1) The set  $A := \{\bar{\varphi} : \varphi \in D[-\alpha, 0]\}$  is a closed subset of  $D[-\beta, 0]$ , which is a Polish space. Indeed, Skorohod convergence implies almost everywhere convergence and we can finish the argument by right continuity of the limit.

(2) As  $d_\beta \leq d_S$ , the  $d_\beta$ -Borel  $\sigma$ -algebra is contained in the  $d_S$ -Borel  $\sigma$ -algebra. For the opposite inclusion it is enough to show that finite-dimensional sets, that is,  $\{\varphi \in D[-\alpha, 0] : (\varphi(s_1), \dots, \varphi(s_n)) \in C\}$ , where  $C \in \mathcal{B}(\mathbb{R}^n)$  and  $-\alpha \leq s_1 \leq \dots \leq s_n \leq 0$ , are in the  $d_\beta$ -Borel  $\sigma$ -algebra (see [3], formula (15.2) on p. 157). This is obvious as  $D[-\alpha, 0], d_\beta$  is a subspace of  $(D[-\beta, 0], d_{D[-\beta, 0]})$ .  $\square$

Next we collect some results on the deterministic delay equation

$$\begin{aligned} x(t) &= \varphi(0) + \int_0^t \left( \int_{[-\alpha, 0]} x(s+a)\mu(da) \right) ds \quad \text{for } t \geq 0, \\ x(a) &= \varphi(a) \quad \text{for } a \in [-\alpha, 0]. \end{aligned} \tag{2}$$

Here  $\alpha > 0$ ,  $\mu$  is a finite signed Borel measure on  $[-\alpha, 0]$ , and the initial condition  $\varphi \in D[-\alpha, 0]$ . The results that we need can be found in a more general framework in [8]. However, we can give these results in a more easily accessible way as follows. According to [7], Theorem (i), p. 972, for each  $\varphi \in D[-\alpha, 0]$ , there exists a unique function  $x : [-\alpha, \infty) \rightarrow \mathbb{R}$  whose restriction to  $[0, \infty)$  is continuous and which satisfies (2). Indeed, the map  $\psi \mapsto \int_{[-\alpha, 0]} \psi(a)\mu(da)$  is a bounded linear map from  $C[-\alpha, 0]$  with the uniform norm into  $\mathbb{R}$ . Therefore it is also bounded on the Sobolev space  $W^{1,1}[-\alpha, 0]$ , as  $W^{1,1}$  is continuously embedded in  $C_b$ . Since each  $\varphi \in D[-\alpha, 0]$  is bounded and measurable, we have  $(\varphi(0), \varphi) \in M^1 := \mathbb{R} \times L^1[-\alpha, 0]$ . Hence we may apply [7], Theorem (i), to obtain a unique solution  $x$  to (2.6) of [7]. By Fubini theorem  $x$  is the unique solution of (2).

To stress the dependence on the initial condition, we denote the solution of (2) by  $x(\cdot, \varphi)$ . The solution corresponding to initial condition  $\varphi(s) = 0$  for  $-\alpha \leq s < 0$  and  $\varphi(0) = 1$  is called the *fundamental solution* and denoted by  $r$ .

The following variation-of-constants formula for  $x(\cdot, \varphi)$  holds,

$$x(t, \varphi) = r(t)\varphi(0) + \int_0^t r(t-s) \int_{[-\alpha, -s)} \varphi(s+a)\mu(da) ds, \quad t \geq 0, \tag{3}$$

where the inner integral is considered to vanish for  $s > \alpha$ , hence the outer integral is actually from 0 to  $t \wedge \alpha$ . Indeed, by Fubini and substitutions one can verify that the right-hand side of (3) satisfies (2) and therefore equals the unique solution  $x(\cdot, \varphi)$ . By similar arguments one can rewrite formula (3) as

$$x(t, \varphi) = \varphi(0)r(t) + \int_{[-\alpha, 0]} \int_s^0 r(t+s-a)\varphi(a) da \mu(ds) \quad \text{for } t \geq 0. \tag{4}$$

The delay Eq. (2) is said to be *stable* if the fundamental solution  $r$  converges to zero as  $t \rightarrow \infty$ . The condition

$$v_0(\mu) := \sup \left\{ \operatorname{Re} \lambda: \lambda \in \mathbb{C}, \lambda - \int_{[-\alpha, 0]} e^{\lambda s} \mu(ds) = 0 \right\} < 0 \tag{5}$$

implies the even stronger property of exponential stability of all solutions, i.e., there exist  $\gamma, K > 0$  such that  $|x(t, \varphi)| \leq Ke^{-\gamma t} \|\varphi\|_\infty$  for all  $t \geq 0$  and for any solution  $x(\cdot, \varphi)$  of (2). Indeed, for the stability of the fundamental solution see the text below Corollary 4.1 on p. 182 of [11], and then the exponential stability of arbitrary solutions with initial condition  $\varphi \in D[-\alpha, 0]$  follows by direct computation from (3).

It is clear from (2) that each of its solutions  $x(\cdot, \varphi)$  is absolutely continuous on  $[0, T]$  for every  $T > 0$  and even continuously differentiable on  $(\alpha, \infty)$ . If (5) holds, then (2) yields the exponential decay of the derivative  $\dot{x}(\cdot, \varphi)$  directly.

### 3. Variation-of-constants formula

This section establishes a variation-of-constants formula for Eq. (1). The major point is to show existence and uniqueness for equations of variation-of-constants form.

Recall that for two local martingales  $M$  and  $N$  their quadratic covariation process is denoted by  $[M, N]$ . Recall also that for  $t \geq 0$ ,  $\int_0^t |dA(s)|$  ( $\int_0^\infty |dA(s)|$ ) denotes the pathwise total variation on  $[0, t]$  ( $[0, \infty)$ ), respectively) of a process  $A \in D[0, \infty)$ . The next two definitions are taken from [20], Section V.2, pp. 250–251.

**Definition 3.1.** Let  $1 \leq p \leq \infty$ . For a process  $X \in \mathbb{D}[0, \infty)$  set

$$\|X\|_{\underline{S}^p} := \left\| \sup_{t \geq 0} |X(t)| \right\|_{L^p}$$

and let  $\underline{S}^p[0, \infty)$  denote the Banach space of  $X \in \mathbb{D}[0, \infty)$  for which  $\|X\|_{\underline{S}^p}$  is finite. Further for a local martingale  $M$  with  $M(0) = 0$  and a càdlàg adapted process  $A$  with paths of finite variation on compact intervals a.s. and  $A(0) = 0$ , set

$$j_p(M, A) := \left\| [N, N]_\infty^{1/2} + \int_0^\infty |dA(s)| \right\|_{L^p}.$$

For a semimartingale  $Z$  with  $Z(0) = 0$  set

$$\|Z\|_{\underline{H}^p} := \inf_{Z=M+A} j_p(M, A),$$

where the infimum is taken over all possible decompositions  $Z = M + A$  where  $M$  is a local martingale with  $M(0) = 0$  and  $A$  is a càdlàg adapted process with paths of finite variation on compact intervals a.s. and  $A(0) = 0$ . Furthermore, let  $\underline{H}^p[0, \infty)$  denote the Banach space of all semimartingales  $Z$  with  $Z(0) = 0$  such that  $\|Z\|_{\underline{H}^p}$  is finite. For  $t_0 > 0$ , the analogous Banach spaces of processes only defined on  $[0, t_0]$  are denoted by  $\underline{S}^p[0, t_0]$  and  $\underline{H}^p[0, t_0]$ .

For  $1 \leq p < \infty$  there exists a constant  $c_p$  such that

$$\|Z\|_{\underline{S}^p} \leq c_p \|Z\|_{\underline{H}^p} \tag{6}$$

for all semimartingales  $Z$  with  $Z(0) = 0$  (see [20], Theorem V.2, p. 252). For a process  $X \in \mathbb{D}[0, \infty)$  and a stopping time  $T$ , the *pre-stopped process*  $X^{T-}$  is given by

$$(X^{T-})(t)(\omega) = X(t)(\omega)\mathbf{1}_{\{0 \leq t < T(\omega)\}} + X(t \wedge T(\omega)-)(\omega)\mathbf{1}_{\{t \geq T(\omega) > 0\}}, \quad \omega \in \Omega, t \geq 0.$$

Observe that  $X^{T-} \in \mathbb{D}[0, \infty)$ .

The following definition from [22] is essentially from [20], p. 256.

**Definition 3.2.** A map  $\Psi : \mathbb{D}[0, \infty) \rightarrow \mathbb{D}[0, \infty)$  is called functional Lipschitz if for any  $X, Y \in \mathbb{D}[0, t_0]$ :

- (a) for any stopping time  $T$ ,  $X^{T-} = Y^{T-}$  implies  $\Psi(X)^{T-} = \Psi(Y)^{T-}$ ;
- (b) there exists a (positive finite) adapted increasing process  $K$  such that

$$|\Psi(X)(\omega, t) - \Psi(Y)(\omega, t)| \leq K(\omega, t) \sup_{s \leq t} |X(\omega, s) - Y(\omega, s)|.$$

**Definition 3.3.** A map  $\Psi : \mathbb{D}[0, \infty) \rightarrow \mathbb{D}[0, \infty)$  is called locally Lipschitz functional with linear growth if it satisfies:

- (a) for all  $X, Y \in \mathbb{D}[0, \infty)$  and all stopping times  $T$

$$X^{T-} = Y^{T-} \implies \Psi(X)^{T-} = \Psi(Y)^{T-};$$

- (b) for each  $n \in \mathbb{N}$  there exists an adapted increasing process  $K_n$  such that for all  $t \geq 0$  and all  $\omega \in \Omega$ ,

$$|\Psi(X)(\omega, t) - \Psi(Y)(\omega, t)| \leq K_n(\omega, t) \sup_{s \leq t} |X(\omega, s) - Y(\omega, s)|$$

whenever  $|X(\omega, s)|, |Y(\omega, s)| \leq n$  for all  $s \leq t$ ;

- (c) there exists a positive increasing adapted process  $\gamma(t)$  such that

$$|\Psi(X)(\omega, t)| \leq \gamma(\omega, t) \left( 1 + \sup_{s \leq t} |X(\omega, s)| \right) \quad \text{for all } X \in \mathbb{D}[0, \infty).$$

Notice that by (b):  $X^{t_0} = Y^{t_0}$  implies  $\Psi(X)^{t_0} = \Psi(Y)^{t_0}$ , for any  $X, Y \in \mathbb{D}[0, \infty)$ . Therefore, for  $U \in \mathbb{D}[0, t_0]$  we can define  $\Psi(U) \in \mathbb{D}[0, t_0]$  unambiguously, simply by extending  $U$  to  $[0, \infty)$ . Notice also that any functional Lipschitz map is a locally Lipschitz functional with linear growth.

In the sequel we will need the following condition on a function  $g : [0, t_0] \rightarrow \mathbb{R}$ .

**Condition 1.** For every  $Y \in \underline{S}^p[0, t_0]$  and  $Z \in \underline{H}^\infty[0, t_0]$ ,

$$\int_0^\cdot g(\cdot - s)Y(s-) dZ(s) \in \underline{H}^p[0, t_0]$$

and

$$\left\| \int_0^\cdot g(\cdot - s)Y(s-) dZ(s) \right\|_{\underline{H}^p[0, t_0]} \leq R \|Y\|_{\underline{S}^p[0, t_0]} \|Z\|_{\underline{H}^\infty[0, t_0]},$$

where  $t_0 > 0$ ,  $R > 0$  and  $1 < p < \infty$  are given constants.

**Lemma 3.4.** *Let  $t_0 > 0$  be given and let  $g$  satisfy Condition 1 for some  $p$  and  $R$ . Assume that  $\Psi$  is locally Lipschitz with linear growth,  $\Psi(0) = 0$ , and such that the processes  $K_n(t)$  and  $\gamma(t)$  are constants. Let  $Z \in \underline{\underline{H}}^\infty[0, t_0]$  with  $\|Z\|_{\underline{\underline{H}}^\infty} < \frac{1}{4c_p R \gamma}$  and let  $J \in \underline{\underline{S}}^p[0, t_0]$ . Then for any stopping time  $T$  the equation*

$$X(t) = J^{T^-}(t) + \left( \int_0^\cdot g(\cdot - s) \Psi(X)(s-) dZ(s) \right)^{T^-}(t), \quad 0 \leq t \leq t_0,$$

has a unique solution  $X \in \underline{\underline{S}}^p[0, t_0]$ . Moreover,  $X$  is a semimartingale if  $J$  is.

**Proof.** For each  $n \in \mathbb{N}$  and  $X \in \mathbb{D}[0, t_0]$  let  $X^{(n)}(t) := (X(t) \wedge n) \vee (-n)$ ,  $t \in [0, t_0]$ . Now define  $\Psi^n : \mathbb{D}[0, t_0] \rightarrow \mathbb{D}[0, t_0]$  by

$$\Psi^n(X) := \Psi(X^{(n)}).$$

Then  $\Psi^n$  is functional Lipschitz: (a) is immediate and for (b) notice that

$$\begin{aligned} |\Psi^n(X)(t) - \Psi^n(Y)(t)| &\leq K_n \sup_{s \leq t} |X^{(n)}(s) - Y^{(n)}(s)| \\ &\leq K_n \sup_{s \leq t} |X(s) - Y(s)|. \end{aligned}$$

By [20], Theorem V.5(ii), p. 254, there is a stopping time  $T^n$  such that  $\mathbb{P}(T^n > t_0) > 1 - 2^{-n}$  and  $Z^{T^n-} \in \mathcal{S}(\frac{1}{2c_p K_n R})$  (notation of [20]), with  $c_p$  as in (6). Then by [22], Lemma 5.6, there is a unique  $X^n \in \underline{\underline{S}}^p[0, t_0]$  such that

$$X^n(t) = J^{T^n-}(t) + \left( \int_0^t g(t-s) \Psi^n(X^n)(s-) dZ^{T^n-}(s) \right)^{T^n-}$$

and we have

$$\begin{aligned} \|X^n\|_{\underline{\underline{S}}^p} &\leq \|J^{T^n-}\|_{\underline{\underline{S}}^p} + \left\| \left( \int_0^\cdot g(\cdot - s) \Psi^n(X^n)(s-) dZ^{T^n-}(s) \right)^{T^n-} \right\|_{\underline{\underline{S}}^p} \\ &\leq 2\|J\|_{\underline{\underline{S}}^p} + 2c_p \left\| \int_0^\cdot g(\cdot - s) \Psi^n(X^n)(s-) dZ^{T^n-}(s) \right\|_{\underline{\underline{H}}^p} \\ &\leq 2\|J\|_{\underline{\underline{S}}^p} + 2c_p R \|\Psi^n(X^n)\|_{\underline{\underline{S}}^p} \|Z^{T^n-}\|_{\underline{\underline{H}}^\infty} \\ &\leq 2\|J\|_{\underline{\underline{S}}^p} + 2c_p R \gamma (1 + \|X^n\|_{\underline{\underline{S}}^p}) 2\|Z\|_{\underline{\underline{H}}^\infty}, \end{aligned}$$

where the second inequality follows from [20], Theorem V.2 and proof of Theorem V.5, the third by Condition 1, and the fourth by (c) of Definition 3.3. Since  $4c_p R \gamma \|Z\|_{\underline{\underline{H}}^\infty} < 1$  by assumption, we obtain

$$\|X^n\|_{\underline{\underline{S}}^p} \leq \frac{2\|J\|_{\underline{\underline{S}}^p} + 4c_p R \gamma \|Z\|_{\underline{\underline{H}}^\infty}}{1 - 4c_p R \gamma \|Z\|_{\underline{\underline{H}}^\infty}} =: C < \infty \quad \text{for all } n.$$

In particular

$$\mathbb{P}\left( \sup_{0 \leq s \leq t_0} |X^n(s)| > n \right) \leq \frac{C^p}{n^p} \quad \text{for all } n \in \mathbb{N}.$$

Let now  $A_n := \{\sup_{0 \leq s \leq t_0} |X^n(s)| \leq n\}$  and  $B_n := \{T^n > t_0\}$ . Then  $\mathbb{P}(A_n \cap B_n) \geq 1 - 2^{-n} - C^p/n^p$ . Set  $\tilde{\Omega}_n := A_n \cap B_n$ ,  $\Omega_n := \bigcap_{k \geq n} \tilde{\Omega}_k$ ,  $\Omega' := \bigcup_n \Omega_n$ , so that  $\Omega_n \subset \Omega_{n+1}$ ,  $\mathbb{P}(\Omega_n) \uparrow \mathbb{P}(\Omega') = 1$ . On  $\Omega_n$  we have that

$$\sup_{0 \leq s \leq t_0} |X^k(s)| \leq k, \quad Z^{T^k-} = Z,$$

and also  $\Psi^{k_1}(X^k) = \Psi^k(X^k) = \Psi(X^k)$  for  $k_1 \geq k \geq n$ .

If we consider measures  $\mathbb{P}^n$  given by  $\mathbb{P}^n(A) := \mathbb{P}(A \cap \Omega_n) / \mathbb{P}(\Omega_n)$  for  $n$  so large that  $\mathbb{P}(\Omega_n) > 0$ , we have for  $k \geq n$  that

$$\begin{aligned} X^k(t) &= J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi^k(X^k)(s-) dZ(s) \right)^{T^-} \\ &= J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi(X^k)(s-) dZ(s) \right)^{T^-}, \\ X^n(t) &= J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi^n(X^n)(s-) dZ(s) \right)^{T^-} \\ &= J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi(X^n)(s-) dZ(s) \right)^{T^-} \\ &= J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi^k(X^n)(s-) dZ(s) \right)^{T^-} \end{aligned}$$

$\mathbb{P}_n$ -a.s., the stochastic integrals being computed according to the probability  $\mathbb{P}_n$ . This is easily seen by applying [20], Theorems II.14 and II.18, and the fact that the stochastic convolutions above are  $\mathbb{P}$ -a.s. càdlàg. Since  $\Psi^k$  is functional Lipschitz in the sense of [22], Definition 5.1, the uniqueness in [22], Proposition 5.8, yields that we have for  $k \geq n$  that  $X^k = X^n$   $\mathbb{P}^n$ -a.s., thus  $\mathbb{P}$ -a.s. on  $\Omega_n$ . Hence there is a process  $X$  such that for almost all  $\omega \in \Omega'$  and all  $t \geq 0$

$$X(t, \omega) = \lim_{n \rightarrow \infty} X^n(t, \omega),$$

which is adapted and càdlàg, since the filtration satisfies the usual conditions. Moreover, we have for each  $n$  that

$$\begin{aligned} X &= X^n = J^{T^-} + \left( \int_0^\cdot g(\cdot-s)\Psi^n(X^n)(s-) dZ^{T^n-}(s) \right)^{T^-} \\ &= J^{T^-} + \left( \int_0^\cdot g(\cdot-s)\Psi(X)(s-) dZ(s) \right)^{T^-} \end{aligned}$$

on  $\Omega_n$  a.s. Hence  $X$  satisfies the equation. We also have that

$$\sup_{0 \leq s \leq t_0} |X(s)|^p = \lim_{n \rightarrow \infty} \sup_{0 \leq s \leq t_0} |X^n(s)|^p \quad \mathbb{P}\text{-a.s.},$$

so that by Fatou's lemma,

$$\|X\|_{\underline{\underline{S}}^p} \leq \sup_n \|X^n\|_{\underline{\underline{S}}^p} < \infty.$$

For uniqueness, suppose  $X'$  is another solution in  $\mathbb{D}[0, t_0]$ . Consider the fundamental sequences  $S^n := \inf\{t: |X(t)| > n\}$  and  $S'_1 := \inf\{t: |X'(t)| > n\}$ . Then on the set  $C_n := \{S^n \wedge S'_1 > t_0\}$  the processes  $X$  and  $X'$  both solve the equation with  $\Psi^n$ , so by uniqueness in [22], Proposition 5.8,  $X = X'$  on  $C_n$ . Hence  $X = X'$  a.s.

Finally, if  $J$  is a semimartingale,  $X^{S^n-}$  is a semimartingale for each  $n$ , by (a) and (c) of Definition 3.3 together with Condition 1, hence  $X$  is a semimartingale as well. □

**Lemma 3.5.** Assume the situation of Lemma 3.4 with  $p > 2$ , but without the condition  $\|Z\|_{\underline{\underline{H}}^\infty[0, t_0]} < \frac{1}{4c_p R \gamma}$ . Then for each stopping time  $T$  the equation

$$X(t) = J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi(X)(s-) dZ(s) \right)^{T^-}, \quad 0 \leq t \leq t_0,$$

has a unique solution in  $\mathbb{D}[0, t_0]$ . If  $J$  is a semimartingale, then  $X$  is a semimartingale as well.

**Proof.** Let  $\Psi^n$  be the functional Lipschitz maps as in the proof of Lemma 3.4. By [22], Proposition 5.8, for each  $n$  there is an  $\tilde{X}^n \in \mathbb{D}[0, t_0]$  such that

$$\tilde{X}^n(t) = J(t) + \int_0^t g(t-s)\Psi^n(\tilde{X}^n)(s-) dZ(s),$$

which is a semimartingale if  $J$  is a semimartingale. (Indeed, before applying [22], Proposition 5.8,  $J$  and  $Z$  can be extended constantly after  $t_0$  to  $[0, \infty)$  and then the solution can be restricted to  $[0, t_0]$ .) Stop  $\tilde{X}^n$  at  $T^-$  to obtain

$$(\tilde{X}^n)^{T^-}(t) = J^{T^-}(t) + \left( \int_0^t g(t-s)\Psi^n((\tilde{X}^n)^{T^-})(s-) dZ(s) \right)^{T^-}$$

and set  $X^n := (\tilde{X}^n)^{T^-}$ . As before, Condition 1 and [20], Theorem V.2, yield

$$\|X^n\|_{\underline{\underline{S}}^p[0, t_0]} \leq 2\|J\|_{\underline{\underline{S}}^p[0, t_0]} + 2c_p R \|\Psi^n(X^n)\|_{\underline{\underline{S}}^p[0, t_0]} \|Z\|_{\underline{\underline{H}}^\infty[0, t_0]}.$$

By the linear growth of  $\Psi$  we have

$$\begin{aligned} |\Psi^n(X^n)(t)| &= |\Psi((X^n \wedge n) \vee (-n))(t)| \\ &\leq \gamma \left( 1 + \sup_{0 \leq s \leq t} |(X^n(s) \wedge n) \vee (-n)| \right) \leq \gamma(1+n) \end{aligned}$$

and obtain

$$\|X^n\|_{\underline{\underline{S}}^p[0, t_0]} \leq 2\|J\|_{\underline{\underline{S}}^p} + 2c_p R \gamma(1+n) \|Z\|_{\underline{\underline{H}}^\infty[0, t_0]}.$$

Let  $A_n := \{\sup_{0 \leq t \leq t_0} |X^n(t)| > n\}$ ,  $n \in \mathbb{N}$ . Then

$$\mathbb{P}(A_n) \leq 2n^{-p} (\|J\|_{\underline{\underline{S}}^p[0, t_0]} + c_p R \gamma(1+n) \|Z\|_{\underline{\underline{H}}^\infty[0, t_0]})$$

by Chebyshev's inequality. Let further  $\Omega_n := \bigcap_{k \geq n} (\Omega \setminus A_k)$ . Notice that  $\Omega_n \subset \Omega_{n+1}$  and  $\mathbb{P}(\Omega_n) \uparrow 1$ , since  $p > 2$ . Define probability measures  $\mathbb{P}^n(A) := \mathbb{P}(\Omega_n \cap A) / \mathbb{P}(\Omega_n)$  for  $n \geq N$ , where  $N$  is such that  $\mathbb{P}(\Omega_N) > 0$ . Arguing as in Lemma 3.4 we obtain for  $k \geq n$  that  $X^k = X^n$  on  $\Omega_n$ , so we can define a process  $X := \lim X^n$ , which is then the unique solution of the equation. If  $J$  is a semimartingale then so is  $X$ .  $\square$

**Proposition 3.6.** *Let  $Z$  be a semimartingale and  $J \in \mathbb{D}[0, \infty)$ . Let  $g$  be a function on  $[0, \infty)$  which satisfies Condition 1 for each  $t_0 > 0$  with some  $R(t_0) > 0$  and some fixed  $p > 2$ . Further, suppose that  $\Psi$  is a locally Lipschitz functional such that the processes  $K_n$  and  $\gamma$  are deterministic. Then the equation*

$$X(t) = J(t) + \int_0^t g(t-s)\Psi(X)(s-) dZ(s), \quad t \geq 0,$$

has a unique solution in  $\mathbb{D}[0, \infty)$ , which is a semimartingale if  $J$  is.

**Proof.** Fix  $t_0 > 0$ . First we show existence and uniqueness on  $[0, t_0]$ . There is a fundamental sequence  $T^\ell$  of stopping times such that  $J^{T^\ell-} \in \underline{\underline{S}}^\infty[0, \infty)$  and  $Z^{T^\ell-} \in \underline{\underline{H}}^\infty[0, \infty)$ . By the linear growth of  $\Psi$  and Condition 1,  $(\int_0^t g(t-s)\Psi(0)(s-) dZ(s))_{t \geq 0}$  is a semimartingale. Hence we may assume that  $\Psi(0) = 0$ .

By the previous lemma, for each  $\ell$  there is an  $X_\ell$  such that

$$X_\ell(t) = J^{T^\ell-}(t) + \left( \int_0^t g(t-s)\Psi(X_\ell)(s-) dZ^{T^\ell-}(s) \right)^{T^\ell-} \quad \text{for all } t \in [0, t_0].$$

Moreover,

$$X_{\ell+k}^{T^{\ell-}}(t) = J^{T^{\ell-}} + \left( \int_0^t g(t-s) \Psi(X_{\ell+k}^{T^{\ell-}})(s-) dZ^{T^{\ell-}}(s) \right)^{T^{\ell-}}.$$

Define  $X := \lim X_\ell$  and argue as in the two previous lemmas to show that  $X$  is the unique solution on  $[0, t_0]$ . Finally, we can use the same techniques to patch the solutions together obtaining a unique global solution.  $\square$

We can now show existence and uniqueness for the general equation of variation-of-constants form.

**Theorem 3.7.** *Let  $\Psi$  be a locally Lipschitz functional with linear growth and let  $g$  be a function on  $[0, \infty)$  which satisfies Condition 1 for each  $t_0 > 0$  with some  $R(t_0) > 0$  and for some fixed  $p > 2$ . Suppose  $Z$  is a semimartingale and  $J \in \mathbb{D}[0, \infty)$ . Then the equation*

$$X(t) = J(t) + \int_0^t g(t-s) \Psi(X)(s-) dZ(s) \quad (7)$$

has a unique solution in  $\mathbb{D}[0, \infty)$ , which is a semimartingale if  $J$  is.

**Proof.** Fix  $t_0 > 0$ . For  $n, k \in \mathbb{N}$  there are constants  $c_{n,k} > 0$ ,  $\gamma_k > 0$  such that

$$\begin{aligned} \mathbb{P}(\Omega_{n,k}) &\geq 1 - 2^{-n-k}, \quad \text{where } \Omega_{n,k} := \{K_n(t_0, \omega) \leq c_{n,k}\}, \\ \mathbb{P}(\tilde{\Omega}_k) &\geq 1 - 2^{-k}, \quad \text{where } \tilde{\Omega}_k := \{\gamma(t_0, \omega) \leq \gamma_k\}. \end{aligned}$$

Set  $\Omega_k := \bigcap_n \Omega_{n,k} \cap \tilde{\Omega}_k$ . Then  $\mathbb{P}(\Omega_k) \geq 1 - 2^{-k+1}$ , and on  $\Omega_k$ ,  $K_n(t_0, \omega)$  and  $\gamma(t_0, \omega)$  are bounded functions. Now apply the last part of the proof of Proposition 5.8 of [22], using existence and uniqueness from the previous proposition.  $\square$

As we mentioned in the [Introduction](#), the fundamental solution  $r$  of the deterministic delay equation is absolutely continuous on compacts and its derivative is bounded on compacts. Hence due to [22], Lemma 4.2,  $r$  satisfies Condition 1 for any  $t_0 > 0$  and  $p \geq 1$  with  $R = 1 + (1 + c_p)t_0 \sup_{0 \leq t \leq t_0} |r'(t)|$ . So there is a unique solution of the variation-of-constants formula (7) with  $g = r$ . This solution also satisfies the stochastic delay differential equation

$$dX(t) = \left( \int_{[-\alpha, 0]} X(t+s) \mu(ds) \right) dt + \Psi(X)(t-) dZ(t). \quad (8)$$

This can be shown by applying the stochastic Fubini theorem. The result is actually Lemma 6.1 in [22]. Although the statement there presupposes  $\Psi$  to be functional Lipschitz, the only property of the functional  $\Psi$  used in the proof is that it is a map  $\mathbb{D}[0, \infty) \rightarrow \mathbb{D}[0, \infty)$ . As (8) has a unique solution (see [13]), it follows that this solution satisfies the variation-of-constants formula. Stated precisely:

**Theorem 3.8.** *Let  $\mu$  be a finite signed Borel measure on  $[-\alpha, 0]$  and let  $r$  be the fundamental solution of the deterministic delay equation. Let  $\Psi$  be a locally Lipschitz functional and let  $Z$  and  $J$  be semimartingales. The unique solution  $X$  of*

$$X(t) = X(0) + J(t) + \int_0^t \int_{[-s, 0]} X(s+a) \mu(da) ds + \int_0^t \Psi(X)(s-) dZ(s), \quad t \geq 0,$$

satisfies

$$X(t) = r(t)X(0) + \int_0^t r(t-s) dJ(s) + \int_0^t r(t-s) \Psi(X)(s-) dZ(s), \quad t \geq 0.$$

#### 4. The equation and the segment process

In the remaining part of the paper we consider (1) and show that it has an invariant measure under suitable conditions.

Our approach is to see the stochastic equation (1) as a perturbation of the deterministic equation (2). If the deterministic part is stable it is plausible to expect existence of an invariant measure under mild conditions on the diffusion part. Therefore we assume that (5) holds. For an analysis of the case where (5) does not hold, see, e.g., [2].

As was shown in [10], Theorem 3.1, even if  $F$  is constant, a necessary condition for the existence of an invariant measure on the jumps of the Lévy process  $L$  is that  $\int_{|x|>1} \log |x| \nu(dx) < \infty$ , where  $\nu$  denotes the Lévy measure of  $L$ . As our situation is even more general, we need this condition as well.

As mentioned in the [Introduction](#), the main point of this paper is to relax the global Lipschitz condition in [21], Assumption 4.1(c), to a locally Lipschitz condition. Our locally Lipschitz condition on  $F$  is the following.

**Definition 4.1.** A map  $F : D[-\alpha, \infty) \rightarrow D[-\alpha, \infty)$  is called a locally Lipschitz functional of deterministic type (in short lolidet) if it satisfies:

(1)  $\forall n \in \mathbb{N} \exists K_n > 0$  such that  $\forall x, y \in D[-\alpha, \infty) \forall t \geq 0$

$$\sup_{s \in [t-\alpha, t]} |x(s)| \vee |y(s)| \leq n \implies |F(x)(t) - F(y)(t)| \leq K_n \sup_{s \in [t-\alpha, t]} |x(s) - y(s)|,$$

(2)  $\exists \gamma > 0$  such that  $\forall x \in D[-\alpha, \infty) \forall t \geq 0$

$$|F(x)(t)| \leq \gamma \left( 1 + \sup_{s \in [t-\alpha, t]} |x(s)| \right).$$

**Definition 4.2.** Let  $(\Phi(s))_{s \in [-\alpha, 0]}$  be an initial condition, that is, a process with càdlàg paths and such that  $\Phi(s)$  is  $\mathcal{F}_0$ -measurable for all  $-\alpha \leq s \leq 0$ . For  $X \in \mathbb{D}[0, \infty)$  or  $X \in \mathbb{D}[-\alpha, \infty)$  define  $X_{-\Phi} \in \mathbb{D}[-\alpha, \infty)$  by

$$X_{-\Phi}(s) := \begin{cases} \Phi(s), & -\alpha \leq s < 0, \\ X(s), & s \geq 0. \end{cases}$$

Here we extend the filtration by setting  $\mathcal{F}_s := \mathcal{F}_0$  for  $s < 0$ . Define further  $\Psi_\Phi : \mathbb{D}[0, \infty) \rightarrow \mathbb{D}[0, \infty)$  by

$$\Psi_\Phi(X)(t, \omega) := F(X_{-\Phi}(\cdot, \omega))(t).$$

By standard arguments,  $\Psi_\Phi$  indeed maps into  $\mathbb{D}[0, \infty)$ .

Set  $\Omega_0 = \emptyset$  and  $\Omega_n := \{\sup_{[-\alpha, 0]} |\Phi(s)| > n\}$  for  $n \geq 1$ . Then  $\Omega_n \uparrow \Omega$ . Define  $C_n : \Omega \rightarrow \mathbb{R}$  and  $\bar{\gamma} : \Omega \rightarrow \mathbb{R}$  by

$$\begin{aligned} C_n(\omega) &:= K_n \quad \text{on } \Omega_n, & C_n(\omega) &:= K_{n+k} \quad \text{on } \Omega_{n+k} \setminus \Omega_{n+k-1}, \\ \bar{\gamma}(\omega) &:= \gamma(1+n) \quad \text{on } \Omega_n \setminus \Omega_{n-1}. \end{aligned}$$

Notice that  $C_n$  and  $\bar{\gamma}$  are  $\mathcal{F}_0$ -measurable for all  $n$ . Moreover,  $\Psi_\Phi$  satisfies:

(1) for  $t \geq 0$  and  $X, Y \in \mathbb{D}[0, \infty)$ ,

$$|\Psi_\Phi(X)(t, \omega) - \Psi_\Phi(Y)(t, \omega)| \leq C_n(\omega) \sup_{s \in [(t-\alpha)^+, t]} |X(s, \omega) - Y(s, \omega)|$$

if  $\sup_{s \in [(t-\alpha)^+, t]} |X(s, \omega)| \vee |Y(s, \omega)| \leq n$ , and

(2)  $|\Psi_\Phi(X)(t, \omega)| \leq \bar{\gamma}(\omega) \left( 1 + \sup_{[(t-\alpha)^+, \alpha]} |X(t, \omega)| \right)$

for all  $X \in \mathbb{D}[0, \infty)$ .

In other words,  $\Psi_\Phi$  satisfies (b) and (c) of Definition 3.3, and (a) is obvious.

Define

$$J_\Phi(t) := \int_0^t \int_{[-\alpha, -s)} \Phi(s+a)\mu(da) ds, \quad t \geq 0.$$

Then  $J_\Phi$  is an adapted (by Fubini arguments) càdlàg process of finite variation. Moreover,  $r(t)\Phi(0)$  is an adapted process of finite variation. Hence by [22], Theorem 4.1,  $r(t)\Phi(0) + \int_0^t r(t-s) dJ(s)$  is a semimartingale.

Let  $X$  be the unique solution (see [13], Theorem 4.5) of

$$\begin{aligned} X(t) &= \Phi(0) + \int_0^t \int_{[-\alpha, 0]} X_{-\Phi}(s+a)\mu(da) ds + \int_0^t F(X_{-\Phi})(s-) dL(s) \\ &= \Phi(0) + J_\Phi(t) + \int_0^t \int_{[-s, 0]} X(s+a)\mu(da) ds + \int_0^t \Psi_\Phi(X)(s-) dL(s). \end{aligned}$$

By Theorem 3.8,  $X$  is also a solution of

$$X(t) = r(t)\Phi(0) + \int_0^t r(t-s) dJ(s) + \int_0^t r(t-s)\Psi_\Phi(X)(s-) dL(s).$$

Because of (3) Theorem 3.8 takes the following form in the current setting.

**Theorem 4.3.** *Let  $F : D[-\alpha, \infty) \rightarrow D[-\alpha, \infty)$  be lolidet. Then for  $X \in \mathbb{D}[0, \infty)$  and  $\Phi \in \mathbb{D}[-\alpha, 0]$  the following two statements are equivalent:*

(1)  $X$  is the unique solution of

$$X(t) = \Phi(0) + \int_0^t \int_{[-\alpha, 0)} X_{-\Phi}(s+a)\mu(da) ds + \int_0^t F(X_{-\Phi})(s-) dZ(s), \quad t \geq 0,$$

(2)  $X$  obeys the variation-of-constants formula

$$X(t) = x(t, \Phi) + \int_0^t r(t-s)F(X_{-\Phi})(s-) dZ(s), \quad t \geq 0.$$

Recall that for a process  $X \in \mathbb{D}[-\alpha, \infty)$  we denote by  $(X_t)_{t \geq 0}$  the segment process, which takes values in  $D[-\alpha, 0]$  for each  $t$ . More precisely,  $X_t(s) = X(t+s)$  for  $-\alpha \leq s \leq 0$ . We wish to show that the segment process is Markov. For this to be true  $F$  obviously has to be autonomous in the sense of the following definition.

**Definition 4.4.** *A map  $F : D[-\alpha, \infty) \rightarrow D[-\alpha, \infty)$  is autonomous if for all  $x \in D[-\alpha, \infty)$  and all  $s, t \geq 0$ ,*

$$F(x(s + \cdot))(t) = F(x)(s+t).$$

Assume that  $F$  is autonomous. For  $u \geq 0$  and  $(Y(s))_{-\alpha \leq s \leq 0}$  càdlàg and  $\mathcal{F}_u$ -measurable, we consider the equation

$$\begin{aligned} X(t) &= Y(0) + \int_0^t \int_{[-\alpha, 0]} X(s+a)\mu(da) ds + \int_0^t F(X)(s-) dL^u(s), \quad t \geq 0, \\ X(t) &= Y(t), \quad -\alpha \leq t < 0, \end{aligned}$$

where  $L^u(t) = L(t+u) - L(u)$ ,  $t \geq 0$ . The underlying filtration  $\mathcal{G}_t^u$  is (the right continuous version of)  $\sigma(L(s+u) - L(u) : 0 \leq s \leq t) \vee \mathcal{F}_u$ . Denote by  $(X_Y^u(t))_{t \geq -\alpha}$  the unique solution of this equation and let  $(X_{Y,t}^u)_{t \geq 0}$  denote the corresponding segment process.

For any  $\mathcal{F}_0$ -adapted initial condition  $\Phi$  and  $t \geq 0$  the process  $X_\Phi := X_\Phi^0$  satisfies

$$\begin{aligned} X_\Phi(u + \cdot)(t) - X_\Phi(u) &= X_\Phi(t + u) - X_\Phi(u) \\ &= \int_u^{t+u} \int_{[-\alpha, 0]} X_\Phi(s + a) \mu(da) ds + \int_u^{t+u} F(X_\Phi)(s-) dL(s) \\ &= \int_0^t \int_{[-\alpha, 0]} X_\Phi(u + \cdot)(s + a) \mu(da) ds + \int_0^t F(X_\Phi(u + \cdot))(s-) dL^u(s), \end{aligned}$$

where the latter equality holds by the fact that  $F$  is autonomous. The process  $L^u$  is a Lévy process relative to the filtration  $\mathcal{F}_{u+}$  and  $\mathcal{G}_t^u \subset \mathcal{F}_{t+u}$  for all  $t \geq 0$ , hence  $X_{X_\Phi, u}^u$  is also a solution of the equation relative to the filtration  $\mathcal{F}_{u+}$ . (see [20], Theorem II.16). Hence

$$X_{X_\Phi, u}^u(t) = X_\Phi(t + u) \tag{9}$$

for all  $t \geq 0$ , due to the strong uniqueness of the equation.

Under additional conditions we will show below that the segment  $(X_{\Phi, t})_t$  is a Markov process, that is,

$$\mathbb{E}[\mathbf{1}_B(X_{X_\Phi, u, t}^u) | \mathcal{F}_u] = \mathcal{A}_B(X_{\Phi, u}), \tag{10}$$

where

$$\mathcal{A}_B = \mathbb{E} \mathbf{1}_B(X_{\varphi, t}^u), \quad \varphi \in D[-\alpha, 0],$$

for every  $B \in \mathcal{B}(D[-\alpha, 0])$ . Notice that  $X_{\Phi, t}$  is  $\mathcal{F}/\mathcal{B}(D[-\alpha, 0])$ -measurable, since the Borel  $\sigma$ -algebra generated by the Skorohod topology  $\mathcal{B}(D[-\alpha, 0])$  equals the  $\sigma$ -algebra generated by the finite-dimensional set [3], formula (15.2) on p. 157. By  $B_b(D[-\alpha, 0])$  we denote the space of bounded Borel (relative to the Skorohod topology) functions on  $D[-\alpha, 0]$ . For  $0 \leq s \leq t$  we define

$$P_{s,t} f(\varphi) := \mathbb{E} f(X_{\varphi, t-s}^s), \quad \varphi \in D[-\alpha, 0], \quad f \in B_b(D[-\alpha, 0]). \tag{11}$$

We will show that  $P_{s,t}$  maps  $B_b(D[-\alpha, 0])$  into  $B_b(D[-\alpha, 0])$ , that  $P_{u,t} = P_{u,s} P_{s,t}$  for  $0 \leq u \leq s \leq t$ , and that  $P_{s,t} = P_{0,t-s}$ .

Then  $X_{\Phi, t}$  is a homogeneous Markov process and the operators

$$P_t := P_{0,t} \tag{12}$$

form a Markovian semigroup. We will also show that  $(P_t)_{t \geq 0}$  is *eventually Feller* in the following sense:

(1) for  $f \in C_b(D[-\alpha, 0])$ ,  $t \geq \alpha$ ,

$$P_t f \in C_b(D[-\alpha, 0]); \tag{13}$$

(2) for  $t \geq \alpha$ ,  $f \in C_b(D[-\alpha, 0])$ ,

$$\lim_{s \downarrow t} P_s f(\varphi) = P_t f(\varphi) \quad \text{uniformly in } \varphi. \tag{14}$$

Since  $\Delta X_\Phi(t) = F(X_\Phi)(t-) \Delta L(t)$ , and  $L$  is stochastically continuous, we have that  $X_\Phi$  is stochastically continuous, hence by [21], Lemma 3.2, the segment process  $X_{\Phi, t}$  is stochastically continuous as well. So by bounded convergence and subsequence arguments, (2) follows.

The proof of (1) will be more involved and is given in Section 7.

In the sequel we will use several assumptions on the input to our Eq. (1), so we list them here.

**Assumption 1.** (1)  $v_0(\mu) < 0$  (see (5)).

(2) The Lévy measure  $\nu$  of  $L$  satisfies  $\int_{|x| > 1} \log |x| d\nu(x) < \infty$ .

- (3)  $F$  is lolidet.
- (4)  $F$  is autonomous and bounded.

We remark that boundedness of  $F$  implies growth condition (2) of Definition 4.1.

### 5. Tightness of segments

In this section we obtain the tightness of the segment process under Assumption 1. We start by showing that for each fixed  $T \geq 0$ , and any uniformly bounded sequence  $\Phi_n$  of initial conditions (i.e.,  $\sup_n \sup_{-\alpha \leq s \leq 0} \sup_{\omega \in \Omega} |\Phi_n(s, \omega)| < \infty$ ), that

$$\sup_n \mathbb{P} \left( \sup_{t \leq T} |X_{\Phi_n}(t)| > K \right) \rightarrow 0 \quad \text{as } K \rightarrow \infty \tag{15}$$

(recall that  $X_\Phi$  denotes the solution of (1) with initial condition  $\Phi$ ), by showing the following stronger result.

**Theorem 5.1.** *Let  $\Phi_n$  be a uniformly bounded sequence of initial conditions. Let further Assumption 1 hold. Then the laws of  $(X_{\Phi_n}(t + s) : s \in [0, \alpha])_{t \geq 0}$ ,  $n \in \mathbb{N}$  form a tight set. Consequently,*

$$\sup_n \sup_{t \geq 0} \mathbb{P} \left( \sup_{0 \leq u \leq \alpha} |X_{\Phi_n}(t + u)| > K \right) \rightarrow 0 \quad \text{as } K \rightarrow \infty. \tag{16}$$

Notice that (16) implies (15). In order to prove Theorem 5.1, we need several lemmas.

**Lemma 5.2.** *Suppose  $v_0(\mu) < 0$ . The fundamental solution  $r$  of the deterministic delay equation is  $C^1$  on  $[\alpha, \infty)$  and its total variation  $TV_{[-\alpha, \infty)} r$  is finite.*

**Proof.** Since  $r$  is absolutely continuous on  $[0, T]$  for every  $T > 0$ , we have that for  $s > t$  and  $s_n \rightarrow s$  that

$$\int_{[-\alpha, 0]} r(s_n + a) \mu(da) =: g(s_n) \rightarrow g(s) := \int_{[-\alpha, 0]} r(s + a) \mu(da),$$

by bounded convergence. So  $h(t) := \int_\alpha^t g(s) ds$ ,  $t > \alpha$ , is an antiderivative of a continuous function, hence  $C^1$ , and  $h'(t) = g(t)$ , which is a continuous function on  $[\alpha, \infty)$ . Moreover, the estimate  $|g(t)| \leq C' \exp(-\beta t)$  holds for some  $C', \beta > 0$ , so that  $TV_{[\alpha, \infty)} r \leq \int_\alpha^\infty C' \exp(-\beta t) dt < \infty$ . Since  $r$  is absolutely continuous on  $[0, \alpha]$  and  $TV_{[-\alpha, 0]} r = 1$ , we obtain that  $TV_{[-\alpha, \infty)} r < \infty$ . □

**Lemma 5.3.** *Suppose  $v_0(\mu) < 0$  and let  $\dot{r}(t) := \int_{[-\alpha, 0]} r(t + a) \mu(da)$ , for  $t \geq 0$ . Then  $\dot{r}$  is almost everywhere on  $[0, \infty)$  equal to the derivative of  $r$  and the total variation  $TV_{[0, \infty)} \dot{r}$  is finite.*

**Proof.** The first claim follows directly. For the second claim, let  $0 \leq t_0 < t_1 < \dots < t_n < \infty$ , then

$$\begin{aligned} \sum_{i=1}^{n-1} |\dot{r}(t_{i+1}) - \dot{r}(t_i)| &\leq \int_{[-\alpha, 0]} \sum_{i=1}^{n-1} |r(t_{i+1} + a) - r(t_i + a)| |\mu|(da) \\ &\leq TV_{[-\alpha, \infty)} r |\mu|. \end{aligned} \tag{17}$$
□

**Lemma 5.4.** *Suppose  $v_0(\mu) < 0$ . Let  $A \subset D[-\alpha, 0]$  such that  $\sup_{\varphi \in A} \sup_{-\alpha \leq s \leq 0} |\varphi(s)| < \infty$ . Then the solutions of the deterministic delay equation satisfy*

$$\sup_{\varphi \in A} \sup_{t \in [-\alpha, \infty)} |x(t, \varphi)| < \infty.$$

**Proof.** By (4), for  $\varphi \in A$ ,

$$|x(t, \varphi)| = \left| \varphi(0)r(t) + \int_{[-\alpha, 0]} \int_s^0 r(t+s-a)\varphi(s) \, ds \, \mu(ds) \right|$$

$$\leq \sup_{\varphi \in A} |\varphi(0)| \sup_{s \in [-\alpha, \infty)} |r(s)| + \sup_{s \in [-\alpha, \infty)} |r(s)| \alpha \sup_{\varphi \in A} \sup_{-\alpha \leq s \leq 0} |\varphi(s)| |\mu| < \infty. \quad \square$$

**Lemma 5.5.** *If Assumption 1 holds,  $\{\mathcal{L}(X_{\Phi_n}(t)) : t \geq 0, n \in \mathbb{N}\}$  is a tight set of laws on  $\mathbb{R}$ , provided that  $\sup_n \sup_{[-\alpha, 0]} \sup_{\omega \in \Omega} |\Phi_n(s, \omega)| < \infty$ .*

**Proof.** Since  $X_{\Phi_n}(t) = x(t, \Phi_n) + \int_0^t r(t-s)F(X_{\Phi_n})(s-) \, dL(s)$  and by Lemma 5.4  $\sup_{t,n} |x(t, \Phi_n)| < \infty$ , we can execute the same proof as in [21], Proposition 4.2, as the only property of  $F(X_{\Phi_n})$  used there is that it is a bounded process.  $\square$

We proceed by showing that the laws of the deducted segments  $X_{\Phi_n}(t + \cdot) - X_{\Phi_n}(t)$ ,  $t \geq 0$ , are tight as well. Define processes

$$I^n(u) := \int_{[-\alpha, 0]} X_{\Phi_n}(u+v)\mu(dv)$$

$$= \int_{[-\alpha, 0]} \left( x(u+v, \Phi_n) + \int_0^{u+v} r(u+v-s)F(X_{\Phi_n})(s-) \, dL(s) \right) \mu(dv)$$

$$= \int_{[-\alpha, 0]} x(u+v, \Phi_n)\mu(dv) + \int_0^u \dot{r}(u-s)F(X_{\Phi_n})(s-) \, dL(s),$$

where we used the stochastic Fubini theorem and that  $r(s) = 0$  for  $s < 0$ . Hence, since  $X_{\Phi_n}$  is càdlàg, the processes

$$V^n(u) := \int_0^u \dot{r}(u-s)F(X_{\Phi_n})(s-) \, dL(s)$$

have càdlàg versions, which we will use in the sequel.

**Lemma 5.6.** *Processes  $V^n$  defined above satisfy*

$$\lim_{K \rightarrow \infty} \sup_{n \in \mathbb{N}} \sup_{t \geq 0} \mathbb{P} \left( \sup_{0 \leq s \leq \alpha} |V^n(t+s)| > K \right) = 0,$$

*in other words,  $\{\sup_{0 \leq s \leq \alpha} |V^n(t+s)| : t \geq 0, n \in \mathbb{N}\}$  is a tight set of laws on  $\mathbb{R}$ .*

**Proof.** First we show that the set of laws  $\{\mathcal{L}(V^n(u)) : u \geq 0, n \in \mathbb{N}\}$  is tight. To do so, we examine the proof of [21], Proposition 4.2. There the authors prove that the family of laws  $\{\mathcal{L}(X(t)) : t \geq 0\}$  is tight, where

$$X(t) = \int_0^t r(t-s)F(X)(s-) \, dL(s), \quad t \geq 0.$$

In the proof they use the boundedness of  $F(X(s-))$ , the fact that  $r(t)$  decays exponentially for  $t \rightarrow \infty$ , and that the Lévy process  $L$  is exactly as in our Assumption 1. Due to our Assumption 1 we have the same bound for  $F(X_{\Phi_n})(s-)$  for all  $n$  simultaneously. As we also have exponential decay of the function  $\dot{r}$ , we can execute the same proof for  $V^n(u)$  and obtain that  $\{\mathcal{L}(V^n(u)) : u \geq 0, n \in \mathbb{N}\}$  is tight.

As

$$\sup_{0 \leq s \leq \alpha} |V^n(t+s)| \leq \sup_{0 \leq s \leq \alpha} |V^n(t+s) - V^n(t)| + |V^n(t)|,$$

and we have that  $\{\mathcal{L}(V^n(t)): t \geq 0, n \in \mathbb{N}\}$  is tight, it is enough to show tightness of the laws of  $\sup_{0 \leq s \leq \alpha} |V^n(t + s) - V^n(t)|$ , where  $n \in \mathbb{N}$  and  $t \geq 0$ .

Our Lévy process  $L$  decomposes into

$$L(t) = bt + M(t) + N(t),$$

where  $N(t) = \sum_{s \leq t} \Delta L(s) \mathbf{1}_{|\Delta L(s)| > 1}$ ,  $M$  is a square integrable Lévy martingale, and  $b \in \mathbb{R}$ . Then

$$\begin{aligned} & \sup_{0 \leq u \leq \alpha} \left| \int_0^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dN(s) \right| \\ & \leq \sum_{s \leq t+\alpha} C' m \exp(-\beta(t-s)) |\Delta N(s)|, \end{aligned}$$

and the last process is bounded in probability by time reversal for compound Poisson processes and [10], Lemma 4.3 (see also [21], Proof of Proposition 4.2). Further,

$$\left| \int_0^{t+u} \dot{r}(t+u-s) b \, ds \right| \leq C' b \int_0^{t+u} \exp(-\beta(t+u-s)) \, ds < C' b.$$

Therefore it is enough to show that the laws of

$$\sup_{0 \leq u \leq \alpha} \left| \int_0^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dM(s) - \int_0^t \dot{r}(t-s) F(X_{\Phi_n})(s-) \, dM(s) \right|,$$

$n \in \mathbb{N}, t \geq 0$ , are a tight family. Now

$$\begin{aligned} & \left| \int_0^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dM(s) - \int_0^t \dot{r}(t-s) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & \leq \left| \int_0^t (\dot{r}(t+u-s) - \dot{r}(t-s)) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & \quad + \left| \int_t^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dM(s) \right|. \end{aligned}$$

For the first term we estimate

$$\begin{aligned} & \left| \int_0^t (\dot{r}(t+u-s) - \dot{r}(t-s)) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & = \left| \int_0^t \int_0^u d\dot{r}(t-s+v) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & = \left| \int_0^t \int_0^{t+u} \mathbf{1}_{t-v \leq s \leq t-v+u} F(X_{\Phi_n})(s-) \, d\dot{r}(v) \, dM(s) \right| \\ & = \left| \int_0^{t+u} \left( \int_{t-v}^{t-v+u} F(X_{\Phi_n})(s-) \, dM(s) \right) d\dot{r}(v) \right| \\ & \leq \int_0^{t+\alpha} \sup_{w \leq \alpha, n \in \mathbb{N}} \left| \int_{t-v}^{t-v+w} F(X_{\Phi_n})(s-) \, dM(s) \right| d|\dot{r}|(v), \end{aligned}$$

by the stochastic Fubini theorem. Hence

$$\begin{aligned} & \mathbb{E} \sup_{u \leq \alpha} \left| \int_0^t (\dot{r}(t+u-s) - \dot{r}(t-s)) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & \leq \int_0^{t+\alpha} \left( \mathbb{E} \sup_{u \leq \alpha} \left| \int_{t-v}^{t-v+u} F(X_{\Phi_n})(s-) \, dM(s) \right|^2 \right)^{1/2} d|\dot{r}|(v) \\ & \leq (TV_{[0, \infty)} \dot{r}) 4m^2 (\mathbb{E} M(\alpha)^2)^{1/2}, \end{aligned}$$

by Doob’s inequality, boundedness of  $F$  and the fact that  $M$  is a Lévy square integrable martingale.

For the second term, we first extend  $\dot{r}$  by  $\dot{r}(s) = \dot{r}(0)$  for  $s < 0$  and compute

$$\begin{aligned} & \left| \int_t^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & = \left| \int_t^{t+u} \left( \int_0^u d\dot{r}(t-s+v) + \dot{r}(t-s) \right) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & \leq \left| \int_t^{t+u} \int_0^u d\dot{r}(t-s+v) F(X_{\Phi_n})(s-) \, dM(s) \right| + |\dot{r}(0) F(X_{\Phi_n})(t-) \Delta M(t)| \\ & \leq \left| \int_t^{t+u} \int_0^u \mathbf{1}_{t-z \leq s \leq t+u-z} F(X_{\Phi_n})(s-) \, d\dot{r}(z) \, dM(s) \right| + |\dot{r}(0)|m \\ & = \left| \int_0^u \left( \int_t^{t+u-z} F(X_{\Phi_n})(s-) \, dM(s) \right) d\dot{r}(z) \right| + |\dot{r}(0)|m, \end{aligned}$$

by applying stochastic Fubini theorem and because  $|\Delta M| \leq 1$ . Hence, arguing as above, we obtain

$$\begin{aligned} & \mathbb{E} \sup_{0 \leq u \leq \alpha} \left| \int_t^{t+u} \dot{r}(t+u-s) F(X_{\Phi_n})(s-) \, dM(s) \right| \\ & \leq (TV_{[0, \infty)} \dot{r}) 4m^2 (\mathbb{E} M(\alpha)^2)^{1/2} + |\dot{r}(0)|m. \end{aligned}$$

Now the proof is complete. □

**Proposition 5.7.** *Let Assumption 1 hold, and let  $\Phi_n$  be a uniformly bounded sequence of initial conditions. Then the laws of  $(X_{\Phi_n}(t+s) - X_{\Phi_n}(t): s \in [0, \alpha], t \geq 0, n \in \mathbb{N})$  are tight in  $D[0, \alpha]$ .*

**Proof.** We use the same strategy as in [21], Proposition 4.3, and only need to show that

$$\lim_{K \rightarrow \infty} \sup_n \sup_{t \geq 0} \mathbb{P} \left( \sup_{0 \leq u \leq \alpha} |I^n(t+u)| > K \right) \rightarrow 0,$$

where

$$\begin{aligned} I^n(u) &= \int_{[-\alpha, 0]} X_{\Phi_n}(u+v) \mu(dv) \\ &= \int_{[-\alpha, 0]} x(u+v, \Phi_n) \mu(dv) + \int_0^u \dot{r}(u-s) F(X_{\Phi_n})(s-) \, dL(s). \end{aligned}$$

Since the first term is bounded in  $n$  and  $t$  by Lemma 5.4, we infer the claim with the aid of Lemma 5.6. □

Now proving Theorem 5.1 is straightforward.

**Proof of Theorem 5.1.** We have  $X_{\phi_n}(t + \cdot) = (X_{\phi_n}(t + \cdot) - X_{\phi_n}(t)\mathbf{1}) + X_{\phi_n}(t)\mathbf{1}$  for all  $t \geq 0$ , where  $\mathbf{1}(s) = 1$  for all  $s \in [0, \alpha]$ . Let  $\varepsilon > 0$ . By Proposition 5.7, there exists a compact set  $K \subset D[0, \alpha]$  such that  $\mathbb{P}(X_{\phi_n}(t + \cdot) - X_{\phi_n}(t)\mathbf{1} \in K) \geq 1 - \varepsilon/2$  for all  $t \geq 0$  and  $n \in \mathbb{N}$ . By Lemma 5.5 there exists a bounded interval  $I \subset \mathbb{R}$  such that  $\mathbb{P}(X_{\phi_n}(t) \in I) \geq 1 - \varepsilon/2$ . Let  $K' := \{\sigma + c\mathbf{1} : \sigma \in K, c \in I\}$ . Then  $\mathbb{P}(X_{\phi_n}(t + \cdot) \in K') \geq 1 - \varepsilon$ . The set  $K'$  has compact closure in  $D[0, \alpha]$ , due to [3], Theorem 12.4. Indeed, as  $K$  is compact it satisfies conditions (12.25) and (12.30) of [3]. Then  $K'$  satisfies these conditions as well, hence it has compact closure by the same theorem. Thus,  $\{\mathcal{L}(X_{\phi_n}(t + \cdot)) : t \geq 0, n \in \mathbb{N}\}$  is tight.  $\square$

### 6. A stability theorem

In this section we prove that  $\Phi_n \rightarrow \Phi$  in  $D[-\alpha, 0]$  w.r.t.  $d_\beta$  in probability implies uniform convergence on compact sets in probability of the corresponding solutions, under Assumption 1 and the following condition:

$$\varphi_n \rightarrow \varphi \text{ in } D[-\alpha, 0] \text{ w.r.t. } d_\beta \implies \int_0^\alpha (F(x_{-\varphi})(t) - F(x_{-\varphi_n})(t))^2 dt \rightarrow 0, \tag{17}$$

where  $x_{-\varphi}(t) := x(t)$  for  $t \geq 0$  and  $x_{-\varphi}(t) := \varphi(t)$  for  $-\alpha \leq t \leq 0$ , and likewise for  $x_{-\varphi_n}$ .

We need the following approximations of  $F$ :

$$F^N(x)(t) := F(x^{(N)})(t), \quad \text{where } x^N := (x \wedge N) \vee (-N), \text{ for } x \in D[-\alpha, \infty), t \geq 0, \text{ and } N > 0.$$

Then  $F^N$  is Lipschitz in the sense of [21], formula (2.5). Indeed, for  $x, y \in D[-\alpha, \infty)$  and  $t \geq 0$  and  $N > 0$ , we have that  $|x^N(t)|, |y^N(t)| \leq N$  for all  $t \geq -\alpha$ , hence by condition (1) of Definition 4.1 there is  $K_N$  such that

$$\begin{aligned} |F^N(x)(t) - F^N(y)(t)| &= |F(x^N)(t) - F(y^N)(t)| \\ &\leq K_N \sup_{s \in [t-\alpha, t]} |x^N(s) - y^N(s)| \leq K_N \sup_{s \in [t-\alpha, t]} |x(s) - y(s)| \end{aligned}$$

for all  $t \geq 0$ .

If (17) holds for  $F$  then it also holds for the approximations  $F^N$ , provided  $N > \sup_{n,s} |\varphi_n(s)| \vee |\varphi(s)|$ , since then for any  $x \in D[0, \infty)$ ,

$$F^N(x_{-\varphi}) = F((x_{-\varphi})^{(N)}) = F((x^{(N)})_{-\varphi})$$

and the same holds for  $\varphi_n$ .

We need the next lemma.

**Lemma 6.1.** *Let  $L$  be a Lévy process with Lévy measure  $\nu$  and let  $T > 0$ . Then for each  $K > 0$  there exist constants  $b$  and  $\sigma$  such that for each stopping time  $R$  with  $|\Delta L^{R-}| < K$  we have*

$$\left\| \int_0^T H(s-) dL^{R-}(s) \right\|_{\underline{H}^2[0, T]}^2 \leq 2 \left( b^2 T + 2\sigma^2 + 2 \int_{(-K, K)} u^2 \nu(du) \right) \int_0^T \mathbb{E} H(t)^2 dt$$

for every predictable process  $H$  with  $\int_0^T \mathbb{E} H(t)^2 dt < \infty$ .

**Proof.** Let  $K > 0$  and let  $R$  be a stopping time such that  $|\Delta L^{R-}| < K$ , and let  $H$  be a predictable process such that  $\int_0^T \mathbb{E} H(t)^2 dt < \infty$ . Consider the Lévy–Itô decomposition of  $L$  (see [1], Theorem 2.4.16),

$$L(t) = bt + \sigma B(t) + \int_{(-K, K)} u \tilde{N}(t, du) + \int_{|u| \geq K} u N(t, du)$$

and set  $L_K(t) = L(t) - \int_{|u| \geq K} u N(t, du)$ . Then  $L^{R-} = L_K^{R-}$ , so we have

$$\left\| \int_0^\cdot H(t) dL^{R-} \right\|_{\underline{\underline{H}}^2[0, T]} = \left\| \int_0^\cdot H(t) dL_K^{R-} \right\|_{\underline{\underline{H}}^2[0, T]} \leq 2 \left\| \int_0^\cdot H(t-) dL_K \right\|_{\underline{\underline{H}}^2[0, T]},$$

arguing as in [20], Proof of Theorem V.5. Denote

$$I_H(t) = \int_0^t \int_{(-K, K)} u H(t) \tilde{N}(dt, du),$$

where the integral is as defined in [1], Section 4.2. Notice that  $I_H(t)$  equals the usual stochastic integral of  $H$  with respect to the Lévy process  $\int_{(-K, K)} u \tilde{N}(t, du)$ , as one can see from the construction of both integrals. Since  $\int_0^T \int_{(-K, K)} \mathbb{E}(uH(t))^2 dt \nu < \infty$  by assumption and Fubini, [1], Theorem 4.2.3, yields that  $I_H$  is a square integrable martingale and  $\mathbb{E}I_H(t)^2 = \int_{(-K, K)} u^2 \nu \int_0^T \mathbb{E}H(t)^2 dt$ . Now

$$\int_0^t H(s) dL_K(s) = b \int_0^t H(s) ds + \sigma \int_0^t H(s) dB(s) + I_H(t),$$

where the first term is a process of bounded variation and the latter two terms are square integrable martingales. Hence by a well-known identity for square integrable martingales (see [20], Corollary 3 to Theorem II.27),

$$\begin{aligned} \left\| \int_0^\cdot H(s) dL(s) \right\|_{\underline{\underline{H}}^2[0, T]}^2 &\leq 2\mathbb{E}\left(TV\left(b \int_0^\cdot H(s) ds\right)\right)^2 + 2\mathbb{E}\left(\sigma \int_0^T H(s) dB(s) + I_H(t)\right)^2 \\ &\leq 2b^2 T \int_0^T \mathbb{E}H(t)^2 dt + 4\sigma^2 \int_0^T \mathbb{E}H(t)^2 dt + 4\mathbb{E}(I_H(T))^2. \end{aligned} \quad \square$$

**Theorem 6.2.** *Let Assumption 1 and (17) hold. If  $\Phi_n \rightarrow \Phi$  in  $D[-\alpha, 0]$  w.r.t.  $d_\beta$  in probability, then  $X_{\Phi_n} \rightarrow X_\Phi$  uniformly on compact subintervals of  $[0, \infty)$  in probability.*

**Proof.** Write  $X = X_\Phi$ ,  $X^n = X_{\Phi_n}$  throughout this proof. Fix  $T > 0$  and  $\varepsilon > 0$ .

Assume first that  $\{\Phi_n, \Phi\}$  is a uniformly bounded family. Hence Theorem 5.1 can be applied and (15) holds, so that there exists  $N_0$  such that for  $N > N_0$ ,

$$\sup_n \mathbb{P}\left(\sup_{t \leq T} |X^n(t)| > N\right), \quad \mathbb{P}\left(\sup_{t \leq T} |X(t)| > N\right) < \varepsilon.$$

Define stopping times  $T^n := \inf\{t: |X^n(t)| > N_0\}$  and  $T^\infty := \inf\{t: |X(t)| > N_0\}$ . Then  $\mathbb{P}(T^n > T) > 1 - \varepsilon$  and  $\mathbb{P}(T^\infty > T) > 1 - \varepsilon$ . Moreover,

$$\begin{aligned} (X^n)^{T^n-}(t) &= \Phi_n(0) + \int_0^{t \wedge T^n-} \int_{[-\alpha, 0]} (X^n)^{T^n-}(s+a) \mu(da) ds + \int_0^t F((X^n)^{T^n-})(s-) dL^{T^n-}(s) \\ &= \Phi_n(0) + \int_0^{t \wedge T^n-} \int_{[-\alpha, 0]} (X^n)^{T^n-}(s+a) \mu(da) ds + \int_0^t F^{N_0}((X^n)^{T^n-})(s-) dL^{T^n-}(s). \end{aligned}$$

So by uniqueness of solutions,  $(X^n)^{T^n-} = (X_{\Phi_n}^{(N_0)})^{T^n-}$ , where  $X_{\Phi_n}^{(N_0)}$  denotes the solution to Eq. (1) with  $F$  replaced by  $F^{N_0}$ . Likewise,  $(X)^{T^\infty-} = (X_\Phi^{(N_0)})^{T^\infty-}$ . Since

$$\sup_{t \leq T} |X^n(t) - X(t)| \leq \sup_{t \leq T} |X^n(t) - X_{\Phi_n}^{(N_0)}(t)| + \sup_{t \leq T} |X_{\Phi_n}^{(N_0)}(t) - X_\Phi^{(N_0)}(t)| + \sup_{t \leq T} |X_\Phi^{(N_0)}(t) - X(t)|,$$

we obtain for  $\delta > 0$ ,

$$\begin{aligned} \mathbb{P}\left(\sup_{t \leq T} |X^n(t) - X(t)| > \delta\right) &\leq \mathbb{P}\left(\sup_{t \leq T} |X^n(t) - X_{\Phi_n}^{(N_0)}(t)| > \delta/3\right) \\ &\quad + \mathbb{P}\left(\sup_{t \leq T} |X_{\Phi_n}^{(N_0)}(t) - X_{\Phi}^{(N_0)}(t)| > \delta/3\right) + \mathbb{P}\left(\sup_{t \leq T} |X_{\Phi}^{(N_0)}(t) - X(t)| > \delta/3\right) \\ &\leq \mathbb{P}(T^n \leq T) + \mathbb{P}\left(\sup_{t \leq T} |X_{\Phi_n}^{(N_0)}(t) - X_{\Phi}^{(N_0)}(t)| > \delta/3\right) + \mathbb{P}(T^\infty \leq T) \\ &\leq 2\varepsilon + \mathbb{P}\left(\sup_{t \leq T} |X_{\Phi_n}^{(N_0)}(t) - X_{\Phi}^{(N_0)}(t)| > \delta/3\right). \end{aligned}$$

Hence in this special case there is no loss of generality by assuming that  $F$  is Lipschitz in the sense of [21], formula (2.5).

Let  $R$  be a stopping time such that  $L^{R-}$  has bounded jumps, is  $\alpha$ -sliceable for suitably small  $\alpha$ , and  $\mathbb{P}(R > T) > 1 - \varepsilon$  (see [20], Theorem V.5). Denote by  $Z$  and  $Z^n$  the solutions of Eq. (1) with  $L$  replaced by  $L^{R-}$  and initial condition  $\Phi$  and  $\Phi_n$ , respectively. By uniqueness of solutions,  $(Z^n)^{R-} = (X^n)^{R-}$  and  $Z^{R-} = X^{R-}$ . Hence for  $\delta > 0$ ,

$$\begin{aligned} &\mathbb{P}\left(\sup_{0 \leq t \leq T} |X^n(t) - X(t)| \geq \delta\right) \\ &\leq \mathbb{P}\left(\sup_{0 \leq t \leq T} |X^n(t) - X(t)| \geq \delta \text{ and } R \geq T\right) + \mathbb{P}(R < T) \\ &\leq \mathbb{P}\left(\sup_{0 \leq t \leq T} |Z^n(t) - Z(t)| \geq \delta\right) + \varepsilon/2, \end{aligned}$$

so it suffices to show that  $Z^n \rightarrow Z$  uniformly on  $[0, T]$  in probability. To show this we introduce some notation:

$$\begin{aligned} Y^n(t) &:= \int_0^t \int_{[-\alpha, 0]} Z_{-\Phi}(s+u) - Z_{-\Phi_n}(s+u) \mu(du) ds + \int_0^t (F(Z_{-\Phi}) - F(Z_{-\Phi_n}))(s-) dL^{R-}(s), \\ \mathcal{G}^n(U)(t) &:= \int_{[-\alpha, 0]} (Z_{-\Phi_n}(t+u) - (Z-U)_{-\Phi_n}(t+u)) \mu(du), \\ \mathcal{H}^n(U)(t) &:= F(Z_{-\Phi_n})(t) - F((Z-U)_{-\Phi_n})(t), \end{aligned}$$

$t \geq 0$ . We obtain for  $U^n := Z - Z^n$  the equation

$$U^n(t) = \Phi(0) - \Phi_n(0) + Y^n(t) + \int_0^t \mathcal{G}^n(U^n)(s) ds + \int_0^t \mathcal{H}^n(U^n)(s-) dL^{R-}(s).$$

Lemma V.3.2 of [20] extended to two driving semimartingales yields  $\|U^n\|_{\underline{\underline{S}}^2[0, T]} \leq C \|\Phi(0) - \Phi_n(0) + Y^n\|_{\underline{\underline{S}}^2[0, T]}$  with a constant  $C > 0$  depending on the process  $L^{R-}$  and the uniform bound for the Lipschitz constants of mappings  $V \mapsto \int_{[-\alpha, 0]} V_{-\Phi}(\cdot + u) \mu(du)$ ,  $V \mapsto \int_{[-\alpha, 0]} V_{-\Phi_n}(\cdot + u) \mu(du)$ ,  $V \mapsto F(V_{-\Phi})$  and  $V \mapsto F(V_{-\Phi_n})$ . (Notice that this bound is finite as  $F$  is assumed to be Lipschitz and we assumed that  $\Phi_n, \Phi$  is a uniformly bounded family. As  $\Phi_n \rightarrow \Phi$  w.r.t.  $d_\beta$  in probability, we have  $\Phi_n(0) \rightarrow \Phi(0)$  in probability, hence  $\|U^n\|_{\underline{\underline{S}}^2[0, T]} \rightarrow 0$  if  $\|Y^n\|_{\underline{\underline{S}}^2[0, T]} \rightarrow 0$ .)

Next we show  $\|Y^n\|_{\underline{\underline{S}}^2[0, T]} \rightarrow 0$ . Due to the continuous embedding of  $\underline{\underline{H}}^2[0, T]$  into  $\underline{\underline{S}}^2[0, T]$  (see [20], Theorem V.2) and Lemma 6.1 this follows if

$$\mathbb{E} \int_0^T \left( \left( \int_{[-\alpha, 0]} (Z_{-\Phi}(t+u) - Z_{-\Phi_n}(t+u)) \mu(du) \right)^2 + (F(Z_{-\Phi})(t) - F(Z_{-\Phi_n})(t))^2 \right) dt \rightarrow 0 \tag{18}$$

as  $n \rightarrow \infty$ . By the boundedness of  $F$  and assumption (17) we have  $\mathbb{E} \int_0^T (F(Z_{-\Phi})(t) - F(Z_{-\Phi_n})(t))^2 dt \rightarrow 0$  as  $n \rightarrow \infty$ . Moreover,

$$\begin{aligned} & \int_0^T \left( \int_{[-\alpha, 0]} Z_{-\Phi}(t+u) - Z_{-\Phi_n}(t+u) \mu(du) \right)^2 dt \\ &= \int_0^T \left( \int_{[-\alpha, 0]} (\Phi(t+u) - \Phi_n(t+u)) \mu(du) \right) \left( \int_{[-\alpha, 0]} (\Phi(t+v) - \Phi_n(t+v)) \mu(dv) \right) dt \\ &= \int_{[-\alpha, 0]} \int_{[-\alpha, 0]} \int_0^T (\Phi(t+u) - \Phi_n(t+u)) (\Phi(t+v) - \Phi_n(t+v)) \mathbf{1}_{[-\alpha, -t]}(u) \\ & \quad \times \mathbf{1}_{[-\alpha, -t]}(v) dt \mu(du) \mu(dv). \end{aligned}$$

This expression converges almost surely to zero and is bounded in  $n$  and  $\omega$ , as convergence in  $d_\beta$  implies almost everywhere convergence on  $[-\alpha, 0]$  and the family  $\Phi_n, \Phi$  is uniformly bounded. Hence (18) holds indeed, and we proved the special case of uniformly bounded initial conditions.

For the general case, notice that since  $\Phi_n \rightarrow \Phi$  w.r.t.  $d_\beta$  in probability, the laws of  $\Phi_n$  converge weakly to the law of  $\Phi$ , and since  $(D[-\alpha, 0], d_\beta)$  is Polish we have by the Prohorov theorem that the family of laws of  $\Phi_n, \Phi$  is tight. Hence for a  $\varepsilon > 0$  there is a set  $K \subset D[-\alpha, 0]$  compact w.r.t.  $d_\beta$  such that  $\mathbb{P}(\Phi_n \in K, \Phi \in K) > 1 - \varepsilon$  for all  $n$ . As convergence w.r.t.  $d_\beta$  is implied by Skorohod convergence,  $K$  is also compact w.r.t.  $d_S$ . Hence all the functions in  $K$  are bounded by some finite constant  $C$ . Consider the truncated initial conditions  $\Phi_n^C$  and  $\Phi^C$  and let  $\bar{X}^n$  and  $\bar{X}$  be the solutions of Eq. (1) with these initial conditions. We have that  $\mathbb{P}(\Phi_n = \Phi_n^C, \Phi = \Phi^C) > 1 - \varepsilon$ , and concentrating  $\mathbb{P}$  on the sets  $\{\Phi_n = \Phi_n^C\}$  and  $\{\Phi = \Phi^C\}$ , with the aid of [20], Theorem IV.23, and the uniqueness of solutions we conclude that

$$\begin{aligned} \bar{X}^n &= X^n \quad \text{a.s. on } \{\Phi_n = \Phi_n^C\}, \\ \bar{X} &= X \quad \text{a.s. on } \{\Phi = \Phi^C\}. \end{aligned}$$

Moreover, it is easy to check that  $\Phi_n^C \rightarrow \Phi^C$  w.r.t.  $d_\beta$  in probability, so that be the special case above,  $\sup_{0 \leq t \leq T} |\bar{X}^n(t) - \bar{X}(t)| \rightarrow 0$  in probability. Finally, for  $\delta > 0$ ,

$$\begin{aligned} \mathbb{P}\left(\sup_{0 \leq t \leq T} |X(t) - X^n(t)| > \delta\right) &\leq \mathbb{P}\left(\sup_{0 \leq t \leq T} |X(t) - \bar{X}(t)| > \delta/3\right) + \mathbb{P}\left(\sup_{0 \leq t \leq T} |\bar{X}(t) - \bar{X}^n(t)| > \delta/3\right) \\ &\quad + \mathbb{P}\left(\sup_{0 \leq t \leq T} |\bar{X}^n(t) - X^n(t)| > \delta/3\right) \\ &\leq 2\varepsilon + \mathbb{P}\left(\sup_{0 \leq t \leq T} |\bar{X}^n(t) - X^n(t)| > \delta/3\right) \end{aligned}$$

and the theorem has been proved. □

**Remark 6.3.**

- (1) By stopping  $X = X_\Phi$  appropriately, we can use similar techniques as before and prove Theorem 6.2 even if  $F$  is not bounded, but merely having linear growth.
- (2) Each of the conditions ‘ $\Phi_n \rightarrow \Phi$  w.r.t.  $d_S$  in probability’, ‘ $\Phi_n \rightarrow \Phi$  w.r.t.  $d_\beta$  a.s.’ and ‘ $\Phi_n \rightarrow \Phi$  w.r.t.  $d_S$  a.s.’ is stronger than the condition of Theorem 6.2.

In the next corollary the use of  $d_\beta$  instead of  $d_S$  is essential; see [21], Section 3.3, for a counterexample with  $d_S$ .

**Corollary 6.4.** *Let Assumption 1 and (17) hold. If  $\Phi_n \rightarrow \Phi$  with respect to  $d_\beta$  in probability, then  $X_{\Phi_n, t} \rightarrow X_{\Phi, t}$  with respect to  $d_\beta$  in probability for every  $t \geq 0$ .*

**Proof.** If  $t \geq \alpha$ , the assertion readily follows from Theorem 6.2. Consider  $0 < t < \alpha$ . Let  $\lambda$  be an increasing homeomorphism from  $[-\beta, 0]$  onto itself. Define  $\rho: [-\beta, 0] \rightarrow [-\beta, 0]$  by  $\rho(s) = s$  for  $s \in [-t, 0]$ ,  $\rho(s) = \lambda(t+s) - t$  for  $s \in [-\alpha, -t]$ , and affine on  $[-\beta, -\alpha]$  with  $\rho(-\beta) = -\beta$ . Then  $\rho$  is an increasing homeomorphism,  $\sup_{s \in [-\beta, 0]} |\rho(s) - s| \leq \sup_{s \in [-\beta, 0]} |\lambda(s) - s|$ , and

$$\begin{aligned} & \sup_{s \in [-\alpha, 0]} |X_{\Phi_n}(t+s) - X_{\Phi}(t+\rho(s))| \\ & \leq \sup_{s \in [-\alpha, -t]} |\Phi_n(t+s) - \Phi(\lambda(t+s))| \vee \sup_{s \in [-t, 0]} |X_{\Phi_n}(t+s) - X_{\Phi}(t+s)|. \end{aligned}$$

Hence  $d_{\beta}(X_{\Phi_n, t}, X_{\Phi, t}) \leq d_{\beta}(\Phi_n, \Phi) + \sup_{s \in [0, \alpha]} |X_{\Phi_n}(s) - X_{\Phi}(s)|$ , and the proof follows.  $\square$

## 7. Markov and eventual Feller property and existence of invariant measure

**Theorem 7.1.** *Let Assumption 1 and (17) hold. The segment process  $(X_{\Phi, t})_{t \geq 0}$  is a Markov process, the transition operators  $P_{s, t}$  defined by (11) map  $B_d(D[-\alpha, 0])$  into  $B_d(D[-\alpha, 0])$  and satisfy*

$$P_{u, t} = P_{u, s} P_{s, t} \quad \text{and} \quad P_{s, t} = P_{0, t-s}$$

for all  $0 \leq u \leq s \leq t$ . Moreover, the Markov semigroup  $(P_t)_{t \geq 0}$  defined by (12) is eventually Feller.

**Proof.** In this proof we endow  $D[-\alpha, 0]$  with the metric  $d_{\beta}$ . Recall that by Lemma 2.1  $d_{\beta}$  and  $d_S$  generate the same Borel  $\sigma$ -algebra  $\mathcal{B}(D[-\alpha, 0])$ .

We begin by showing (10). Let  $0 \leq u \leq t$  and  $B \in \mathcal{B}(D[-\alpha, 0])$ . Observe that  $\mathbf{1}_B(X_{X_{\Phi, t-u}}^u)$  is measurable, as  $X_{X_{\Phi, t-u}}^u = X_{\Phi}(t)$ , by (9). Let  $C_b$  denote the space of bounded functions  $f: D[-\alpha, 0] \rightarrow \mathbb{R}$  that are continuous with respect to  $d_{\beta}$ . Let  $f \in C_b$  and let  $\xi$  be an  $\mathcal{F}_u$ -measurable random variable with values in  $D[-\alpha, 0]$ . For a  $\mathcal{F}_u$ -measurable random variable  $\xi$  with values in  $D[-\alpha, 0]$ , let  $A(\xi, \omega) := f(X_{\xi, t}^u(\omega))$ . Let further  $\mathcal{A}(\varphi) := \mathbb{E}A(\varphi, \cdot)$  for  $\varphi \in D[-\alpha, 0]$ . Assume first that

$$\xi = \sum_{i=1}^n a_i \mathbf{1}_{C_i} \tag{19}$$

with  $a_i \in D[-\alpha, 0]$  and  $C_i \in \mathcal{F}_u$ , and  $C_i$  mutually disjoint,  $\bigcup_i C_i = \Omega$ , and  $\mathbb{P}(C_i) > 0$  for all  $i$ . Then  $A(\xi, \omega) = \sum_i A(a_i, \omega) \mathbf{1}_{C_i}(\omega)$  (as before we rescale  $\mathbb{P}$  to  $C_i$  and use [20], Theorem IV.23, and uniqueness of solutions), so

$$\mathbb{E}[A(\xi(\cdot), \cdot) | \mathcal{F}_u] = \sum_i \mathbf{1}_{C_i} \mathbb{E}A(a_i, \cdot) = \sum_i \mathcal{A}(a_i) \mathbf{1}_{C_i} = \mathcal{A}(\xi(\cdot)).$$

If  $\xi$  is an arbitrary  $\mathcal{F}_u$ -measurable random variable with values in  $D[-\alpha, 0]$ , then there are  $\xi_m$  of the form (19) such that  $d_{\beta}(\xi_m(\omega), \xi(\omega)) \rightarrow 0$  as  $m \rightarrow \infty$  for a.e.  $\omega$  (see [24], Proposition I.1.9). Due to the continuity of  $f$  and Corollary 6.4 we have  $A(\xi_m, \cdot) \rightarrow A(\xi, \cdot)$  in probability, so that

$$\mathcal{A}(\xi_m) = \mathbb{E}[A(\xi_m, \cdot) | \mathcal{F}_u] \rightarrow \mathbb{E}[A(\xi, \cdot) | \mathcal{F}_u],$$

as  $f$  is bounded. Again by Corollary 6.4 we have  $\mathcal{A}(\varphi_n) \rightarrow \mathcal{A}(\varphi)$  whenever  $d_{\beta}(\varphi_n, \varphi) \rightarrow 0$ , so that  $\mathcal{A}(\xi_m) \rightarrow \mathcal{A}(\xi)$  a.s. Hence  $\mathbb{E}[A(\xi, \cdot) | \mathcal{F}_u] = \mathcal{A}(\xi)$  a.s.

Next let  $C$  be a closed subset of  $D[-\alpha, 0]$  and choose  $f_n \in C_b$  such that  $f_n \downarrow \mathbf{1}_C$  pointwise. Let  $A_n(\omega) := f_n(X_{\varphi, t}^u(\omega))$  and  $A(\varphi, \omega) = \mathbf{1}_C(X_{\varphi, t}^u(\omega))$ ,  $\omega \in \Omega$ , and  $\mathcal{A}_n(\varphi) = \mathbb{E}A_n(\varphi, \cdot)$  and  $\mathcal{A}(\varphi) = \mathbb{E}A(\varphi, \cdot)$ ,  $\varphi \in D[-\alpha, 0]$ . Then  $A_n \downarrow A$  and  $\mathcal{A}_n \downarrow \mathcal{A}$  pointwise, so

$$\mathbb{E}[A(\xi, \cdot) | \mathcal{F}_u] = \lim_{n \rightarrow \infty} \mathbb{E}[A_n(\xi, \cdot) | \mathcal{F}_u] = \lim_{n \rightarrow \infty} \mathcal{A}_n(\xi) = \mathcal{A}(\xi) \quad \text{a.s.}$$

By a monotone class argument we can extend the above identity to any  $C \in \mathcal{B}(D[-\alpha, 0])$ , that is, we have proved (10).

We show that  $P_{s,t}$  maps  $B_b(D[-\alpha, 0])$  into  $B_b(D[-\alpha, 0])$ . Indeed, if  $f \in C_b$ , then Corollary 6.4 yields that  $P_{s,t}f \in C_b$ . If  $C$  is a closed subset of  $D[-\alpha, 0]$ , then there are  $f_n \in C_b$  such that  $f_n \downarrow \mathbf{1}_C$  pointwise and then  $P_{s,t}f_n \downarrow P_{s,t}\mathbf{1}_C$  pointwise, so  $P_{s,t}\mathbf{1}_C \in B_b(D[-\alpha, 0])$ . By a monotone class argument we obtain  $P_{s,t}\mathbf{1}_C \in B_b(D[-\alpha, 0])$  for any  $F \in \mathbb{B}(D[-\alpha, 0])$  and then it follows that  $P_{s,t}f \in B_b(D[-\alpha, 0])$  for any  $f \in B_b(D[-\alpha, 0])$ .

The Markov property (10) yields for  $0 \leq u \leq s \leq t$  that

$$\begin{aligned} P_{u,t}f(\varphi) &= \mathbb{E}f(X_{\varphi,t-u}^u) = \mathbb{E}(\mathbb{E}[f(X_{\varphi,t-u}^u)|\mathcal{F}_s]) \\ &= \mathbb{E}(\mathbb{E}[f(X_{X_{\varphi,s-u}^u,t-s}^s)|\mathcal{F}_s]) = \mathbb{E}(\mathbb{E}f(X_{\psi,t-s}^s)|X_{\varphi,s-u}^u=\psi) \\ &= P_{u,s}P_{s,t}f(\varphi). \end{aligned}$$

By uniqueness in law [14], Section IX.6c, we have that  $(X_{\varphi,t}^u)_{t \geq 0}$  has the same law as  $(X_{\varphi,t})_{t \geq 0}$  for each  $u \geq 0$ , since  $L^u$  and  $L$  have the same law. Hence  $P_{s,t} = P_{0,t-s}$ .

Finally, we establish that  $(P_t)_t$  is eventually Feller. By Proposition 6.2 we have that for each  $t \geq \alpha$ ,  $\varphi_n \rightarrow \varphi$  in  $D[-\alpha, 0]$  implies  $X_{\varphi_n,t} \rightarrow X_{\varphi,t}$  in  $D[-\alpha, 0]$  in probability, so  $P_t f(\varphi_n) \rightarrow P_t f(\varphi)$ , hence (13) holds. Property (14) has already been shown.  $\square$

**Remark 7.2.** *The condition (17) is rather mild as the following examples show:*

- (1) *Let  $\rho$  be a finite signed Borel measure on  $[-\alpha, 0]$  and let  $f$  be a locally Lipschitz function on  $\mathbb{R}$  with linear growth. Define*

$$\begin{aligned} F(x)(t) &:= f\left(\int_{[-\alpha,0]} x(t+v)\rho(dv)\right), \quad t \geq 0, \\ F(x)(t) &:= 0, \quad -\alpha \leq t \leq 0, \end{aligned}$$

*$x \in D[-\alpha, \infty)$ . Then if  $\varphi_n \rightarrow \varphi$  in  $D[-\alpha, 0]$  and  $x \in D[0, \infty)$ ,*

$$\begin{aligned} &\int_0^\alpha (F(x_{-\varphi}(t)) - F(x_{-\varphi_n}(t)))^2 dt \\ &= \int_0^\alpha \left( f\left(\int_{[-\alpha,0]} x_{-\varphi_n}(t+v)\rho(dv)\right) - f\left(\int_{[-\alpha,0]} x_{-\varphi}(t+v)\rho(dv)\right) \right)^2 dt \\ &\leq C \int_0^\alpha \left( \int_{[-\alpha,0]} (x_{-\varphi_n}(t+v) - x_{-\varphi}(t+v))\rho(dv) \right)^2 dt \\ &= C \int_0^\alpha \left( \int_{[-\alpha,0]} (x_{-\varphi_n}(t+v) - x_{-\varphi}(t+v))\rho(dv) \right) \left( \int_{[-\alpha,0]} (x_{-\varphi_n}(t+u) - x_{-\varphi}(t+u))\rho(du) \right) dt \\ &= C \int_0^\alpha \int_{[-\alpha,0]} \int_{[-\alpha,0]} (\varphi_n(t+v) - \varphi(t+v))(\varphi_n(t+u) - \varphi(t+u)) \mathbf{1}_{u < -t} \mathbf{1}_{v < -t} \rho(du) \rho(dv) dt \end{aligned}$$

*for some  $C$  depending only on  $f$ ,  $x$ ,  $\rho$  and  $(\varphi_n)$ , where the latter equality follows by Fubini theorem and the fact that  $x_{-\varphi_n}(t+w) = x_{-\varphi}(t+w)$  whenever  $t+w \geq 0$ . Now by the Fubini theorem and dominated convergence  $F$  satisfies (17) for each  $x \in D[0, \infty)$ .*

*Moreover,  $F$  is lolidet: for  $t \geq 0$  and  $x, y \in D[-\alpha, \infty)$  such that  $\sup_{s \in [1-\alpha, t]} |x(s)| \vee |y(s)| \leq n$  we have*

$$\left| \int_{[-\alpha,0]} x(t+v)\rho(dv) \right|, \quad \left| \int_{[-\alpha,0]} y(t+v)\rho(dv) \right| \leq n|\rho|.$$

Hence as  $f$  is locally Lipschitz there is a  $C_n > 0$  such that

$$\begin{aligned} |F(x)(t) - F(y)(t)| &\leq C_n \left| \int_{[-\alpha, 0]} (x(t+v) - y(t+v)) \rho(dv) \right| \\ &\leq C_n |\rho| \sup_{v \in [-\alpha, 0]} |x(t+v) - y(t+v)|. \end{aligned}$$

Since  $f$  has linear growth, it follows that  $F$  is lolidet.

However,  $F$  need not be Lipschitz in the sense of [21], formula (2.5), if  $f$  is not Lipschitz. To see this, take  $f(t) = \sin(t^2)$ ,  $\rho$  the Lebesgue measure on  $[-\alpha, 0]$ , and evaluate  $F(x_n)(t) = \sin(\alpha^2 n^2)$ , where  $x_n \equiv n$ .

(2) Likewise we can take  $\rho_1, \dots, \rho_d$  signed Borel measures on  $[-\alpha, 0]$  and  $f$  a locally Lipschitz on  $\mathbb{R}^d$ . In particular, we may take for  $F$  combinations of finitely many point evaluations.

As above,  $F$  is lolidet but need not be Lipschitz in the sense of [21], formula (2.5).

(3) Let  $f$  be a locally Lipschitz function on  $\mathbb{R}$ . Let for  $x \in D[-\alpha, \infty)$ ,

$$F(x)(t) := f\left(\sup_{t-\alpha \leq s \leq t} |x(s)|\right) \quad \text{for } t \geq 0,$$

$$F(x)(t) := 0 \quad \text{for } -\alpha \leq t < 0.$$

Then if  $\varphi_n \rightarrow \varphi$  in  $D[-\alpha, 0]$ , for  $x \in D[0, \infty)$ ,

$$\int_0^\alpha (F(x_{\varphi_n})(t) - F(x_\varphi)(t))^2 dt \leq C \int_0^\alpha \left( \sup_{t-\alpha \leq s \leq t} |x_{\varphi_n}(s)| - \sup_{t-\alpha \leq s \leq t} |x_\varphi(s)| \right)^2 dt$$

for some  $C$  depending only on  $f, x$ , and  $(\varphi_n)$ , hence by bounded convergence the last expression above will tend to zero if for a.e.  $t \in [0, \alpha]$

$$\sup_{t-\alpha \leq s \leq t} |x_{\varphi_n}(s)| - \sup_{t-\alpha \leq s \leq t} |x_\varphi(s)| \rightarrow 0. \tag{20}$$

Let us show that (20) holds for  $t$  such that  $\varphi$  is continuous at  $t - \alpha$ . Let  $(\lambda_n)$  be a sequence of increasing homeomorphisms on  $[-\alpha, 0]$  such that  $\|\varphi_n - \varphi \circ \lambda_n\|_\infty + \|\lambda_n - I\|_\infty \rightarrow 0$ . Then

$$\begin{aligned} &\left| \sup_{t-\alpha \leq s \leq 0} |\varphi_n(s)| - \sup_{t-\alpha \leq s \leq 0} |\varphi(s)| \right| \\ &\leq \left| \sup_{t-\alpha \leq s \leq 0} |\varphi_n(s)| - \sup_{t-\alpha \leq s \leq 0} |\varphi \circ \lambda_n(s)| \right| + \left| \sup_{t-\alpha \leq s \leq 0} |\varphi \circ \lambda_n(s)| - \sup_{t-\alpha \leq s \leq 0} |\varphi(s)| \right|. \end{aligned}$$

The first term converges to 0 as  $\|\varphi_n - \varphi \circ \lambda_n\|_\infty$ . For the second term, let  $\varepsilon > 0$  and let  $\delta > 0$  be such that  $t - \alpha - \delta < s < t - \alpha + \delta$  implies  $\varphi(t - \alpha) - \varepsilon < \varphi(s) < \varphi(t - \alpha) + \varepsilon$ . Fix  $N$  such that  $\|\lambda_n - I\|_\infty < \delta$  for all  $n \geq N$ . Then

$$\sup_{t-\alpha \leq s \leq 0} |\varphi \circ \lambda_n(s)| = \sup_{\lambda_n(t-\alpha) \leq s \leq 0} |\varphi(s)| \leq \sup_{t-\alpha-\delta \leq s \leq 0} |\varphi(s)| \leq \sup_{t-\alpha \leq s \leq 0} |\varphi(s)| + \varepsilon$$

and

$$\sup_{t-\alpha \leq s \leq 0} |\varphi(s)| - 2\varepsilon \leq \sup_{t-\alpha+\delta \leq s \leq 0} |\varphi(s)| \leq \sup_{\lambda_n(t-\alpha) \leq s \leq 0} |\varphi(s)|,$$

so that the second term is less than  $2\varepsilon$  whenever  $n \geq N$ . Now since  $\sup_{t-\alpha \leq s \leq t} |x_\varphi(s)| = \sup_{t-\alpha \leq s \leq 0} |\varphi(s)| \vee \sup_{0 \leq s \leq t} |x(s)|$  and the same with  $\varphi_n$ , (20) holds if  $\varphi$  is continuous at  $t - \alpha$ . Hence  $F$  satisfies (17) for each  $x \in D[0, \infty)$ .

If  $x, y \in D[-\alpha, \infty)$  are bounded by  $n$  on  $[t - \alpha, t]$ , then  $|\sup_{[t-\alpha, t]} |x(s)| - \sup_{[t-\alpha, t]} |y(s)|| \leq n$ , so by the assumption on  $f$  there is a  $C_n > 0$  such that

$$|F(X)(t) - F(y)(t)| \leq C_n \left| \sup_{[t-\alpha, t]} |x(s)| - \sup_{[t-\alpha, t]} |y(s)| \right|.$$

As  $\sup_{s \in [t-\alpha, t]} |x(s)| \leq \sup_{s \in [t-\alpha, t]} |x(s) - y(s)| + \sup_{s \in [t-\alpha, t]} |y(s)|$ , we obtain by symmetry  $|\sup_{s \in [t-\alpha, t]} |x(s)| - \sup_{s \in [t-\alpha, t]} |y(s)|| \leq \sup_{s \in [t-\alpha, t]} |x(s) - y(s)|$ . Since  $f$  has linear growth, it follows that  $F$  is lolidet. Again,  $F$  need not be Lipschitz in the sense of [21], formula (2.5), if  $f$  is not Lipschitz, as we see by taking  $f(t) = \sin(t^2)$  and evaluating  $F$  on the sequence  $x_n \equiv n$ .

(4) Similar arguments as in (3) can be given for functionals like  $f(\sup_{t-\alpha \leq s \leq t} x(s))$ ,  $f(\inf_{t-\alpha \leq s \leq t} x(s))$  and  $f(\inf_{t-\alpha \leq s \leq t} |x(s)|)$ .

Notice that all functionals  $F$  in the previous remark are autonomous in the sense of Definition 4.4. If  $f$  is bounded, then  $F$  satisfies all conditions of Assumption 1 and (17).

Finally, we consider existence of an invariant measure. Denote by  $\mathcal{P}$  the set of Borel probability measures on  $D[-\alpha, 0]$  endowed with the topology of weak convergence of measures. Let  $B_b$  denote the space of all real valued bounded Borel functions on  $D[-\alpha, 0]$  and denote  $\langle \zeta, f \rangle = \int f d\zeta$ ,  $f \in B_b$ ,  $\zeta \in \mathcal{P}$ . The adjoint of the Markov semigroup defined in (12) is given by

$$\langle P_t^* \zeta, f \rangle = \langle \zeta, P_t f \rangle, \quad f \in B_b, \zeta \in \mathcal{P}.$$

A measure  $\eta \in \mathcal{P}$  is called an *invariant measure* for (1) if

$$P_t^* \eta = \eta \quad \text{for all } t \geq 0.$$

If  $\eta$  is the distribution of an initial segment  $\Phi$ , then  $P_t^* \eta$  is the distribution of the segment  $X_{t, \Phi}$ . Therefore if  $\Phi$  is an  $\mathcal{F}_0$ -measurable random variable with values in  $D[-\alpha, 0]$  whose law is an invariant measure, the segment process corresponding to the solution  $X$  of (1) with initial condition  $\Phi$  is constant in law. In this case the solution  $X$  itself is also constant in law.

**Theorem 7.3.** *Grant Assumption 1 and assume that (17) holds for every  $x \in D[0, \infty)$ . Then Eq. (1) has an invariant measure.*

**Proof.** It follows from Theorem 7.1 that  $P_t$  maps  $C_b = C_b(D[-\alpha, 0])$  into  $C_b$  for  $t \geq \alpha$  and that  $t \mapsto P_t^* \zeta$  is a continuous map from  $[\alpha, \infty)$  to  $\mathcal{P}$ . Moreover,  $P_{s+t}^* = P_s^* P_t^*$  for all  $s, t \geq 0$ . Theorem 5.1 yields that the set  $\{P_t^* \zeta : t \geq \alpha\}$  is tight, where, for instance,  $\zeta$  is the distribution of the initial condition  $\varphi \equiv 0$ .

Next, proceeding as in [21], Section 4.2, the invariant measure  $\eta$  is obtained by means of the Krylov–Bogoliubov method. □

**Example 7.4.** *Let us illustrate by means of an example how transformation of an equation with unbounded globally Lipschitz coefficients may lead to an equation with bounded coefficients that are locally but not globally Lipschitz. Consider the equation*

$$dX(t) = -ax(t) dt + f(X_t) dt + g(X_t) dW(t), \quad t \geq 0,$$

where  $a > 0, \alpha > 0$ , and  $f, g : D[-\alpha, 0] \rightarrow \mathbb{R}$  are Lipschitz with respect to  $\|\cdot\|_\infty$  and continuous with respect to the Skorohod metric  $d_\beta$  and such that

$$|f(y)| \vee |g(y)| \leq C(1 + |y(0)|^r) \quad \text{for all } y \in D[-\alpha, 0]$$

for some  $C > 0$  and  $0 < r < 1$ . The process  $W$  is a real valued Wiener process. With a sufficiently smooth strictly increasing  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\phi(x) = \text{sgn}(x)|x|^s \quad \text{for large } |x|$$

for some  $0 < s < 1 - r$ , the process

$$Y(t) = \phi(X(t))$$

satisfies

$$dY(t) = -asY(t) dt + \tilde{f}(Y_t) dt + \tilde{g}(Y_t) dW(t),$$

due to Ito's formula. The coefficients  $\tilde{f}$  and  $\tilde{g}$  are given by

$$\begin{aligned} \tilde{f}(y) &= asy(0) - as\phi'(\phi^{-1}(y(0)))\phi^{-1}(y(0)) + \phi'(\phi^{-1}(y(0)))f(t \mapsto \phi^{-1}(y(t))) \\ &\quad + \frac{1}{2}\phi''(\phi^{-1}(y(0)))g(t \mapsto \phi^{-1}(y(t)))^2, \\ \tilde{g}(y) &= \phi'(\phi^{-1}(y(0)))g(t \mapsto \phi^{-1}(y(t))) \end{aligned}$$

for  $y \in D[-\alpha, 0]$ . A computation reveals that  $\tilde{f}$  and  $\tilde{g}$  are bounded. Indeed, choose  $R \geq 1$  such that  $\phi(x) = \text{sgn}(x)|x|^s$  for  $|x| \geq R^{1/s}$ . Then  $\phi'(\phi^{-1}(x))\phi^{-1}(x) = sx$ ,  $|\phi'(\phi^{-1}(x))| = s|x|^{1-1/s}$ , and  $|\phi''(\phi^{-1}(x))| = s(1-s)|x|^{1-2/s}$  for  $|x| \geq R$ , so that

$$|\phi'(\phi^{-1}(x))|(1 + |x|^{r/s}) \leq 2s \quad \text{and} \quad |\phi''(\phi^{-1}(x))|(1 + |x|^{r/s})^2 \leq 4s(1-s)$$

for  $|x| \geq R$  and therefore

$$\begin{aligned} |\tilde{f}(y)| &\leq |asy(0) - a\phi'(\phi^{-1}(y(0)))\phi^{-1}(y(0))| + C|\phi'(\phi^{-1}(y(0)))|(1 + |\phi^{-1}(y(0))|^r) \\ &\quad + \frac{1}{2}C|\phi''(\phi^{-1}(y(0)))|(1 + |\phi^{-1}(y(0))|^r)^2 \\ &\leq M \vee 2C(s + s(1-s)) \end{aligned}$$

for all  $y \in D[-\alpha, 0]$ , where

$$\begin{aligned} M &= \sup_{|x| \leq R} |asx - a\phi'(\phi^{-1}(x))\phi^{-1}(x)| + C|\phi'(\phi^{-1}(x))|(1 + |\phi^{-1}(x)|^r) \\ &\quad + \frac{1}{2}C|\phi''(\phi^{-1}(x))|(1 + |\phi^{-1}(x)|^r)^2, \end{aligned}$$

and, similarly,

$$|\tilde{g}(y)| \leq M \vee 2s \quad \text{for all } y \in D[-\alpha, 0].$$

It easily follows from the local Lipschitz continuity of  $f$ ,  $g$ ,  $\phi'$ ,  $\phi''$  and  $\phi^{-1}$  that  $\tilde{f}$  and  $\tilde{g}$  are locally Lipschitz continuous. Since  $\phi^{-1}$  is not Lipschitz,  $\tilde{f}$  and  $\tilde{g}$  need not be globally Lipschitz, even if  $f$  and  $g$  are globally Lipschitz.

## References

- [1] D. Applebaum. *Lévy Processes and Stochastic Calculus*. Cambridge Univ. Press, Cambridge, 2004. [MR2072890](#)
- [2] J. Appleby and X. Mao. Stochastic stabilisation of functional differential equations. *Systems Control Lett.* **54** (2005) 1069–1081. [MR2170288](#)
- [3] P. Billingsley. *Convergence of Probability Measures*, 2nd edition. Wiley, New York, 1999. [MR1700749](#)
- [4] S. Cerrai. Stochastic reaction-diffusion systems with multiplicative noise and non-Lipschitz reaction term. *Probab. Theory Related Fields* **125** (2003) 271–304. [MR1961346](#)
- [5] F. Confortola. Dissipative backward stochastic differential equations with locally Lipschitz nonlinearity. *Stochastic Process. Appl.* **117** (2007) 613–628. [MR2320952](#)
- [6] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. Cambridge Univ. Press, Cambridge, 1992. [MR1207136](#)
- [7] M. C. Delfour. The largest class of hereditary systems defining a  $C_0$  semigroup on the product space. *Canad. J. Math.* **32** (1980) 969–978. [MR0590659](#)

- [8] O. Diekmann, S. A. van Gils, S. M. Verduyn Lunel and H.-O. Walther. *Delay Equations. Functional-, Complex-, and Nonlinear Analysis*. Springer, New York, 1995. [MR1345150](#)
- [9] A. Es-Sarhir and W. Stannat. Invariant measures for semilinear SPDE's with local Lipschitz drift coefficients and applications. *J. Evol. Equ.* **8** (2008) 129–154. [MR2383485](#)
- [10] A. Gushchin and U. Küchler. On stationary solutions of delay differential equations driven by a Lévy process. *Stochastic Process. Appl.* **88** (2000) 195–211. [MR1767844](#)
- [11] J. Hale. *Theory of Functional Differential Equations*. Springer, New York, 1977. [MR0508721](#)
- [12] E. Hausenblas. SPDEs driven by Poisson random measures with non Lipschitz coefficients: Existence results. *Probab. Theory Related Fields* **137** (2007) 161–200. [MR2278455](#)
- [13] J. Jacod and J. Memin. Weak and strong solutions of stochastic differential equations: Existence and stability. In *Stochastic Integrals, Proceedings, LMS Durham Symposium, 1980* 169–212. D. Williams (Ed.). *Springer LNM* **851**. Springer, Berlin, 1981. [MR0620991](#)
- [14] J. Jacod and A. N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer, Berlin, 2003. [MR1943877](#)
- [15] X. Mao. *Stochastic Differential Equations and Their Applications*. Horwood Publishing, Chichester, 1997. [MR1475218](#)
- [16] S.-E. A. Mohammed. *Stochastic Functional Differential Equations*. Pitman Advanced Publishing Program, Melbourne, 1984. [MR0754561](#)
- [17] S.-E. A. Mohammed and M. K. R. Scheutzow. Lyapunov exponents of linear stochastic functional differential equations. II: Examples and case studies. *Ann. Probab.* **25** (1997) 1210–1240. [MR1457617](#)
- [18] L. Mytnik, E. Perkins and A. Sturm. On pathwise uniqueness for stochastic heat equations with non-Lipschitz coefficients. *Ann. Probab.* **34** (2006) 1910–1959. [MR2271487](#)
- [19] M. Ondreját. Uniqueness for stochastic non-linear wave equations. *Nonlinear Anal.* **67** (2007) 3287–3310. [MR2350886](#)
- [20] P. E. Protter. *Stochastic Integration and Differential Equations*, 2nd edition. Springer, Berlin, 2005. [MR2273672](#)
- [21] M. Reiß, M. Riedle and O. van Gaans. Delay differential equations driven by Lévy processes: Stationarity and Feller properties. *Stochastic Process. Appl.* **116** (2006) 1409–1432. [MR2260741](#)
- [22] M. Reiß, M. Riedle and O. van Gaans. On Émery's inequality and a variation-of-constants formula. *Stoch. Anal. Appl.* **25** (2007) 353–379. [MR2303092](#)
- [23] R. Stelzer. Multivariate COGARCH(1, 1) processes. *Bernoulli* **16** (2010) 80–115. [MR2648751](#)
- [24] N. N. Vakhania, V. I. Tarieladze and S. A. Chobanyan. *Probability Distributions on Banach Spaces*. D. Reidel Publishing Company, Dordrecht, 1987. [MR1435288](#)
- [25] S. J. Wolfe. On a continuous analogue of the stochastic difference equation  $X_n = \rho X_{n-1} + B_n$ . *Stochastic Process. Appl.* **12** (1982) 301–312. [MR0656279](#)