

## Prediction of future failures for generalized exponential distribution under Type-I or Type-II hybrid censoring

R. Valiollahi<sup>a</sup>, A. Asgharzadeh<sup>b</sup> and D. Kundu<sup>c</sup>

<sup>a</sup>*Semnan University*

<sup>b</sup>*University of Mazandaran*

<sup>c</sup>*Indian Institute of Technology*

**Abstract.** In this paper, we consider the prediction of a future observation based on either Type-I or Type-II hybrid censored samples when the lifetime distribution of the experimental units is assumed to be a generalized exponential random variable. Different point and interval predictors are obtained using classical and Bayesian approaches. Monte Carlo simulations are performed to compare the performances of the different methods, and the analysis of one data set has been presented for illustrative purposes.

### 1 Introduction

The generalized exponential (GE) distribution is one of the most widely used distributions in reliability and survival analysis. It has been used quite successfully to analyze lifetime data in place of Weibull or gamma distribution. Because of its explicit expression of the cumulative distribution function (CDF), it has been used quite conveniently for analyzing censored data. The probability density function (PDF) and hazard function can take variety of shapes. The PDF of the GE distribution can be a decreasing or an unimodal function, and the hazard function can be either increasing, decreasing or constant, depending on the shape parameter.

In this manuscript, it is assumed that the two-parameter generalized exponential distribution has the following PDF and CDF

$$f(x; \alpha, \lambda) = \begin{cases} \alpha \lambda e^{-\lambda x} (1 - e^{-\lambda x})^{\alpha-1} & \text{if } x > 0, \\ 0 & \text{otherwise} \end{cases} \quad (1.1)$$

and

$$F(x; \alpha, \lambda) = \begin{cases} (1 - e^{-\lambda x})^\alpha & \text{if } x > 0, \\ 0 & \text{otherwise,} \end{cases} \quad (1.2)$$

respectively. Here  $\alpha > 0$  is the shape parameter and  $\lambda > 0$  is the scale parameter. From now on, a two-parameter generalized exponential distribution with the PDF (1.1) and CDF (1.2) will be denoted by  $GE(\alpha, \lambda)$ .

Censoring is very common in any reliability or lifetesting experiment. The complete survival times may not be observed by the experimenter most of times. The

---

*Key words and phrases.* Type-I hybrid censoring, Type-II hybrid censoring, generalized exponential distribution, predictor, prediction interval, Monte Carlo simulation.

Received August 2015; accepted October 2015.

two most common censoring schemes are namely Type-I and Type-II censoring. They can be briefly described as follows. Consider a sample of  $n$  units is placed on a life testing experiment at the time point zero. In Type-I censoring scheme, a time  $T$ , independent of the failure times, is pre-fixed so that beyond this time no failures will be observed, that is, the experiment terminates at the time point  $T$ . In Type-II censoring scheme, the number of observed failures is fixed, say  $r$  ( $r \leq n$ ), and the experiment stops when the  $r$ th failure takes place. Epstein (1954) introduced a new censoring scheme which is a mixture of Type-I and Type-II censoring schemes, and called it as the hybrid censoring scheme. From now on, we will call it as the Type-I hybrid censoring scheme (Type-I HCS). Type-I hybrid censoring scheme can be described as follows. Suppose  $n$  items are put on a life testing experiment at the time point zero. Let  $X_{1:n} < \dots < X_{n:n}$  denote the ordered lifetime of the experimental units. In Type-I HCS, the experiment stops at  $T_1^* = \min\{X_{r:n}, T\}$ , where  $r$  and  $T$  are pre-fixed.

The main advantage of Type-I HCS is that the experiment will not last more than  $T$  units of time, and also more than  $r$  units will not fail. The major disadvantage of a Type-I HCS is that there is a possibility that very few failures may occur before time  $T$ . Hence, the inference of the lifetime distribution of the experimental units will not be very accurate. For this reason, Childs et al. (2003) proposed a new censoring scheme, known as the Type-II hybrid censoring scheme (Type-II HCS), which guarantees a fixed number of failures. In this case, the termination point is  $T_2^* = \max\{X_{r:n}, T\}$ . For a detailed comparison of Type-I and Type-II HCS, the readers are referred to Childs et al. (2003).

Recently, the hybrid censoring scheme has received considerable interest among the statisticians. Epstein (1954) first introduced Type-I HCS and analyzed the data under the assumption of exponential lifetime distribution of the experimental units. Since the introduction of Type-I HCS by Epstein (1954), extensive work has been done on hybrid censoring and many different variations of it. See, for example, Fairbanks, Madison and Dykstra (1982), Draper and Guttman (1987), Chen and Bhattacharya (1988), Ebrahimi (1986, 1992), Jeong, Park and Yum (1996), Kundu and Gupta (1988), Childs et al. (2003), Kundu (2007), Kundu and Banerjee (2008) and the recent review article by Balakrishnan and Kundu (2013) on this topic.

One of the most important problems in life testing is predicting future failures given a record of observed failures. Information regarding future observations can tell us at an early stage of testing how costly the testing is and whether actions should be taken to redesign the test. Extensive work has been done on prediction problem based on frequentist and Bayesian framework. Kaminsky and Rhodin (1985) have applied maximum likelihood to the joint prediction of a future random variable and unknown parameter. Smith (1997, 1999) discussed the properties of the different predictors based on Bayes and frequentist procedures for a class of parametric family of distribution functions, under smooth loss functions. Dellaportas and Wright (1991) used a numerical method to Bayesian prediction for the two-parameter Generalized exponential distribution under the squared error loss function. Raqab, Asgharzadeh and Valiollahi (2010), Balakrishnan, Beut-

ner and Cramer (2010) and Basak, Basak and Balakrishnan (2006) have discussed different predictors of times to failure of units censored in progressively Type-II censored samples. Kundu and Raqab (2012) considered the prediction of future observation from a Type-II censored data for the two-parameter Weibull distribution under fairly flexible priors on the shape and scale parameters. Recently, Asgharzadeh, Valiollahi and Kundu (2015) discussed the prediction of future observation based on a Type-I hybrid censored data from the two-parameter Weibull distribution.

The main aim of this paper is to consider the prediction of future observation based on Type-I or Type-II hybrid censored samples from a two-parameter Generalized exponential distribution. We obtain the maximum likelihood predictor, best unbiased predictor, conditional median predictor and Bayesian predictor. We also obtain different prediction intervals. It is difficult to compare the performances of the different methods theoretically. We perform some Monte Carlo simulations to compare the performances of the different methods. Finally one data analysis has been performed for illustrative purposes.

Rest of the paper is organized as follows. In Section 2, we provide the preliminaries and notations. Different classical predictors are presented in Section 3. In Section 4, we discuss Bayesian prediction of future observations. In Section 5, one data analysis has been presented. Monte Carlo simulation results are presented in Section 6. Finally, we conclude the paper in Section 7.

## 2 Notation and preliminaries

Consider  $\mathbf{X} = (X_{1:n}, X_{2:n}, \dots, X_{D:n})$  as a sample from model (1.1) which is obtained based on a Type-I HCS or Type-II HCS. Here  $D$  is the number of failures during the experiment. In Type-I HCS,  $D = r$  if  $X_{r:n} \leq T$ , and  $D = J < r$  if  $X_{r:n} > T$ . For Type-II HCS,  $D = r$  if  $X_{r:n} \geq T$ , and  $D = J \geq r$  if  $X_{r:n} < T$ . In both the cases,  $J$  is determined such that  $X_{J:n} < T < X_{J+1:n}$  where  $J = 0, \dots, n$ .

For notational simplicity, we will write  $(X_1, X_2, \dots, X_D)$  for  $(X_{1:n}, X_{2:n}, \dots, X_{D:n})$ . Based on the observed data, the likelihood function for  $\alpha$  and  $\lambda$  without the normalizing constant is

$$L(\alpha, \lambda) = \alpha^D \lambda^D e^{-\lambda \sum_{i=1}^D x_i} [1 - (1 - e^{-\lambda T_0})^\alpha]^{n-D} \left\{ \prod_{i=1}^D (1 - e^{-\lambda x_i})^{\alpha-1} \right\}, \quad (2.1)$$

where  $T_0$  denotes the time when the experiment is stopped. In other words, for Type-I HCS,  $T_0 = X_{r:n}$  if  $X_{r:n} \leq T$  and  $T_0 = T$  if  $X_{r:n} > T$ . For Type-II HCS,  $T_0 = X_{r:n}$  if  $X_{r:n} \geq T$  and  $T_0 = T$  if  $X_{r:n} < T$ .

From (2.1), the maximum likelihood estimators (MLEs) of  $\alpha$  and  $\lambda$  can be obtained as solutions of the following equations

$$\frac{D}{\alpha} + \sum_{i=1}^D \ln(1 - e^{-\lambda x_i}) - (n - D) \frac{(1 - e^{-\lambda T_0})^\alpha \ln(1 - e^{-\lambda T_0})}{1 - (1 - e^{-\lambda T_0})^\alpha} = 0, \quad (2.2)$$

and

$$\frac{D}{\lambda} - \sum_{i=1}^D x_i + (\alpha - 1) \frac{x_i e^{-\lambda x_i}}{1 - e^{-\lambda x_i}} - (n - D) \frac{\alpha T_0 e^{-\lambda T_0} (1 - e^{-\lambda T_0})^{\alpha-1}}{1 - (1 - e^{-\lambda T_0})^\alpha}. \quad (2.3)$$

The main aim of this paper is to discuss different methods of prediction of  $Y = X_{s+D:n}$  ( $s = 1, 2, \dots, n - D$ ) of all the  $n - D$  censored units based on observed data  $\mathbf{X} = (X_1, \dots, X_D)$ . Due to the Markovian property of censored-order statistics, the conditional distribution of  $Y$  given  $\mathbf{X} = \mathbf{x}$  is just the distribution of  $X_{s+D:n}$  given  $X_D = x_D$ . This implies that the density of  $Y$  given  $\mathbf{X} = \mathbf{x}$  is the same as the density of the  $s$ th order statistic out of  $n - D$  units from the population with density  $f(y)/(1 - F(T_0))$ ,  $y \geq T_0$  (left truncated density at  $T_0$ ). Therefore, the conditional density of  $Y = X_{s+D:n}$  given  $\mathbf{X} = \mathbf{x}$ , for  $y \geq T_0$ , is given by

$$f(y|\mathbf{x}) = s \binom{n-D}{s} f(y) [F(y) - F(T_0)]^{s-1} \times [1 - F(y)]^{n-D-s} [1 - F(T_0)]^{-(n-D)}, \quad (2.4)$$

where  $s = 1, 2, \dots, n - D$ . For model (1.1), (2.4) reduces to

$$f(y|\mathbf{x}, \alpha, \lambda) = s \binom{n-D}{s} \alpha \lambda e^{-\lambda y} (1 - e^{-\lambda y})^{\alpha-1} [(1 - e^{-\lambda y})^\alpha - (1 - e^{-\lambda T_0})^\alpha]^{s-1} \times [1 - (1 - e^{-\lambda y})^\alpha]^{n-D-s} [1 - (1 - e^{-\lambda T_0})^\alpha]^{-(n-D)}, \quad (2.5)$$

which is the conditional density of  $Y = X_{s+D:n}$  given  $\mathbf{X} = (X_1, \dots, X_D)$ .

### 3 Classical prediction of future observations

In this section, we obtain several predictors of  $Y = X_{s+D:n}$  on the basis of  $\mathbf{X} = (X_1, \dots, X_D)$  via classical approaches.

#### 3.1 Likelihood prediction approach

In likelihood prediction approach, the principle of maximum likelihood is applied to the joint prediction and estimation of a future random variable and unknown parameters. We assume dependence between present and future, and the approach is non-Bayesian. The predictive likelihood function (PLF) of  $Y$  and  $(\alpha, \lambda)$  is given by

$$L(y, \alpha, \lambda|\mathbf{x}) = f(y|\mathbf{x}, \alpha, \lambda) f(\mathbf{x}|\alpha, \lambda). \quad (3.1)$$

Suppose  $\hat{Y} = u(\mathbf{X})$ ,  $\hat{\alpha} = v_1(\mathbf{X})$  and  $\hat{\lambda} = v_2(\mathbf{X})$  are statistics for which

$$L(u(\mathbf{x}), v_1(\mathbf{x}), v_2(\mathbf{x})|\mathbf{x}) = \sup_{(y, \alpha, \lambda)} L(y, \alpha, \lambda|\mathbf{x}),$$

then  $u(\mathbf{X})$  is said to be the maximum likelihood predictor (MLP) of  $Y$  and  $v_1(\mathbf{x})$  and  $v_2(\mathbf{x})$  the predictive maximum likelihood estimators (PMLEs) of  $\alpha$  and  $\lambda$ , respectively.

For the GE model, using (2.1) and (2.5), the predictive likelihood function (PLF) of  $Y$ ,  $\alpha$  and  $\lambda$ , is given by

$$L(y, \alpha, \lambda) = s \binom{n-D}{s} (\alpha\lambda)^{d+1} (1 - e^{-\lambda y})^{\alpha-1} \left\{ \prod_{i=1}^D (1 - e^{-\lambda x_i})^{\alpha-1} \right\} \\ \times e^{-\lambda[y + \sum_{i=1}^D x_i]} [(1 - e^{-\lambda y})^\alpha - (1 - e^{-\lambda T_0})^\alpha]^{s-1} \quad (3.2) \\ \times [1 - (1 - e^{-\lambda y})^\alpha]^{n-D-s}.$$

Apart from a constant term, the predictive log-likelihood function is

$$\ln L(y, \alpha, \lambda) = (D + 1)[\ln(\alpha) + \ln(\lambda)] \\ + (\alpha - 1) \left[ \ln[1 - e^{-\lambda y}] + \sum_{i=1}^D \ln[1 - e^{-\lambda x_i}] \right] \quad (3.3) \\ - \lambda \left[ y + \sum_{i=1}^D x_i \right] + (s - 1) \ln[(1 - e^{-\lambda y})^\alpha - (1 - e^{-\lambda T_0})^\alpha] \\ + (n - D - s) \ln[1 - (1 - e^{-\lambda y})^\alpha].$$

By using (3.3), the predictive likelihood equations (PLEs) for  $y$ ,  $\alpha$  and  $\lambda$  are given, respectively, by

$$\frac{\partial \ln L(y, \alpha, \lambda)}{\partial y} \\ = -\lambda + (\alpha - 1) \frac{\lambda e^{-\lambda y}}{1 - e^{-\lambda y}} + (s - 1) \frac{\alpha \lambda e^{-\lambda y} (1 - e^{-\lambda y})^{\alpha-1}}{(1 - e^{-\lambda y})^\alpha - (1 - e^{-\lambda T_0})^\alpha} \quad (3.4) \\ - (n - D - s) \frac{\alpha \lambda e^{-\lambda y} (1 - e^{-\lambda y})^{\alpha-1}}{1 - (1 - e^{-\lambda y})^\alpha}$$

$$= 0,$$

$$\frac{\partial \ln L(y, \alpha, \lambda)}{\partial \alpha} \\ = \ln[1 - e^{-\lambda y}] + \sum_{i=1}^D \ln[1 - e^{-\lambda x_i}] \\ - (n - D - s) \frac{(1 - e^{-\lambda y})^\alpha \ln[1 - e^{-\lambda y}]}{1 - (1 - e^{-\lambda y})^\alpha} + \frac{D + 1}{\alpha} \quad (3.5)$$

$$\begin{aligned}
& + (s-1) \frac{(1-e^{-\lambda y})^\alpha \ln[1-e^{-\lambda y}] - (1-e^{-\lambda T_0})^\alpha \ln[1-e^{-\lambda T_0}]}{(1-e^{-\lambda y})^\alpha - (1-e^{-\lambda T_0})^\alpha} \\
& = 0, \\
& \frac{\partial \ln L(y, \alpha, \lambda)}{\partial \lambda} \\
& = \frac{D+1}{\lambda} - \left[ y + \sum_{i=1}^D x_i \right] + (\alpha-1) \left[ \frac{ye^{-\lambda y}}{1-e^{-\lambda y}} + \sum_{i=1}^D \frac{x_i e^{-\lambda x_i}}{1-e^{-\lambda x_i}} \right] \\
& + (s-1) \frac{\alpha y e^{-\lambda y} (1-e^{-\lambda y})^{\alpha-1} - \alpha T_0 e^{-\lambda T_0} (1-e^{-\lambda T_0})^{\alpha-1}}{(1-e^{-\lambda y})^\alpha - (1-e^{-\lambda T_0})^\alpha} \quad (3.6) \\
& - (n-D-s) \frac{\alpha y e^{-\lambda y} (1-e^{-\lambda y})^{\alpha-1}}{1 - (1-e^{-\lambda y})^\alpha} \\
& = 0.
\end{aligned}$$

From (3.4) and (3.6), we obtain the MLP of  $Y$  as

$$\begin{aligned}
\hat{Y}_{\text{MLP}} = & -\frac{1}{\tilde{\lambda}} \ln \left[ 1 - \left( \frac{(s-1)\tilde{\alpha} T_0 e^{-\tilde{\lambda} T_0} (1-e^{-\tilde{\lambda} T_0})^{\tilde{\alpha}-1}}{(D+1)/\tilde{\lambda} - \sum_{i=1}^D x_i (1-\tilde{\alpha} e^{-\tilde{\lambda} x_i}) / (1-e^{-\tilde{\lambda} x_i})} \right. \right. \\
& \left. \left. + (1-e^{-\tilde{\lambda} T_0})^{\tilde{\alpha}} \right)^{1/\tilde{\alpha}} \right], \quad (3.7)
\end{aligned}$$

where  $(\tilde{\alpha}, \tilde{\lambda})$  is PMLE of  $(\alpha, \lambda)$  that can be obtained numerically from (3.5) and (3.6).

### 3.2 Conditional prediction approach

In conditional prediction approach, the conditional distribution of  $Y = X_{s+D:n}$  given  $\mathbf{X} = (X_1, \dots, X_D)$  is applied to derive the predictors of  $Y$ . A statistic  $\hat{Y}$  which is used to predict  $Y = X_{s+D:n}$  is called a best unbiased predictor (BUP) of  $Y$ , if the predictor error  $\hat{Y} - Y$  has a mean zero and its prediction error variance  $\text{Var}(\hat{Y} - Y)$  is less than or equal to that of any other unbiased predictor of  $Y$ .

The BUP of  $Y$  is

$$\hat{Y}_{\text{BUP}} = E(Y|\mathbf{X}) = \int_{T_0}^{\infty} y f(y|\mathbf{x}, \alpha, \lambda) dy.$$

Using (2.5) and the binomial expansion

$$\begin{aligned}
& [(1-e^{-\lambda y})^\alpha - (1-e^{-\lambda T_0})^\alpha]^{s-1} \\
& = \sum_{j=0}^{s-1} \binom{s-1}{j} (-1)^{s-j-1} [1 - (1-e^{-\lambda T_0})^\alpha]^j [1 - (1-e^{-\lambda y})^\alpha]^{s-j-1} \quad (3.8)
\end{aligned}$$

we obtain

$$\begin{aligned} \hat{Y}_{\text{BUP}} &\equiv I(T_0; \alpha, \lambda) \\ &= s \binom{n-D}{s} \sum_{j=0}^{s-1} \binom{s-1}{j} (-1)^{s-j-1} [1 - (1 - e^{-\lambda T_0})^\alpha]^{-(n-D-j)} h_1(T_0), \end{aligned} \quad (3.9)$$

where

$$\begin{aligned} h_1(T_0) &= \int_{T_0}^{\infty} y \alpha \lambda e^{-\lambda y} (1 - e^{-\lambda y})^{\alpha-1} [1 - (1 - e^{-\lambda y})^\alpha]^{n-D-j-1} dy \\ &= -\frac{\alpha}{\lambda} \int_0^{e^{-\lambda T_0}} \ln(t) (1-t)^{\alpha-1} [1 - (1-t)^\alpha]^{n-D-j-1} dt. \end{aligned} \quad (3.10)$$

Because the parameters  $\alpha$  and  $\lambda$  are unknown, they have to be estimated. Thus, one would replace them by their corresponding MLEs and obtain the BUP of  $Y$ .

Let us now consider conditional median predictor (CMP) which was first suggested by Raqab and Nagaraja (1995). A predictor  $\hat{Y}$  is called the CMP of  $Y$ , if it is the median of the conditional distribution of  $Y$  given  $\mathbf{X} = \mathbf{x}$ , that is

$$P_\theta(Y \leq \hat{Y} | \mathbf{X} = \mathbf{x}) = P_\theta(Y \geq \hat{Y} | \mathbf{X} = \mathbf{x}). \quad (3.11)$$

Using the relation

$$\begin{aligned} P_{\alpha,\lambda}(Y \leq \hat{Y} | \mathbf{X} = \mathbf{x}) \\ = P_{\alpha,\lambda} \left( 1 - \frac{1 - (1 - e^{-\lambda Y})^\alpha}{1 - (1 - e^{-\lambda T_0})^\alpha} \leq 1 - \frac{1 - (1 - e^{-\lambda \hat{Y}})^\alpha}{1 - (1 - e^{-\lambda T_0})^\alpha} \middle| \mathbf{X} = \mathbf{x} \right), \end{aligned} \quad (3.12)$$

and using the fact that the distribution of  $1 - \frac{1 - (1 - e^{-\lambda Y})^\alpha}{1 - (1 - e^{-\lambda T_0})^\alpha}$  given  $\mathbf{X} = \mathbf{x}$  is a Beta( $s, n - D - s + 1$ ) distribution, we obtain the CMP of  $Y$  as

$$\hat{Y}_{\text{CMP}} = -\frac{1}{\lambda} \ln[1 - (1 - [1 - \text{Med}(B)][1 - (1 - e^{-\lambda T_0})^\alpha])^{1/\alpha}], \quad (3.13)$$

where  $B$  has Beta( $s, n - D - s + 1$ ) distribution and  $\text{Med}(B)$  stands for median of  $B$ . By substituting  $\alpha$  and  $\lambda$  with their corresponding MLEs, we obtain the CMP of  $Y$ .

### 3.3 Classical prediction intervals

In this section, we construct two prediction intervals (PI's) for  $Y = X_s$  based on the Type I or Type-II censored sample  $\mathbf{X} = (X_1, X_2, \dots, X_D)$ .

Let us define the random variable  $Z$  as

$$Z = 1 - \frac{1 - (1 - e^{-\lambda Y})^\alpha}{1 - (1 - e^{-\lambda T_0})^\alpha}.$$

As mentioned before, the distribution of  $Z$  given  $\mathbf{X} = \mathbf{x}$  is a  $\text{Beta}(s, n - D - s + 1)$  distribution. So, we can consider  $Z$  as a pivotal quantity to obtain the prediction interval for  $Y$ . Now, a  $(1 - \gamma)100\%$  PI for  $Y$  is  $(L_1(\mathbf{X}), U_1(\mathbf{X}))$  where

$$L_1(\mathbf{X}) = -\frac{1}{\lambda} \ln[1 - (1 - [1 - B_{\gamma/2}][1 - (1 - e^{-\lambda T_0})^\alpha])^{1/\alpha}], \quad (3.14)$$

$$U_1(\mathbf{X}) = -\frac{1}{\lambda} \ln[1 - (1 - [1 - B_{1-\gamma/2}][1 - (1 - e^{-\lambda T_0})^\alpha])^{1/\alpha}], \quad (3.15)$$

where  $B_\gamma$  stands for  $100\gamma$ th percentile of  $\text{Beta}(s, n - D - s + 1)$  distribution. When  $\alpha$  and  $\lambda$  are unknown, the parameters in (3.14) and (3.15), have to be estimated. For example, by replacing  $\alpha$  and  $\lambda$  with their corresponding MLEs, the prediction limits for  $Y$  can be obtained.

Now let us consider another prediction interval for  $Y = X_s$ . The distribution of  $Z$  given  $\mathbf{X} = \mathbf{x}$  is a  $\text{Beta}(s, n - D - s + 1)$  distribution with PDF

$$g(z) = \frac{z^{s-1}(1-z)^{n-D-s}}{\text{Beta}(s, n - D - s + 1)}, \quad 0 < z < 1,$$

which is a unimodal function of  $z$ , for  $1 < s < n - D$ . Therefore, the  $(1 - \gamma)100\%$  highest conditional density (HCD) prediction limits for  $Y$  are given by

$$L_2(\mathbf{X}) = -\frac{1}{\lambda} \ln[1 - (1 - [1 - w_1][1 - (1 - e^{-\lambda T_0})^\alpha])^{1/\alpha}], \quad (3.16)$$

$$U_2(\mathbf{X}) = -\frac{1}{\lambda} \ln[1 - (1 - [1 - w_2][1 - (1 - e^{-\lambda T_0})^\alpha])^{1/\alpha}], \quad (3.17)$$

where  $w_1$  and  $w_2$  are the simultaneous solutions of the following equations:

$$\int_{w_1}^{w_2} g(z) dz = 1 - \gamma \quad (3.18)$$

and

$$g(w_1) = g(w_2). \quad (3.19)$$

Now, we simplify Equations (3.18) and (3.19) as

$$B_{w_2}(s, n - D - s + 1) - B_{w_1}(s, n - D - s + 1) = 1 - \gamma, \quad (3.20)$$

and

$$\left(\frac{1 - w_2}{1 - w_1}\right)^{n-D-s} = \left(\frac{w_1}{w_2}\right)^{s-1}, \quad (3.21)$$

where

$$B_t(a, b) = \frac{1}{B(a, b)} \int_0^t x^{a-1} (1-x)^{b-1} dx,$$

is the incomplete beta function. It is clear from (3.21), that the method cannot be used to construct the prediction interval when  $s = 1$ .

#### 4 Bayesian prediction of future observations

In this section, we obtain the Bayesian predictors of  $Y = X_{s+D}$  ( $s = 1, 2, \dots, n - D$ ) based on the observed hybrid censored sample  $\mathbf{x} = (x_1, \dots, x_D)$ . For finding Bayesian predictors, it is assumed that  $\alpha$  and  $\lambda$  each have independent  $\text{Gamma}(a_1, b_1)$  and  $\text{Gamma}(a_2, b_2)$  priors, respectively. Based on these priors, we can obtain the joint posterior density function of  $\alpha$  and  $\lambda$  given the data as

$$\pi(\alpha, \lambda|\mathbf{x}) \propto g_1(\alpha|\lambda, \mathbf{x})g_2(\lambda|\mathbf{x}). \quad (4.1)$$

Here  $g_2(\lambda|\mathbf{x})$  is a gamma density function with the shape and scale parameters as  $D + a_2$  and  $\sum_{i=1}^D x_i + b_2$ , respectively. Further,

$$g_1(\alpha|\lambda, \mathbf{x}) \propto \frac{e^{-b_1\alpha}\alpha^{D+a_1-1}}{(\sum_{i=1}^D x_i + b_2)^{D+a_2}} [1 - (1 - e^{-\lambda T_0})^\alpha]^{n-D} \prod_{i=1}^D (1 - e^{-\lambda x_i})^{\alpha-1}, \quad (4.2)$$

which is a proper density function.

The Bayesian predictors are obtained from the predictive density function. The predictive density of  $Y = X_{s+D}$  given  $\mathbf{x}$  is

$$f_s^*(y|\mathbf{x}) = \int_0^\infty \int_0^\infty f(y|\mathbf{x}, \alpha, \lambda)\pi(\alpha, \lambda|\mathbf{x}) d\alpha d\lambda, \quad (4.3)$$

where  $f(y|\mathbf{x}, \alpha, \lambda)$  is the conditional density of  $Y = X_{s+D}$  given  $\mathbf{x}$ . By replacing (4.1) in (4.3), we obtain the predictive density function  $f_s^*(y|\mathbf{x})$  as

$$f_s^*(y|\mathbf{x}) = \int_0^\infty \int_0^\infty f(y|\mathbf{x}, \alpha, \lambda)g_1(\alpha|\lambda, \mathbf{x})g_2(\lambda|\mathbf{x}) d\alpha d\lambda. \quad (4.4)$$

Now, the Bayesian point (BP) predictor of  $Y$  under a squared error loss is

$$\widehat{Y}_{BP} = \int_{T_0}^\infty y f_s^*(y|\mathbf{x}) dy. \quad (4.5)$$

It is not possible to compute (4.5) explicitly. Therefore, we propose here Gibbs sampling procedure to obtain the Bayes predictor. In the Gibbs sampling procedure, we can generate samples from the posterior density function  $\pi(\alpha, \lambda|\mathbf{x})$  and in turn obtain the Bayes predictor and also the corresponding prediction intervals based on the generated posterior samples. For later use, we need the following result.

**Theorem 1.** *The conditional distribution of  $\alpha$  given the data,  $g_1(\alpha|\lambda, \mathbf{x})$  is log-concave.*

**Proof.** See the [Appendix](#). □

Now, using Theorem 1 and following the idea of [Geman and Geman \(1984\)](#), we propose the following algorithm to generate  $(\alpha, \lambda)$  from the posterior density function.

1. Generate  $\lambda_1$  from  $g_2(\cdot|\mathbf{x})$ .
2. Generate  $\alpha_1$  from  $g_1(\cdot|\lambda_1, \mathbf{x})$  using the method proposed by Devroye (1984).
3. Repeat steps 1 and 2,  $M$  times and obtain  $(\alpha_1, \lambda_1), \dots, (\alpha_M, \lambda_M)$ .

Now, using the generated samples  $(\alpha_1, \lambda_1), \dots, (\alpha_M, \lambda_M)$ , the simulation consistent estimator of  $f_s^*(y|\mathbf{x})$  is

$$\widehat{f}_s^*(y|\mathbf{x}) = \frac{1}{M} \sum_{i=1}^M f(y|\mathbf{x}, \alpha_i, \lambda_i). \quad (4.6)$$

Therefore, by replacing (4.6) in (4.5), the BP of  $Y$  can be approximated as

$$\widehat{Y}_{\text{BP}} = \int_{T_0}^{\infty} y \left( \frac{1}{M} \sum_{i=1}^M f(y|\mathbf{x}, \alpha_i, \lambda_i) \right) dy \quad (4.7)$$

$$= \frac{1}{M} \sum_{i=1}^M \int_{T_0}^{\infty} y f(y|\mathbf{x}, \alpha_i, \lambda_i) dy \quad (4.8)$$

$$= \frac{1}{M} \sum_{i=1}^M I(T_0; \alpha_i, \lambda_i), \quad (4.9)$$

where  $I(T_0, \alpha, \lambda)$  is defined in (3.9).

Bayesian prediction intervals are obtained from the Bayes predictive density  $f^*(y|\mathbf{x})$ . For some positive  $u$ , the survivor function of  $Y$  given  $\mathbf{x}$  is

$$P(Y > u|\mathbf{x}) = \int_u^{\infty} f^*(y|\mathbf{x}) dy.$$

which by using (4.6) can be approximated as

$$\begin{aligned} P(Y > u|\mathbf{x}) &= \frac{1}{M} \sum_{i=1}^M \int_u^{\infty} f(y|\mathbf{x}, \alpha_i, \lambda_i) dy \\ &= s \binom{n-D}{s} \frac{1}{M} \sum_{i=1}^M \sum_{j=0}^{s-1} \binom{s-1}{j} \frac{(-1)^{s-j-1}}{n-D-j} \left[ \frac{1 - (1 - e^{-\lambda_i u})^{\alpha_i}}{1 - (1 - e^{-\lambda_i T_0})^{\alpha_i}} \right]^{n-D-j}. \end{aligned}$$

Therefore, the  $100(1 - \gamma)\%$  Bayesian prediction interval for  $Y$  is  $(L_3(\mathbf{X}), U_3(\mathbf{X}))$ , where the prediction limits  $L_3(\mathbf{X})$  and  $U_3(\mathbf{X})$  can be obtained by solving the non-linear equations

$$P(Y > L_3(\mathbf{x})|\mathbf{x}) = 1 - \frac{\gamma}{2}, \quad P(Y > U_3(\mathbf{x})|\mathbf{x}) = \frac{\gamma}{2}.$$

## 5 Real data analysis

In this section, we consider one real data set mainly to illustrate how the different methods perform when the underlying distribution is known to be GE. The following data below (Lawless, 1982, p. 267) are from an accelerator life test of 59 conductors. The data are

2.997 4.137 4.288 4.531 4.700 4.706 5.009 5.381 5.434  
 5.459 5.589 5.640 5.807 5.923 6.033 6.071 6.087 6.129  
 6.352 6.369 6.479 6.492 6.515 6.522 6.538 6.545 6.573  
 6.725 6.869 6.923 6.948 6.956 6.958 7.024 7.224 7.365  
 7.398 7.459 7.489 7.495 7.496 7.543 7.683 7.937 7.945  
 7.974 8.120 8.336 8.532 8.591 8.687 8.799 9.218 9.254  
 9.289 9.663 10.092 10.491 11.038.

A subset of the data set we will be using as samples (training data) and the rest will be used for verification purposes. Before progressing further, first we want to see whether GE model provides a good fit to the data set or not. We obtain the MLEs of the GE parameters as  $\hat{\alpha} = 52.411$  and  $\hat{\lambda} = 0.642$ . It is observed that the Kolmogorov–Smirnov (K–S) distance and the corresponding  $p$ -value are, respectively,

$$\text{K-S} = 0.103 \quad \text{and} \quad p = 0.532.$$

Since  $p$ -value is quite high, we can say that the GE provides a good fit to the data set.

**Scheme 1.** Let us consider the following sampling scheme:  $r = 20$  and  $T = 6$ . Based on Type-I HCS, the sample is:

2.997 4.137 4.288 4.531 4.700 4.706 5.009 5.381 5.434  
 5.459 5.589 5.640 5.807 5.923.

In this case  $D = 14$  and  $T_0 = 6$ . Now based on the above sample, we want to predict  $X_{(s)}$ , for different values of  $s$ , namely for  $s = 1, 3, 5, 8$ . The predicted values and the prediction intervals using different methods are computed and they are reported in Table 1. Note that for computing Bayesian predictions, since we do not have any prior information, we assumed that the priors on  $\alpha$  and  $\lambda$  are improper, that is,  $a_1 = b_1 = a_2 = b_2 = 0$ .

**Scheme 2.** In this case, we consider Type-II HCS with the same  $r = 20$  and  $T = 6$ . The observed sample becomes

2.997 4.137 4.288 4.531 4.700 4.706 5.009 5.381 5.434  
 5.459 5.589 5.640 5.807 5.923 6.033 6.071 6.087 6.129  
 6.352 6.369.

**Table 1** The values of point predictions and 95% PIs for  $Y = X_s$ 

	Exact value	Point prediction				Interval prediction		
		MLP	BUP	CMP	BP	Pivotal method	HCD method	Bayesian method
Type-I HCS								
$s = 1$	6.033	5.923	6.030	5.999	6.048	(5.926, 6.307)	–	(5.602, 6.352)
$s = 3$	6.087	6.131	6.241	6.213	6.301	(5.993, 6.651)	(5.958, 6.578)	(5.461, 6.725)
$s = 5$	6.352	6.334	6.448	6.423	6.604	(6.106, 6.937)	(6.067, 6.875)	(5.988, 7.205)
$s = 8$	6.492	6.633	6.966	6.734	7.203	(6.313, 7.331)	(6.275, 7.280)	(6.163, 7.464)
Type-II HCS								
$s = 1$	6.479	6.369	6.464	6.436	6.521	(6.372, 6.710)	–	(6.065, 6.929)
$s = 3$	6.515	6.557	6.653	6.626	6.852	(6.430, 7.028)	(6.403, 6.961)	(6.386, 7.094)
$s = 5$	6.538	6.745	6.843	6.819	7.162	(6.532, 7.299)	(6.498, 7.239)	(6.284, 7.744)
$s = 8$	6.725	7.030	7.156	7.112	7.802	(6.721, 7.685)	(6.690, 7.636)	(6.069, 8.785)

Here  $D = 20$  and  $T_0 = 6.369$ . Now based on the above Type-II HCS, the predicted value of  $X_{(s)}$  and the prediction intervals are reported in Table 1.

From Table 1, it is observed that different point predictors are quite close to the true observations. Also, we observe that different prediction intervals are containing the true observations.

## 6 Numerical comparisons

It is difficult to compare the performances of the different predictors and predictive intervals theoretically, as proposed in the previous sections. In this section, we present a simulation study to compare the performance of the different methods. The simulation is performed based on two types of hybrid censoring schemes, and for different  $T$  and different parameter values. We have kept  $r = 10$  and  $n = 20$  to be fixed throughout. The experiment can be briefly describe as follows. For given  $n$  and the parameter values, we generate a sample of size  $n$  from a  $GE(\alpha, \lambda)$ . Based on the sampling scheme we obtain the (training) sample from the whole sample as  $\{x_1, \dots, x_D\}$ . Now based on the sample  $\{x_1, \dots, x_D\}$ , we predict  $Y = X_{s+D:n}$ , and also obtain different predictive intervals. For computing Bayesian point and interval predictors, we assume two priors as follows:

$$\text{Prior 1: } a_j = 0.0001, \quad b_j = 0.0001, \quad j = 1, 2,$$

$$\text{Prior 2: } a_j = 1, \quad b_j = 3, \quad j = 1, 2.$$

The two priors have the same means. But, the variance of Prior 2 is smaller than that of Prior 1. So, the Prior 2 is more informative than Prior 1.

We compare the performances of the MLP, BUP, CMP and Bayes predictors in terms of biases, mean squares prediction errors (MSPE) and Pitman closeness (PC) measure, based on 10,000 replications. It may be mentioned the measure of Pitman closeness has received considerable attention in recent years, see for example Balakrishnan et al. (2011) and the references cited therein. The bias and MSPE of the different predictors are reported in Table 2, and the PC measures of the different predictors are reported in Table 3.

We have calculated the Bias, MSPE and PC as follows. Suppose  $\hat{Y}_i$  is the prediction of  $Y = X_{s+D:n}$  derived in the  $i$ th iteration of simulation, where  $i = 1, \dots, N = 10,000$ , then the Bias and the MSPE computed as:

$$\text{Bias} = \frac{1}{N} \sum_{i=1}^N (\hat{Y}_i - Y) \tag{6.1}$$

and

$$\text{MSPE} = \frac{1}{N} \sum_{i=1}^N (\hat{Y}_i - Y)^2. \tag{6.2}$$

If we consider,  $\hat{Y}_i^{(1)}$  and  $\hat{Y}_i^{(2)}$  as the predictors of  $Y = X_{s+D:n}$ , obtained based on the first and second method, respectively, then PC for  $\hat{Y}_i^{(1)}$  against  $\hat{Y}_i^{(2)}$ , can be calculated as follows:

$$\text{PC} = P(|\hat{Y}^{(1)} - Y| < |\hat{Y}^{(2)} - Y|) = \frac{1}{N} \sum_{i=1}^N I(|\hat{Y}_i^{(2)} - Y| - |\hat{Y}_i^{(1)} - Y|), \tag{6.3}$$

where  $I(\cdot)$  is the indicator function, and  $I(x) = 1$  if  $x > 0$  and 0, otherwise. We say that  $\hat{Y}^{(1)}$  competes with  $\hat{Y}^{(2)}$  if  $\text{PC} > 0.5$ .

We also compute three prediction intervals obtained using pivotal, HCD and Bayesian methods as discussed in Sections 3 and 4. We compared the performances of the three intervals in terms of their average lengths and coverage percentages based on 10,000 replications. The results are reported in Table 4.

Some of the points are quite clear from this simulation experiments.

- From Table 2, it is observed the MSPEs increase as  $s$  increases for all the prediction methods.
- For the same  $\alpha$ ,  $\lambda$  and  $T$ , the MSPEs of the different predictors for Type-II HCS are smaller than the corresponding Type-I HCS.
- In terms of MSPEs, BUP is slightly better than the MLP and CMP.
- In terms of MSPEs, the Bayes predictors under Prior 2 work better than the Bayes predictor under Prior 1.
- From Table 3, it is observed that based on PC measure BUP is better than the MLP and CMP. Also based on PC measure, the BP predictor under Prior 2 works better than the BP predictor under Prior 1.

**Table 2** The biases and MSPEs of point predictors for  $n = 20$  and  $r = 10$ 

		$\alpha = 0.75, \lambda = 1$									
		$T = 0.40$					$T = 0.60$				
		BUP	MLP	CMP	BP		BUP	MLP	CMP	BP	
					Prior 1	Prior 2				Prior 1	Prior 2
		Type-I HCS									
$s = 1$	Bias	0.016	0.021	0.018	0.024	0.020	0.019	0.024	0.021	0.034	0.030
	MSPE	0.003	0.004	0.004	0.006	0.004	0.002	0.002	0.002	0.006	0.003
$s = 2$	Bias	0.024	0.026	0.019	0.016	0.017	0.024	0.026	0.022	0.038	0.028
	MSPE	0.004	0.005	0.006	0.009	0.006	0.002	0.003	0.003	0.007	0.004
$s = 3$	Bias	0.026	0.030	0.023	0.026	0.024	0.031	0.029	0.028	0.028	0.026
	MSPE	0.006	0.007	0.009	0.010	0.008	0.004	0.006	0.006	0.009	0.006
$s = 4$	Bias	0.028	0.027	0.031	0.041	0.028	0.042	0.024	0.029	0.029	0.027
	MSPE	0.008	0.008	0.010	0.012	0.010	0.005	0.006	0.008	0.011	0.009
$s = 5$	Bias	0.034	0.029	0.025	0.027	0.019	0.038	0.034	0.026	0.019	0.017
	MSPE	0.010	0.012	0.013	0.015	0.013	0.007	0.009	0.010	0.014	0.010
		Type-II HCS									
$s = 1$	Bias	0.016	0.026	0.027	0.029	0.024	0.015	0.020	0.021	0.013	0.016
	MSPE	0.001	0.002	0.003	0.005	0.002	0.001	0.001	0.001	0.003	0.001
$s = 2$	Bias	0.026	0.027	0.021	0.016	0.019	0.019	0.015	0.024	0.013	0.019
	MSPE	0.002	0.004	0.005	0.008	0.005	0.001	0.002	0.002	0.006	0.004
$s = 3$	Bias	0.027	0.029	0.026	0.042	0.046	0.016	0.024	0.019	0.024	0.035
	MSPE	0.004	0.006	0.007	0.009	0.007	0.002	0.004	0.005	0.007	0.006
$s = 4$	Bias	0.019	0.015	0.022	0.031	0.037	0.019	0.027	0.030	0.064	0.050
	MSPE	0.005	0.007	0.008	0.011	0.010	0.003	0.005	0.006	0.010	0.007
$s = 5$	Bias	0.029	0.025	0.020	0.030	0.026	0.020	0.024	0.021	0.024	0.014
	MSPE	0.008	0.010	0.012	0.014	0.012	0.005	0.007	0.009	0.012	0.009

**Table 2** (Continued)

		$\alpha = 1.5, \lambda = 1$									
		$T = 0.80$					$T = 1.20$				
		BUP	MLP	CMP	BP		BUP	MLP	CMP	BP	
					Prior 1	Prior 2				Prior 1	Prior 2
Type-I HCS											
$s = 1$	Bias	0.030	0.036	0.039	0.049	0.037	0.035	0.029	0.041	0.024	0.028
	MSPE	0.005	0.007	0.007	0.012	0.009	0.004	0.006	0.007	0.010	0.008
$s = 2$	Bias	0.033	0.028	0.046	0.037	0.029	0.024	0.028	0.037	0.028	0.027
	MSPE	0.008	0.010	0.011	0.016	0.012	0.006	0.008	0.008	0.014	0.011
$s = 3$	Bias	0.045	0.056	0.028	0.049	0.050	0.064	0.052	0.034	0.042	0.046
	MSPE	0.011	0.014	0.016	0.019	0.013	0.009	0.010	0.012	0.021	0.018
$s = 4$	Bias	0.059	0.067	0.065	0.034	0.022	0.029	0.034	0.035	0.075	0.046
	MSPE	0.013	0.016	0.019	0.025	0.021	0.010	0.013	0.016	0.021	0.018
$s = 5$	Bias	0.062	0.067	0.046	0.024	0.029	0.049	0.057	0.061	0.075	0.060
	MSPE	0.016	0.018	0.022	0.028	0.026	0.013	0.019	0.021	0.027	0.023
Type-II HCS											
$s = 1$	Bias	0.058	0.064	0.046	0.072	0.068	0.042	0.038	0.048	0.067	0.062
	MSPE	0.005	0.006	0.007	0.010	0.008	0.003	0.004	0.006	0.009	0.008
$s = 2$	Bias	0.041	0.049	0.046	0.034	0.037	0.045	0.036	0.047	0.045	0.054
	MSPE	0.007	0.008	0.008	0.011	0.010	0.005	0.007	0.008	0.010	0.009
$s = 3$	Bias	0.033	0.037	0.035	0.028	0.026	0.056	0.034	0.045	0.060	0.054
	MSPE	0.009	0.010	0.013	0.017	0.015	0.008	0.009	0.011	0.016	0.013
$s = 4$	Bias	0.051	0.052	0.046	0.057	0.053	0.078	0.082	0.064	0.067	0.072
	MSPE	0.011	0.013	0.014	0.020	0.016	0.009	0.012	0.015	0.018	0.014
$s = 5$	Bias	0.067	0.066	0.055	0.058	0.056	0.034	0.038	0.064	0.052	0.042
	MSPE	0.012	0.018	0.020	0.025	0.022	0.022	0.016	0.018	0.024	0.021

**Table 3** PC comparison of different predictors for  $n = 20$  and  $r = 10$ 

$\alpha = 0.75, \lambda = 1$								
$T = 0.40$				$T = 0.60$				
	BUP vs MLP	BUP vs CMP	MLP vs CMP	BP (Prior 2) vs BP (Prior 1)	BUP vs MLP	BUP vs CMP	MLP vs CMP	BP (Prior 2) vs BP (Prior 1)
	Type-I HCS							
$s = 1$	0.545	0.568	0.526	0.649	0.584	0.604	0.529	0.684
$s = 2$	0.612	0.649	0.586	0.682	0.648	0.682	0.629	0.719
$s = 3$	0.671	0.697	0.641	0.729	0.706	0.729	0.687	0.780
$s = 4$	0.712	0.748	0.730	0.772	0.738	0.759	0.720	0.816
$s = 5$	0.746	0.759	0.741	0.791	0.784	0.798	0.758	0.842
	Type-II HCS							
$s = 1$	0.527	0.558	0.521	0.613	0.542	0.579	0.534	0.640
$s = 2$	0.549	0.584	0.531	0.663	0.576	0.597	0.564	0.687
$s = 3$	0.604	0.627	0.589	0.705	0.637	0.684	0.610	0.729
$s = 4$	0.642	0.681	0.624	0.734	0.685	0.705	0.659	0.779
$s = 5$	0.693	0.713	0.673	0.751	0.725	0.746	0.706	0.792
$\alpha = 1.5, \lambda = 1$								
$T = 0.80$				$T = 1.20$				
	BUP vs MLP	BUP vs CMP	MLP vs CMP	BP (Prior 2) vs BP (Prior 1)	BUP vs MLP	BUP vs CMP	MLP vs CMP	BP (Prior 2) vs BP (Prior 1)
	Type-I HCS							
$s = 1$	0.546	0.568	0.537	0.629	0.564	0.579	0.543	0.613
$s = 2$	0.576	0.592	0.550	0.671	0.591	0.628	0.571	0.649
$s = 3$	0.608	0.631	0.582	0.703	0.638	0.662	0.616	0.682
$s = 4$	0.628	0.654	0.603	0.741	0.652	0.689	0.662	0.726
$s = 5$	0.659	0.682	0.631	0.768	0.694	0.726	0.672	0.752
	Type-II HCS							
$s = 1$	0.528	0.549	0.520	0.608	0.534	0.550	0.528	0.597
$s = 2$	0.549	0.573	0.534	0.614	0.552	0.576	0.534	0.621
$s = 3$	0.568	0.593	0.551	0.629	0.581	0.608	0.571	0.657
$s = 4$	0.586	0.616	0.572	0.684	0.620	0.657	0.604	0.703
$s = 5$	0.634	0.652	0.612	0.711	0.659	0.689	0.639	0.726

**Table 4** Average lengths and coverage probabilities of 95% PIs for  $n = 20$  and  $r = 10$

		$\alpha = 0.75, \lambda = 1$							
		$T = 0.40$				$T = 0.60$			
		Pivotal method	HCD method	Bayesian Prior 1	Bayesian Prior 2	Pivotal method	HCD method	Bayesian Prior 1	Bayesian Prior 2
		Type-I HCS							
$s = 1$	Length	0.125	–	0.112	0.103	0.116	–	0.107	0.091
	CP	0.934	–	0.935	0.938	0.933	–	0.935	0.937
$s = 2$	Length	0.203	0.180	0.195	0.188	0.161	0.134	0.153	0.146
	CP	0.936	0.941	0.938	0.940	0.935	0.940	0.936	0.937
$s = 3$	Length	0.247	0.219	0.235	0.230	0.199	0.164	0.189	0.173
	CP	0.938	0.941	0.938	0.941	0.937	0.941	0.936	0.938
$s = 4$	Length	0.298	0.269	0.286	0.276	0.234	0.206	0.222	0.216
	CP	0.942	0.945	0.942	0.944	0.940	0.944	0.940	0.941
$s = 5$	Length	0.352	0.329	0.348	0.337	0.279	0.242	0.266	0.254
	CP	0.943	0.947	0.945	0.945	0.942	0.945	0.943	0.944
		Type-II HCS							
$s = 1$	Length	0.133	–	0.124	0.116	0.108	–	0.096	0.088
	CP	0.933	–	0.936	0.934	0.929	–	0.934	0.937
$s = 2$	Length	0.196	0.169	0.181	0.175	0.148	0.119	0.138	0.130
	CP	0.935	0.938	0.937	0.934	0.930	0.934	0.933	0.931
$s = 3$	Length	0.264	0.231	0.254	0.242	0.194	0.160	0.175	0.168
	CP	0.938	0.941	0.938	0.940	0.935	0.938	0.936	0.936
$s = 4$	Length	0.306	0.281	0.297	0.291	0.254	0.224	0.244	0.238
	CP	0.939	0.943	0.943	0.943	0.938	0.943	0.940	0.941
$s = 5$	Length	0.336	0.310	0.328	0.321	0.266	0.238	0.255	0.247
	CP	0.941	0.946	0.942	0.945	0.941	0.944	0.941	0.943

**Table 4** (Continued)

		$\alpha = 1.5, \lambda = 1$							
		$T = 0.80$				$T = 1.20$			
		Pivotal method	HCD method	Bayesian Prior 1	Bayesian Prior 2	Pivotal method	HCD method	Bayesian Prior 1	Bayesian Prior 2
		Type-I HCS							
$s = 1$	Length	0.167	–	0.154	0.147	0.142	–	0.132	0.125
	CP	0.937	–	0.939	0.940	0.935	–	0.936	0.938
$s = 2$	Length	0.266	0.241	0.258	0.252	0.206	0.174	0.194	0.187
	CP	0.938	0.942	0.939	0.941	0.936	0.940	0.938	0.939
$s = 3$	Length	0.322	0.301	0.320	0.311	0.261	0.238	0.250	0.244
	CP	0.938	0.943	0.940	0.941	0.938	0.942	0.939	0.941
$s = 4$	Length	0.405	0.371	0.394	0.382	0.317	0.284	0.310	0.301
	CP	0.944	0.949	0.947	0.949	0.944	0.947	0.945	0.945
$s = 5$	Length	0.470	0.451	0.464	0.460	0.344	0.318	0.329	0.324
	CP	0.947	0.955	0.951	0.954	0.945	0.952	0.950	0.950
		Type-II HCS							
$s = 1$	Length	0.161	–	0.150	0.142	0.138	–	0.126	0.120
	CP	0.935	–	0.938	0.937	0.933	–	0.934	0.936
$s = 2$	Length	0.231	0.212	0.224	0.219	0.226	0.201	0.219	0.211
	CP	0.938	0.942	0.939	0.941	0.935	0.938	0.935	0.937
$s = 3$	Length	0.298	0.268	0.284	0.278	0.287	0.261	0.280	0.274
	CP	0.941	0.945	0.943	0.945	0.937	0.941	0.938	0.938
$s = 4$	Length	0.394	0.367	0.388	0.379	0.311	0.294	0.304	0.300
	CP	0.942	0.946	0.945	0.945	0.940	0.945	0.942	0.944
$s = 5$	Length	0.461	0.449	0.459	0.455	0.339	0.312	0.330	0.328
	CP	0.944	0.949	0.946	0.948	0.942	0.947	0.945	0.946

- From Table 4, it is observed that the coverage probabilities for all prediction intervals are very close to the nominal level.
- For the same  $\alpha$ ,  $\lambda$  and  $T$ , the average lengths and coverage percentages increase as  $s$  increases for all the methods.
- From Table 4, the HCD method works better than the pivotal and Bayesian methods in terms of both average lengths and coverage probabilities. Also, the Bayesian prediction intervals under Prior 2 are shorter than the Bayesian prediction intervals under Prior 1.

Therefore, based on the simulation results here, we propose the following: for prediction of future observation BUP may be used, and for prediction interval, for  $s > 1$ , we propose to use the HCD method, and for  $s = 1$ , Bayesian method may be used.

## 7 Conclusions

In this paper, we have considered the prediction of future observations and the associated prediction interval based on Type-I HCS or Type-II HCS samples, when the lifetime distributions of the experimental units follow GE distribution. We have used three different classical predictors namely BUP, MLP and CMP. We have also considered Bayesian predictor under squared error loss function. We then compared these different point predictors using biases, MSPE and PC. Comparing all the points, we recommend to use BUP for predicting the future observation. To calculate the prediction intervals, we have used three methods namely HCD, Pivotal and Bayesian methods. It is observed HCD method cannot be used when  $s = 1$ . Three proposed methods maintain the coverage percentages. HCD method performs slightly better than the Pivotal method and Bayesian method for  $s > 1$ .

In this paper, we have mainly considered the prediction of the future failures under Type-I or Type-II hybrid censoring. Throughout the study, it has been assumed that the time point  $T$  is pre-fixed. It may be of interest if it is assumed that  $T$  is also a random variable. Both Bayesian and frequentist analysis are possible in this case. More work is needed along that direction.

## Appendix

The conditional density of  $\alpha$  given  $\lambda$  and the data is

$$g_1(\alpha|\lambda, \mathbf{x}) \propto \frac{e^{-b_1\alpha}\alpha^{D+a_1-1}}{(\sum_{i=1}^D x_i + b_2)^{D+a_2}} [1 - (1 - e^{-\lambda T_0})^\alpha]^{n-D} \prod_{i=1}^D (1 - e^{-\lambda x_i})^{\alpha-1}.$$

The log-likelihood of  $g_1(\alpha|\lambda, \mathbf{x})$  is

$$\begin{aligned} \ln g_1(\alpha|\lambda, \mathbf{x}) \propto & -b_1\alpha + (D + a_1 - 1) \ln(\alpha) + (\alpha - 1) \sum_{i=1}^D \ln[1 - e^{-\lambda x_i}] \\ & + (n - D) \ln[1 - (1 - e^{-\lambda T_0})^\alpha]. \end{aligned}$$

Since we have

$$\frac{d^2 \ln g_1(\alpha|\lambda, \mathbf{x})}{d\alpha^2} = -\frac{D + a_1 - 1}{\alpha^2} - (n - D) \frac{\ln^2[1 - e^{-\lambda T_0}](1 - e^{-\lambda T_0})^\alpha}{(1 - (1 - e^{-\lambda T_0})^\alpha)^2} \leq 0,$$

the result follows.

## Acknowledgments

The authors would like to thank the referees and the associate editor for their very constructive suggestions.

## References

- Asgharzadeh, A., Valiollahi, R. and Kundu, D. (2015). Prediction for future failures in Weibull distribution under hybrid censoring. *J. Stat. Comput. Simul.* **85**, 824–838. [MR3284645](#)
- Balakrishnan, N., Davies, K. F., Keating, J. P. and Mason, R. L. (2011). Pitman closeness, monotonicity and consistency of best linear unbiased and in variant estimators for exponential distribution under type-II censoring. *J. Stat. Comput. Simul.* **81**, 985–999. [MR2820061](#)
- Balakrishnan, N., Beutner, E. and Cramer, E. (2010). Exact two-sample non-parametric confidence, prediction, and tolerance intervals based on ordinary and progressively type-II right censored data. *TEST* **19**, 68–91. [MR2610918](#)
- Basak, I., Basak, P. and Balakrishnan, N. (2006). On some predictors of times to failures of censored items in progressively censored sample. *Comput. Statist. Data Anal.* **50**, 1313–1337. [MR2224374](#)
- Balakrishnan, N. and Kundu, D. (2013). Hybrid censoring: Models, inferential results and applications (with discussion). *Comput. Statist. Data Anal.* **57**, 166–209. [MR2981081](#)
- Chen, S. and Bhattacharya, G. K. (1988). Exact confidence bounds for an exponential parameter under hybrid censoring. *Comm. Statist. Theory Methods* **17**, 1857–1870. [MR0945789](#)
- Childs, A., Chandrasekhar, B., Balakrishnan, N. and Kundu, D. (2003). Exact likelihood inference based on type-I and type-II hybrid censored samples from the exponential distribution. *Ann. Inst. Statist. Math.* **55**, 319–330. [MR2001867](#)
- Dellaportas, P. and Wright, D. E. (1991). Numerical prediction for the two-parameter Weibull distribution. *The Statistician* **40**, 365–372.
- Devroye, L. (1984). A simple algorithm for generating random variates with a logconcave density. *Computing* **33**, 247–257. [MR0773927](#)
- Draper, N. and Guttman, I. (1987). Bayesian analysis of hybrid life tests with exponential failure times. *Ann. Inst. Statist. Math.* **39**, 219–225. [MR0886518](#)
- Ebrahimi, N. (1986). Estimating the parameter of an exponential distribution from hybrid life test. *J. Statist. Plann. Inference* **14**, 255–261. [MR0852530](#)
- Ebrahimi, N. (1992). Prediction intervals for future failures in exponential distribution under hybrid censoring. *IEEE Transactions on Reliability* **41**, 127–132.
- Epstein, B. (1954). Life tests in the exponential case. *Ann. Math. Stat.* **25**, 555–564. [MR0064372](#)
- Fairbanks, K., Madison, R. and Dykstra, R. (1982). A confidence interval for an exponential parameter from a hybrid life test. *J. Amer. Statist. Assoc.* **77**, 137–140.
- Geman, S. and Geman, A. (1984). Stochastic relaxation, Gibbs distributions and the Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **6**, 721–740.
- Jeong, H. S., Park, J. I. and Yum, B. J. (1996). Development of  $(r, T)$  hybrid sampling plans for exponential lifetime distributions. *J. Appl. Stat.* **23**, 601–607.

- Kaminsky, K. S. and Rhodin, L. S. (1985). Maximum likelihood prediction. *Ann. Inst. Statist. Math.* **37**, 507–517. [MR0818048](#)
- Kundu, D. (2007). On hybrid censored Weibull distribution. *J. Statist. Plann. Inference* **137**, 2127–2142. [MR2325420](#)
- Kundu, D. and Banerjee, A. (2008). Inference based on type-II hybrid censored data from a Weibull distribution. *IEEE Transaction on Reliability* **57**, 369–378.
- Kundu, D. and Gupta, R. D. (1988). Hybrid censoring schemes with exponential failure distribution. *Comm. Statist. Theory Methods* **27**, 3065–3083.
- Kundu, D. and Raqab, M. Z. (2012). Bayesian inference and prediction of order statistics for a type-II censored Weibull distribution. *J. Statist. Plann. Inference* **142**, 41–47. [MR2827128](#)
- Lawless, J. F. (1982). *Statistical Models and Methods for Life Time Data*. New York: Wiley. [MR0640866](#)
- Raqab, M. Z. and Nagaraja, H. N. (1995). On some predictors of future order statistics. *Metron* **53**, 185–204. [MR1380909](#)
- Raqab, M. Z., Asgharzadeh, A. and Valiollahi, R. (2010). Prediction for Pareto distribution based on progressively type-II censored samples. *Comput. Statist. Data Anal.* **54**, 1732–1743. [MR2608969](#)
- Smith, R. L. (1997). Statistics for exceptional athletics record: Letter to the editor. *Appl. Statist.* **46**, 123–127.
- Smith, R. L. (1999). Bayesian and frequentist approaches to parametric predictive inference (with discussion). In *Bayesian Statistics, Vol. 6* (J. M. Bernardo, J. O. Berger, A. P. Dawid and A. F. M. Smith, eds.) 589–612. Oxford: Oxford University Press. [MR1724875](#)

R. Valiollahi  
Department of Mathematics, Statistics  
and Computer Science  
Semnan University  
Semnan  
Iran

A. Asgharzadeh  
Department of Statistics  
Faculty of Mathematical Sciences  
University of Mazandaran  
Babolsar  
Iran  
E-mail: [a.asgharzadeh@umz.ac.ir](mailto:a.asgharzadeh@umz.ac.ir)

D. Kundu  
Department of Mathematics and Statistics  
Indian Institute of Technology  
Kanpur  
India