

HIGH ORDER EXPANSIONS FOR RENEWAL FUNCTIONS AND APPLICATIONS TO RUIN THEORY¹

BY DOMBRY CLÉMENT AND RABEHASAINA LANDY

Université de Bourgogne Franche-Comté

A high order expansion of the renewal function is provided under the assumption that the inter-renewal time distribution is light tailed with finite moment generating function g on a neighborhood of 0. This expansion relies on complex analysis and is expressed in terms of the residues of the function $1/(1 - g)$. Under the assumption that g can be extended into a meromorphic function on the complex plane and some technical conditions, we obtain even an exact expansion of the renewal function. An application to risk theory is given where we consider high order expansion of the ruin probability for the standard compound Poisson risk model. This precises the well-known Crámer-Lundberg approximation of the ruin probability when the initial reserve is large.

1. Introduction. Let $(X_k)_{k \in \mathbb{N}}$ be an i.i.d. sequence of nonnegative random variables with common cumulative distribution F . The arrival times $(S_n)_{n \in \mathbb{N}}$ are defined by $S_0 = 0$ and, for $n \geq 1$, $S_n = \sum_{k=1}^n X_k$. We consider the counting process N defined by

$$N(x) := \sum_{n \geq 0} 1_{\{S_n \leq x\}}, \quad x \geq 0,$$

and the associated renewal function

$$U(x) := \mathbb{E}[N(x)] = \sum_{n=0}^{\infty} F^{*n}(x), \quad x \geq 0.$$

The renewal theorem states that if the inter-arrival distribution F has a finite first moment $\mu := \mathbb{E}(X_1)$, then

$$U(x) \sim \frac{x}{\mu} \quad \text{as } x \rightarrow +\infty.$$

Recall that the distribution F is called lattice if F is supported by $h\mathbb{N} = \{0, h, 2h, \dots\}$ for some mesh $h > 0$. It is well known (see, e.g., Asmussen [1],

Received June 2016; revised November 2016.

¹Supported in part by the French National Agency of Research ANR Grant, project AMMSI, number ANR-2011-BS01-021.

MSC2010 subject classifications. Primary 60K05; secondary 60K10.

Key words and phrases. Renewal process, renewal function, ruin probability, Crámer-Lundberg approximation, compound Poisson model.

Proposition 6.1) that if F has a finite second moment $\mu_2 := \mathbb{E}(X_1^2)$, then

$$(1) \quad U(x) = \begin{cases} \frac{x}{\mu} + \frac{\mu_2 + \mu}{2\mu^2} + o(1), & \text{if } F \text{ is lattice,} \\ \frac{x}{\mu} + \frac{\mu_2}{2\mu^2} + o(1), & \text{otherwise} \end{cases} \quad \text{as } x \rightarrow +\infty.$$

When F has infinite first or second moment, Teugels [25] provides asymptotics for the renewal function U under some regular variation conditions. In this paper, we focus on the case when F is light-tailed and we assume that X_1 has some finite exponential moment so that

$$(2) \quad R := \sup \left\{ r \geq 0; \int_0^\infty e^{rx} dF(x) < \infty \right\} > 0.$$

Then the moment generating function

$$g(z) := \int_0^\infty e^{zx} dF(x)$$

is well-defined and holomorphic on

$$S_R = \{z \in \mathbb{C}, \Re(z) < R\}.$$

Using complex analysis, Stone [22] proved that under the strong nonlattice condition:

$$(3) \quad \limsup_{\theta \rightarrow \pm\infty} \left| \frac{1}{1 - g(i\theta)} \right| < \infty,$$

there exists some $r > 0$ such that

$$(4) \quad U(x) = \frac{x}{\mu} + \frac{\mu_2}{2\mu^2} + o(e^{-rx}) \quad \text{as } x \rightarrow \infty.$$

Up to now, relatively few results concern either expansions or closed form expressions for $U(x)$. It appears that the only case where closed form expansions are available are when F is Matrix exponential distributed, in which case an expression of $U(x)$ is given in Asmussen and Bladt [3]. Mitov and Omey [18] provide heuristics on many terms asymptotics of $U(x)$ that are verified on the already known cases. However, as the authors point out, those interesting expansions are only given formally and are not proved. Other expansions are available in [8], Theorem 4, in the context of potential densities of Lévy processes in terms of the associated Lévy jump distribution. The approach by Stone [22] for obtaining Expansion (4) is mainly based on complex analysis and proved fruitful for obtaining expansions involving survival functions of random sums, see Blanchet and Glynn [6]. The approach in [22] was later generalized for spread out distributions in [24]; note that the generalization of our results to spread out distribution is not available in the present paper because the main technical assumption that enable us to obtain higher expansions [namely, Assumption (6) thereafter], which is stronger

than the nonlattice condition (3), is not verified for spread out distributions. This is not really surprising, as it is on the other hand known that a spreadout distribution is strongly nonlattice (see, e.g., Proposition 1.6 page 189 of [1]), which is why many results in renewal theory that hold for strongly nonlattice distributions also hold for spreadout distributions.

We consider in this paper both the lattice and the nonlattice cases and, in view of equation (1), we introduce the function:

$$v(x) := \begin{cases} U(x) - \frac{x}{\mu} - \frac{\mu_2 + \mu}{2\mu^2}, & \text{if } F \text{ is lattice,} \\ U(x) - \frac{x}{\mu} - \frac{\mu_2}{2\mu^2}, & \text{if } F \text{ is nonlattice.} \end{cases}$$

Following Stone’s approach but with more detailed computations, we obtain higher order expansions for the function v both in the lattice and nonlattice cases (Theorems 1 and 3, resp.). When g has a meromorphic extension to the whole complex plane and under some technical conditions, we are even able to provide exact expansions for v (Corollary 4).

The paper is structured as follows. Section 2 presents our results on high order expansions for the renewal function as well as some examples. Section 3 is devoted to applications to ruin theory: we provide asymptotics of the ruin probability in the setting of continuous or discrete time risk processes and also consider a two-dimensional risk process. Proofs are gathered in Sections 4 and 5.

2. High order expansions for the renewal function.

2.1. *Main results.* In the sequel, the solutions of the equation $g(z) = 1$ play a major role. Note that 0 is the unique real solution in S_R and that other solutions satisfy $\Re(z) \geq 0$ and come in pairs, that is, if z is a solution then so is \bar{z} .

We consider first the lattice case and we assume without loss of generality that the mesh of the distribution is equal to $h = 1$, that is, F is supported by \mathbb{N} . In this case, the moment generating function $z \mapsto g(z)$ is $2i\pi$ -periodic on $S_R = \{z \in \mathbb{C}, \Re(z) < R\}$ and we introduce the fundamental domain $S_R^f = \{z \in \mathbb{C}; \Re(z) < R, -\pi \leq \Im(z) \leq \pi\}$.

THEOREM 1. *Let $R_0 \in (0, R)$ be such that equation $g(z) = 1$ has no root verifying $\Re(z) = R_0$. Let us denote by $z_0 = 0, z_1, \dots, z_N$ the solutions of this equation in $S_{R_0}^f$. Then $v(k)$ has the asymptotic expansion:*

$$(5) \quad v(k) = \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-kz}}{(e^z - 1)(1 - g(z))}; z_j\right) + o(e^{-R_0k}), \quad k \rightarrow +\infty, k \in \mathbb{N},$$

where the notation $\operatorname{Res}(f(z); z_j)$ denotes the residue of the meromorphic function f at pole z_j .

If z_j is a simple zero of $g - 1$, that is, $g'(z_j) \neq 0$, then the j th term in (5) has the simple form:

$$-\frac{e^{-kz_j}}{(e^{z_j} - 1)g'(z_j)}.$$

Next, we consider the case when F is nonlattice. It is well known that a distribution is nonlattice if and only if $\theta = 0$ is the unique real solution of the equation $g(i\theta) = 1$. We will need here the following stronger technical assumption: for all $R_0 < R$:

$$(6) \quad \limsup_{\theta \rightarrow \pm\infty} \sup_{0 \leq r \leq R_0} \left| \frac{1}{1 - g(r + i\theta)} \right| < \infty.$$

Assumption (6) is stronger than the strong lattice condition (3) from Stone [22]. It is however not too restrictive and satisfied by a large class of distributions as shown by the following proposition.

PROPOSITION 2. *Suppose that the distribution F is absolutely continuous with respect to the Lebesgue measure. Then Assumption (6) is satisfied for all $R_0 < R$.*

Our main result in the nonlattice case is the following theorem.

THEOREM 3. *Consider a strictly nonlattice distribution F satisfying assumption (6). Then for all $R_0 \in (0, R)$, the equation $g(z) = 1$ has a finite number of solutions in $S_{R_0} = \{z \in \mathbb{C}; \Re(z) < R_0\}$ that we denote by $z_0 = 0, z_1, \dots, z_N$. Then, supposing that R_0 is such that there is no solution to that equation verifying $\Re(z) = R_0$, $v(x)$ has the asymptotic expansion:*

$$(7) \quad v(x) = \sum_{j=1}^N \operatorname{Res} \left(\frac{e^{-xz}}{z(1 - g(z))}; z_j \right) + o(e^{-R_0x}) \quad \text{as } x \rightarrow +\infty.$$

If z_j is simple zero of $g - 1$, that is, $g'(z_j) \neq 0$, then the j th term in (7) has the simple form:

$$-\frac{e^{-xz_j}}{z_j g'(z_j)}.$$

It is worth noting that Theorems 1 and 3 can be extended to obtain asymptotics of higher order, that is, of order e^{-rx} with $r > R$, if we assume that the moment generating function g has a meromorphic extension to $S_{\bar{R}}$ for some $\bar{R} > R$. Theorems 1 and 3 and their proofs extend in a straightforward way, but not Proposition 2. In the case when the moment generating function g has a meromorphic extension to the whole complex plane, that is, $\bar{R} = +\infty$, it is even possible under some technical assumption to get an exact expansion for the renewal function.

COROLLARY 4. Assume that $g(z)$ has a meromorphic extension to the whole complex plane:

- In the lattice case, we assume furthermore that

$$(8) \quad \liminf_{r \rightarrow +\infty} \sup_{-\pi \leq \theta \leq \pi} \left| \frac{1}{e^r (1 - g(r + i\theta))} \right| = 0.$$

We denote by $z_0 = 0, z_1, \dots, z_N$ (possibly $N = +\infty$) the solution of $g(z) = 1$ in the fundamental domain $S^f = \{z \in \mathbb{C}; -\pi \leq \Im(z) \leq \pi\}$ with $\Re(z_0) \leq \Re(z_1) \leq \dots$. Then we have the exact expansion

$$(9) \quad v(k) = \sum_{j=1}^N \operatorname{Res} \left(\frac{e^{-kz}}{(e^z - 1)(1 - g(z))}; z_j \right), \quad k \geq 0.$$

- In the nonlattice case, we suppose that assumption (6) holds for all $R_0 > 0$. Let $x > 0$. Let us furthermore suppose that:

- One has an infinite number of roots $(z_n)_{n \in \mathbb{N}}$ of equation $g(z) = 1$, all of which are simple, real, and such that the following series converges:

$$(10) \quad \sum_{j=1}^{\infty} \left| \operatorname{Res} \left(\frac{e^{-xz}}{z(1 - g(z))}; z_j \right) \right| = \sum_{j=1}^{\infty} \left| \frac{e^{-xz_j}}{z_j g'(z_j)} \right| < +\infty.$$

- The following holds:

$$(11) \quad \liminf_{r \rightarrow +\infty} \sup_{\theta \in \mathbb{R}} \left| \frac{1}{r(1 - g(r + i\theta))} \right| = 0.$$

Then we have the exact expansion

$$(12) \quad v(x) = \sum_{j=1}^{\infty} \operatorname{Res} \left(\frac{e^{-xz}}{z(1 - g(z))}; z_j \right).$$

REMARK 5. One of the assumptions of Corollary 4 in the nonlattice case is that there is an infinite number of roots of equation $g(z) = 1$. One may wonder what happens in the case when those roots are in finite number N . In fact, one shows in this case, and thanks to Condition (11), that the moment generating function $g(z)$ is rational, in which case a finite expansion for $v(x)$ can be deduced almost straightforwardly.

In the previous results, a kind of dichotomy arises between the lattice and nonlattice cases. Interestingly, a unified statement can be deduced for the renewal mass function or the renewal density function in the nonlattice and lattice case, respectively.

COROLLARY 6. Under the same assumptions as Corollary 4:

- In the lattice case, the renewal measure has mass function:

$$u(k) = - \sum_{j=0}^N \operatorname{Res} \left(\frac{e^{-kz}}{1-g(z)}; z_j \right), \quad k \geq 0.$$

As a particular case, if all the poles are simple,

$$u(k) = \sum_{j=0}^N \frac{e^{-kz_j}}{g'(z_j)}, \quad k \geq 0.$$

- In the nonlattice case, the renewal measure has density function:

$$(13) \quad u(x) = - \sum_{j=0}^N \operatorname{Res} \left(\frac{e^{-xz}}{1-g(z)}; z_j \right), \quad x > 0.$$

As a particular case, if all the poles are simple,

$$(14) \quad u(x) = \sum_{j=0}^N \frac{e^{-xz_j}}{g'(z_j)}, \quad x > 0.$$

To conclude this subsection, we present an informal argument leading to (and motivating) Expansion (7) and that may lead to some better comprehension of proof of Theorem 3 given in Section 4.3. One verifies, using Fubini, that the Laplace transform of $U(\cdot)$ is

$$\begin{aligned} \int_0^\infty e^{-xz} U(x) dx &= \int_0^\infty e^{-xz} \left[\sum_{k=0}^\infty \mathbb{P}(S_k \leq x) \right] dx \\ &= \sum_{k=0}^\infty \mathbb{E} \left[\int_{S_k}^\infty e^{-xz} dx \right] \\ &= \sum_{k=0}^\infty \frac{g(-z)^k}{z} = \frac{1}{z(1-g(-z))}, \end{aligned}$$

so that, the inversion formula for the Laplace transform leads formally to

$$(15) \quad U(x) = \frac{1}{2i\pi} \int_{c-i\infty}^{c+i\infty} \frac{e^{xz}}{z(1-g(-z))} dz,$$

where $c > 0$ is such that all singularities of $\frac{e^{xz}}{z(1-g(-z))}$ are on the left of vertical line $c + i\mathbb{R}$. The right-hand side integral of (15) does not necessarily converge, however, let us suppose that this is the case. The poles of $\frac{e^{xz}}{z(1-g(-z))}$ satisfying $\Re(z) > -R_0$ are $-z_0 = 0, \dots, -z_N$, and one can verify that the residue of $\frac{e^{xz}}{z(1-g(-z))}$ at $-z_0 = 0$

is $\frac{x}{\mu} + \frac{\mu_2}{2\mu^2}$. Thus, using a rectangular contour and the theorem of residues leads to the following:

$$\begin{aligned}
 U(x) &= \sum_{j=0}^N \operatorname{Res}\left(\frac{e^{xz}}{z(1-g(-z))}; -z_j\right) + \frac{1}{2i\pi} \int_{-R_0+i\infty}^{-R_0-i\infty} \frac{e^{xz}}{z(1-g(-z))} dz \\
 (16) \quad &= \frac{x}{\mu} + \frac{\mu_2}{2\mu^2} + \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))}; z_j\right) \\
 &\quad + \frac{1}{2i\pi} \int_{-R_0+i\infty}^{-R_0-i\infty} \frac{e^{xz}}{z(1-g(-z))} dz.
 \end{aligned}$$

The last integral is an $o(e^{-R_0x})$, so that one would obtain Expansion (7). However, the main failing points in this reasoning are first that the integral in (15) is not convergent, and that the contour argument leading to (16) is more delicate than it seems. The convergence issue will be solved by introducing a Gaussian kernel (an idea already introduced by Stone [22, 23]) which, by inversion, will make the corresponding integral converge; see Step 2 in the proof in forthcoming Section 4.3. The contour argument will involve Assumption (6), which will enable two of the pieces of the contour to vanish in the proof; see again Step 2 in Section 4.3 as well as the corresponding Figure 3.

2.2. *Examples.* We provide some examples that illustrate the results above.

EXAMPLE 7. In the lattice case, we consider the negative binomial distribution with parameters $p \in (0, 1)$ and $n \geq 1$ defined by

$$F(dx) = \sum_{k \geq 0} \binom{k+n-1}{k} p^k (1-p)^n \delta_k(dx), \quad p \in (0, 1), n \geq 1.$$

Its moment generating function is given by

$$g(z) = \left(\frac{1-p}{1-pe^z}\right)^n, \quad \Re(z) < R = -\log p,$$

so that

$$\frac{1}{1-g(z)} = \frac{(1-pe^z)^n}{(1-pe^z)^n - (1-p)^n}$$

defines a meromorphic function on \mathbb{C} . The poles are the solutions of

$$e^z = \frac{1 - (1-p)e^{2i\pi \frac{j}{n}}}{p}, \quad 0 \leq j \leq n-1.$$

For $n = 1$, the only pole in the fundamental domain $S^f_{\mathbb{R}}$ is $z_0 = 0$ so that Theorem 1 implies $v(x) = o(e^{-rx})$ for all $r > 0$. In the general case $n \geq 1$, there are exactly n poles in the fundamental domain S^f given by $z_j = r_k e^{i\alpha_j}$ with

$$r_j = \log \left| \frac{1 - (1 - p)e^{2i\pi \frac{j}{n}}}{p} \right| \quad \text{and} \quad \alpha_j = \arg(1 - (1 - p)e^{2i\pi \frac{j}{n}}).$$

Furthermore, Assumption (8) is easily satisfied (comparison between an exponential and a power growth) so that Corollaries 4 and 6 apply. We obtain that the renewal measure has mass function:

$$u(k) = \sum_{j=0}^{n-1} \frac{1 - pe^{z_j}}{np(1 - p)e^{z_j}} e^{-kz_j} \quad \text{with } e^{z_j} = \frac{1 - (1 - p)e^{2i\pi \frac{j}{n}}}{p}.$$

EXAMPLE 8. In the nonlattice case let us consider *Matrix exponential distributions* with parameters (α, T) , where α is an $\mathbb{R}^{1 \times (N+1)}$ probability row vector and T is an $\mathbb{R}^{(N+1) \times (N+1)}$ subintensity matrix. The definition and principal properties of phase-type distributions can be found in Asmussen and Albrecher [2], Chapter IX. The moment generating function $g(z)$ is a rational function given by

$$(17) \quad g(z) = \alpha(-zI - T)^{-1}s,$$

where $s := -Te$ and $e = (1, \dots, 1)^T$ (see [2], Theorem 1.5). The equation $g(z) = 1$ is a polynomial equation with $N + 1$ solutions $z_0 = 0, \dots, z_N$ and Condition (6) is satisfied with $R_0 = +\infty$ and Assumption (11) is easily satisfied (comparison between an exponential and a power growth). From Corollaries 4 and 6, we obtain a closed formula for the renewal distribution U and the renewal density u . On the other hand, Asmussen and Bladt [3], Theorem 3.1, provide the simple expression:

$$(18) \quad u(x) = \alpha e^{(s\alpha + T)x} s, \quad x > 0.$$

Let us check that this formula agrees with Corollary 6 in the case when the roots z_0, \dots, z_N are simple. It is easy to check that the $-z_j$'s are exactly the eigenvalues of the matrix $s\alpha + T$. Denoting by P the $(N + 1) \times (N + 1)$ matrix made up with the eigenvectors $v_1, \dots, v_j \in \mathbb{R}^{N+1}$ corresponding to eigenvalues z_0, \dots, z_N , we have

$$s\alpha + T = P\Delta P^{-1} \quad \text{with } \Delta := \text{Diag}(-z_j, j = 0, \dots, N).$$

Denoting by J_k the $\mathbb{R}^{(N+1) \times (N+1)}$ matrix with 1 at the (k, k) th position and 0 elsewhere, we compute

$$(19) \quad u(x) = \alpha e^{(s\alpha + T)x} s = \alpha P e^{\Delta x} P^{-1} s = \sum_{j=1}^{N+1} e^{-z_{j-1}x} \alpha P J_j P^{-1} s.$$

On the other hand, we deduce from (17)

$$\begin{aligned} g'(z) &= \alpha(-zI - T)^{-2}s = \alpha(-zI + s\alpha - P\Delta P^{-1})^{-2}s \\ &= \alpha P(-zI + P^{-1}s\alpha P - \Delta)^{-2}P^{-1}s. \end{aligned}$$

Hence, in order to prove that (19) and (13) agree, we need to prove

$$\alpha P J_j P^{-1} s = \frac{1}{g'(z_{j-1})} \quad \text{for all } j = 1, \dots, N + 1$$

or equivalently

$$\alpha P J_j P^{-1} s \times \alpha P(-z_{j-1}I + P^{-1}s\alpha P - \Delta)^{-2}P^{-1}s = 1.$$

This can be easily verified with elementary algebra [the relation $J_j(\Delta + z_{j-1}I) = 0$ is useful].

EXAMPLE 9. Let us consider the simple case of the *uniform distribution* on $[0, 1]$, that is, $X \sim \mathcal{U}[0, 1]$. In that case $g(z) = \frac{e^z - 1}{z}$ and the equation $g(z) = 1$ is equivalent to $e^z = z + 1$. The solutions are $z_j = -W_j(-e^{-1}) - 1$, $j \in \mathbb{Z}$, where $W_j(\cdot)$ is the j th generalized Lambert function. Proposition 2 together with Theorem 3 provide an asymptotic expansion of v . Using the relations $z_{-j} = \bar{z}_j$ and $g'(z_j) = 1$, we obtain, for all $N \geq 1$,

$$v(x) = -2 \sum_{j=1}^N \Re \left(\frac{1}{z_j} e^{-xz_j} \right) + o(e^{-r_N x}) \quad \text{as } x \rightarrow +\infty,$$

with $z_j = -W_j(-e^{-1}) - 1$ and $r_N = \Re(z_N)$. As $N \rightarrow +\infty$, $r_N \rightarrow +\infty$ so that the expansion has arbitrary high order.

At this point, there still lacks an example of distribution X such that a meromorphic extension $g(z)$ exists, equation $g(z) = 1$ admits an infinite number of solution and infinite expansion (12) holds. An example of such an infinite expansion of $v(x)$ will be given in upcoming Example 14, in the context of meromorphic Lévy processes.

3. Application to ruin theory. As an application of Theorem 3, we provide asymptotic expansions for the ruin probability in risk theory. We consider both a continuous setting (compound Poisson risk process) and a discrete setting (binomial risk process). Estimation of the ruin probability in a two-dimensional model is also investigated.

3.1. *Ruin theory in continuous time.* We consider the following classical continuous time risk process:

$$R_t^x = x + ct - \sum_{k=1}^{N_t} Z_k = x + Y_t, \quad t \geq 0,$$

with $\{N_t, t \geq 0\}$ a Poisson process with intensity $\alpha > 0$ and $(Z_k)_{k \in \mathbb{N}}$ an i.i.d. sequence of nonnegative random variables with common distribution G and finite expectation m , independent from $\{N_t, t \geq 0\}$. Such a process models the capital of an insurance company with premium rate $c > 0$, initial reserve x , and incoming claims $(Z_k)_{k \in \mathbb{N}}$; see, for example, Asmussen and Albrecher [2]. We define $\bar{G} = 1 - G$ the tail function and assume the following on the moment generating function:

$$(20) \quad s \mapsto \int_0^\infty e^{sx} dG(x) \quad \text{is finite for all } s > 0.$$

We are interested in the ruin probability

$$(21) \quad \psi(x) := \mathbb{P}\left(\inf_{t \geq 0} R_t^x < 0\right), \quad x \geq 0,$$

and its asymptotic when the initial reserve x is large. It is well known that $\psi(x) < 1$ if and only if the safety loading is positive, that is,

$$(22) \quad \mathbb{E}(Y_1) = c - \alpha m > 0.$$

In the asymptotic analysis, a key role is played by the Lundberg equation

$$(23) \quad \int_0^\infty e^{zy} \frac{\alpha}{c} \bar{G}(y) dy = 1, \quad z \in \mathbb{C}.$$

Under Assumption (22), this equation restricted to real numbers admits a unique real solution denoted by $\kappa > 0$. The Lundberg inequality states that

$$\psi(x) \leq e^{-\kappa x} \quad \text{for all } x > 0,$$

while the Cramér–Lundberg approximation provides the asymptotic behavior as $x \rightarrow +\infty$:

$$(24) \quad \psi(x) \sim Ce^{-\kappa x} \quad \text{with } C = \frac{c - \alpha m}{\mathbb{E}(Ze^{\kappa Z}) - c}.$$

We provide high order asymptotic expansions for the ruin probability $\psi(x)$. Similar considerations as well as exact expansions have been proved with different methods by Kuznetsov and Morales [14] for a so-called meromorphic risk process and by Roynette, Vallois and Volpi [20].

Using the fact that $\kappa > 0$ solves the Lundberg equation (23), one can define the probability measure F on $[0, +\infty)$ by

$$F(dx) := e^{\kappa x} \frac{\alpha}{c} \bar{G}(x) dx.$$

The moment generating function

$$g(z) = \int_0^\infty e^{zx} F(dx), \quad z \in \mathbb{C}$$

is well-defined and holomorphic on the complex plane. For future reference, note that the Lundberg equation (23) is equivalent to

$$(25) \quad g(z - \kappa) = 1, \quad z \in \mathbb{C}.$$

Using a renewal equation solved by the ruin probability function ψ and the asymptotic behavior of the renewal function provided by Theorem 3, we can deduce an asymptotic expansion for $\psi(x)$ as $x \rightarrow +\infty$.

THEOREM 10. *Assume conditions (20) and (22) are satisfied. Let $r > 0$ be fixed and $z_0 = 0, z_1, \dots, z_N$ the solutions of $g(z) = 1$ in $S_r = \{z \in \mathbb{C}; \Re(z) \leq r\}$. Then the ruin probability $\psi(x)$ has the asymptotic expansion:*

$$(26) \quad \psi(x) = \sum_{j=0}^N \operatorname{Res} \left(\frac{(\alpha m - c)e^{-x(z+\kappa)}}{c(1-g(z))(z+\kappa)}; z_j \right) + o(e^{-(r+\kappa)x}) \quad \text{as } x \rightarrow +\infty.$$

If z_j is a simple zero of $g - 1$, that is, $g'(z_j) \neq 0$, then the j th term in (26) has the simple form:

$$(27) \quad -\frac{(\alpha m - c)}{cg'(z_j)(z_j + \kappa)} e^{-x(z_j+\kappa)} = \frac{c - \alpha m}{\alpha \mathbb{E}(Ze^{(z_j+\kappa)Z}) - c} e^{-(z_j+\kappa)x}.$$

The term $j = 0$ of the asymptotic expansion (39) is exactly the Cramer–Lundberg approximation (24).

EXAMPLE 11. Let us consider the case where the claims are of the form $X = \min(V, d)$ where $d > 0$ and V has an exponential distribution with parameter $\lambda > 0$. This models a reinsurance scenario where a reinsurance company covers the excess of claim above d only, that is, according to a stop loss contract with priority $d > 0$. In that case, the Lundberg equation (23) reads

$$(28) \quad e^{(z-\lambda)d} = 1 + \frac{c}{\alpha}(z - \lambda), \quad z \in \mathbb{C},$$

and the Lundberg exponent is $\kappa = \lambda$. Solutions z_j satisfy

$$z_j = -\frac{\alpha}{c} - \frac{1}{d} W_j \left(-\frac{\alpha d}{c} e^{-\frac{\alpha d}{c}} \right), \quad j \in \mathbb{Z},$$

where $W_j(\cdot)$ is the j th generalized Lambert function. It is easy to check that

$$g'(z_j) = d + \frac{d\alpha/c - 1}{z_j + \kappa - \lambda} \neq 0,$$

so that Theorem 10 entails the asymptotic expansion (26) with j th term given by (27). Since claims here are bounded, the expression of the ruin probability can in fact be made more precise. Indeed, $\psi(x) = 1 - E[Y_1]W(x)$ where $W(x)$ is the so-called scale function associated to Lévy process $\{Y_t, t \geq 0\}$ (see Expression (8.7) page 215 of [16]). The expression of $W(x)$ is available in Theorem 3 of [15] as an infinite series; it yields that (26) can in fact be written as an infinite series, that is,

$$\psi(x) = - \sum_{j=0}^{\infty} \frac{(\alpha m - c)}{c g'(z_j)(z_j + \kappa)} e^{-x(z_j + \kappa)}.$$

REMARK 12. It is worth comparing Theorem 10 with the results of Kuznetsov and Morales [14]. They consider a so-called meromorphic risk process $\{R_t, t \geq 0\}$, which amounts to assume that the claims Z_k 's have density:

$$\frac{\mathbb{P}[Z_k \in dx]}{dx} = \sum_{m=1}^{\infty} b_m e^{-\rho_m x}, \quad x \geq 0,$$

for some positive coefficients $(b_m)_{m \geq 1}$ and increasing sequence $(\rho_m)_{m \geq 1}$ satisfying $\rho_m \rightarrow +\infty$. Corollary 1 of [14] states that the Laplace exponent $\Lambda(z) := \log \mathbb{E}(e^{zY_1})$ of the Lévy process $\{R_t, t \geq 0\}$ admits a meromorphic extension on $z \in \mathbb{C}$ and that all the solutions of the (extended) Lundberg equation (25) are real, negative and simple. Furthermore, denoting these solutions by $(-\zeta_n)_{n \geq 1}$, the ruin probability (21) has expansion

$$(29) \quad \psi(x) = \sum_{n=1}^{\infty} \frac{-\mathbb{E}[Y_1]}{\Lambda'(-\zeta_n)} e^{-\zeta_n x}.$$

In this framework, condition (20) is not satisfied but we check below that these results are still consistent with Theorem 10. The Laplace exponent satisfies

$$\Lambda(z) = cz + \alpha[\mathbb{E}(e^{-zZ_1}) - 1] \quad \text{and} \quad \Lambda'(z) = c - \alpha \mathbb{E}(Z_1 e^{-zZ_1}).$$

Elementary computations reveal that

$$(30) \quad \begin{aligned} g(z) &= \frac{\alpha}{c} \frac{1}{z + \kappa} \mathbb{E}(e^{(z+\kappa)Z} - 1) = \frac{\Lambda(-z - \kappa)}{c(z + \kappa)} + 1 \quad \text{and} \\ g'(z) &= \frac{\alpha}{c} \frac{\mathbb{E}(Z e^{(z+\kappa)Z})}{z + \kappa} - \frac{\alpha}{c} \frac{\mathbb{E}(e^{(z+\kappa)Z} - 1)}{(z + \kappa)^2}. \end{aligned}$$

One can check that the solutions of $g(z) = 1$ satisfy $z_n + \kappa = -\zeta_n$ and that

$$g'(z_j) = \frac{1}{z_j + \kappa} \left[\frac{\alpha}{c} \mathbb{E}(Z e^{(z_j + \kappa)Z}) - 1 \right],$$

so that Expansion (26), with corresponding terms given by (27), corresponds to the $N + 1$ first terms of (29).

REMARK 13. With some more effort, Theorem 10 can be extended to more general Gerber–Shiu functions, for example, of the form:

$$\psi(x, \theta, b, a) := \mathbb{E}_x(e^{-\theta\tau} \mathbb{1}_{\{R_{\tau-} \geq b; R_{\tau} \geq a; \tau < +\infty\}}),$$

where θ, b, a are nonnegative, $\tau := \inf\{t \geq 0 \mid R_t < 0\}$ is the ruin time of the risk process and $\underline{R}_t = \inf_{0 \leq s \leq t} R_s$ is the running minimum at time t ; see Theorem 2.8 of [20] as well as Theorem 1 of [14] for example of such expansions. For ease of presentation, we stick in this paper to $\psi(x)$ as defined by (21).

EXAMPLE 14. We give an example of an infinite expansion in the nonlattice case of $v(x)$ as in Corollary 4. Conditions (10) and (11) may look hard to verify in practice. To exhibit such an X , we again use the theory of meromorphic Lévy processes. As in Remark 12, we pick spectrally negative process $\{Y_t, t \geq 0\}$, $Y_t = ct - \sum_{k=1}^{N_t} Z_k$ where $\{N_t, t \geq 0\}$ is a Poisson process with intensity $\alpha > 0$, such that Laplace exponent is of the form:

$$\Lambda(z) = \tilde{\mu}z + z^2 \sum_{m=1}^{\infty} \frac{b_m}{\rho_m^2(\rho_m + z)},$$

for some $\tilde{\mu} > 0$, where sequences of positive real numbers $(b_m)_{m \in \mathbb{N}^*}$ and (strictly) increasing $(\rho_m)_{m \in \mathbb{N}^*}$ are such that series $\sum_{m=1}^{\infty} \frac{b_m}{\rho_m}$ converges so that Lévy process $\{Y_t, t \geq 0\}$ is indeed a compound Poisson process; see (3.2) in [14]. We will additionally suppose that sequence $(\rho_m)_{m \in \mathbb{N}^*}$ grows like a polynomial (in addition to being increasing), that is, there exists $a \geq 1$ such that

$$(31) \quad \rho_m \sim Cm^a, \quad m \rightarrow \infty$$

for some $C > 0$. Remembering that G is the c.d.f. of the Z_k 's, we then consider r.v. X with descending ladder height distribution of Lévy process $\{Y_t, t \geq 0\}$, with corresponding moment generating function:

$$g(z) := \int_0^{\infty} e^{(\kappa+z)y} \frac{\alpha}{c} \bar{G}(y) dy,$$

where $\kappa > 0$ is solution to Lundberg equation (23); see Relation (5.7) page 87 of [2]. We proceed to show that an infinite expansion for the corresponding function $v(x)$ is available. The sole condition for this expansion is (31), which is not too stringent and covers a wide range of processes. The relation between $g(z)$ and $\Lambda(z)$ is given by (30). This has two important consequences. The first one is that $z \mapsto g(z)$ is meromorphic, as $\Lambda(z)$ is. The second one is that z is a solution to $g(z) = 1$ iff $\Lambda(-z - \kappa) = 0$. By Properties (v) and (vi) of [14] (see also Theorem 1 (7) of [13]), one deduces that roots $(z_n)_{n \in \mathbb{N}}$ are *real nonnegative* and verify

$$(32) \quad z_0 = 0 < \rho_1 - \kappa < z_1 < \rho_2 - \kappa < z_2 < \dots$$

We now turn back to Conditions (10) and (11). We start by (10). We compute from (30)

$$(33) \quad g(z) = -\frac{\tilde{\mu}}{c} + 1 + (z + \kappa) \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m^2(\rho_m - z - \kappa)},$$

hence, for all $j \in \mathbb{N}$, $g'(z_j) = \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m(\rho_m - z_j - \kappa)^2}$, which happens to be positive. We now write

$$z_j^2 g'(z_j) \geq \sum_{m=1}^j \frac{b_m}{c\rho_m(1 + \frac{\rho_m - \kappa}{z_j})^2} = \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m(1 + \frac{\rho_m - \kappa}{z_j})^2} \mathbb{1}_{[m \leq j]}.$$

Using (32) and the dominated convergence theorem, one easily shows that $\sum_{m=1}^{\infty} \frac{b_m}{c\rho_m(1 + \frac{\rho_m - \kappa}{z_j})^2} \mathbb{1}_{[m \leq j]} \rightarrow \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m}$ as $j \rightarrow \infty$. One then deduces from the above inequality that

$$(34) \quad \liminf_{j \rightarrow \infty} z_j^2 g'(z_j) \geq \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m} > 0.$$

Now, (31) and (32) implies that $\sum_{m=1}^{\infty} e^{-xz_m}$ is a convergent series for all $x > 0$ which, combined with (34), implies the convergence (10).

We now prove (11), by establishing that $\lim_{n \rightarrow \infty} \frac{1}{r_n(1 - g(r_n + i\theta))} = 0$ with $r_n := \rho_n - \kappa$. Using (33) and $g(0) = 1$ implies for all $\theta \in \mathbb{R}$

$$\begin{aligned} 1 - g(r_n + i\theta) &= g(0) - g(r_n + i\theta) \\ &= \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m^2} \left[\frac{\kappa}{\rho_m - \kappa} - \frac{\rho_n + i\theta}{\rho_m - \rho_n - i\theta} \right] \\ &= \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m^2} \left[\frac{\kappa}{\rho_m - \kappa} + 1 - \rho_m \frac{\rho_m - \rho_n + i\theta}{(\rho_m - \rho_n)^2 + \theta^2} \right] \\ &:= R_n(\theta) + i I_n(\theta). \end{aligned}$$

Let us set $R_{n,1}(\theta) := \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m^2} \left[\frac{\kappa}{\rho_m - \kappa} + 1 \right] + \sum_{m=1}^n \frac{b_m}{c\rho_m^2} \rho_m \frac{\rho_n - \rho_m}{(\rho_m - \rho_n)^2 + \theta^2}$ and $R_{n,2}(\theta) := \sum_{m=n+1}^{\infty} \frac{b_m}{c\rho_m^2} \rho_m \frac{\rho_n - \rho_m}{(\rho_m - \rho_n)^2 + \theta^2}$. As $(\rho_n)_{n \in \mathbb{N}}$ is increasing one gets the following inequalities:

$$\begin{aligned} R_{n,1}(\theta) &\geq \sum_{m=1}^{\infty} \frac{b_m}{c\rho_m^2} \left[\frac{\kappa}{\rho_m - \kappa} + 1 \right] := \xi > 0, \\ 0 &\geq R_{n,2}(\theta) \geq \sum_{m=n+1}^{\infty} \frac{b_m}{c\rho_m^2} \rho_m \frac{\rho_n - \rho_m}{(\rho_m - \rho_n)^2} = \sum_{m=n+1}^{\infty} \frac{b_m}{c\rho_m} \frac{1}{\rho_n - \rho_m} := \chi_n, \end{aligned}$$

so that the real part of $1 - g(r_n + i\theta)$ verifies the inequality

$$(35) \quad |R_n(\theta)| \geq |R_{n,1}(\theta)| - |R_{n,2}(\theta)| = R_{n,1}(\theta) + R_{n,2}(\theta) \geq \xi + \chi_n.$$

(31) entails that $|\chi_n| \leq \frac{1}{\rho_{n+1}-\rho_n} \sum_{m=n+1}^{\infty} \frac{b_m}{c\rho_m} \sim \frac{1}{Cana-1} \sum_{m=n+1}^{\infty} \frac{b_m}{c\rho_m} \rightarrow 0$ as $n \rightarrow \infty$. One then deduces that $\xi + \chi_n > 0$ for n large enough, and

$$\sup_{\theta \in \mathbb{R}} \frac{1}{r_n |1 - g(r_n + i\theta)|} \leq \sup_{\theta \in \mathbb{R}} \frac{1}{r_n |R_n(\theta)|} \leq \frac{1}{r_n (\xi + \chi_n)} \rightarrow 0, \quad n \rightarrow \infty,$$

proving (11). Hence, infinite expansion (12) holds.

3.2. *Skip-free random walks on \mathbb{Z} .* Quite unlike its continuous time counterpart, risk theory in discrete time seems to have been less studied. We refer to [17] for an overview of such processes, as well as [2], Chapter XVI. This type of process is but a skip-free random walk, that is, a random walk with at most unit upward movement, and is in fact studied in many fields of applied probability. We consider here the so-called binomial discrete time risk model defined by

$$R_n = x + n - \sum_{j=1}^n Z_j = x + Y_n, \quad n \in \mathbb{N},$$

where $x \in \mathbb{N}$ is the initial reserve, the premium rate is assumed w.l.o.g. to be equal to 1, the claims $(Z_j)_{j \in \mathbb{N}}$ form an i.i.d. sequence taking values in \mathbb{N} . We let $m_1 = \mathbb{E}[Z_1]$ and assume that $m_1 \in (0, 1)$.

The discrete ruin probability is defined by

$$(36) \quad \psi(x) := \mathbb{P}\left(\inf_{n \in \mathbb{N} \setminus \{0\}} R_n \leq 0\right), \quad x \in \mathbb{N}.$$

This corresponds to the probability that a \mathbb{Z} -valued random walk starting from $x \in \mathbb{N}$ eventually becomes nonpositive. The condition $m_1 \in (0, 1)$ ensures that the random walk has a positive drift so that $\psi(x) < 1$.

A closed form expression for $\psi(x)$ may be found in [11]; however, this expression requires computing an infinite number of convolutions of distribution of Z_1 . We are here interested in finding a simple expansion of $\psi(x)$ as $x \rightarrow \infty$. Similar to Condition (20) in the continuous case, we assume here that

$$(37) \quad \text{the moment generating function } \mathbb{E}[e^{sZ}] \text{ is finite for all } s > 0.$$

In this discrete setting, the Lundberg equation writes

$$(38) \quad \sum_{k=0}^{\infty} e^{zk} \mathbb{P}[Z > k] = 1, \quad z \in \mathbb{C}.$$

Restricted to the real numbers, this equation has, thanks to convexity of mean generating function of Z , a unique solution $\kappa > 0$. We define the probability mass

function f defined by $f(k) = e^{\kappa k} \mathbb{P}[Z > k]$, $k \in \mathbb{N}$, with moment generating function:

$$g(z) = \sum_{k=0}^{\infty} e^{z k} f(k) = \frac{1 - \mathbb{E}(e^{(z+\kappa)Z})}{1 - e^{z+\kappa}}, \quad z \in \mathbb{C}.$$

The following theorem provides an asymptotic expansion of $\psi(x)$ as $x \rightarrow \infty$, $x \in \mathbb{N}$ and is the discrete analog of Theorem 10.

THEOREM 15. *Assume conditions (37) holds. Let $r > 0$ be fixed and $z_0 = 0, z_1, \dots, z_N$ the solutions to equation $g(z) = 1$ in $S_r^f = \{z \in \mathbb{C}; \Re(z) < r, -\pi \leq \Im(z) \leq \pi\}$. Then the ruin probability $\psi(x)$ has the asymptotic expansion:*

$$(39) \quad \psi(x) = - \sum_{j=0}^N \text{Res} \left[\frac{1}{1-g(z)} \frac{m-g(z)e^{\kappa+z}}{1-e^{\kappa+z}} e^{-(z+\kappa)x}; z_j \right] + o(e^{-(r+\kappa)x})$$

as $x \rightarrow \infty$.

If z_j is a simple zero of $g - 1$, that is, $g'(z_j) \neq 0$, then the j th term in (39) has the simple form:

$$(40) \quad \frac{m - e^{\kappa+z_j}}{e^{\kappa+z_j} - \mathbb{E}(Z e^{(\kappa+z_j)Z})} e^{-(z_j+\kappa)x}.$$

3.3. A two-dimensional ruin problem. We consider a two-dimensional ruin problem motivated by applications in reinsurance. The capitals of two insurance companies are modeled by the risk processes:

$$(41) \quad R_t^j = x_j + c_j t - \sum_{k=1}^{N_t^j} Z_k^j = x_j + Y_t^j, \quad t \geq 0, j = 1, 2,$$

where, $x_j \geq 0, c_j > 0$ are the respective initial reserves and premium rates, $\{N_t^j, t \geq 0\}$ are Poisson processes with intensities $\alpha_j > 0$, and $(Z_k^j)_{k \in \mathbb{N}}$ are the corresponding claims with mean m^j . For each $j = 1, 2$, independence between $\{N_t^j, t \geq 0\}$ and $(Z_k^j)_{k \in \mathbb{N}}$ is assumed. However, no independence is required between processes $\{R_t^1, t \geq 0\}$ and $\{R_t^2, t \geq 0\}$. We suppose that the mean drifts $\mathbb{E}(Y_j) = c_j - \alpha_j m^j, j = 1, 2$, are positive, and then define the eventual ruin probabilities for each company

$$\psi_j(x_j) := \mathbb{P} \left(\inf_{t \geq 0} R_t^j < 0 \right), \quad j = 1, 2.$$

We also consider the probability that (at least) one of the companies is eventually ruined:

$$\psi_{\text{or}}(x_1, x_2) := \mathbb{P} \left(\inf_{t \geq 0} R_t^1 < 0 \text{ or } \inf_{t \geq 0} R_t^2 < 0 \right),$$

that is, the probability that the two-dimensional process $\{(R_t^1, R_t^2), t \geq 0\}$ exits the first quadrants $[0, +\infty)^2$.

We are interested here in the asymptotics of $\psi_{\text{or}}(x_1, x_2)$ as (x_1, x_2) tend to infinity along a fixed direction $x_2/x_1 = q \in (0, +\infty)$. We refer to [4, 19] for related results that concern light tailed claims, or [5, 12] for models featuring heavy tailed claims.

For $j = 1, 2$, letting $g_j(z) := \mathbb{E}(e^{zZ_1^j})$, we suppose that equation $g_j(z) = 1$ has solutions $\overline{z_0^j} = 0, z_1^j, \overline{z_1^j}$ in $S_r = \{z \in \mathbb{C}, 0 \leq \Re(z) < r\}$ for some $r > 0$, and that those $z_1^j, \overline{z_1^j}$ are simple zeros of $g - 1$. Thus, $\psi_j(x_j)$ has the following 2 terms expansion from Theorem 10:

$$(42) \quad \psi_j(x_j) = C_0^j e^{-\kappa_j x_j} + \Re[C_1^j e^{-(\kappa_j + z_1^j)x_j}] + \varepsilon_j(x_j) e^{-(r + \kappa_j)x_j},$$

$$x_j \rightarrow +\infty, j = 1, 2,$$

where $\varepsilon_j(x_j) \rightarrow 0$ as $x_j \rightarrow +\infty$, and

$$C_k^j := \frac{c_j - \alpha_j m^j}{\alpha_j \mathbb{E}(Z^j e^{(z_j + \kappa)Z^j}) - c_j}, \quad k = 0, 1, j = 1, 2.$$

The main result of this subsection is the following theorem.

PROPOSITION 16. *A two term asymptotic for $\psi_{\text{or}}(x_1, x_2)$ as (x_1, x_2) tend to infinity along the fixed direction $x_2/x_1 = q \in (0, +\infty)$, is given by*

$$(43) \quad \psi_{\text{or}}(x, qx) = \Re[D_0 e^{-d_0(q)x}] + \Re[D_1 e^{-d_1(q)x}] + \eta_q(x) e^{-\Re(d_1(q))x},$$

$$x \rightarrow \infty,$$

where $d_0(q) > 0, \Re(d_1(q)) > d_0(q), 0 \leq \limsup_{x \rightarrow \infty} |\eta_q(x)| \leq 1$ and where $x \mapsto \Re[D_0 e^{-d_0(q)x}]$ and $x \mapsto \Re[D_1 e^{-d_1(q)x}]$ are the two first dominant functions among $x \mapsto C_0^1 e^{-\kappa_1 x}, x \mapsto C_0^2 e^{-q\kappa_2 x}, x \mapsto \Re[C_1^1 e^{-(\kappa_1 + z_1^1)x}]$ and $x \mapsto \Re[C_1^2 e^{-q(\kappa_2 + z_1^2)x}]$.

Four different cases occur in the asymptotic described in Proposition 16, depending on the asymptotic direction $\vec{u} = (1, q)$:

$$\Re(\kappa_1 + z_1^1) > q\kappa_2, \quad \kappa_1 > q\kappa_2 > \Re(\kappa_1 + z_1^1),$$

$$q\kappa_2 > \kappa_1 > q\Re(\kappa_2 + z_1^2), \quad \text{or} \quad q\Re(\kappa_2 + z_1^2) > \kappa_1.$$

To each case corresponds a different two terms expansion for $\psi_{\text{or}}(x, qx)$ as summarized in Figure 1. Proposition 16 generalizes the one term expansion given in Theorem 3 of [4]. The last term in (43) is only $O(e^{-\Re(d_1(q))x})$ but the condition $\limsup_{x \rightarrow \infty} |\eta_q(x)| \leq 1$ provides information on how fast this term tends to 0.




FIG. 1. Two term asymptotic expansions on the four different regions.

EXAMPLE 17. Let us consider the Stop Loss contract scenario with priority $d > 0$. We assume that $\{R_t^1, t \geq 0\}$ is the capital of an insurance company with claims $(Z_n^1)_{n \in \mathbb{N}}$ distributed as $\min(V, d)$ where V is exponentially distributed with parameter λ . The second risk process $\{R_t^2, t \geq 0\}$ corresponds to the capital of a reinsurance company which covers the excess of claims with priority d , that is, claims $(Z_n^2)_{n \in \mathbb{N}}$ are distributed as $(V - d)^+$, as described in Example 11. In that case, the two risk processes R_t^1 and R_t^2 are dependent. Because of the memoryless property of the exponential distribution, the Z_n^2 's are exponentially distributed with parameter λ and the intensity of the Poisson process $\{N_t^2, t \geq 0\}$ is given by

$$\alpha_2 = \alpha_1 \mathbb{P}(V > d) = \alpha_1 e^{-\lambda d}.$$

As the Z_n^2 's are exponentially distributed,

$$(44) \quad \psi_2(x_2) = C_0^2 e^{-\kappa_2 x_2}, \quad \kappa_2 = \lambda - \alpha_2/c_2, \quad C_0^2 = \frac{\alpha_2}{c_2 \lambda},$$

see Corollary 3.2 page 78 of [2]. The two terms expansion for $\psi_1(x_1)$ is given by

$$\psi_1(x_1) = C_0^1 e^{-\kappa_1 x_1} + \Re[C_1^1 e^{-(\kappa_1 + z_1^1)x}] + o(e^{-(r + \kappa_1)x_1}),$$

where z_1^1 is the solution to equation (28) with smallest real part $\Re(z_1^1) > \kappa_1$ and $r > \Re(z_1^1)$. Note that even though $\{R_t^1, t \geq 0\}$ and $\{R_t^2, t \geq 0\}$ are corre-




FIG. 2. Two term asymptotic expansions for the stop loss model.

lated, Proposition 16 may be applied and, since $C_1^2 = 0$, only three cases occur:

$$\Re(\kappa_1 + z_1^1) > q\kappa_2, \quad \kappa_1 > q\kappa_2 > \Re(\kappa_1 + z_1^1) \quad \text{and} \quad q\kappa_2 > \kappa_1.$$

This is summarized in Figure 2.

4. Proofs for Section 2.

4.1. *Proof of Theorem 1.* For the proof of Theorem 1, we need the following lemma.

LEMMA 18. *Let $f : [-\pi, \pi] \rightarrow \mathbb{C}$ be a continuous function satisfying $f(0) \in \mathbb{R}$ and $\Im(f(\theta)) = O(\theta)$ as $\theta \rightarrow 0$. Then we have the following convergence as $r \rightarrow 1^-$:*

$$(45) \quad \int_{-\pi}^{\pi} \Re\left(f(\theta) \frac{1}{1 - rg(i\theta)}\right) d\theta \rightarrow \int_{-\pi}^{\pi} \Re\left(f(\theta) \frac{1}{1 - g(i\theta)}\right) d\theta + \pi \frac{f(0)}{\mu}.$$

The above lemma is akin to the preliminary result of Stone [23], page 330; see also Breiman [7] and Feller and Orey [10]. In these references, only the case of a real valued function f is considered. In the complex case, we give a short proof inspired by Lemma 10.11 page 221 of [7].

PROOF OF LEMMA 18. Inspecting the proof of [7], one can see that it is sufficient to check that $\theta \mapsto \Re(f(\theta) \frac{1}{1-g(i\theta)})$ is integrable at $\theta = 0$. The rest of the proof may be applied similarly (with minor modification) in order to prove (45). Since

$$\Re\left(f(\theta) \frac{1}{1-g(i\theta)}\right) = \frac{\theta^2}{|1-g(i\theta)|^2} \frac{\Re(f(\theta)(1-\overline{g(i\theta)}))}{\theta^2}$$

and

$$\frac{\theta^2}{|1-g(i\theta)|^2} \rightarrow \frac{1}{\mu^2} \quad \text{as } \theta \rightarrow 0,$$

it is sufficient to prove local integrability of $\frac{\Re(f(\theta)(1-\overline{g(i\theta)}))}{\theta^2}$ at $\theta = 0$. We compute further

$$\Re(f(\theta)(1-\overline{g(i\theta)})) = \Re(f(\theta))\Re(1-\overline{g(i\theta)}) - \Im(f(\theta))\Im(1-\overline{g(i\theta)}).$$

The first term is integrable at 0 since

$$\begin{aligned} \int_{-\varepsilon}^{\varepsilon} \frac{|\Re(f(\theta))\Re(1-\overline{g(i\theta)})|}{\theta^2} d\theta &\leq \sup_{\theta \in [-\pi, \pi]} |f(\theta)| \times \int_{-\varepsilon}^{\varepsilon} \frac{1 - \mathbb{E}(\cos(\theta X_1))}{\theta^2} d\theta \\ &< +\infty. \end{aligned}$$

For the integrability of the second term, we need the assumption $\Im(f(\theta)) = O(\theta)$ which implies the existence of a constant $C > 0$ such that $|\Im(f(\theta))| \leq C|\theta|$ for $|\theta| \leq \varepsilon$. We use also the inequality $|\sin(x)| \leq x, x \in \mathbb{R}$. Using this, we have

$$\begin{aligned} \int_{-\varepsilon}^{\varepsilon} \left| \frac{\Im(f(\theta))\Im(1-\overline{g(i\theta)})}{\theta^2} \right| d\theta &\leq \mathbb{E} \int_{-\varepsilon}^{\varepsilon} \left| \frac{\Im(f(\theta))}{\theta} \frac{\sin(\theta X_1)}{\theta} \right| d\theta \\ &\leq \mathbb{E} \int_{-\varepsilon}^{\varepsilon} C|X_1| d\theta = 2\varepsilon C \mathbb{E}(|X_1|) < +\infty. \quad \square \end{aligned}$$

PROOF OF THEOREM 1. Let us define

$$u_k := U(k) - U(k-1) = \sum_{n=0}^{\infty} \mathbb{P}(S_n = k), \quad k \in \mathbb{N},$$

with the convention $U(-1) = 0$. We use the basic fact that S_n has Fourier transform $g(i\theta)^n$ and that the probabilities $\mathbb{P}(S_n = k)$'s are linked to the Fourier transform by

$$\mathbb{P}(S_n = k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ik\theta} g(i\theta)^n d\theta, \quad n \in \mathbb{N}, k \in \mathbb{N},$$

which can be verified by writing $g(i\theta)^n = \mathbb{E}(e^{i\theta S_n})$ and using Fubini. Hence, by Lebesgue’s dominated convergence and Fubini’s theorems,

$$\begin{aligned} u_k &= \lim_{r \rightarrow 1^-} \sum_{n=0}^{\infty} r^n \mathbb{P}(S_n = k) \\ &= \lim_{r \rightarrow 1^-} \sum_{n=0}^{\infty} r^n \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ik\theta} g(i\theta)^n d\theta \\ &= \lim_{r \rightarrow 1^-} \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ik\theta} \frac{1}{1 - rg(i\theta)} d\theta. \end{aligned}$$

Note that $\Im(u_k) = 0$. We deduce, thanks to Lemma 18,

$$u_k = \Re(u_k) = \frac{1}{2\mu} + \frac{1}{2\pi} \int_{-\pi}^{\pi} \Re\left(e^{-ik\theta} \frac{1}{1 - g(i\theta)}\right) d\theta.$$

We apply the same argument to the i.i.d. r.v. $(X'_n)_{n \in \mathbb{N}}$ with distribution $X'_n \sim \delta_1$. This yields, for all $k \in \mathbb{N}$,

$$1 = \frac{1}{2} + \frac{1}{2\pi} \int_{-\pi}^{\pi} \Re\left(e^{-ik\theta} \frac{1}{1 - e^{i\theta}}\right) d\theta,$$

whence we deduce

$$\begin{aligned} (46) \quad u_k - \frac{1}{\mu} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \Re\left[e^{-ik\theta} \left(\frac{1}{1 - g(i\theta)} - \frac{1}{\mu} \frac{1}{1 - e^{i\theta}}\right)\right] d\theta \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ik\theta} \left(\frac{1}{1 - g(i\theta)} - \frac{1}{\mu} \frac{1}{1 - e^{i\theta}}\right) d\theta, \quad k \in \mathbb{N}, \end{aligned}$$

the last line justified by the fact that the integral is convergent. The integrand function

$$z \mapsto e^{-kz} \left(\frac{1}{1 - g(z)} - \frac{1}{\mu} \frac{1}{1 - e^z}\right)$$

is meromorphic on the domain $\{z \in \mathbb{C}; -\pi \leq \Im(z) \leq \pi, 0 \leq \Re(z) \leq r\}$, $r < R_0$. For $R_0 - r$ small enough, the poles inside this domain are exactly z_1, \dots, z_N (the pole at $z_0 = 0$ has been removed). Cauchy’s residue theorem with contour given in the left panel of Figure 3 implies

$$\begin{aligned} (47) \quad u_k - \frac{1}{\mu} &= - \sum_{j=1}^N \text{Res}\left(\frac{e^{-kz}}{1 - g(z)}; z_j\right) \\ &\quad + \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-k(r+i\theta)} \left(\frac{1}{1 - g(r+i\theta)} - \frac{1}{\mu} \frac{1}{1 - e^{r+i\theta}}\right) d\theta. \end{aligned}$$




FIG. 3. Contours used for the application of Cauchy's residue theorem in the lattice case (left) and the nonlattice case (right).

Using this, we obtain finally

$$\begin{aligned}
 v(k) &= \sum_{m=0}^{\infty} [v(k+m) - v(k+m+1)] \\
 &= \sum_{m=0}^{\infty} [1/\mu - u(k+m+1)] \\
 &= \sum_{m=0}^{\infty} \sum_{j=1}^N \text{Res} \left(\frac{e^{-(k+m+1)z}}{1-g(z)}; z_j \right) \\
 (48) \quad &\quad - \frac{1}{2\pi} \sum_{m=0}^{\infty} \int_{-\pi}^{\pi} e^{-(k+m+1)(r+i\theta)} \left(\frac{1}{1-g(r+i\theta)} - \frac{1}{\mu} \frac{1}{1-e^{r+i\theta}} \right) d\theta \\
 &= \sum_{j=1}^N \text{Res} \left(\frac{e^{-kz}}{(e^z-1)(1-g(z))}; z_j \right) \\
 &\quad - \frac{e^{-kr}}{2\pi} \int_{-\pi}^{\pi} \frac{e^{-ik\theta}}{e^{(r+i\theta)}-1} \left(\frac{1}{1-g(r+i\theta)} - \frac{1}{\mu} \frac{1}{1-e^{r+i\theta}} \right) d\theta.
 \end{aligned}$$

Here, we have used that all residues $\text{Res}(\frac{e^{-(k+m+1)z}}{1-g(z)}; z_j)$ are obtained by integrating $\frac{e^{-(k+m+1)z}}{1-g(z)}$ on a compact contour around z_j , so that exchanging $\sum_{m=0}^{\infty}$ and $\text{Res}(\cdot)$ is indeed justified by Fubini's theorem. By the Lebesgue lemma, the last term in (48) is $o(e^{-rk})$ and this proves equation (5). \square

4.2. Proof of Proposition 2. We prove that if F is absolutely continuous, then for all $R_0 < R$,

$$(49) \quad \lim_{\theta \rightarrow \pm\infty} \sup_{0 \leq r \leq R_0} |g(r+i\theta)| = 0.$$

Clearly, equation (49) implies Proposition 2. It is worth noting that this is a uniform version of the Riemann–Lebesgue lemma. Let us set for all $\theta \in \mathbb{R}$, $f_\theta(z) := g(z + i\theta)$, $z \in \mathbb{C}$. Since F admits a density, the Riemann–Lebesgue lemma implies that $f_\theta(z)$ converges pointwise to 0 as $\theta \rightarrow \infty$ when $z \in \Omega := \{z \in \mathbb{C} \mid 0 \leq \Re(z) \leq R_0\}$. Let us now note that we have the uniform bound

$$|f_\theta(z)| = |\mathbb{E}[e^{(z+i\theta)X}]| \leq \mathbb{E}[e^{R_0X}] < +\infty, \quad \forall z \in \Omega.$$

By Theorem 1.6.4, page 26 of [21], $f_\theta(z)$ converges uniformly towards 0 as $\theta \rightarrow \infty$ when z lies in any compact subset $K \subset \Omega$. Picking in particular $K := \{z \in \mathbb{C} \mid \Re(z) \in [0, R_0] \text{ and } \Im(z) = 0\}$ yields

$$\sup_{z \in K} |f_\theta(z)| = \sup_{r \in [0, R_0]} |g(r + i\theta)| \longrightarrow 0, \quad \theta \rightarrow \infty,$$

which we were set to prove.

4.3. *Proof of Theorem 3.* The proof of Theorem 3 follows the same lines as the proof of the main theorem in Stone [22].

For the sake of clarity, we divide the proof into several steps.

Step 1: We prove that Condition (6) implies that the equation $g(z) = 1$ has only a finite number of solutions in S_{R_0} . Condition (6) entails the existence of $M > 0$ such that the equation $g(r + i\theta) = 1$ has no solution with $0 \leq r \leq R_0$ and $|\theta| > M$. Since furthermore the obvious bound $|g(z)| < 1$ if $\Re(z) < 0$ exclude solutions in the half-plane $\Re(z) < 0$, the only possible solutions of $g(z) = 1$ in S_{R_0} belong to the compact set $K = \{z \in \mathbb{C}; 0 \leq \Re(z) \leq R_0, -M \leq \Im(z) \leq M\}$. The function $g(z) - 1$ being holomorphic, its zeros are isolated. Hence, the equation $g(z) = 1$ has finitely many solution in the compact set K .

Step 2: Following Stone [22], we introduce for positive a and h ,

$$V(x, h, a) := \mathbb{E}[U(x + h - aZ) - U(x - aZ)] = \mathbb{E}[U(x - aZ, h)]$$

with

$$U(x, h) = U(x + h) - U(x),$$

and Z a random variable with standard Gaussian distribution. Using Fourier analysis, one can show (similarly to equation (7) in [22])

$$\begin{aligned} (50) \quad V(x, h, a) &= \frac{h}{2\mu} + \frac{h}{2\pi} \int_{-\infty}^{+\infty} \Re \left\{ e^{-ix\theta} \frac{1 - e^{-ih\theta}}{ih\theta} e^{-a^2\theta^2/2} \frac{1}{1 - g(i\theta)} \right\} d\theta \\ &= \frac{h}{2\mu} + I(x, h, a). \end{aligned}$$

Setting $T(h, z) := \frac{1-e^{-hz}}{hz}$ and $\varphi(z) := \frac{1}{1-g(z)} + \frac{1}{\mu z}$, we then split the integral $I(x, h, a)$ into

$$\begin{aligned}
 I(x, h, a) &= I_1(x, h, a) - I_2(x, h, a), \\
 (51) \quad I_1(x, h, a) &= \frac{h}{2\pi} \int_{-\infty}^{+\infty} \Re\{e^{-ix\theta} T(h, i\theta) e^{-a^2\theta^2/2} \varphi(i\theta)\} d\theta, \\
 I_2(x, h, a) &= \frac{h}{2\pi} \int_{-\infty}^{+\infty} \Re\left\{e^{-ix\theta} T(h, i\theta) e^{-a^2\theta^2/2} \frac{1}{i\mu\theta}\right\} d\theta.
 \end{aligned}$$

The analysis of the first term $I_1(x, h, a)$ relies on Cauchy’s residue theorem. In the domain S_{R_0} , the function $\varphi(\cdot)$ is meromorphic with poles z_1, \dots, z_N (note that the pole at $z_0 = 0$ has been removed). We apply Cauchy’s residue theorem with the contour given in the right panel of Figure 3 and we let the parameter L tend to $+\infty$. Thanks to condition (6), the contribution of the horizontal parts of the contour vanishes as $L \rightarrow +\infty$ and we get, for all $r \in (\Re(z_N), R_0)$,

$$\begin{aligned}
 (52) \quad &\frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ix\theta} T(h, i\theta) e^{-a^2\theta^2/2} \varphi(i\theta) d\theta \\
 &= - \sum_{j=1}^N \text{Res}(e^{-xz} T(h, z) e^{a^2z^2/2} \varphi(z); z_j) + \Re \tilde{I}_1(x, h, a, r)
 \end{aligned}$$

with

$$(53) \quad \tilde{I}_1(x, h, a, r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-x(r+i\theta)} T(h, r+i\theta) e^{a^2(r+i\theta)^2/2} \varphi(r+i\theta) d\theta.$$

Note that the sum of residues is a real number because for conjugate poles z and \bar{z} , the residues at z and \bar{z} are also conjugate so that the sum is real. Similarly for the second term $I_2(x, h, a)$, Cauchy’s residue theorem yields

$$(54) \quad \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ix\theta} T(h, i\theta) e^{-a^2\theta^2/2} \frac{1}{i\mu\theta} d\theta = -\frac{1}{2\mu} + \tilde{I}_2(x, h, a, r)$$

with

$$(55) \quad \tilde{I}_2(x, h, a, r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-x(r+i\theta)} T(h, r+i\theta) e^{a^2(r+i\theta)^2/2} \frac{1}{\mu(r+i\theta)} d\theta.$$

Equations (50), (51), (52) and (54) together yield

$$\begin{aligned}
 (56) \quad V(x, h, a) &= \frac{h}{\mu} - h \sum_{j=1}^N \text{Res}(e^{-xz} T(h, z) e^{a^2z^2/2} \varphi(z); z_j) \\
 &\quad + h\Re\{\tilde{I}_1(x, h, a, r)\} - h\Re\{\tilde{I}_2(x, h, a, r)\}.
 \end{aligned}$$

Note that the Riemann–Lebesgue lemma implies that the second term $\tilde{I}_2(x, h, a, r)$ satisfies, for all $r > 0$,

$$(57) \quad \tilde{I}_2(x, h, a, r) = o(e^{-rx}) \quad \text{uniformly for } a \text{ and } h \text{ in bounded sets.}$$

Step 3: We analyze here the term $\tilde{I}_1(x, h, a, r)$ and, similarly as equation (8) from [22], we show that

$$(58) \quad \tilde{I}_1(x, h, a, r) = o(e^{-rx}(1 + |\ln(a)|))$$

uniformly for a and h in bounded sets.

The argument for this statement is given almost without proof in [22] but can be adapted from a similar argument in [23] in the following way. One notices that

$$\varphi(z) = \frac{g(z) - 1 - \mu z}{\mu^2 z^2} + \frac{(g(z) - 1 - \mu z)^2}{\mu^2 z^2(1 - g(z))},$$

so that we have, for $a < 1$,

$$(59) \quad \tilde{I}_1(x, h, a, r) = J_1 + J_2 + J_3 + J_4,$$

with

$$J_1 = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-x(r+i\theta)} T(h, r+i\theta) e^{a^2(r+i\theta)^2/2} \frac{g(r+i\theta) - 1 - \mu(r+i\theta)}{\mu^2(r+i\theta)^2} d\theta,$$

$$J_2 = \frac{1}{2\pi} \int_{|\theta| \leq 1} e^{-x(r+i\theta)} T(h, r+i\theta) \times e^{a^2(r+i\theta)^2/2} \frac{(g(r+i\theta) - 1 - \mu(r+i\theta))^2}{\mu^2(r+i\theta)^2(1 - g(r+i\theta))} d\theta.$$

$$J_3 = \frac{1}{2\pi} \int_{1 < |\theta| < a^{-2}} e^{-x(r+i\theta)} T(h, r+i\theta) \times e^{a^2(r+i\theta)^2/2} \frac{(g(r+i\theta) - 1 - \mu(r+i\theta))^2}{\mu^2(r+i\theta)^2(1 - g(r+i\theta))} d\theta.$$

$$J_4 = \frac{1}{2\pi} \int_{|\theta| \geq a^{-2}} e^{-x(r+i\theta)} T(h, r+i\theta) \times e^{a^2(r+i\theta)^2/2} \frac{(g(r+i\theta) - 1 - \mu(r+i\theta))^2}{\mu^2(r+i\theta)^2(1 - g(r+i\theta))} d\theta.$$

Since $\frac{g(z)-1-\mu z}{\mu^2 z^2}$ is analytic on S_R (the pole at $z = 0$ has been removed), Cauchy's residue theorem and Lebesgue's lemma yield

$$J_1 = o(e^{-rx}) \quad \text{uniformly for } a \text{ and } h \text{ in bounded sets.}$$

For the other terms, we use the fact that the function

$$F(\theta) := T(h, r+i\theta) e^{a^2(r+i\theta)^2/2} \frac{(g(r+i\theta) - 1 - \mu(r+i\theta))^2}{\mu^2(r+i\theta)^2(1 - g(r+i\theta))}$$

is bounded for $|\theta| \leq 1$ and satisfies

$$F'(\theta) = O\left(\frac{1}{\theta}\right) \quad \text{as } |\theta| \geq 1$$

and

$$F(\theta) = O(e^{-a^2\theta^2/2}) \quad \text{as } |\theta| \geq 1.$$

This justifies the following estimates:

$$\begin{aligned} |J_2| &= \frac{1}{2\pi} \left| \int_{|\theta| \leq 1} e^{-x(r+i\theta)} F(\theta) d\theta \right| = O(e^{-rx}), \\ |J_3| &= \frac{1}{2\pi} \left| \int_{1 < |\theta| < a^{-2}} e^{-x(r+i\theta)} F(\theta) d\theta \right| = O(e^{-rx} |\log(a)|), \\ |J_4| &= \frac{1}{2\pi} \left| \int_{|\theta| \geq a^{-2}} e^{-x(r+i\theta)} F(\theta) d\theta \right| = O(e^{-rx}). \end{aligned}$$

Gathering the above inequalities in (59), we obtain (58). The term $O(e^{-rx} |\log(a)|)$ can be replaced by $o(e^{-rx} |\log(a)|)$ because $r \in (\mathfrak{N}(z_N), R_0)$ is arbitrary. Equations (56), (57) and (58) together yield

$$\begin{aligned} (60) \quad V(x, h, a) &- \frac{h}{\mu} \\ &= -h \sum_{j=1}^N \text{Res}(e^{-xz} T(h, z) e^{a^2 z^2/2} \varphi(z); z_j) + o(e^{-rx} (1 + |\ln(a)|)). \end{aligned}$$

Step 4: We prove the following inequality:

$$\begin{aligned} (61) \quad V(x + e^{-rx}, 1 - 2e^{-rx}, e^{-rx}/x) &- Me^{-rx} \leq U(x, 1) \\ &\leq (1 - e^{-rx})^{-1} V(x - e^{-rx}, 1 + 2e^{-rx}, e^{-rx}/x). \end{aligned}$$

Recall the definition $V(x, h, a) = \mathbb{E}[U(x - aZ, h)]$ with $U(x, h) = U(x + h) - U(x)$ and Z a standard Gaussian random variable. Using the fact that $U(\cdot)$ is non-decreasing, we check, for $|y| \leq e^{-rx}$,

$$U(x + e^{-rx} - y, 1 - 2e^{-rx}) \leq U(x, 1) \leq U(x - e^{-rx} - y, 1 + 2e^{-rx}).$$

Let $x_0 > 0$ be such that, for all $x \geq x_0$, $\mathbb{P}(|Z| \geq x) \leq e^{-rx} \leq 1/2$. Besides, the fact that renewal function $x \mapsto U(x)$ is sub-additive implies that there exists some constant M independent from h and x such that

$$(62) \quad 0 \leq U(x, h) \leq M, \quad \text{for all } x > 0 \text{ and } h \text{ in a bounded set.}$$

By the definition $V(x, h, a) = \mathbb{E}[U(x - aZ, h)]$, both sides of equation (61) are obtained by splitting

$$\begin{aligned} V(x \pm e^{-rx}, h, a) &= \mathbb{E}[U(x \pm e^{-rx} - aZ, h) \mathbb{1}_{\{|Z| \leq x\}}] + \mathbb{E}[U(x \pm e^{-rx} - aZ, h) \mathbb{1}_{\{|Z| > x\}}] \end{aligned}$$

with the values $a = a(x) = e^{-rx}/x$ and $h = h(x) = 1 \pm 2e^{-rx}$.

Step 5: Let us now prove the following estimate:

$$(63) \quad U(x, 1) - \frac{1}{\mu} = - \sum_{j=1}^N \text{Res}(e^{-xz}T(1, z)\varphi(z); z_j) + o(e^{-rx}).$$

This is roughly obtained from (60) by setting

$$a = a(x) = e^{-rx}/x \quad \text{and} \quad h = h_{\pm}(x) = 1 \pm 2e^{-rx}.$$

This can be made rigorous thanks to equation (61). We have, uniformly in the neighborhood of z_j , $j = 1, \dots, N$,

$$\begin{aligned} e^{-(x \pm e^{-rx})z}T(h(x), z)e^{-a(x)^2z^2/2} \\ = e^{-xz}[1 + o(e^{-rx})][T(1, z) + o(e^{-rx})][1 + o(e^{-rx})] \\ = e^{-xz}T(1, z) + o(e^{-rx}) \end{aligned}$$

so that

$$(64) \quad \begin{aligned} \text{Res}(e^{-(x \pm e^{-rx})z}T(h, z)e^{a^2z^2/2}\varphi(z); z_j) \\ = \text{Res}(e^{-xz}T(1, z)\varphi(z); z_j) + o(e^{-rx}). \end{aligned}$$

Besides, left-hand side of equation (61) and equation (64) entail

$$(65) \quad \begin{aligned} U(x, 1) - \frac{1}{\mu} + \sum_{j=1}^N \text{Res}(e^{-xz}T(1, z)\varphi(z); z_j) \\ \geq V(x + e^{-rx}, 1 - 2e^{-rx}, e^{-rx}/x) - Me^{-rx} - \frac{1}{\mu} \\ + \sum_{j=1}^N \text{Res}(e^{-xz}T(1, z)\varphi(z); z_j) \\ = V(x + e^{-rx}, h(x), a(x)) - Me^{-rx} \\ - \frac{1}{\mu} + \sum_{j=1}^N \text{Res}(e^{-(x \pm e^{-rx})z}T(h, z)e^{a^2z^2/2}\varphi(z); z_j) + o(e^{-rx}). \end{aligned}$$

Now from (60), we have

$$\begin{aligned} V(x + e^{-rx}, h(x), a(x)) - \frac{h(x)}{\mu} \\ + h(x) \sum_{j=1}^N \text{Res}(e^{-(x \pm e^{-rx})z}T(h, z)e^{a^2z^2/2}\varphi(z); z_j) \\ = o(xe^{-rx}). \end{aligned}$$

Together with (65) and since $h(x) = 1 + O(e^{-rx})$, this yields

$$(66) \quad U(x, 1) - \frac{1}{\mu} + \sum_{j=1}^N \operatorname{Res}(e^{-xz}T(1, z)\varphi(z); z_j) \geq o(xe^{-rx}).$$

A similar upper bound is proved in the same way, using the right-hand side of equation (61). We deduce

$$U(x, 1) - \frac{1}{\mu} + \sum_{j=1}^N \operatorname{Res}(e^{-xz}T(1, z)\varphi(z); z_j) = o(xe^{-rx}).$$

We can replace $o(xe^{-rx})$ by $o(e^{-rx})$ because r is arbitrary in $(\Re(z_N), R_0)$. This proves equation (63).

Step 6: We finally prove equation (7). Since $\lim_{x \rightarrow \infty} v(x) = 0$, we have

$$(67) \quad v(x) = \sum_{k=0}^{\infty} [v(x+k) - v(x+k+1)] = \sum_{k=0}^{\infty} [-U(x+k, 1) + 1/\mu].$$

Using equation (63), we deduce, for all $r \in (\Re(z_N), R_0)$,

$$(68) \quad \begin{aligned} v(x) &= \sum_{k=0}^{\infty} \sum_{j=1}^N \operatorname{Res}(e^{-(x+k)z}T(1, z)\varphi(z); z_j) + o(e^{-r(x+k)}) \\ &= \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-xz}}{1 - e^{-z}}T(1, z)\varphi(z); z_j\right) + o(e^{-rx}) \\ &= \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-xz}}{z(1 - g(z))}; z_j\right) + o(e^{-rx}). \end{aligned}$$

Equation (7) follows easily.

4.4. Proof of Corollaries 4 and 6.

PROOF OF COROLLARY 4. We consider first the lattice case. According to Condition (8), one can consider $(r_n)_{n \geq 1}$ an increasing sequence such that $r_n \rightarrow +\infty$ and

$$\lim_{n \rightarrow +\infty} \sup_{-\pi \leq \theta \leq \pi} \left| \frac{1}{e^{r_n}(1 - g(r_n + i\theta))} \right| = 0.$$

According to equation (48),

$$\begin{aligned} v(k) &= \sum_{j=1}^{N(r_n)} \operatorname{Res}\left(\frac{e^{-kz}}{(e^z - 1)(1 - g(z))}; z_j\right) \\ &\quad - \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{e^{-k(r_n+i\theta)}}{e^{r_n+i\theta} - 1} \left(\frac{1}{1 - g(r_n + i\theta)} - \frac{1}{\mu} \frac{1}{1 - e^{r_n+i\theta}} \right) d\theta \end{aligned}$$

with $N(r_n)$ the number of solutions of the equation $g(z) = 1$ in $S_{r_n}^f$. Lebesgue’s dominated convergence theorem implies, for $k \geq 1$,

$$\lim_{n \rightarrow +\infty} \int_{-\pi}^{\pi} \frac{e^{-k(r_n+i\theta)}}{e^{r_n+i\theta} - 1} \left(\frac{1}{1 - g(r_n + i\theta)} - \frac{1}{\mu} \frac{1}{1 - e^{r_n+i\theta}} \right) d\theta = 0.$$

This yields the result (9).

We now consider the nonlattice case. Thanks to Assumption (11), let $(r_n)_{n \geq 1}$ be such that $r_n \rightarrow +\infty$ and

$$\sup_{\theta \in \mathbb{R}} \left| \frac{1}{r_n(1 - g(r_n + i\theta))} \right| \rightarrow 0, \quad n \rightarrow \infty.$$

We will prove below that setting $h = 1$, $r = r_n$ and $a = a_n(x) = e^{-r_n x}/x$ in equation (56) and letting $n \rightarrow +\infty$, we obtain

$$(69) \quad U(x, 1) = \frac{1}{\mu} - \sum_{j=1}^{\infty} \text{Res}(e^{-xz} T(1, z) \varphi(z); z_j) = \frac{1}{\mu} - \sum_{j=1}^{\infty} \frac{e^{-xz_j} (e^{-z_j} - 1)}{z_j g'(z_j)},$$

remembering that roots $(z_j)_{j \in \mathbb{N}}$ are simple. This is justified as follows:

– Condition (62) together with Lebesgue’s dominated convergence theorem implies

$$(70) \quad V(x, h, a) = V(x, 1, e^{-r_n x}/x) \rightarrow U(x, h) \quad \text{as } n \rightarrow +\infty.$$

– Recalling that $\tilde{I}_1(x, h, a, r)$ and $\tilde{I}_2(x, h, a, r)$ are respectively given by (53) and (55), the integral term $\tilde{I}_1(x, h, a, r) - \tilde{I}_2(x, h, a, r)$ with $h = 1$, $r = r_n$ and $a = e^{-r_n x}/x$ converge to 0 as $n \rightarrow +\infty$. Indeed, condition (8) and the upper bound $|T(h, r + i\theta)| \leq 2/r$ imply

$$\begin{aligned} & |\tilde{I}_1(x, h, a, r) - \tilde{I}_2(x, h, a, r)| \\ &= \frac{1}{2\pi} \left| \int_{-\infty}^{+\infty} e^{-x(r+i\theta)} T(h, r + i\theta) e^{a^2(r+i\theta)^2/2} \frac{1}{1 - g(r + i\theta)} d\theta \right| \\ &\leq \frac{e^{-xr}}{\pi} \int_{-\infty}^{+\infty} \left| \frac{1}{r(1 - g(r + i\theta))} \right| e^{a^2 r^2/2 - a^2 \theta^2/2} d\theta \\ &\leq \sup_{\theta \in \mathbb{R}} \left| \frac{1}{r(1 - g(r + i\theta))} \right| \times \frac{e^{-xr}}{\pi} \frac{e^{a^2 r^2/2} \sqrt{2\pi}}{a} \\ &\leq \sup_{\theta \in \mathbb{R}} \left| \frac{1}{r_n(1 - g(r_n + i\theta))} \right| \times \frac{\sqrt{2x}}{\sqrt{\pi}} \exp\left(\frac{r_n^2 e^{-2r_n x}}{2x}\right). \end{aligned}$$

This last quantity goes to 0 as $r_n \rightarrow +\infty$.

– Let us recall inequality $|e^z - 1| \leq e|z|$ for all $|z| \leq 1$. As roots $(z_n)_{n \in \mathbb{N}}$ are simple and real, one has for all $j = 1, \dots, N(r_n)$, and for n large enough,

$$\begin{aligned} & \left| \operatorname{Res}(e^{-(x \pm e^{-r_n x})z} T(1, z) e^{a_n(x)^2 z^2 / 2} \varphi(z); z_j) - \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right| \\ &= \left| \operatorname{Res}([e^{\pm e^{-r_n x} z} e^{a_n(x)^2 z^2 / 2} - 1] e^{-xz} T(1, z) \varphi(z); z_j) \right| \\ &= \left| [e^{\pm e^{-r_n x} z_j} e^{a_n(x)^2 z_j^2 / 2} - 1] \right| \cdot \left| \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right| \\ &\leq e \left| \pm e^{-r_n x} z_j + a_n(x)^2 z_j^2 / 2 \right| \cdot \left| \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right| \\ &\leq M e^{-2r_n x} r_n^2 \cdot \left| \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right|, \end{aligned}$$

where $M = M(x)$ is a constant independent from n and j , but which may depend on $x > 0$. One thus deduces

$$\begin{aligned} & \sum_{j=1}^{N(r_n)} \left| \operatorname{Res}(e^{-(x \pm e^{-r_n x})z} T(1, z) e^{a_n(x)^2 z^2 / 2} \varphi(z); z_j) \right. \\ (71) \quad & \left. - \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right| \\ & \leq M e^{-2r_n x} r_n^2 \sum_{j=1}^{N(r_n)} \left| \operatorname{Res}(e^{-xz} T(1, z) \varphi(z); z_j) \right| \end{aligned}$$

which tends to zero as $n \rightarrow \infty$ because of Assumption (10).

– The fact that $\tilde{I}_1(x, h, a_n(x), r_n) - \tilde{I}_2(x, h, a_n(x), r_n)$ tends to 0 as $n \rightarrow \infty$ and convergence (70) (with $h = 1$) as well as inequality (71) implies (69) from (56).

The end of the proof follows easily, as in Step 6 of the proof of Theorem 3: equations (67) and (69) imply (12). \square

PROOF OF COROLLARY 6. In the lattice case, we deduce from Corollary 4 that

$$\begin{aligned} u(k) &= U(k) - U(k - 1) = v(k) - v(k - 1) + \frac{1}{\mu} \\ &= \frac{1}{\mu} + \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-kz} - e^{-(k-1)z}}{(e^z - 1)(1 - g(z))}; z_j\right) = \frac{1}{\mu} - \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-kz}}{1 - g(z)}; z_j\right). \end{aligned}$$

Finally, note that $z_0 = 0, g(0) = 1, g'(0) = \mu$ so that

$$\frac{1}{\mu} = \operatorname{Res}\left(\frac{e^{-kz}}{1 - g(z)}; z_0\right).$$

In the nonlattice case, we deduce from Corollary 4 that

$$u(x) = \frac{dU(x)}{dx} = \frac{1}{\mu} + \frac{dv(x)}{dx} = \frac{1}{\mu} - \sum_{j=1}^N \operatorname{Res}\left(\frac{e^{-kz}}{1 - g(z)}; z_j\right).$$

The result follows since $\frac{1}{\mu} = \text{Res}\left(\frac{e^{-kz}}{1-g(z)}; z_0\right)$. \square

5. Proofs for Section 3.

5.1. *Proof of Theorem 10.* Assumption (20) that the moment generating function of G is finite everywhere is equivalent to the fact that the tail function $\bar{G} = 1 - G$ decreases super-exponentially fast, that is, \bar{G} is in \mathcal{D} with

$$(72) \quad \mathcal{D} := \{f : [0, +\infty) \rightarrow \mathbb{C} \mid \forall r > 0, f(x) = o(e^{-rx}) \text{ as } x \rightarrow \infty\}.$$

The following lemma can be verified easily.

LEMMA 19. *Let $f \in \mathcal{D}$, then:*

1. $x \mapsto e^{zx} f(x)$ belongs to \mathcal{D} for all $z \in \mathbb{C}$;
2. $x \mapsto \int_x^\infty f(y) dy$ belongs to \mathcal{D} .

PROOF OF THEOREM 10. It is well known that $\psi(x)$ satisfies the following defective renewal equation:

$$(73) \quad \psi(x) = \bar{L}(x) + \int_0^x \psi(x - y) dL(y),$$

where

$$(74) \quad L(x) := \int_0^x \frac{\alpha}{c} \bar{G}(y) dy, \quad x \geq 0,$$

can be seen as the c.d.f. of a defective distribution [because $L(+\infty) = \frac{\alpha}{c} m < 1$]; see, for example, Corollary 3.3, page 79 of Asmussen and Albrecher [2] or equation (7.2) page 377 of Feller [9].

We introduce the probability distribution $dF(x) = e^{\kappa x} dL(x)$ and use the notation:

$$(75) \quad \begin{aligned} g(z) &:= \int_0^\infty e^{zx} dF(x) = \frac{\alpha}{c} \frac{1}{z + \kappa} \mathbb{E}(e^{(z+\kappa)Z} - 1), \\ \mu &:= \int_0^\infty x dF(x) \quad \text{and} \quad \mu_2 := \int_0^\infty x^2 dF(x) \end{aligned}$$

for the moment generating function and the first and second moments of F . Elementary calculation yields

$$(76) \quad g'(z_j) = \frac{1}{z_j + \kappa} \left[\frac{\alpha}{c} \mathbb{E}(Z e^{(z_j+\kappa)Z}) - 1 \right].$$

Thanks to the assumption $\bar{G} \in \mathcal{D}$ and Lemma 19, the tail function of F belongs also to \mathcal{D} whence the moment generating function $g(z)$ is defined for all $z \in \mathbb{C}$. Setting $\psi_\kappa(x) := e^{\kappa x} \psi(x)$, (73) entails the nondefective renewal equation:

$$(77) \quad \psi_\kappa(x) = z(x) + \int_0^x \psi_\kappa(x - y) dF(y), \quad z(x) := e^{\kappa x} \bar{L}(x),$$

see Feller [9] equation (6.12), page 376. The distribution F is nonlattice so that Smith’s renewal theorem entails the classical Cramer–Lundberg asymptotics:

$$(78) \quad \psi_\kappa(x) \rightarrow C := \frac{\int_0^\infty z(y) dy}{\int_0^\infty y dF(y)} = \frac{c - \alpha m}{c - \alpha \mathbb{E}(Z_1 e^{\kappa Z_1})} \quad \text{as } x \rightarrow +\infty,$$

the last equality being obtained by direct computation; see, for example, Theorem 5.3 page 86 of [2].

We now wish to provide an expansion of $\psi_\kappa(x)$ and provide extra terms in (78). Let us introduce

$$U(x) = \sum_{n=0}^\infty F^{*(n)}(x), \quad v(x) = U(x) - \frac{x}{\mu} - \frac{\mu_2}{2\mu^2}, \quad x \geq 0.$$

The solution to (77) is given by

$$(79) \quad \psi_\kappa(x) = \int_0^x z(x - y) dU(y) = \int_0^x z(x - y) \frac{dy}{\mu} + \int_0^x z(x - y) dv(y).$$

Using the fact that $\int_0^x z(x - y) dy = \int_0^x z(y) dy$, an integration by parts and the definition of C , we obtain

$$(80) \quad \begin{aligned} \psi_\kappa(x) - C &= - \int_x^\infty z(y) \frac{dy}{\mu} + \int_0^x z'(x - y)v(y) dy - v(0-)z(x) + \frac{\alpha}{c}mv(x) \\ &:= I_1(x) + I_2(x) + I_3(x) + I_4(x). \end{aligned}$$

We recall that $x \mapsto \bar{G}(x)$ belongs to \mathcal{D} defined in (72), so that one easily sees that $z(x)$ defined in (77) belongs to \mathcal{D} , and that $I_1(x)$ and $I_3(x)$ are $o(e^{-rx})$ from Lemma 19. Thus, we only need to study asymptotics of the terms $I_2(x)$ and $I_4(x)$ as $x \rightarrow \infty$ in (80).

We start with $I_4(x)$. Let us note that, as $g(z)$ is defined for all $z \in \mathbb{C}$, R defined in (2) is equal to $+\infty$, as well as R_0 in Theorem 3. Expansion (7) in Theorem 3 yields

$$(81) \quad I_4(x) = \sum_{j=1}^N \frac{\alpha}{c} m \operatorname{Res} \left(\frac{e^{-xz}}{z(1 - g(z))}; z_j \right) + o(e^{-rx}).$$

Now turning to $I_2(x)$. By a change of variable one has that $I_2(x) = \int_0^x z'(y)v(x - y) dy$, so that, using Expansion (7) in Theorem 3 for $v(x - y)$ yields

$$(82) \quad \begin{aligned} I_2(x) &= \sum_{j=1}^N \int_0^x z'(y) \operatorname{Res} \left(\frac{e^{-(x-y)z}}{z(1 - g(z))}; z_j \right) dy \\ &\quad + \int_0^x z'(y)e^{-r(x-y)} \varepsilon(x - y) dy \end{aligned}$$

for some function $\varepsilon(\cdot)$ vanishing at $+\infty$. Furthermore, it is easy to check that $x \mapsto z'(x) \in \mathcal{D}$. One then verifies that $\int_x^\infty z'(y) \operatorname{Res} \left(\frac{e^{-(x-y)z}}{z(1 - g(z))}; z_j \right) dy$ is an $o(e^{-rx})$,

by writing the residue as an integral over a fixed contour including z_j then using Lemma 19. One also checks that the last term on the right-hand side of (82) is $o(e^{-rx})$ by a dominated convergence argument. One may thus write in (82)

$$(83) \quad I_2(x) = \sum_{j=1}^N \int_0^\infty z'(y) \operatorname{Res}\left(\frac{e^{-(x-y)z}}{z(1-g(z))}; z_j\right) dy + o(e^{-rx}).$$

Deriving $z(y) = e^{\kappa y} \bar{L}(y)$ with $L(y)$ given in (74), expressing the residue as an integral over a fixed contour and using Fubini, thus yields the following expression for the terms in the summation on the right-hand side of (83):

$$\begin{aligned} & \int_0^\infty z'(y) \operatorname{Res}\left(\frac{e^{-(x-y)z}}{z(1-g(z))}; z_j\right) dy \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \int_0^\infty z'(y) e^{yz} dy; z_j\right) \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \left[\int_0^\infty \kappa e^{(z+\kappa)y} \int_y^\infty \frac{\alpha}{c} \bar{G}(v) dv dy \right. \right. \\ &\quad \left. \left. - \int_0^\infty e^{(z+\kappa)y} \frac{\alpha}{c} \bar{G}(y) dy \right]; z_j\right) \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \left[\frac{\alpha}{c} \mathbb{E}\left(\int_0^\infty \int_0^\infty \kappa e^{(z+\kappa)y} \mathbb{1}_{[Z \geq v \geq y]} dy dv\right) \right. \right. \\ &\quad \left. \left. - \frac{\alpha}{c} \mathbb{E}\left(\int_0^\infty e^{(z+\kappa)y} \mathbb{1}_{[Z \geq y]} dy\right) \right]; z_j\right) \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \left[\frac{\alpha}{c} \frac{\kappa}{z+\kappa} \mathbb{E}\left(\frac{e^{(z+\kappa)Z} - 1}{z+\kappa} - Z\right) \right. \right. \\ &\quad \left. \left. - \frac{\alpha}{c} \frac{1}{z+\kappa} \mathbb{E}(e^{(z+\kappa)Z} - 1) \right]; z_j\right), \end{aligned}$$

which, since z_j satisfies $g(z_j) = 1$ with $g(z)$ given by (75), yields

$$(84) \quad \begin{aligned} & \int_0^\infty z'(y) \operatorname{Res}\left(\frac{e^{-(x-y)z}}{z(1-g(z))}; z_j\right) dy \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \left[\frac{\kappa}{z+\kappa} - \frac{\alpha}{c} \frac{\kappa}{z+\kappa} m - 1 \right]; z_j\right) \\ &= \operatorname{Res}\left(\frac{e^{-xz}}{z(1-g(z))} \frac{[-z - \frac{\alpha}{c} \kappa m]}{z+\kappa}; z_j\right). \end{aligned}$$

Plugging (84) in (83) and summing $I_2(x)$ and Expression (81) of $I_4(x)$ then yields from (80)

$$\psi_k(x) - C = \sum_{j=1}^N \operatorname{Res} \left(\frac{e^{-xz}}{1-g(z)} \frac{[-1 + \frac{\alpha}{c}m]}{z+\kappa}; z_j \right) + o(e^{-rx}).$$

Let us now note that in the case where $g'(z_j) \neq 0$, the residue in the j th term in the above can be explicitly given and, thanks to (76), is equal to (27). Let us also note that expression of C provided in (78) is but $-\frac{-1+\frac{\alpha}{c}m}{\frac{\alpha}{c}\mathbb{E}(Ze^{(z_0+\kappa)Z})-1}$ with $z_0 = 0$ and that $g'(z_0) \neq 0$, so that the above yields Expansion (39). \square

5.2. *Proof of Theorem 15.* We let $b(k) := \mathbb{P}(Z_1 = k)$, $k \in \mathbb{N}$, the probability mass distribution of Z_1 and denote its survival function by $l(k) := \mathbb{P}[Z_1 \geq k + 1] = \sum_{j=k+1}^\infty b(j)$. Let us in particular note that $b(0)$ can be positive. In other words, there is a possibility that there is no claim (or equivalently, a claim of size 0) at time n . We define

$$\mathcal{D}_d := \{f : \mathbb{N} \longrightarrow \mathbb{C} \mid f(k) = o(e^{-sk}) \forall s > 0\}$$

and suppose here that the Z_k 's verify that $k \mapsto l(k) \in \mathcal{D}_d$. we let $\kappa > 0$ its unique real positive solution.

The following lemma is the discrete analog of Lemma 19.

LEMMA 20. *Let $f \in \mathcal{D}_d$, then:*

- 1. $k \in \mathbb{N} \mapsto e^{zk} f(k)$ belongs to \mathcal{D}_d for all $z \in \mathbb{C}$,
- 2. $k \in \mathbb{N} \mapsto \sum_{j=k}^\infty f(j)$ belongs to \mathcal{D}_d .

Similarly to (74), (23) and (75) we define the following discrete measures $F(\cdot)$, associated probability mass function $f(\cdot)$ and complex valued function $g(\cdot)$ which we recall here:

$$F(x) := \sum_{k=0}^x f(k), \quad x \in \mathbb{N}, \tag{85}$$

$$f(k) := e^{\kappa k} l(k) = e^{\kappa k} \mathbb{P}(Z \geq k + 1), \quad k \in \mathbb{N},$$

$$g(z) := \int_0^\infty e^{zx} dF(x) = \sum_{k=0}^\infty e^{(z+\kappa)k} \mathbb{P}(Z \geq k + 1) \tag{86}$$

$$= \frac{1 - \mathbb{E}(e^{(z+\kappa)Z})}{1 - e^{z+\kappa}}, \quad z \in \mathbb{C}.$$

$g(z)$ is thus defined for all $z \in \mathbb{C}$ and is equal to $\mathbb{E}(e^{zX})$ for some integer valued r.v. X with probability mass function $f(\cdot)$, and with expectation and second moment $\mathbb{E}(X) := \mu$ and $\mathbb{E}(X^2) := \mu_2$.

PROOF OF THEOREM 15. We proceed along the lines of proof of Proposition 10. Setting $\bar{L}(x) := \sum_{y=x}^{\infty} l(y)$, $x \in \mathbb{N}$, the discrete time analog of (73) is

$$(87) \quad \psi(x) = \sum_{k=0}^x \psi(x-k)l(k) + \bar{L}(x), \quad x \in \mathbb{N};$$

see, for example, Proposition 1.2, page 488 in [2]. Letting $\psi_\kappa(x) := e^{\kappa x} \psi(x)$ and $z(x) := e^{\kappa x} \bar{L}(x)$, and by the definition of $f(k)$ for all k in (85), we obtain from (87) the renewal equation:

$$(88) \quad \psi_\kappa(x) = z(x) + \sum_{k=0}^x \psi_\kappa(x+1-k)f(k).$$

Since X is lattice, the corresponding Smith’s renewal theorem implies that

$$(89) \quad \psi_\kappa(x) \longrightarrow C := \frac{1}{\mu} \sum_{k=0}^{\infty} z(k) = \frac{m - e^\kappa}{e^\kappa - \mathbb{E}(Ze^{\kappa Z})}, \quad x \rightarrow \infty,$$

the last equality can be verified by direct computation. Let us note in passing that (89) provides the Cramer–Lundberg asymptotics $\psi(x) \sim \frac{m - e^\kappa}{e^\kappa - \mathbb{E}(Ze^{\kappa Z})} e^{-\kappa x}$ for $x \rightarrow \infty$, $x \in \mathbb{N}$, in the discrete case. Defining now

$$U(j) := \sum_{n=0}^{\infty} F^{*n}(j), \quad u_j := U(j) - U(j-1),$$

$$v(j) := U(j) - \frac{j}{\mu} - \frac{\mu_2 + \mu^2}{2\mu^2}, \quad j \in \mathbb{N}$$

with $U(-1) = 0$, solution to (88) is given by

$$\begin{aligned} \psi_\kappa(x) &= \sum_{k=0}^x z(x-k)u_k = \sum_{k=0}^x z(x-k) \left[\frac{1}{\mu} + v(k) - v(k-1) \right] \\ &= \frac{1}{\mu} \sum_{k=0}^x z(x-k) + \sum_{k=0}^x z(x-k)v(k) - \sum_{k=-1}^{x-1} z(x-k-1)v(k) \\ &= \frac{1}{\mu} \sum_{k=0}^x z(k) + \sum_{k=0}^{x-1} [z(x-k) - z(x-k-1)]v(k) \\ &\quad + z(0)v(x) - z(x)v(-1). \end{aligned}$$

Subtracting in the above constant C given by (89) yields

$$(90) \quad \begin{aligned} \psi_\kappa(x) - C &= -\frac{1}{\mu} \sum_{k=x+1}^{\infty} z(k) + \sum_{k=0}^{x-1} [z(x-k) - z(x-k-1)]v(k) \\ &\quad + z(0)v(x) - z(x)v(-1). \end{aligned}$$

The first and last term on the right-hand side of (90) are $o(e^{-rk})$ by Lemma 20, hence we are interested in the following quantities:

$$I_1(x) := \sum_{k=0}^{x-1} [z(x-k) - z(x-k-1)]v(k) = \sum_{k=1}^x [z(k) - z(k-1)]v(x-k),$$

$$I_2(x) := z(0)v(x) = \bar{L}(0)v(x) = mv(x).$$

Thanks to Theorem 1, one has that

$$(91) \quad I_2(x) = \sum_{j=0}^N \text{Res} \left(\frac{e^{-kz}}{(e^z - 1)(1 - g(z))}; z_j \right) + o(e^{-rx})$$

and we then turn to $I_1(x)$. Writing

$$(92) \quad \begin{aligned} z(k) - z(k-1) &= e^{\kappa k} \bar{L}(k) - e^{\kappa(k-1)} \bar{L}(k-1) \\ &= [1 - e^{-\kappa}] e^{\kappa k} \bar{L}(k) + e^{\kappa(k-1)} [\bar{L}(k) - \bar{L}(k-1)] \\ &= [1 - e^{-\kappa}] e^{\kappa k} \bar{L}(k) - e^{\kappa(k-1)} l(k-1) \\ &= [1 - e^{-\kappa}] z(k) - e^{\kappa(k-1)} l(k-1), \end{aligned}$$

then writing, in view of expansion (5),

$$v(x-k) = \sum_{j=1}^N \text{Res} \left(\frac{e^{-(x-k)z}}{(e^z - 1)(1 - g(z))}; z_j \right) + \varepsilon_{x-k} e^{-r(x-k)}$$

for some $(\varepsilon_n)_{n \in \mathbb{N}}$ vanishing at ∞ , one obtains

$$(93) \quad I_1(x) = \sum_{j=1}^N \text{Res} \left[\frac{1}{(e^z - 1)(1 - g(z))} \sum_{k=1}^x [1 - e^{-\kappa}] e^{\kappa k} \bar{L}(k) e^{-(x-k)z}; z_j \right]$$

$$(94) \quad - \sum_{j=1}^N \text{Res} \left[\frac{1}{(e^z - 1)(1 - g(z))} \sum_{k=1}^x e^{\kappa(k-1)} l(k-1) e^{-(x-k)z}; z_j \right]$$

$$(95) \quad + \sum_{k=1}^x [z(k) - z(k-1)] \varepsilon_{x-k} e^{-r(x-k)}.$$

Using Lemma 20 as well as a dominated convergence theorem, it is not hard to see that (95) is an $o(e^{-rx})$ as $x \rightarrow \infty$, $x \in \mathbb{N}$. We then study (93) and (94). One verifies that $x \in \mathbb{N} \mapsto \text{Res}[\frac{1}{(e^z-1)(1-g(z))} \sum_{k=x+1}^{\infty} e^{\kappa(k-1)} l(k-1) e^{kz}; z_j] \in \mathcal{D}_d$ by writing the residue as an integral over a fixed contour including z_j then using Lemma 20, so that in (94) we compute

$$\begin{aligned}
 & \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} e^{-xz} \sum_{k=1}^x e^{\kappa(k-1)} l(k-1) e^{kz}; z_j \right] \\
 &= \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} e^{-xz} e^z \sum_{k=0}^{\infty} e^{(z+\kappa)k} \mathbb{P}(Z \geq k+1); z_j \right] \\
 (96) \quad & - \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} e^{-xz} \sum_{k=x+1}^{\infty} e^{\kappa(k-1)} l(k-1) e^{kz}; z_j \right] \\
 &= \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} e^{-xz} e^z g(z); z_j \right] + o(e^{-rx}), \quad j = 1, \dots, N.
 \end{aligned}$$

Next, we compute, for all z ,

$$\begin{aligned}
 \sum_{k=1}^{\infty} e^{(\kappa+z)k} \bar{L}(k) &= \sum_{k=1}^{\infty} \sum_{n=k}^{\infty} e^{(\kappa+z)k} l(n) = \sum_{n=1}^{\infty} \left[\sum_{k=1}^n e^{(\kappa+z)k} \right] l(n) \\
 &= \sum_{n=1}^{\infty} \left[e^{\kappa+z} \frac{1 - e^{(\kappa+z)n}}{1 - e^{\kappa+z}} \right] l(n) \\
 (97) \quad &= \frac{e^{\kappa+z}}{1 - e^{\kappa+z}} \sum_{n=1}^{\infty} l(n) - \frac{1}{1 - e^{\kappa+z}} e^{\kappa+z} \sum_{n=1}^{\infty} e^{(\kappa+z)n} l(n) \\
 &= \frac{e^{\kappa+z}}{1 - e^{\kappa+z}} [m - l(0)] - \frac{1}{1 - e^{\kappa+z}} e^{\kappa+z} [g(z) - l(0)] \\
 &= \frac{e^{\kappa+z}}{1 - e^{\kappa+z}} [m - g(z)].
 \end{aligned}$$

One verifies thanks to Lemma 20 and writing the residue as a contour around z_j that $x \in \mathbb{N} \mapsto \text{Res}[\frac{1}{(e^z-1)(1-g(z))} \sum_{k=x+1}^{\infty} e^{(\kappa+z)k} \bar{L}(k); z_j] \in \mathcal{D}_d$, so that inserting (97) in (93) yields

$$\begin{aligned}
 & \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} \sum_{k=1}^x [1 - e^{-\kappa}] e^{\kappa k} \bar{L}(k) e^{-(x-k)z}; z_j \right] \\
 (98) \quad &= \text{Res} \left[\frac{1}{(e^z-1)(1-g(z))} e^{-xz} \frac{e^{\kappa+z}}{1 - e^{\kappa+z}} [m - g(z)]; z_j \right] + o(e^{-rx}).
 \end{aligned}$$

Respectively inserting (96) and (98) in (94) and (93) in $I_1(x)$, and adding $I_2(x)$ in (91), yields

$$\begin{aligned}
 I_1(x) + I_2(x) &= \sum_{j=1}^N \operatorname{Res} \left[\frac{1 - e^{-\kappa}}{(e^z - 1)(1 - g(z))} \frac{e^{\kappa+z}}{1 - e^{\kappa+z}} [m - g(z)] e^{-xz}; z_j \right] \\
 &\quad - \sum_{j=1}^N \operatorname{Res} \left[\frac{g(z)}{(e^z - 1)(1 - g(z))} e^z e^{-xz}; z_j \right] \\
 (99) \quad &\quad + \sum_{j=0}^N \operatorname{Res} \left[\frac{m}{(e^z - 1)(1 - g(z))} e^{-xz}; z_j \right] + o(e^{-rx}) \\
 &= - \sum_{j=1}^N \operatorname{Res} \left[\frac{1}{1 - g(z)} \frac{m - g(z)e^{\kappa+z}}{1 - e^{\kappa+z}} e^{-xz}; z_j \right] + o(e^{-rx}).
 \end{aligned}$$

Plugging (99) into (90) yields (39), provided that we prove that residue of $-\frac{1}{1-g(z)} \frac{m-g(z)e^{\kappa+z}}{1-e^{\kappa+z}} e^{-xz}$ at $z = z_0 = 0$ equals $Ce^{-\kappa x}$, C given by (89). This computation is in fact included in the case where the $g'(z_j)$'s are nonzero, in which case j th term of (99) is equal to (40), which we proceed to consider now. Direct computation from (86) coupled to the fact that $g(z_j) = 1$ gives

$$\begin{aligned}
 g'(z_j) &= \frac{-\mathbb{E}(Ze^{(\kappa+z_j)Z})[1 - e^{\kappa+z_j}] + [1 - \mathbb{E}(e^{(\kappa+z_j)Z})]e^{\kappa+z_j}}{(1 - e^{\kappa+z_j})^2} \\
 &= -\frac{\mathbb{E}(Ze^{(\kappa+z_j)Z})}{1 - e^{\kappa+z_j}} + \frac{e^{\kappa+z_j}}{1 - e^{\kappa+z_j}}
 \end{aligned}$$

so that it is easy to verify that $\operatorname{Res}[\frac{1}{1-g(z)} \frac{m-g(z)e^{\kappa+z}}{1-e^{\kappa+z}} e^{-xz}; z_j]$ is (40) times $e^{-(\kappa+z_j)x}$. \square

5.3. *Proof of Proposition 16.* We recall that two functions f and h satisfy $f(x) \gg h(x)$ iff $h(x)/f(x)$ tends to 0 as $x \rightarrow \infty$. In the following, we consider cases where $q > 0$ is such that functions respectively satisfy

$$(100) \quad C_0^1 e^{-\kappa_1 x} \gg \Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] \gg \{C_0^2 e^{-q\kappa_2 x}, \Re[C_1^2 e^{-q(\kappa_2+z_1^2)x}]\},$$

$$(101) \quad C_0^1 e^{-\kappa_1 x} \gg C_0^2 e^{-q\kappa_2 x} \gg \{\Re[C_1^1 e^{-(\kappa_1+z_1^1)x}], \Re[C_1^2 e^{-q(\kappa_2+z_1^2)x}]\},$$

all other cases being treated similarly. We will use the following inequalities (an easy and direct consequence of Lemma 2 (i) of [4]):

$$(102) \quad \psi_1(x_1) \leq \psi_{\text{or}}(x_1, x_2) \leq \psi_1(x_1) + \psi_2(x_2).$$

Let us first consider the case (100). Using expansions (42) with $x_1 = x$, $x_2 = qx$, as well as (102) we get

$$\begin{aligned}
 \varepsilon_1(x)e^{-(r+\kappa_1)x} &\leq \psi_{\text{or}}(x, qx) - C_0^1 e^{-\kappa_1 x} - \Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] \\
 (103) \qquad \qquad \qquad &\leq C_0^2 e^{-q\kappa_2 x} + \Re[C_1^2 e^{-q(\kappa_2+z_1^2)x}] + \varepsilon_1(x)e^{-(r+\kappa_1)x} \\
 &\qquad \qquad \qquad + \varepsilon_2(qx)e^{-(r+\kappa_1)qx}.
 \end{aligned}$$

We now note that the terms on the far left- and right-hand side of (103) divided by $e^{-\Re(\kappa_1+z_1^1)x}$ tend to 0 because of Assumption (100), which proves (43). We now consider case (101). Using again (102) we get, similarly to (103),

$$\begin{aligned}
 \varepsilon_1(x)e^{-(r+\kappa_1)x} + \Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] - C_0^2 e^{-q\kappa_2 x} \\
 (104) \qquad \qquad \qquad &\leq \psi_{\text{or}}(x, qx) - C_0^1 e^{-\kappa_1 x} - C_0^2 e^{-q\kappa_2 x} \\
 &\leq \Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] + \Re[C_1^2 e^{-q(\kappa_2+z_1^2)x}] \\
 &\qquad \qquad \qquad + \varepsilon_1(x)e^{-(r+\kappa_1)x} + \varepsilon_2(qx)e^{-(r+\kappa_1)qx}.
 \end{aligned}$$

Because of Assumption (101), we get from (104) that

$$\begin{aligned}
 -1 &= \lim_{x \rightarrow \infty} \frac{\varepsilon_1(x)e^{-(r+\kappa_1)x} + \Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] - C_0^2 e^{-q\kappa_2 x}}{C_0^2 e^{-q\kappa_2 x}} \\
 &\leq \liminf_{x \rightarrow \infty} \frac{\psi_{\text{or}}(x, qx) - C_0^1 e^{-\kappa_1 x} - C_0^2 e^{-q\kappa_2 x}}{C_0^2 e^{-q\kappa_2 x}} \\
 &\leq \limsup_{x \rightarrow \infty} \frac{\psi_{\text{or}}(x, qx) - C_0^1 e^{-\kappa_1 x} - C_0^2 e^{-q\kappa_2 x}}{C_0^2 e^{-q\kappa_2 x}} \\
 &\leq \lim_{x \rightarrow \infty} (\Re[C_1^1 e^{-(\kappa_1+z_1^1)x}] + \Re[C_1^2 e^{-q(\kappa_2+z_1^2)x}] + \varepsilon_1(x)e^{-(r+\kappa_1)x} \\
 &\qquad \qquad \qquad + \varepsilon_2(qx)e^{-(r+\kappa_1)qx}) / (C_0^2 e^{-q\kappa_2 x}) \\
 &= 0,
 \end{aligned}$$

which proves (43).

Acknowledgment. The authors wish to thank an anonymous referee for pointing out the informal proof given at the end of Section 2.1, comments leading to Remark 5, as well as reference [15] leading to the infinite expansion for $\psi(x)$ in Example 11.

REFERENCES

[1] ASMUSSEN, S. (2003). *Applied Probability and Queues: Stochastic Modelling and Applied Probability*, 2nd ed. Springer, New York. MR1978607

- [2] ASMUSSEN, S. and ALBRECHER, H. (2010). *Ruin Probabilities*, 2nd ed. World Scientific, Hackensack, NJ. [MR2766220](#)
- [3] ASMUSSEN, S. and BLADT, M. (1997). Renewal theory and queueing algorithms for matrix-exponential distributions. In *Matrix-Analytic Methods in Stochastic Models (Flint, MI). Lecture Notes in Pure and Applied Mathematics* **183** 313–341. Dekker, New York. [MR1427279](#)
- [4] AVRAM, F., PALMOWSKI, Z. and PISTORIUS, M. R. (2008). Exit problem of a two-dimensional risk process from the quadrant: Exact and asymptotic results. *Ann. Appl. Probab.* **18** 2421–2449. [MR2474542](#)
- [5] BIARD, R. (2013). Asymptotic multivariate finite-time ruin probabilities with heavy-tailed claim amounts: Impact of dependence and optimal reserve allocation. *Bull. Fr. Actuarial* **13** 79–92.
- [6] BLANCHET, J. and GLYNN, P. (2007). Uniform renewal theory with applications to expansions of random geometric sums. *Adv. in Appl. Probab.* **39** 1070–1097. [MR2381589](#)
- [7] BREIMAN, L. (1968). *Probability*. Addison-Wesley, Reading, MA. [MR0229267](#)
- [8] DÖRING, L. and SAVOV, M. (2011). (Non)differentiability and asymptotics for potential densities of subordinators. *Electron. J. Probab.* **16** 470–503. [MR2781843](#)
- [9] FELLER, W. (1965). *An Introduction to Probability Theory and Its Applications*. Wiley, New York.
- [10] FELLER, W. and OREY, S. (1961). A renewal theorem. *J. Math. Fluid Mech.* **10** 619–624.
- [11] GERBER, H. U. (1988). Mathematical fun with compound binomial process. *Astin Bull.* **18** 161–168.
- [12] HU, Z. and JIANG, B. (2013). On joint ruin probabilities of a two-dimensional risk model with constant interest rate. *J. Appl. Probab.* **50** 309–322.
- [13] KUZNETSOV, A., KYPRIANOU, A. E. and PARDO, J. C. (2012). Meromorphic Lévy processes and their fluctuation identities. *Ann. Appl. Probab.* **22** 1101–1135.
- [14] KUZNETSOV, A. and MORALES, M. (2014). Computing the finite-time expected discounted penalty function for a family of Lévy risk processes. *Scand. Actuar. J.* **1** 1–31. [MR3176015](#)
- [15] KUZNETSOV, A. and PENG, X. (2012). On the Wiener–Hopf factorization for Lévy processes with bounded positive jumps. *Stochastic Process. Appl.* **122** 2610–2638.
- [16] KYPRIANOU, A. E. (2006). *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer, Berlin. [MR2250061](#)
- [17] LI, S., LU, Y. and GARRIDO, J. (2009). A review of discrete-time risk models. *Journal Serie A Matemáticas (RACSAM)* **103** 321–337.
- [18] MITOV, K. V. and OMEY, E. (2014). Intuitive approximations for the renewal function. *Statist. Probab. Lett.* **84** 72–80.
- [19] RABEHASAINA, L. (2012). A Markov additive risk process in dimension 2 perturbed by a fractional Brownian motion. *Stochastic Process. Appl.* **122** 2925–2960. [MR2931347](#)
- [20] ROYNETTE, B., VALLOIS, P. and VOLPI, A. (2008). Asymptotic behavior of the hitting time, overshoot and undershoot for some Lévy processes. *ESAIM Probab. Stat.* **12** 58–93. [MR2367994](#)
- [21] SCHIFF, J. L. (1993). *Normal Families*. Springer, New York. [MR1211641](#)
- [22] STONE, C. (1965). On moment generating functions and renewal theory. *Ann. Math. Stat.* **36** 1298–1301.
- [23] STONE, C. (1965). On characteristic functions and renewal theory. *Trans. Amer. Math. Soc.* **120** 327–342.
- [24] STONE, C. (1966). On absolutely continuous components and renewal theory. *Ann. Math. Stat.* **37** 271–275. [MR0196795](#)

- [25] TEUGELS, J. (1968). Renewal theorems when the first or the second moment is infinite. *Ann. Math. Stat.* **39** 1210–1219.

LABORATOIRE DE MATHÉMATIQUES
UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ
CNRS UMR 6623
UFR SCIENCES ET TECHNIQUES
16 ROUTE DE GRAY
25030 BESANÇON CEDEX
FRANCE
E-MAIL: Clement.Dombry@univ-fcomte.fr
E-MAIL: Landy.Rabehasaina@univ-fcomte.fr