

## When does the minimum of a sample of an exponential family belong to an exponential family?

Shaul K. Bar-Lev\*      Gérard Letac†

### Abstract

It is well known that if  $(X_1, \dots, X_n)$  are i.i.d. r.v.'s taken from either the exponential distribution or the geometric one, then the distribution of  $\min(X_1, \dots, X_n)$  is, with a change of parameter, is also exponential or geometric, respectively. In this note we prove the following result. Let  $F$  be a natural exponential family (NEF) on  $\mathbb{R}$  generated by an arbitrary positive Radon measure  $\mu$  (not necessarily confined to the Lebesgue or counting measures on  $\mathbb{R}$ ). Consider  $n$  i.i.d. r.v.'s  $(X_1, \dots, X_n)$ ,  $n \geq 2$ , taken from  $F$  and let  $Y = \min(X_1, \dots, X_n)$ . We prove that the family  $G$  of distributions induced by  $Y$  constitutes an NEF if and only if, up to an affine transformation,  $F$  is the family of either the exponential distributions or the geometric distributions. The proof of such a result is rather intricate and probabilistic in nature.

**Keywords:** exponential family; exponential distribution; geometric distribution; order statistics; Radon measure.

**AMS MSC 2010:** 62E10.

Submitted to ECP on 2015-08-03, final version accepted on 2015-12-24.

## 1 Introduction

Both distributions, the geometric distribution supported on  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$  and the exponential distribution supported on  $[0, \infty)$ , possess similar properties. We outline only some of them:

- Like its continuous analogue (the exponential distribution), the geometric distribution is memoryless.
- If a r.v.  $X$  has an exponential distribution with mean  $1/\lambda$  then  $\lfloor X \rfloor$ , where  $\lfloor x \rfloor$  denotes the floor function of a real number  $x$ , is geometrically distributed with parameter  $p = 1 - e^{-\lambda}$ .
- If  $(X_1, \dots, X_n)$  are i.i.d. r.v.'s taken from either the exponential distribution or the geometric one, then the distribution of  $\min(X_1, \dots, X_n)$  is, with a change of parameter, also exponential or geometric, respectively.
- Both families of distributions belong to the class of natural exponential families (NEF's).

Indeed, the present note incorporates the last two properties in the following sense. Let  $F$  be an NEF on  $\mathbb{R}$  generated by an arbitrary positive Radon measure  $\mu$  (not necessarily confined to the Lebesgue or counting measures on  $\mathbb{R}$ ). Consider  $n$  i.i.d. r.v.'s

---

\*University of Haifa, Israel. E-mail: barlev@stat.haifa.ac.il

†Université Paul Sabatier, France. E-mail: gerard.letac@math.univ-toulouse.fr

$(X_1, \dots, X_n)$ ,  $n \geq 2$ , taken from  $F$  and let  $Y = \min(X_1, \dots, X_n)$ . Then we prove that the family  $G$  of distributions induced by  $Y$  constitutes an NEF if and only if, up to an affine transformation,  $F$  is the family of either the exponential distributions or the geometric distributions.

A similar, but rather more restrictive, problem has been treated by Bar-Lev and Bshouty (2008) in which they considered the case where  $\mu$  has the form  $\mu(dx) = h(x)dx$ . Then under some restrictive conditions on  $h$  (as differentiability) they showed that the family of distributions induced by  $Y$  is an NEF if and only if the distribution of the  $X_i$ 's is an exponential one (up to an affinity  $x \mapsto ax + b$ ). In their concluding remarks, Bar-Lev and Bshouty (2008) indicated the mathematical difficulties arising for proving that when  $\mu$  is a counting measure on  $\mathbb{N}_0$  then the family  $G$  is an NEF iff  $F$  is the family geometric distributions. It should be noted, however, that for the restricted case  $\mu(dx) = h(x)dx$ , Bar-Lev and Bshouty (2008) treated the question of when  $G_r$ , the family of distributions induced by the  $r$ -th order statistic  $X_{(r)}$  (out  $(X_1, \dots, X_n)$ ), is an NEF. They showed that necessarily  $r = 1$  in which case the NEF  $F$  must be that of the exponential distributions.

As already indicated, we consider here the case  $r = 1$  and prove in Theorem 1 a more general result for an arbitrary measure  $\mu$  (which includes the Lebesgue measure and counting measure as special cases).

In Section 2 we introduce some required preliminaries on NEF's. In Section 3 we present and prove our main result. The style of the result and the methods of the proof are close to the celebrated Balkema-de Haan-Pickands theorem on extreme values (see [1] and [5]).

## 2 Some preliminaries on NEF's

For proving our main result we shall need the definition of an NEF (for a detailed description of NEF's on  $\mathbb{R}$  see Letac and Mora, 1990).

Let  $\mu$  be a positive non-Dirac Radon measure on  $\mathbb{R}$ . The Laplace transform of  $\mu$  is

$$L_\mu(\theta) = \int_{-\infty}^{\infty} e^{\theta x} \mu(dx) \leq \infty.$$

Let

$$D(\mu) = \{\theta \in \mathbb{R} : L_\mu(\theta) < \infty\}, \quad \Theta(\mu) = \text{int } D(\mu)$$

and denote  $k_\mu(\theta) = \log L_\mu(\theta)$ ,  $\theta \in \Theta(\mu)$ . Also, let  $\mathcal{M}(\mathbb{R})$  denote the set of positive measures  $\mu$  on  $\mathbb{R}$  not concentrated on one point such that  $\Theta(\mu) \neq \emptyset$ . Then, the family of probabilities

$$F = F(\mu) = \{P(\theta, \mu) : \theta \in \Theta(\mu)\}$$

where

$$P(\theta, \mu)(dx) = e^{\theta x - k_\mu(\theta)} \mu(dx)$$

is called the NEF generated by  $\mu$ .

The two special cases of the geometric and exponential families have the following NEF features:

- Geometric:

$$\mu(dx) = \sum_{k=0}^{\infty} \delta_k(dx), \quad L_\mu(\theta) = (1 - e^\theta)^{-1}, \quad k_\mu(\theta) = -\ln(1 - e^\theta), \quad \Theta(\mu) = (-\infty, 0),$$

where  $\delta_k$  is the Dirac mass on  $k$ . In this case

$$P(\theta, \mu)(dx) = \sum_{x \in \mathbb{N}_0} (1 - q)q^x \delta_x$$

where  $q = e^\theta < 1$ . Let  $X_1, \dots, X_n$  be i.i.d. r.v.'s with common geometric distribution with parameter  $q$ , then the p.d.f. of  $Y = \min(X_1, \dots, X_n)$  is geometric with parameter  $q^n$ , or in its NEF p.d.f. form with  $\theta \mapsto n\theta$ .

- Exponential:

$$\mu(dx) = \mathbf{1}_{(0,\infty)}(dx), L_\mu(\theta) = (-\theta)^{-1}, k_\mu(\theta) = -\ln(-\theta), \Theta(\mu) = (-\infty, 0),$$

where its known p.d.f. form is

$$\lambda e^{-\lambda x} \mathbf{1}_{(0,\infty)}, \lambda > 0,$$

in which case

$$\theta = -\lambda.$$

If  $X_1, \dots, X_n$  be i.i.d. r.v.'s with common exponential distribution with parameter  $\lambda$  then the p.d.f. of  $Y = \min(X_1, \dots, X_n)$  is again exponential with parameter  $n\lambda$ , or in its NEF p.d.f. form with  $\theta \mapsto n\theta$ .

### 3 The main result

**Theorem 3.1.** *Let  $\mu \in \mathcal{M}(\mathbb{R})$  and  $n \geq 2$  be an integer. Let  $X_1, \dots, X_n$  be i.i.d. r.v.'s with common distribution  $P(\theta, \mu)$  and denote by  $Q_\theta$  the distribution of  $Y = \min(X_1, \dots, X_n)$ . Then there exist a measure  $\nu \in \mathcal{M}(\mathbb{R})$ , an NEF  $F(\nu)$  and a differentiable mapping  $\theta \mapsto \alpha(\theta)$  from  $\Theta(\mu)$  to  $\Theta(\nu)$  such that  $Q_\theta = P(\alpha(\theta), \nu)$  for all  $\theta \in \Theta(\mu)$  if and only if  $F(\mu)$  is a positive affine transformation of either the NEF of geometric distributions or the NEF of exponential distributions.*

*Proof.* The statement  $\Leftarrow$  is simple as can be seen from the remarks at the end of Section 2. Indeed, with the choices of  $\mu$  made there, we have for both, the geometric and exponential cases, that  $\mu = \nu$  and  $\alpha(\theta) = n\theta$ .

We prove the statement  $\Rightarrow$  in six steps. In the first step we derive the functional equation (3.3) which provides a necessary condition for  $Q_\theta \sim Y = \min(X_1, \dots, X_n)$  to belong to some NEF  $F(\nu)$ . The second step proves that the support of  $\mu$  is bounded on the left, while the third step shows that such a support is unbounded on the right. The fourth step further analyzes the functional equation (3.3) and provides an important equation (3.7) associated with the measure  $\mu$ . More specifically the problem is then being reduced to the case where the support interval (i.e., the convex hull of the support) of  $\mu$  is exactly  $[0, \infty)$ . If we denote by  $\mu_x$  the translate of  $\mu(dt)$  by  $t \mapsto t - x$  and then truncate at zero, the equality (3.7) is  $k'_{\mu_x} = k'_\mu$  for  $\mu$  almost all  $x$ . This equality reduces the characterization problem to the problem of whether  $\mu$  possesses at least one atom or not. If  $\mu$  has at least one atom the fifth step proves that  $\mu$  generates the geometric NEF. Otherwise, the sixth step shows that  $\mu$  generates the exponential NEF. Such six steps then conclude the proof.

**First step.** This step is devoted to the setting of the functional equation (3.3) below. For simplicity, we write  $k = k_\mu, \Theta = \Theta(\mu)$  and so on. In the sequel we write

$$\int_{a^-}^{b^+} f(t)\mu(dt) \text{ for } \int_{[a,b]} f(t)\mu(dt) \text{ and } \int_{a^+}^{b^+} f(t)\mu(dt) \text{ for } \int_{(a,b]} f(t)\mu(dt).$$

If the law of  $Y$  belongs to an NEF  $F(\nu)$  then for  $\theta \in \Theta$  and real  $y$ , the number  $P(Y \geq y)$  can be represented in two different ways, by which one gets the following equality

$$e^{-nk(\theta)} \left( \int_{y^-}^{\infty} e^{\theta t} \mu(dt) \right)^n = e^{-k\nu(\alpha(\theta))} \int_{y^-}^{\infty} e^{\alpha(\theta)t} \nu(dt),$$

and hence the following equality, between two probability measures, holds:

$$n e^{-nk(\theta)} \left( \int_{y^-}^{\infty} e^{\theta t} \mu(dt) \right)^{n-1} e^{\theta y} \mu(dy) = e^{-k_{\nu}(\alpha(\theta))} e^{\alpha(\theta)y} \nu(dy).$$

This proves that the measures  $\nu$  and  $\mu$  are equivalent and we can introduce the Radon Nikodym derivative  $g(y) = \frac{d\nu}{d\mu}(y)$ . Hence, the following equality which holds  $\mu$  almost everywhere:

$$n e^{-nk(\theta) + k_{\nu}(\alpha(\theta))} \left( \int_{y^-}^{\infty} e^{\theta t} \mu(dt) \right)^{n-1} e^{(\theta - \alpha(\theta))y} = g(y).$$

By denoting  $g_n(y) = \left( \frac{g(y)}{n} \right)^{1/(n-1)}$  and

$$A(\theta) = \frac{-\theta + \alpha(\theta)}{n-1}, \quad B(\theta) = \frac{nk(\theta) - k_{\nu}(\alpha(\theta))}{n-1}, \tag{3.1}$$

and elevating to the power  $1/(n-1)$ , we get the following equality which holds  $\mu$  almost everywhere:

$$e^{-yA(\theta) - B(\theta)} \int_{y^-}^{\infty} e^{\theta t} \mu(dt) = g_n(y). \tag{3.2}$$

Assume, without loss of generality, that  $\mu$  and  $\nu$  are probability measures. Then, the Hölder inequality, applied to the pair of functions  $(g, 1)$  and to  $(p, q) = (n-1, (n-1)/(n-2))$ , shows that  $\int_{-\infty}^{\infty} g_n(y) \mu(dy) < \infty$ . Integrating (3.2) on  $[x, \infty)$  with respect to  $\mu(dy)$  yields for all  $\theta \in \Theta$

$$\int_{x^-}^{\infty} \left( e^{-yA(\theta) - B(\theta)} \int_{y^-}^{\infty} e^{\theta t} \mu(dt) \right) \mu(dy) = \int_{x^-}^{\infty} g_n(y) \mu(dy).$$

Now, by differentiating, with respect to  $\theta$ , of both sides of the latter equality, we obtain

$$\int_{x^-}^{\infty} \left( e^{-yA(\theta) - B(\theta)} \int_{y^-}^{\infty} e^{\theta t} (t - yA'(\theta) - B'(\theta)) \mu(dt) \right) \mu(dy) = 0.$$

Since the latter equality holds for all  $x$ , it follows that for each fixed  $\theta \in \Theta$ ,

$$\int_{x^-}^{\infty} e^{\theta t} (t - xA'(\theta) - B'(\theta)) \mu(dt) = 0, \tag{3.3}$$

which holds  $\mu(dx)$  almost everywhere. The equality (3.3) holds in particular for any element  $x$  of the support  $S$  of the measure  $\mu$ . To prove this statement, we denote by  $H(x)$  the left hand side of (3.3). Then locally,  $H$  has a bounded variation (i.e., it is the difference of two non-increasing functions) and its discontinuity points are the atoms of  $\mu$ . Therefore  $H(x) = 0$  if  $x$  is an atom of  $\mu$ . If  $x \in S$  and is not an atom of  $\mu$  then there exists a sequence  $(x_k)$  such that  $H(x_k) = 0$  for all  $k$  and such that  $x_k \rightarrow x$ . Since  $H$  is continuous in  $x$  it follows that  $H(x) = 0$  for all  $x \in S$ .

**Second step.** We prove that the support of  $\mu$  is bounded on the left. If not, the equality (3.3) holds for some fixed  $\theta \in \Theta$  and for some sequence  $(x_k)$  such that  $\lim_{k \rightarrow \infty} x_k = -\infty$ . This implies that  $A'(\theta) = 0$  and  $B'(\theta) = k'(\theta)$ . But then clearly the equality

$$\int_{x_k^-}^{\infty} e^{\theta t} (t - k'(\theta)) \mu(dt) = 0$$

cannot hold for all  $k$ . Indeed, if  $k_0$  is such that  $x_{k_0} \leq k'(\theta)$  then such an equality would imply that for any  $k > k_0$

$$0 = \int_{x_k^-}^{x_{k_0}^-} e^{\theta t} (t - k'(\theta)) \mu(dt),$$

while the right hand side is negative for  $k$  large enough.

**Third step.** This step proves that the support of  $\mu$  is unbounded on the right. It relies on the following lemma, which has its own interest with its characterisation of the distribution  $B(1, a)$  up to a dilation by  $b$  :

**Lemma 1.** Let  $P$  be a non-Dirac probability on  $[0, \infty)$  and  $K > 0$  such that for  $P$  almost all  $x$  we have

$$\int_{0-}^{x+} tP(dt) = Kx \int_{0-}^{x+} P(dt). \tag{3.4}$$

Then  $K < 1$  and there exists  $b > 0$  such that  $P(dt) = \frac{a}{b^a} t^{a-1} 1_{(0,b)}(t)dt$ , where  $a = K/(1 - K)$ .

**Proof.** If  $K > 1$  then for at least one  $x > 0$  we have

$$\int_{0-}^{x+} P(dt) = \frac{1}{K} \int_{0-}^{x+} \frac{t}{x} P(dt) < \int_{0-}^{x+} P(dt),$$

which is a contradiction. If  $K = 1$  then  $0 = \int_{0-}^{x+} (t - x)P(dt)$  for  $P$  almost all  $x$ . This implies that  $t - x = 0$  for  $P(dt)P(dx)$  almost all  $(t, x)$ , which is possible only if  $P$  is a Dirac measure, a contradiction. The probability measure  $P$  has no atom on  $t_0 > 0$  since (3.4) implies  $t_0P(\{t_0\}) = Kt_0P(\{t_0\})$  which contradicts that  $K < 1$ . Similarly,  $P$  has no atom on zero. If not, since for at least one  $x > 0$  one has

$$\int_{0+}^x P(dt) \geq \int_{0+}^x \frac{t}{x} P(dt) = \int_{0-}^x \frac{t}{x} P(dt) = KP(\{0\}) + \int_{0+}^x P(dt) > \int_{0+}^x P(dt),$$

we get a contradiction.

The support  $S$  of  $P$  contains 0. If not, there exists  $b$  in  $S$  such that  $P([0, b)) = 0$ . Since  $P$  is not Dirac there exists a sequence  $x_n \searrow b$  such that

$$\int_b^{x_n} \frac{t}{x_n} P(dt) = A \int_b^{x_n} P(dt).$$

Now consider the conditional probability  $P_n$  which is  $P(dt)$  conditioned on  $b < t < x_n$ . Then,  $P_n$  converges weakly to  $\delta_b$  (the simplest way to prove this is to use the distribution function of  $P_n$ ). Since  $\int_b^{x_n} tP_n(dt) = Kx_n$ , then by passing to the limit we get the contradiction for  $K = 1$ .

The support  $S$  of  $P$  is an interval containing zero. If not, and since  $0 \in S$ , there exist  $0 < x_1 < x_2$  such that  $P((x_1, x_2)) = 0$ ,  $x_1, x_2 \in S$  and  $\int_0^{x_1} P(dt) > 0$ . Hence, from (3.4), we get the following contradiction

$$Kx_1 \int_0^{x_1} P(dt) \stackrel{(a)}{=} \int_0^{x_1} tP(dt) \stackrel{(b)}{=} \int_0^{x_2} tP(dt) \stackrel{(c)}{=} Kx_2 \int_0^{x_2} P(dt) \stackrel{(d)}{=} Kx_2 \int_0^{x_1} P(dt),$$

where the equalities (a) and (c) stem from (3.4) and the fact that  $x_1$  and  $x_2$  are in  $S$ . The equalities (b) and (d) come from the fact that  $P((x_1, x_2)) = 0$ .

Now, since  $P$  has no atoms, the function

$$f(x) = \int_0^x tP(dt) - Kx \int_0^x P(dt)$$

is continuous. Furthermore,  $f$  is zero  $P$  almost everywhere. This implies that  $f$  is zero on the support  $S$  of  $P$ . If not, there exists  $x_0 \in S$  such that  $|f(x_0)| > 0$  and an open interval  $(x_0 - h, x_0 + h)$  such that  $|f(x)| > 0$  if  $|x - x_0| < h$ . However,

$$\int_{x_0-h}^{x_0+h} |f(t)|\mu(dt) = 0$$

and thus  $(x_0 - h, x_0 + h)$  and  $S$  are disjoint (recall that  $S$  is the complementary set of the largest open set with  $P$  zero measure). Hence, a contradiction follows.

Accordingly,  $S = [0, b]$  for some real  $b$  or  $S = [0, \infty)$ . Denote by  $S_0$  the interior of  $S$ . We have seen that for all  $x \in S$ ,  $f(x) = 0$ . We rewrite this fact as

$$\int_0^x tP(dt) = Kx \int_0^x P(dt).$$

Differentiating this equality (in the Stieltjes sense) we get (on  $S_0$ ) that

$$xP(dx) = a \left( \int_0^x P(dt) \right) dx,$$

where  $a = \frac{K}{1-K}$ . This shows that  $P(dx) = g(x)dx$  is absolutely continuous. In fact, from  $xg(x) = a \int_0^x g(t)dt$ , it follows that the function  $g$  is continuous and even differentiable on  $S_0$ . This leads to the differential equation  $g'(x)/g(x) = (a - 1)/x$  on  $S_0$  and  $g(x) = Cx^{a-1}$ , where  $C > 0$ . If  $S$  is unbounded then  $g$  cannot be a probability density. Therefore  $S = [0, b]$  is bounded and the lemma is proved. ■

We now prove the claim of Step 3 that the support of  $\mu$  is unbounded on the right. If not, from Step 2, we may assume without loss of generality that the support interval of  $\mu$  is exactly  $[0, b]$  with  $b > 0$ . Substituting  $x = 0$  in (3.3) gives  $B'(\theta) = k'(\theta)$ . We now show that  $A'(\theta) = 1 - \frac{1}{b}k'(\theta)$ . To see this we rewrite (3.3) as follows

$$\frac{\int_{x-}^{b+} e^{\theta t} t \mu(dt)}{\int_{x-}^{b+} e^{\theta t} \mu(dt)} = xA'(\theta) + k'(\theta) \tag{3.5}$$

and we do  $x \nearrow b$  in (3.5). The left hand side converges to  $b$  and  $A'(\theta) = 1 - \frac{1}{b}k'(\theta)$  is proved. This now leads to the equation

$$\frac{\int_{x-}^{b+} e^{\theta t} (t - x) \mu(dt)}{\int_{x-}^{b+} e^{\theta t} \mu(dt)} = \left(1 - \frac{x}{b}\right) k'(\theta). \tag{3.6}$$

Fix  $\theta$ , consider the change of variable  $t \mapsto b - t$  and apply Lemma 1 to the image  $P(dt)$  of the probability  $e^{\theta t - k(\theta)} \mu(dt)$  and to  $A = k'(\theta)/b$ . Then, it follows that the  $a$  of Lemma 1 is  $a(\theta) = k'(\theta)/(b - k'(\theta))$ . Since the support interval of  $P$  is also  $[0, b]$  we can claim that

$$e^{\theta t - k(\theta)} \mu(dt) = a(\theta)(b - t)^{a(\theta)-1} 1_{(0,b)}(t) dt,$$

an equality which cannot hold for all  $\theta$ . One may realize this as follows. Since

$$\mu(dt) = e^{-t\theta + (a(\theta)-1) \log(b-t) + c(\theta)} 1_{(0,b)}(t) dt,$$

where  $c(\theta) = k(\theta) + \log a(\theta)$ , we have, by differentiating by  $\theta$ , that for all  $(\theta, t) \in \Theta \times (0, b)$ ,

$$-t + a'(\theta) \log(b - t) + c'(\theta) = 0.$$

Then, differentiating by  $t$ , we get  $b - t = a'(\theta)$ , which is clearly impossible.

**Fourth step.** From Steps 2 and 3, we may assume throughout the sequel that the support interval of  $\mu$  is exactly  $[0, \infty)$ . This assumption implies that we are allowed to substitute  $x = 0$  in (3.3) to obtain

$$\int_{0-}^{\infty} e^{\theta t} (t - B'(\theta)) \mu(dt) = 0,$$

which shows that  $B' = k'$ . By the definition of  $B$  in (3.1), this implies that  $k(\theta) - k_\nu(\alpha(\theta))$  is a constant. We denote by  $\mu_x(du)$  the image of the measure  $\mu$  by the map  $t \mapsto u = t - x$  multiplied by the function  $\mathbf{1}_{[0, \infty)}(u)$ . The equality (3.3) can then be reformulated as

$$k'_{\mu_x}(\theta) = k'(\theta), \tag{3.7}$$

for  $\mu(dx)$  almost everywhere. We now analyze (3.7) according to whether  $\mu$  has at least one atom (Fifth Step), an assumption that will lead to the geometric NEF, or not (Sixth Step), a fact that will lead to the exponential NEF.

**Fifth step.** Assume that  $\mu$  has an atom  $x_0$ . We prove that there exists a countable additive subgroup  $G$  of  $\mathbb{R}$  and a real character  $\chi$  of  $G$  such that

$$\mu(dt) = \mu(0) \sum_{x \in G \cap [0, \infty)} e^{\chi(x)} \delta_x(dt),$$

where  $\mu(x)$  denotes the mass of the atom  $x$ .

This assumption implies that (3.7) is true for  $x = x_0$  and thus that  $\mu$  has an atom on  $\mathbb{R}$  (and thus are all the measures  $\mu_x$  for which (3.7) is true). This implies that  $\mu$  is purely atomic. Denote by  $S$  the set of atoms of  $\mu$ . From (3.7) we infer that for all  $x \in S$  we have

$$S = (S - x) \cap [0, \infty).$$

Denote  $G = S \cup (-S)$ . Then  $G$  is an additive group with  $S = G \cap [0, \infty)$ . Write  $\mu(dt) = \sum_{x \in S} \mu(x) \delta_x(dt)$ , then (3.7) implies that for all  $x \in S$  we have

$$\mu_x(dt) = \frac{\mu(x)}{\mu(0)} \mu(dt).$$

Calculating the mass of this measure on  $s \in S$  we get

$$\mu(s) = \frac{\mu(0)}{\mu(x)} \mu_x(s) = \frac{\mu(0)}{\mu(x)} \mu(x + s).$$

For  $x \in S$ , denote  $\chi(x) = \log \mu(x) - \log \mu(0)$  and for  $x \in -S$  denote  $\chi(x) = -\chi(-x)$ . Then the latter equality implies

$$\chi(x + s) = \chi(x) + \chi(s),$$

that is,  $\chi$  is a real character of  $G$ .

We now prove that  $G$  is  $a\mathbb{Z}$  for some for some  $a > 0$ . If not, then  $G$  is a dense in  $\mathbb{R}$ . Then, either any pair  $(x, x')$  of  $G \setminus \{0\}$  is such that  $x/x'$  is rational, or there exists a pair such that  $x/x'$  is irrational. Without loss of generality, we may assume for the latter two cases that  $1 \in G$ . In the first case (where  $x/x'$  is rational) there exist arbitrary small rational numbers  $x \in G$  such that  $\chi(x) = x\chi(1)$ . Thus, for  $A > 0$ , the family  $\{e^{\chi(x)} : x \in G \cap [0, A]\}$  cannot be summable and  $\mu$  is not a Radon measure. Similarly, for the second case ( $x/x'$  is irrational),  $G$  contains a subgroup  $\mathbb{Z}(\alpha)$  for some irrational number  $\alpha$  (where  $\mathbb{Z}(\alpha)$  is the set of  $a + b\alpha$  with  $a, b$  in  $\mathbb{Z}$ ). By denoting  $p_1 = e^{\chi(1)}$  and  $p_2 = e^{\chi(\alpha)}$  we obtain that  $p_1^a p_2^b = e^{\chi(a+b\alpha)}$ . We now need to prove that

$$\sum \{p_1^a p_2^b : 0 \leq a + b\alpha \leq A\} = \infty. \tag{3.8}$$

This can be accomplished by a tedious discussion and analysis of the nine cases  $0 < p_1 < 1, p_1 = 1$  and  $p_1 > 1$  combined with  $0 < p_2 < 1, p_2 = 1$  and  $p_2 > 1$  (we omit details for brevity). This, however, would finally show that  $\mu$  cannot be a Radon measure.

Thus we conclude the case where  $\mu$  has at least one atom by stating that for this case there exist  $a > 0$  and numbers  $p = e^{\chi(a)} > 0$  and  $q = \mu(0)$  such that

$$\mu(dt) = \mu(0) \sum_{n=0}^{\infty} qp^n \delta_{na}(dt).$$

This is equivalent to saying that  $F(\mu)$  is the image of the geometric distributions by the dilation  $n \mapsto an$ .

**Sixth step.** We assume that  $\mu$  has no atoms. Denote by  $X \subset [0, \infty)$  the set of  $x$  such that (3.7) holds. We prove that the closure  $\bar{X}$  of  $X$  is the support  $S$  of  $\mu$ . To see that  $S \subset \bar{X}$ , we choose  $x_0 \in S$ . If there is no sequence  $(x_n)$  of  $X$  converging to  $x_0$ , this would imply the existence of  $\epsilon > 0$  such that  $\mu([x_0 - \epsilon, x_0 + \epsilon]) = 0$  and thus contradict the fact that  $x_0 \in S$ . To see that  $X \subset S$  we choose  $x_0 \in X$ . If  $x_0 \notin S$  then this would imply the existence of  $\epsilon > 0$  such that  $\mu([x_0 - \epsilon, x_0 + \epsilon]) = 0$ . Since  $0 \in S$ , the measure  $\mu_{x_0}$  cannot be equivalent to  $\mu$ . Thus, the statement that  $S = \bar{X}$  is proved.

Now, the fact that  $\mu$  has no atoms implies that  $x \mapsto \mu_x$  is a continuous function on  $\mathbb{R}$  for the vague topology of Radon measures. The equality (3.7) is thus equivalent to the existence of a function  $\chi$  on  $X$  such that

$$\mu_x(dt) = e^{\chi(x)} \mu(dt), \quad (3.9)$$

and the preceding remark implies that  $\chi$  is a continuous function on  $X$  and is extendable in a continuous function to  $\bar{X}$ . Thus (3.7) and (3.8) hold on  $S$ . Now we observe that (3.7) implies that for all  $x \in S$  we have  $S = (S - x) \cap [0, \infty)$ . Thus  $G = S \cup (-S)$  is an additive subgroup of  $\mathbb{R}$ . Since  $G$  is closed, then either  $G = \{0\}$ , or there exists  $a > 0$  such that  $G = a\mathbb{Z}$  or  $G = \mathbb{R}$ . Such two cases can be excluded since  $\mu$  has no atoms, and thus we get  $S = [0, \infty)$ .

We now show that  $\chi(x+s) = \chi(x) + \chi(s)$  for all  $x \geq 0$  and  $s \geq 0$ . For this we observe that (3.7) implies that for all  $x \geq 0$  the measure  $\mu_x$  generates the NEF  $F(\mu)$ . Thus  $\mu_x$  must share with  $\mu$  the property (3.7), and for  $s \geq 0$  we therefore have

$$\mu_{x+s}(dt) = e^{\chi(x)} \mu(dt).$$

Since  $\mu_x$  and  $\mu$  are proportional, the factor  $e^{\chi(x)}$  is the same. Since we also have  $\mu_{x+s}(dt) = e^{\chi(x+s)} \mu_x(dt)$ , the equality  $\chi(x+s) = \chi(x) + \chi(s)$  follows.

As  $\chi$  is continuous, it is simple to see that there exists  $b \in \mathbb{R}$  such that  $\chi(x) = bx$ . One can consult Bingham, Teugels and Goldie for reference to this Cauchy functional equation. By introducing the measure  $\tilde{\mu}(dt) = e^{-bt} \mu(dt)$ , we have  $F(\tilde{\mu}) = F(\mu)$ . Furthermore (3.9) implies that for all  $x \geq 0$  we have

$$\tilde{\mu}_x(dt) = \tilde{\mu}(dt).$$

This implies that for all intervals  $I \subset [0, \infty)$ , we have  $\tilde{\mu}(x+I) = \tilde{\mu}(I)$ . Thus  $\tilde{\mu}$  is proportional to the restriction of the Lebesgue measure to  $[0, \infty)$  and the theorem is proved.  $\square$

**Acknowledgement.** We thank the referee for a very careful reading of the original version of the paper.

## References

- [1] Balkema, A.A. and de Haan, L. (1974). Residual life time at great age. *Ann. Probab.* **2**(5), 792–804. MR-0359049
- [2] Bar-Lev, S.K. and Bshouty, D. (2008). Exponential families are not preserved by the formation of order statistics. *Statist. Probab. Lett.* **78**, 2787–2792. MR-2465123
- [3] Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1989) *Regular Variation*. Cambridge University Press. MR-1015093
- [4] Letac, G. and Mora, M. (1990). Natural real exponential families with cubic variance functions. *Ann. Statist.* **18**, 1–37. MR-1041384
- [5] Pickands, J. (1975). Statistical inference using extreme order statistics. *Ann. Statist.* **3**, 119–131. MR-0423667