

An inequality for the heat kernel on an Abelian Cayley graph

Thomas McMurray Price*

Abstract

We demonstrate a relationship between the heat kernel on a finite weighted Abelian Cayley graph and Gaussian functions on lattices. This can be used to prove a new inequality for the heat kernel on such a graph: when $t \leq t'$,

$$\frac{H_t(u, v)}{H_t(u, u)} \leq \frac{H_{t'}(u, v)}{H_{t'}(u, u)}.$$

This was an open problem posed by Regev and Shinkar.

Keywords: heat kernel; cayley graph.

AMS MSC 2010: 60J27.

Submitted to ECP on March 7, 2017, final version accepted on August 29, 2017.

1 Introduction

Let $\Gamma = (V, E)$ be a finite undirected weighted graph. Let L be the graph laplacian of Γ , given by $L = D - A$, with D the degree matrix and A the adjacency matrix. We then define the **heat kernel** H_t as the matrix exponential e^{-tL} , for any $t \in \mathbb{R}_{>0}$. The value $H_t(u, v)$ can be thought of as the probability that a continuous-time random walk starting at u ends up at v after time t . See Section 1.1 of [RS16] for more detail on this interpretation. We will say that G has **monotonic diffusion** if the following holds whenever $t' \geq t$, for all vertices v and u :

$$\frac{H_t(u, v)}{H_t(u, u)} \leq \frac{H_{t'}(u, v)}{H_{t'}(u, u)}.$$

We will say that Γ is a **weighted Abelian Cayley graph** if there is an Abelian group structure on V under which Γ is translation-invariant. In this note we will prove the following theorem:

Theorem 1.1. *Any weighted Abelian Cayley graph has monotonic diffusion.*

For arbitrary weighted undirected graphs, monotonic diffusion often fails to hold. See the appendix of [RS16] for a simple case. It has previously been asked whether vertex-transitive graphs necessarily have monotonic diffusion; this question was attributed to Peres in [RS16], and was resolved in the negative in the same paper. It also makes sense to ask whether a Riemannian manifold has monotonic diffusion, since there is a uniquely determined heat kernel in this setting, provided the manifold is complete and the Ricci curvature is bounded from below [C94]. We present an argument in Appendix A, due to

*Toronto, Canada. E-mail: tom.price.math@gmail.com

Jeff Cheeger, that monotonicity does not always hold for Riemannian manifolds. In the other direction, it was shown in [RS15] that all flat tori have monotonic diffusion. Since weighted Abelian Cayley graphs are the closest thing we have to flat tori in the world of graph theory, it is natural to ask whether they have monotonic diffusion. This problem was posed in [RS16] and is resolved, in the affirmative, in this note.

We will use the following strategy: we show that, if v is a nonnegative, even, real-valued function on a finite Abelian group, then the convolutional exponential $\exp^*(v)$ can be represented as a pushforward of a Gaussian function on a lattice. This will allow us to apply an inequality in [RS15] regarding Gaussian sums on lattice cosets.

2 Gaussian pushforwards

Throughout this paper, G will refer to an arbitrary finite Abelian group.

Definition 2.1. A **lattice** is a discrete subgroup of a finite-dimensional inner product space. If L is a lattice, we define the function $\rho : L \rightarrow \mathbb{R}$ as $\rho(x) = e^{-\pi\langle x, x \rangle}$. When $S \subseteq L$, we will use the notation $\rho(S)$ to mean $\sum_{x \in S} \rho(x)$.

Definition 2.2. We will say that a function $\chi : G \rightarrow \mathbb{R}$ is a **Gaussian pushforward** if there exists a lattice L and a group homomorphism $h : L \rightarrow G$ such that $\chi(g) = \rho(h^{-1}(g))$ for all $g \in G$. We will use X to denote the set of all Gaussian pushforwards on G .

Proposition 2.3. X is closed under convolution.

Proof. Suppose that $\chi_1 \in X$ and $\chi_2 \in X$. Then we have lattices L_1 and L_2 , as well as homomorphisms $h_1 : L_1 \rightarrow G$ and $h_2 : L_2 \rightarrow G$, such that $\chi_1(g) = \rho(h_1^{-1}(g))$ and $\chi_2(g) = \rho(h_2^{-1}(g))$ for all $g \in G$. Take L_3 to be the orthogonal direct sum of L_1 and L_2 . We have the homomorphism $h_3 : L_3 \rightarrow G$ given by $h_3((x_1, x_2)) = h_1(x_1) + h_2(x_2)$. Define $\chi_3 : G \rightarrow \mathbb{R}$ as $\chi_3(g) = \rho(h_3^{-1}(g))$. Then $\chi_3 = \chi_1 * \chi_2$. We also clearly have that $\chi_3 \in X$. This yields the result. \square

Remark 2.4. Curiously, we also have that X is closed under multiplication, if we instead take L_3 to be the sublattice of $L_1 \oplus L_2$ of points (x_1, x_2) with $h_1(x_1) = h_2(x_2)$, and $h_3((x_1, x_2))$ to be $h_1(x_1)$.

Proposition 2.5. If $\chi \in X$, then for all $g_1, g_2 \in G$, we have

$$\frac{\chi(g_1)\chi(g_2)}{\chi(0)} \leq \frac{\chi(g_1 + g_2) + \chi(g_1 - g_2)}{2}.$$

Proof. This follows immediately from (4c) of corollary 2.2 of [RS15], if we take the lattice \mathcal{L} to be the kernel of a homomorphism from a lattice to G whose corresponding Gaussian pushforward function is χ . \square

From continuity, we immediately have the following stronger statement:

Corollary 2.6. If χ is in the topological closure \bar{X} of X then for all $g_1, g_2 \in G$, we have

$$\frac{\chi(g_1)\chi(g_2)}{\chi(0)} \leq \frac{\chi(g_1 + g_2) + \chi(g_1 - g_2)}{2}.$$

Proposition 2.7. If χ is in \bar{X} , v is a nonnegative, even, real-valued function on G , and $\omega = \chi * v$, then for all $g \in G$,

$$\frac{\chi(g)}{\chi(0)} \leq \frac{\omega(g)}{\omega(0)}.$$

Proof. For all $g \in G$, we have

$$\omega(g) = \sum_{g' \in G} \chi(g - g')v(g')$$

$$= \sum_{g' \in G} \chi(g + g')v(-g')$$

$$\sum_{g' \in G} \chi(g + g')v(g').$$

Taking the mean of the first and third sums above, we have:

$$\omega(g) = \sum_{g' \in G} \frac{\chi(g + g') + \chi(g - g')}{2} v(g').$$

Applying 2.6,

$$\omega(g) \geq \sum_{g' \in G} \frac{\chi(g)\chi(g')v(g')}{\chi(0)}$$

$$= \frac{\chi(g)}{\chi(0)} \sum_{g' \in G} \chi(g')v(-g')$$

$$= \frac{\chi(g)\omega(0)}{\chi(0)}.$$

Dividing by $\omega(0)$ yields the result. □

3 The convolutional exponential of nonnegative even functions

Definition 3.1. Suppose v is a real-valued function on G . We define $\exp^*(v)$, the **convolutional exponential** of v , by

$$\exp^*(v) = \sum_{n=0}^{\infty} \frac{v^{*n}}{n!},$$

where v^{*n} is the n th convolutional power of v . In general, we have the following relationship between convolution, multiplication, and Fourier transform for real-valued functions on finite groups:

$$\widehat{x * y} = \hat{x}\hat{y}.$$

We therefore have the following equivalent definition of $\exp^*(v)$:

$$\widehat{\exp^*(v)} = \exp(\hat{v}),$$

where \exp is the pointwise exponential operator. From this perspective, it is clear that $\exp^*(a + b) = \exp^*(a) * \exp^*(b)$.

In this section, we will show that the convolutional exponential of any nonnegative even function on G is a Gaussian pushforward.

Definition 3.2. We will use Φ to refer to the set of functions on G of the form $\phi(g) = \delta(g - g_0) + \delta(g + g_0)$ for some $g_0 \in G$. Here we have $\delta(0) = 1$ and $\delta(g) = 0$ for $g \neq 0$. We clearly have that Φ is a basis for the even functions on G .

Definition 3.3. We will say that a sequence of real-valued functions v_n on G is $O(f(n))$ if, for any $g \in G$, we have that $v_n(g)$ is $O(f(n))$. Since all norms on a finite-dimensional vector space are equivalent, this is the same as saying that $\|v_n\|$ is $O(f(n))$, for any choice of norm on \mathbb{R}^G . We also have that v_n is $O(f(n))$ if and only if \hat{v}_n is $O(f(n))$, by the Plancherel theorem.

The next lemma is elementary and so the proof will be skipped.

Lemma 3.4. If a and b are both in $[0, C]$, then for all $n \in \mathbb{N}$, we have $|a^n - b^n| \leq nC^{n-1}|a - b|$.

Lemma 3.5. *If a_n and b_n are both $1 + O(1/n)$, and $a_n - b_n$ is $O(f(n))$, then $a_n^n - b_n^n$ is $O(nf(n))$.*

Proof. We have, for some C , that $a_n \leq 1 + C/n$ and $b_n \leq 1 + C/n$. By applying 3.4, we have

$$|a_n^n - b_n^n| \leq n(1 + C/n)^{n-1}|a_n - b_n| \leq ne^C|a_n - b_n| \in O(nf(n)). \quad \square$$

Lemma 3.6. *Suppose $\phi \in \Phi$ and $\alpha \in \mathbb{R}_{>0}$. Then we can find a sequence of Gaussian pushforwards (χ_n) such that $\delta + \alpha\phi/n - \chi_n$ is $O(1/n^4)$.*

Proof. We will choose χ_n arbitrarily when $n \leq \alpha$, and the rest of this proof will be concerned with the tail $(\chi_n)_{n>\alpha}$. Choose a g_0 such that $\phi(g) = \delta(g + g_0) + \delta(g - g_0)$ for all $g \in G$. Let $r_n = \sqrt{\ln(n/\alpha)/\pi}$, so that $\alpha/n = \rho(r_n)$. Let L_n be the lattice in \mathbb{R} of reals of the form kr_n , with $k \in \mathbb{Z}$. We then have a unique group homomorphism h_n from L_n to G that sends r_n to g_0 . We define χ_n by $\chi_n(g) = \rho(h_n^{-1}(g))$. Let $A_n = \{r_n, 0, -r_n\}$ and $B_n = L_n \setminus A_n$. We can break χ_n into two smaller sums:

$$\chi_n(g) = \rho(h_n^{-1}(g) \cap A_n) + \rho(h_n^{-1}(g) \cap B_n).$$

The left term is equal to $\delta + \alpha\phi/n$. The right term is $O(1/n^4)$. This yields the result. \square

Lemma 3.7. *If $\phi \in \Phi$ and $\alpha \in \mathbb{R}_{>0}$, then $\exp^*(\alpha\phi)$ is in the topological closure \bar{X} of X .*

Proof. Let $\psi = \exp^*(\alpha\phi)$. From the convolution theorem, we have that $\hat{\psi} = \exp(\alpha\hat{\phi})$. Therefore, we have

$$\hat{\psi} = \lim_{n \rightarrow \infty} (1 + \frac{\alpha\hat{\phi}}{n})^n.$$

From 3.6, we have a sequence of Gaussian pushforwards (χ_n) such that $\delta + \alpha\phi/n - \chi_n$ is $O(1/n^4)$. Taking the Fourier transform, we have that $1 + \alpha\hat{\phi}/n - \hat{\chi}_n$ is $O(1/n^4)$. We then have from 3.5 that $(1 + \alpha\hat{\phi}/n)^n - \hat{\chi}_n^n$ is $O(1/n^3)$. Combining this with the formula for $\hat{\psi}$ above, we have:

$$\lim_{n \rightarrow \infty} \hat{\chi}_n^n = \hat{\psi}.$$

Applying the convolution theorem then yields:

$$\lim_{n \rightarrow \infty} \chi_n^{*n} = \psi.$$

From 2.3, we know that χ_n^{*n} is a Gaussian pushforward, so this last limit proves the result. \square

Theorem 3.8. *Suppose v is a nonnegative even function on G . Then $\exp^* v$ is in \bar{X} .*

Proof. We can clearly represent v as a sum $\alpha_1\phi_1 + \dots + \alpha_n\phi_n$, with α_i being positive scalars and $\phi_i \in \Phi$. Then, applying the convolution theorem, we can represent $\exp^*(v)$ as the convolution $\exp^*(\alpha_1\phi_1) * \dots * \exp^*(\alpha_n\phi_n)$. From 3.7, we know that each individual term in this convolution is in the closure of X , so the result follows from 2.3 and the continuity of the convolution operation. \square

4 The heat kernel on weighted Abelian Cayley graphs

In this section, we prove 1.1. The hard work has already been done in [RS15] and in the previous section; now we just need to translate these results into the language of graph theory.

Proof of Theorem 1.1. We define τ to be the element of \mathbb{R}^V obtained by applying the graph laplacian to $-\delta$. In other words, we have $\tau : V \rightarrow \mathbb{R}$ and $\tau(v) = -L(0, v)$, with L the graph laplacian, given by $L = D - A$, with D the degree matrix and A the adjacency matrix. Then τ is an even function and is nonnegative everywhere except at 0. We also have that the linear operator determined by the graph laplacian is just convolution with $-\tau$; it follows that the heat kernel matrix H_t corresponds to convolution with $\exp^*(t\tau)$, and therefore $H_t(u, v) = \exp^*(t\tau)(u - v)$. So, to prove 1.1, it suffices to show that, for $t \leq t'$, and for any $v \in V$,

$$\frac{\exp^*(t\tau)(v)}{\exp^*(t\tau)(0)} \leq \frac{\exp^*(t'\tau)(v)}{\exp^*(t'\tau)(0)}.$$

Let $\tau' : V \rightarrow \mathbb{R}$ be given by $\tau'(0) = 0$ and $\tau'(v) = \tau(v)$ for $v \neq 0$. Then τ' is a nonnegative even function. We also have that $\exp^*(t\tau') = e^{-\tau(0)} \exp^*(t\tau)$. It therefore suffices to show that, for $t \leq t'$, and for any $v \in V$,

$$\frac{\exp^*(t\tau')(v)}{\exp^*(t\tau')(0)} \leq \frac{\exp^*(t'\tau')(v)}{\exp^*(t'\tau')(0)}.$$

Since we have $\exp^*(t'\tau') = \exp^*(t\tau') * \exp^*((t' - t)\tau')$, the result follows immediately from 3.8 and 2.7. \square

5 Directions for further work

In order to prove 1.1, we needed the inequality 2.6 for χ a Gaussian pushforward:

$$\frac{\chi(g_1)\chi(g_2)}{\chi(0)} \leq \frac{\chi(g_1 + g_2) + \chi(g_1 - g_2)}{2},$$

which itself follows from the stronger inequality

$$\chi(g_1)^2\chi(g_2)^2 \leq \chi(g_1 + g_2)\chi(g_1 - g_2)\chi(0)^2, \tag{5.1}$$

which is a special case of Theorem 2.1 of [RS15].

The proof of monotonic diffusion on flat tori also uses (5.1), except $\chi(g)$ is replaced by $H_t(0, g)$, with H the heat kernel on the torus in question.

So, if we want to see where else we have monotonic diffusion, it is natural to ask: where else does (5.1) hold? On the surface, it looks we need group structure to even state (5.1). However, (5.1) can be reformulated in the language of Riemannian symmetric spaces. Indeed, suppose M is a Riemannian symmetric space, that is, M is a connected Riemannian manifold equipped with an isometric automorphism s_p for each point p , such that $s_p(p) = p$ and the derivative of s_p at p is the negation map on T_p . For flat tori then, we have a symmetric space structure with $s_x(y) = 2x - y$, and so the following is equivalent to (5.1):

$$H_t(a, b)^2 H_t(b, c)^2 \leq H_t(a, c) H_t(a, s_b(c)) H_t(a, a)^2 \tag{5.2}$$

To see this, substitute 0 for a , x for b , and $x + y$ for c . Then $s_b(c)$ is $x - y$, and we recover the original inequality.

It is shown Appendix B that any symmetric space which satisfies (5.2) has monotonic diffusion. This inequality does not hold for arbitrary symmetric spaces; it is shown

in Appendix C that it fails on \mathbb{H}^3 . However, we already know from [RS15] that (5.2) holds for flat tori, and the author has numerically tested (5.2) for the 2-sphere and real projective 2-space, using the spherical harmonic functions of SciPy to approximate the heat kernel. The inequality appears to hold for these spaces. It seems natural then to ask: for which Riemannian symmetric spaces does (5.2) hold? Perhaps it holds on all Riemannian symmetric spaces of compact type.

It can also be readily seen from equation (1.6) of [GN98] that the heat kernel has monotonic diffusion on \mathbb{H}^3 , despite the fact that (5.2) fails on this space. Perhaps monotonic diffusion holds on other hyperbolic spaces as well.

A Appendix: lack of monotonic diffusion on arbitrary manifolds

We present here an argument that not all Riemannian manifolds have monotonic diffusion. This argument is due to Jeff Cheeger and was relayed to the author by Oded Regev [R17]. It is very similar to the argument given in the appendix of [RS16], also due to Cheeger, which deals with diffusion on graphs rather than manifolds.

It is shown in [CV87] that, on a compact manifold M of dimension at least 3, any finite sequence $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ of positive real numbers appears as the first nonzero eigenvalues, counted with multiplicity, of the Laplacian on M for some choice of metric. In particular, there exist compact Riemannian manifolds on which the first nonzero eigenvalue of the Laplacian has multiplicity 1. Let M be such a manifold, with λ_1 the first eigenvalue and λ_2 the second. Let $f : M \rightarrow \mathbb{R}$ be an eigenfunction of the Laplacian corresponding to the eigenvalue λ_1 . Choose points x and y in M so that $f(y) > f(x) > 0$. We have the following, for some positive constant C :

$$H_t(x, x) = 1 + Ce^{-\lambda_1 t} f(x) + O(e^{-\lambda_2 t})$$

$$H_t(x, y) = 1 + Ce^{-\lambda_1 t} f(y) + O(e^{-\lambda_2 t}).$$

Since $\lambda_2 > \lambda_1$, we have that $H_t(x, y) > H_t(x, x)$ for sufficiently large t . Since the quotient $H_t(x, y)/H_t(x, x)$ is eventually greater than 1, but it converges to 1 as $t \rightarrow \infty$, it can not be monotonically increasing. Therefore, we do not have monotonic diffusion on M .

B Appendix: monotonic diffusion on Riemannian symmetric spaces

Here we show that, if the heat kernel of a Riemannian symmetric space M satisfies the inequality

$$H_t(a, b)^2 H_t(b, c)^2 \leq H_t(a, c) H_t(a, s_b(c)) H_t(a, a)^2 \tag{2}$$

then it has monotonic diffusion. First of all, note that (5.2) implies the following analogue of 2.5:

$$\frac{H_t(a, b) H_t(b, c)}{H_t(a, a)} \leq \frac{H_t(a, c) + H_t(a, s_b(c))}{2} \tag{B.1}$$

This follows from dividing both sides of (5.2) by $H_t(a, a)^2$, and then applying the inequality of arithmetic and geometric means. This is essentially the same as the proof of 2.5.

From this, we can derive the monotonic diffusion inequality in a way that parallels the proof of 2.7. We have, for $t' \geq t$:

$$\begin{aligned} H_{t'}(x, y) &= \int_M H_{t'-t}(x, z) H_t(z, y) dz \\ &= \int_M H_{t'-t}(x, s_x(z)) H_t(s_x(z), y) dz \end{aligned}$$

$$= \int_M H_{t'-t}(x, z) H_t(s_x(z), y) dz.$$

Averaging the first and third lines above, we have:

$$\begin{aligned} H_{t'}(x, y) &= \int_M H_{t'-t}(x, z) \frac{H_t(z, y) + H_t(s_x(z), y)}{2} dz \\ &= \int_M H_{t'-t}(x, z) \frac{H_t(y, z) + H_t(y, s_x(z))}{2} dz. \end{aligned}$$

Applying (B.1) then gives

$$\begin{aligned} H_{t'}(x, y) &\geq \int_M H_{t'-t}(x, z) \frac{H_t(y, x) H_t(x, z)}{H_t(y, y)} dz \\ &= \int_M H_{t'-t}(x, z) H_t(z, x) dz \frac{H_t(x, y)}{H_t(y, y)} \\ &= \frac{H_{t'}(x, x) H_t(x, y)}{H_t(y, y)}. \end{aligned}$$

Dividing by $H_{t'}(x, x)$ gives the result.

C Appendix: failure of (5.2) on \mathbb{H}^3

Here we present an argument, due to Oded Regev [R17], that the inequality (5.2) fails on hyperbolic 3-space. In this section we realize \mathbb{H}^3 through the hyperboloid model, as described in [CFKP97]. In other words, we see \mathbb{H}^3 as the set of points (x_0, x_1, x_2, x_3) in \mathbb{R}^4 with $x_0^2 - x_1^2 - x_2^2 - x_3^2 = 1$. The geodesic distance between two points $x = (x_0, x_1, x_2, x_3)$ and $y = (y_0, y_1, y_2, y_3)$ in \mathbb{H}^3 is then given by

$$d(x, y) = \operatorname{arccosh}(x_0 y_0 - x_1 y_1 - x_2 y_2 - x_3 y_3).$$

Suppose $d_1 \in \mathbb{R}_{>0}$. We define the following points in \mathbb{H}^3 :

$$\begin{aligned} a &= (\cosh(d_1), \sinh(d_1), 0, 0) \\ b &= (1, 0, 0, 0) \\ c &= (\cosh(d_1), 0, \sinh(d_1), 0). \end{aligned}$$

Then we have $d_1 = d(a, b) = d(b, c)$, and we define d_2 as $d(a, c) = \operatorname{arccosh}(\cosh(d_1)^2)$. We also fix some arbitrary $t \in \mathbb{R}_{>0}$. We have a formula for the heat kernel on \mathbb{H}^3 , given in [GN98]. When the geodesic distance between x and y is given by d , we have:

$$H_t(x, y) = \frac{1}{(4\pi t)^{3/2}} \frac{d}{\sinh(d)} \exp\left(-t - \frac{d^2}{4t}\right).$$

Using this formula, the inequality (5.2) reduces to the following, for our choices of a, b, c , and t :

$$\frac{d_1^2}{\sinh(d_1)^2} \exp\left(-\frac{d_1^2}{2t}\right) \leq \frac{d_2}{\sinh(d_2)} \exp\left(-\frac{d_2^2}{4t}\right).$$

Let LS and RS be the left and right sides of the above inequality. We have, as d_1 and d_2 grow:

$$\begin{aligned} \ln(LS) &= \frac{-d_1^2}{2t} + O(d_1) \\ \ln(RS) &= \frac{-d_2^2}{4t} + O(d_2). \end{aligned}$$

However, it can be readily seen that $d_2 = 2d_1 + O(1)$. We therefore have

$$\ln(RS) = \frac{-d_1^2}{t} + O(d_1),$$

and therefore $RS < LS$ for sufficiently large d_1 . This violates (5.2).

References

- [CV87] Y. Colin de Verdière. Construction de laplaciens dont une partie finie du spectre est donnée. *Annales Scientifiques de l'École Normale Supérieure*, 20:599–615, 1987. MR-0932800
- [C94] I. Chavel. *Eigenvalues in Riemannian Geometry*. Pure and applied mathematics. Academic Press, Orlando, 1994. MR-0768584
- [CFKP97] James W. Cannon, William J. Floyd, Richard Kenyon, and Walter R. Parry. Hyperbolic Geometry. *Flavors of geometry*, pp. 59–115, Cambridge University Press, 1997. MR-1491098
- [GN98] A. Grigor'yan and M. Noguchi. The heat kernel on hyperbolic space. *Bull. London Math. Soc.* 30 (1998), No. 6. MR-1642767
- [P14] T. Price. Is the heat kernel more spread out with a smaller metric? *MathOverflow*. <http://mathoverflow.net/questions/186428/>, 2014.
- [R17] O. Regev, 2016. Private Communication.
- [RS15] O. Regev and N. Stephens-Davidowitz, An Inequality for Gaussians on Lattices, [arXiv:1502.04796v2](https://arxiv.org/abs/1502.04796v2) [math.PR]. MR-3632259
- [RS16] O. Regev and I. Shinkar. A counterexample to monotonicity of relative mass in random walks. *Electron. Commun. Probab.* 21 (2016), paper no. 8, 8 pp. doi:10.1214/16-ECP4392. <http://projecteuclid.org/euclid.ecp/1454682824>.

Acknowledgments. The train of thought that lead to the main idea of this paper began with a discussion with Alexander Borisov. He had noticed that strongly positive-definite functions obey the main inequality of [RS15]. It was through trying to relate this to the fact that Gaussian pushforwards obey the same inequality that I came across the construction used in section 3. So I thank him for the stimulating discussion, as well as for helpful comments on a draft. I thank Oded Regev as well for helpful comments and discussion.

This paper was written while I was visiting the University of Münster. I thank Christopher Deninger for making this possible.