

OPTIMAL INVESTMENT AND CONSUMPTION WITH LABOR INCOME IN INCOMPLETE MARKETS

BY OLEKSII MOSTOVYI¹ AND MIHAI SÎRBU²

¹*Department of Mathematics, University of Connecticut, oleksii.mostovyi@uconn.edu*

²*Department of Mathematics, University of Texas at Austin, sirbu@math.utexas.edu*

We consider the problem of optimal consumption from labor income and investment in a general incomplete semimartingale market. The economic agent cannot borrow against future income, so the total wealth is required to be positive at (all or some) previous times. Under very general conditions, we show that an optimal consumption and investment plan exists and is unique, and provide a dual characterization in terms of an optional strong supermartingale deflator and a decreasing part, which charges only the times when the no-borrowing constraint is binding. The analysis relies on the infinite-dimensional parametrization of the income/liability streams and, therefore, provides the first-order dependence of the optimal investment and consumption plans on future income/liabilities (as well as a pricing rule).

1. Introduction. Optimal investment with intermediate consumption and a stream of labor income (or liabilities) is one of the central problems in mathematical economics. If borrowing against the future income is prohibited, the main technical difficulty lies in the fact that there are infinitely many constraints. Even in the deterministic case of no stocks and nonrandom income, a classical approach is based on the *convexification of the constraints* that leads to a nontrivial dual problem formulated over decreasing nonnegative functions.

Borrowing constraints imposed at all times not only affect the notion of admissibility, leading to more difficult mathematical analysis, but also change the meaning to fundamental concepts of mathematical finance such as replicability and completeness. The latter is formulated via the attainability of every (bounded) contingent claim by a portfolio of traded assets. For a labor income/liability streams that pays off dynamically, there is no a priori guarantee that such a replicating portfolio (if it exists at all) is admissible, that is, satisfies the constraints. Thus, in the terminology of [18], even a complete market becomes dynamically incomplete under the borrowing constraints. The analysis of such a problem (in otherwise complete Brownian settings with a corresponding unique risk-neutral measure), is performed in [18] and later in [16]. The nonnegative decreasing *processes* (that parametrize the dynamic incompleteness mentioned above) play an important role in the characterizations of optimal investment and consumption plans. The analysis in [18] and [16] is connected with optimal stopping techniques from [22].

Incomplete markets with no-borrowing constraints have been analyzed only in specific Markovian models in [14] and [13] based on partial differential equations techniques. The goal of the present paper is to study the problem of consumption and investment with no-borrowing constraints in general (so, non-Markovian) incomplete models. This leads to having, simultaneously, two layers of incompleteness. One comes from the many martingale measures, the other from a similar class of nondecreasing processes (as above) that describe the dynamic incompleteness. We refer to [28] for the examples of market incompleteness in finance and macroeconomics.

Received June 2018; revised January 2019.

MSC2010 subject classifications. Primary 91G10, 93E20; secondary 91B16.

Key words and phrases. Utility maximization, labor income, semimartingale incomplete market, duality theory, optional supermartingale deflator.

In contrast to [18] and [16], our model not only allows for incompleteness, but also for jumps. Mathematically, this means we choose to work in a general semimartingale framework. As in [18] and [16], our approach is based on duality. One of the principal difficulties is in the construction of the dual feasible set and the dual value function. It is well known that the martingale measures drive the dual domain in many problems of mathematical finance. On the other hand, the convexification of constraints leads to the decreasing processes as the central dual object as well. We show that the dual elements in the incomplete case can be approximated by products of the densities of martingale measures and such nonnegative decreasing processes. This is one of the primary results of this work (see Section 5) that leads to the complementary slackness characterization of optimal wealth in Section 6, where it is shown that the approximating sequence for the dual minimizer leads to a nonincreasing process, which decreases at most when the constraints are attained. In turn, the dual minimizer can be written as a product of such a nondecreasing process and an optional strong supermartingale deflator. In the case of complete Brownian markets, a similar result is proved in [18] and [16].

In order to implement the approach, we *increase the dimensionality* of the problem and treat as arguments of the indirect utility not only the initial wealth, but also the function that specifies the number of units of labor income (or the stream of liabilities) at any later time. This parametrization has the spirit of [19], however, unlike [19], we go into (infinite-dimensional) nonreflexive spaces, which gives both novelty and technical difficulties to our analysis. Also, our formulation permits to price by marginal rate of substitution the whole labor income process. This is done through the subdifferentiability results in Section 4. Note that the subdifferential elements (prices) are time-dependent, so infinite-dimensional, unlike in [19].

Another contribution of the paper lies in the *unified framework of admissibility* outlined in Section 2. More precisely, we assume that no-borrowing constraints are imposed starting from some prespecified stopping time and hold up to the terminal time horizon. This framework allows us to treat in one formulation both the problem of no-borrowing constraints at all times (described above) and the one where borrowing against the future income is permitted with a constraint only at the end. The latter is well studied in the literature; see [7, 19, 24, 37] and [32]. In such a formulation, the constraints reduce to a single inequality and the decreasing processes in the dual feasible set become constants.

Among the many possibilities of constraints, [6] considers the problem of investment and consumption with labor income and no-borrowing constraints in Brownian market, even allowing for incompleteness. The dual problem cannot be solved directly (in part because for these constraints the dual space considered is too small) but the primal can be solved with direct methods. An approximate dual sequence can then be recovered from the primal. We generalize [18] and [16] (complete Brownian markets) and [6] (possibly incomplete Brownian markets) to the case of general semi-martingale incomplete markets. Our dual approach allows us, at the same time to obtain a dual characterization (complementary slackness) of the optimal consumption plan (not present in [6]), similar to the complete case in [18] and [16] and the possibility to study the dependence on labor income streams, through the parametrization of such streams.

Embedding path dependent problems into the convex duality framework have been analyzed in [29, 36], [30], Section 3.3, whereas without duality but with random endowment it is considered in [34], in the abstract singular control setting the duality approach is investigated in [1]. Our embedding does not require any condition on labor income replicability, which becomes highly technical in the presence of extra admissibility constraints. Even in the case where the only constraint is imposed at maturity, in this part our approach differs from the one in [19], where nonreplicability of the endowment (in the appropriate sense) is used in the

proofs as it ensures that the effective domain of the dual problem has the same dimensionality as the primal domain. Note that even if the labor income is spanned by the same sources of randomness as the stocks, the idea of replicating the labor income and then reducing the problem to the one without it, does not necessarily work under the borrowing constraints; see the discussion in [18], pages 671–673.

Some of the more specific technical contributions of this paper can be summarized as follows:

- We analyze the *boundary behavior* of the value functions. Note that the value functions are defined over infinite-dimensional spaces.
- The finiteness of the indirect utilities *without* labor income is imposed only, as a necessary and sufficient condition that allows for the standard conclusions of the utility maximization theory; see [31].
- We show existence-uniqueness results for the *unbounded* labor income both from above and below.
- We observe that the “Snell envelope proposition” ([27], Proposition 4.3) can be extended to the envelope over all stopping times that exceed a given initial stopping time θ_0 .
- We represent the dual value function in terms of *uniformly integrable* densities of martingale measures, that is, the densities of martingale measures under which the maximal wealth process of a self-financing portfolio that superreplicates the labor income, is a uniformly integrable martingale; see Lemma 3.11 below.

Organization of the paper. In Section 2, we specify the model. We state and prove existence, uniqueness, semicontinuity and biconjugacy results in Section 3, subdifferentiability is proven in Section 4. Structure of the dual domain is analyzed in Section 5 and complimentary slackness is established in Section 6.

2. Model. We consider a financial market model with finite time horizon $[0, T]$ and a zero interest rate. The price process $S = (S^i)_{i=1}^d$ of the stocks is assumed to be a semimartingale on a complete stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$, where \mathcal{F}_0 is trivial.

Let $(e)_{t \in [0, T]}$ be an optional process that specifies the labor income rate, which is assumed to follow a certain stochastic clock, that we specify below. Both processes S and e are given exogenously.

We define a *stochastic clock* as a nondecreasing, càdlàg, adapted process such that

$$(2.1) \quad \kappa_0 = 0, \quad \mathbb{P}[\kappa_T > 0] > 0 \quad \text{and} \quad \kappa_T \leq A$$

for some finite constant A . We note that the stochastic clock allows to include multiple standard formulations of the utility maximization problem in one formulation; see, for example, [31], Examples 2.5–2.9. Let us define

$$(2.2) \quad K_t := \mathbb{E}[\kappa_t], \quad t \in [0, T].$$

REMARK 2.1. The function K defined in (2.2) is right continuous with left limits and takes values in $[0, A]$.

We assume the income and consumption are given in terms of the clock κ . Define a portfolio Π as a quadruple (x, q, H, c) , where the constant x is the initial value of the portfolio, the function $q : [0, T] \rightarrow \mathbb{R}$ is a bounded and Borel measurable function, which specifies the amount of labor income rate, $H = (H_i)_{i=1}^d$ is a predictable S -integrable process that corresponds to the amount of each stock in the portfolio, and $c = (c_t)_{t \in [0, T]}$ is the consumption rate, which we assume to be optional and nonnegative.

The *wealth process* $V = (V_t)_{t \in [0, T]}$ generated by the portfolio is

$$V_t = x + \int_0^t H_s dS_s + \int_0^t (q_s e_s - c_s) d\kappa_s, \quad t \in [0, T].$$

A portfolio Π with $c \equiv 0$ and $q \equiv 0$ is called *self-financing*. The collection of nonnegative wealth processes generated by self-financing portfolios with initial value $x \geq 0$ is denoted by $\mathcal{X}(x)$, that is,

$$\mathcal{X}(x) := \left\{ X \geq 0 : X_t = x + \int_0^t H_s dS_s, t \in [0, T] \right\}, \quad x \geq 0.$$

A probability measure \mathbb{Q} is an *equivalent local martingale measure* if \mathbb{Q} is equivalent to \mathbb{P} and every $X \in \mathcal{X}(1)$ is a local martingale under \mathbb{Q} . We denote the family of equivalent local martingale measures by \mathcal{M} and assume that

$$(2.3) \quad \mathcal{M} \neq \emptyset.$$

This condition is equivalent to the absence of arbitrage opportunities in the market; see [9, 11] as well as [23] for the exact statements and further references.

To rule out doubling strategies in the presence of random endowment, we need to impose additional restrictions. Following [10], we say that a nonnegative process in $\mathcal{X}(x)$ is *maximal* if its terminal value cannot be dominated by that of any other process in $\mathcal{X}(x)$. As in [10], we define an *acceptable* process to be a process of the form $X = X' - X''$, where X' is a nonnegative wealth process generated by a self-financing portfolio and X'' is maximal.

Our unified framework of admissibility is given by a fixed stopping time θ_0 . The no-borrowing constraints will hold starting at this stopping time until the end. Let Θ be the set of stopping times that are greater or equal than θ_0 .

LEMMA 2.2. *Let q^1 and q^2 be bounded, Borel measurable functions on $[0, T]$, such that $q^1 = q^2, dK$ -a.e. Then the cumulative labor income processes $\int_0^\tau q_s^i e_s d\kappa_s, i = 1, 2$, are indistinguishable.*

The proof of this lemma is given in Section 3.1. Following [19], we denote by $\mathcal{X}(x, q)$ the set of acceptable processes with initial values x , that dominate the labor income on Θ :

$$\mathcal{X}(x, q) := \left\{ \text{acceptable } X : X_0 = x \text{ and } X_\tau + \int_0^\tau q_s e_s d\kappa_s \geq 0, \mathbb{P}\text{-a.s. for every } \tau \in \Theta \right\}.$$

Let us set

$$\mathcal{K} := \{(x, q) : \mathcal{X}(x, q) \neq \emptyset\}.$$

Let $\overset{\circ}{\mathcal{K}}$ denote the interior of \mathcal{K} in the $\mathbb{R} \times \mathbb{L}^\infty(dK)$ -norm topology. We characterize \mathcal{K} in Lemma 3.2 below that in particular asserts that under Assumption 2.5, $\overset{\circ}{\mathcal{K}} \neq \emptyset$.

The set of admissible consumptions is defined as

$$\mathcal{A}(x, q) := \left\{ \text{optional } c \geq 0 : \text{there exists } X \in \mathcal{X}(x, q), \text{ such that } \int_0^\tau c_s d\kappa_s \leq X_\tau + \int_0^\tau q_s e_s d\kappa_s, \text{ for every } \tau \in \Theta \right\}, \quad (x, q) \in \mathcal{K}.$$

Note that $c \equiv 0$ belongs to $\mathcal{A}(x, q)$ for every $(x, q) \in \mathcal{K}$.

REMARK 2.3. The no-borrowing constraints can also be written as

$$\mathbb{P}\left(\int_0^t c_s d\kappa_s \leq X_t + \int_0^t q_s e_s d\kappa_s, \text{ for every } \theta_0 \leq t \leq T\right) = 1.$$

We write the constraints in terms of stopping times $\tau \in \Theta$ as we use the stopping times $\tau \in \Theta$ (and the corresponding decreasing processes that jump from one to zero at these times) as the building blocks of our analysis.

REMARK 2.4. It follows from Lemma 2.2, for every $x \in \mathbb{R}$, we have

$$\mathcal{X}(x, q^1) = \mathcal{X}(x, q^2) \quad \text{and} \quad \mathcal{A}(x, q^1) = \mathcal{A}(x, q^2),$$

where some of these sets might be empty

Hereafter, we shall impose the following conditions on the endowment process.

ASSUMPTION 2.5. There exists a maximal wealth process X' such that

$$X'_t \geq |e_t|, \quad \text{for every } t \in [0, T], \mathbb{P}\text{-a.s.}$$

Moreover, Assumption 2.5 and (2.3) imply that all the assertions of Lemma 3.2 hold.

REMARK 2.6. If $\theta_0 = \{T\}$, then Assumption 2.5 is equivalent to the assumptions on endowment in [19] (for the case of one-dimensional random endowment).

The preferences of an economic agent are modeled with a *utility stochastic field* $U = U(t, \omega, x) : [0, T] \times \Omega \times [0, \infty) \rightarrow \mathbb{R} \cup \{-\infty\}$. We assume that U satisfies the conditions below.

ASSUMPTION 2.7. For every $(t, \omega) \in [0, T] \times \Omega$, the function $x \rightarrow U(t, \omega, x)$ is strictly concave, increasing, continuously differentiable on $(0, \infty)$ and satisfies the Inada conditions:

$$\lim_{x \downarrow 0} U'(t, \omega, x) = \infty \quad \text{and} \quad \lim_{x \rightarrow \infty} U'(t, \omega, x) = 0,$$

where U' denotes the partial derivative with respect to the third argument. At $x = 0$ we suppose, by continuity, $U(t, \omega, 0) = \lim_{x \downarrow 0} U(t, \omega, x)$, which may be $-\infty$. For every $x \geq 0$, the stochastic process $U(\cdot, \cdot, x)$ is optional. Below, following the standard convention, we will not write ω in U .

The agent can control investment and consumption. The goal is to maximize expected utility. The value function u is defined as

$$(2.4) \quad u(x, q) := \sup_{c \in \mathcal{A}(x, q)} \mathbb{E}\left[\int_0^T U(t, c_t) d\kappa_t\right], \quad (x, q) \in \mathcal{K}.$$

In (2.4), we use the convention

$$\mathbb{E}\left[\int_0^T U(t, c_t) d\kappa_t\right] := -\infty \quad \text{if} \quad \mathbb{E}\left[\int_0^T U^-(t, c_t) d\kappa_t\right] = \infty.$$

Here and below, W^- and W^+ denote the negative and positive parts of a stochastic field W , respectively.

We employ duality techniques to obtain the standard conclusions of the utility maximization theory. We first define the convex conjugate stochastic field

$$(2.5) \quad V(t, y) := \sup_{x > 0} (U(t, x) - xy), \quad (t, y) \in [0, T] \times [0, \infty),$$

and then observe that $-V$ satisfies Assumption 2.7. In order to construct the feasible set of the dual problem, we define the set \mathcal{L} as the polar cone of $-\mathcal{K}$:

$$(2.6) \quad \mathcal{L} := \left\{ (y, r) \in \mathbb{R} \times \mathbb{L}^1(dK) : xy + \int_0^T q_s r_s dK_s \geq 0 \text{ for every } (x, q) \in \mathcal{K} \right\}.$$

REMARK 2.8. Under the conditions (2.1), (2.3) and Assumption 2.5, the set \mathcal{L} is nonempty. By definition, it is closed in $\mathbb{R} \times \mathbb{L}^1(dK)$ -norm and $\sigma(\mathbb{R} \times \mathbb{L}^1(dK), \mathbb{R} \times \mathbb{L}^\infty(dK))$ topologies. Also, as shown later, the set $\mathcal{L} = (-\mathring{\mathcal{K}})^o$, that is, the polar of $-\mathring{\mathcal{K}}$.

By \mathcal{Z} , we denote the set of càdlàg densities of equivalent local martingale measures:

$$(2.7) \quad \mathcal{Z} := \left\{ \text{càdlàg} \left(\frac{d\mathbb{Q}_t}{d\mathbb{P}_t} \right)_{t \in [0, T]} : \mathbb{Q} \in \mathcal{M} \right\}.$$

Let us denote by $\mathbb{L}^0 = \mathbb{L}^0(d\kappa \times \mathbb{P})$ the linear space of (equivalence classes of) real-valued optional processes on the stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ which we equip with the topology of convergence in measure $(d\kappa \times \mathbb{P})$. For each $y \geq 0$, we define

$$(2.8) \quad \mathcal{Y}(y) := \text{cl} \{ Y : Y \text{ is càdlàg adapted and } 0 \leq Y \leq yZ(d\kappa \times \mathbb{P}) \text{ a.e. for some } Z \in \mathcal{Z} \},$$

where the closure is taken in \mathbb{L}^0 . Now we are ready to set the domain of the dual problem:

$$(2.9) \quad \mathcal{Y}(y, r) := \left\{ Y : Y \in \mathcal{Y}(y) \text{ and } \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq xy + \int_0^T q_s r_s dK_s, \right. \\ \left. \text{for every } (x, q) \in \mathcal{K} \text{ and } c \in \mathcal{A}(x, q) \right\}.$$

Note that the definition (2.9) requires that every element of $\mathcal{Y}(y, r)$ is in $\mathcal{Y}(y)$, $y \geq 0$. Also, for every $(y, r) \in \mathcal{L}$, $\mathcal{Y}(y, r) \neq \emptyset$, since $0 \in \mathcal{Y}(y, r)$.

We can now state the dual optimization problem:

$$(2.10) \quad v(y, r) := \inf_{Y \in \mathcal{Y}(y, r)} \mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right], \quad (y, r) \in \mathcal{L},$$

where we use the convention:

$$\mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right] := \infty \quad \text{if } \mathbb{E} \left[\int_0^T V^+(t, Y_t) d\kappa_t \right] = \infty.$$

Also, we set

$$(2.11) \quad v(y, r) := \infty \quad \text{for } (y, r) \in \mathbb{R} \times \mathbb{L}^1(dK) \setminus \mathcal{L} \quad \text{and} \\ u(x, q) := -\infty \quad \text{for } (x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK) \setminus \mathcal{K}.$$

With this definition, it will be shown below in Theorem 3.1 that $u < \infty$ and $v > -\infty$ everywhere, so u and v are proper functions in the language of convex analysis. Let us recall that in the absence of random endowment, the dual value function is defined as

$$\tilde{w}(y) := \inf_{Y \in \mathcal{Y}(y)} \mathbb{E} \left[\int_0^T V(t, Y_s) d\kappa_s \right], \quad y > 0,$$

whereas the primal value function is given by

$$(2.12) \quad w(x) := u(x, 0), \quad x > 0.$$

3. Existence, uniqueness and biconjugacy.

THEOREM 3.1. *Let (2.1) and (2.3), Assumptions 2.5 and 2.7 hold true and*

$$(3.1) \quad w(x) > -\infty \text{ for every } x > 0 \text{ and } \tilde{w}(y) < \infty \text{ for every } y > 0.$$

Then we have:

(i) *u is finite-valued on \mathring{K} and $u < \infty$ on $\mathbb{R} \times \mathbb{L}^\infty(dK)$. The dual value function v satisfies $v > -\infty$ on $\mathbb{R} \times \mathbb{L}^1(dK)$, and the set $\{v < \infty\}$ is a nonempty convex subset of \mathcal{L} , whose closure in $\mathbb{R} \times \mathbb{L}^1(dK)$ equals to \mathcal{L} .*

(ii) *u is concave, proper and upper semicontinuous with respect to the norm-topology of $\mathbb{R} \times \mathbb{L}^\infty(dK)$ and the weak-star topology $\sigma(\mathbb{R} \times \mathbb{L}^\infty(dK), \mathbb{R} \times \mathbb{L}^1(dK))$. For every $(x, q) \in \{u > -\infty\}$, there exists a unique solution to (2.4). Likewise, v is convex, proper, and lower semicontinuous with respect to the norm-topology of $\mathbb{R} \times \mathbb{L}^1(dK)$ and the the weak topology $\sigma(\mathbb{R} \times \mathbb{L}^1(dK), \mathbb{R} \times \mathbb{L}^\infty(dK))$. For every $(y, r) \in \{v < \infty\}$, there exists a unique solution to (2.10).*

(iii) *The functions u and v satisfy the biconjugacy relations*

$$u(x, q) = \inf_{(y,r) \in \mathcal{L}} \left(v(y, r) + xy + \int_0^T r_s q_s dK_s \right), \quad (x, q) \in \mathcal{K},$$

$$v(y, r) = \sup_{(x,q) \in \mathcal{K}} \left(u(x, q) - xy - \int_0^T r_s q_s dK_s \right), \quad (y, r) \in \mathcal{L}.$$

LEMMA 3.2. *Let (2.3) and Assumption 2.5 hold. Then we have:*

- (i) *for every $x > 0$, $(x, 0)$ belongs to \mathring{K} (in particular, $\mathring{K} \neq \emptyset$),*
- (ii) *for every $q \in \mathbb{L}^\infty(dK)$, there exists $x > 0$ such that $(x, q) \in \mathring{K}$,*
- (iii) $\sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}[\int_0^T |e_s| d\kappa_s] < \infty$,
- (iv) *there exists a nonnegative maximal wealth process X'' , such that*

$$X''_T \geq \int_0^T |e_s| d\kappa_s, \quad \mathbb{P}\text{-a.s.},$$

- (v) *there exists a nonnegative maximal wealth process X'' , such that*

$$X''_t \geq \int_0^t |e_s| d\kappa_s, \quad t \in [0, T], \mathbb{P}\text{-a.s.}$$

PROOF. First, via [21] and (2.1), Proposition I.4.49, we get

$$\begin{aligned} \int_0^T |e_s| d\kappa_s &\leq \int_0^T X'_s d\kappa_s = - \int_0^T \kappa_{s-} dX'_s + \kappa_T X'_T \\ &\leq - \int_0^T \kappa_{s-} dX'_s + AX'_T = AX'_0 + \int_0^T (A - \kappa_{s-}) dX'_s. \end{aligned}$$

Therefore, there exists a self-financing wealth process \bar{X} , such that

$$\int_0^T |e_s| d\kappa_s \leq \bar{X}_T.$$

Consequently, [10], Theorem 2.3, asserts the existence of a nonnegative maximal process X'' , such that

$$\int_0^T |e_s| d\kappa_s \leq X''_T,$$

that is, (iv) holds. Therefore, (iii) is valid as well, by [11], Theorem 5.12.

Relation (iv) and (2.3) imply (v). To prove (i) and (ii), without loss of generality, we will suppose that in (v), $X''_0 > 0$. Let $q \in \mathbb{L}^\infty(dK)$ and $\varepsilon > 0$ be fixed. Let us define

$$x(q) := (\|q\|_{\mathbb{L}^\infty(dK)} + 2\varepsilon)X''_0.$$

We claim that $(x(q), q) \in \mathring{\mathcal{K}}$. Let us consider arbitrary

$$(3.2) \quad |x'| \leq \varepsilon \text{ and } q' \in \mathbb{L}^\infty(dK) : \quad \|q'\|_{\mathbb{L}^\infty(dK)} \leq \varepsilon,$$

and set

$$(3.3) \quad \tilde{X}_t := (\|q\|_{\mathbb{L}^\infty(dK)} + 2\varepsilon + x')X''_t, \quad t \in [0, T].$$

Then, by item (v), we have

$$\begin{aligned} \tilde{X}_t &\geq (\|q\|_{\mathbb{L}^\infty(dK)} + \varepsilon)X''_t \geq (\|q\|_{\mathbb{L}^\infty(dK)} + \varepsilon) \int_0^t |e_s| d\kappa_s \\ &\geq - \int_0^t (q_s + q'_s)|e_s| d\kappa_s. \end{aligned}$$

We deduce that $\tilde{X} \in \mathcal{X}((\|q\|_{\mathbb{L}^\infty(dK)} + 2\varepsilon + x')X''_0, q + q')$. In particular,

$$\mathcal{X}((\|q\|_{\mathbb{L}^\infty(dK)} + 2\varepsilon + x')X''_0, q + q') \neq \emptyset.$$

As x' and q' are arbitrary elements satisfying (3.2), we deduce that $(x(q), q) \in \mathring{\mathcal{K}}$. This proves (ii).

In order to show (i), first we observe that $\mathring{\mathcal{K}}$ is a convex cone. Therefore, it suffices to prove that, for a given $\varepsilon > 0$, we have

$$(3.4) \quad (2\varepsilon X''_0, 0) \in \mathring{\mathcal{K}}.$$

Again, let us consider x' and q' satisfying (3.2) and \tilde{X} satisfying (3.3) for $q \equiv 0$. Then for every $t \in [0, T]$, we have

$$\tilde{X}_t \geq \varepsilon X''_t \geq \varepsilon \int_0^t |e_s| d\kappa_s \geq - \int_0^t q'_s |e_s| d\kappa_s.$$

Thus, $\mathcal{X}((2\varepsilon + x')X''_0, q') \neq \emptyset$. Consequently, as x' and q' are arbitrary elements satisfying (3.2), (3.4) holds and so is (i). This completes the proof of the lemma. \square

REMARK 3.3. A close look at the proofs shows that the conclusions of Theorem 3.1 also hold if instead of Assumption 2.5, we impose any of the equivalent assertions (iii)–(v) of Lemma 3.2.

3.1. *Characterization of the primal and dual domains.* The polar, A° , of a nonempty subset A of $\mathbb{R} \times \mathbb{L}^\infty(dK)$, is the subset of $\mathbb{R} \times \mathbb{L}^1(dK)$, defined by

$$A^\circ := \left\{ (y, r) \in \mathbb{R} \times \mathbb{L}^1(dK) : xy + \int_0^T q_s r_s dK_s \leq 1, \text{ for every } (x, q) \in A \right\}.$$

The polar of a subset of $\mathbb{R} \times \mathbb{L}^1(dK)$ is defined similarly.

PROPOSITION 3.4. *Under Assumption 2.5 and conditions (2.1) and (2.3), we have:*

(i) *Let $(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK)$. Then $c \in \mathcal{A}(x, q)$ (thus, $\mathcal{A}(x, q) \neq \emptyset$ so $(x, q) \in \mathcal{K}$) if and only if*

$$\mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq xy + \int_0^T q_s r_s dK_s \quad \text{for every } (y, r) \in \mathcal{L} \text{ and } Y \in \mathcal{Y}(y, r).$$

(ii) Likewise, for $(y, r) \in \mathbb{R} \times \mathbb{L}^1(dK)$ we have $(y, r) \in \mathcal{L}$ and $Y \in \mathcal{Y}(y, r)$ if and only if

$$\mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq xy + \int_0^T q_s r_s dK_s \quad \text{for every } (x, q) \in \mathcal{K} \text{ and } c \in \mathcal{A}(x, q).$$

We also have $\mathcal{K} = (-\mathcal{L})^o = \text{cl} \mathring{\mathcal{K}}$, where the closure is taken both in norm $\mathbb{R} \times \mathbb{L}^1(dK)$ and $\sigma(\mathbb{R} \times \mathbb{L}^\infty(dK), \mathbb{R} \times \mathbb{L}^1(dK))$ topologies.

REMARK 3.5. It follows from Proposition 3.4 that for every $(x, q) \in \mathcal{K}$, $\mathcal{A}(x, q) \neq \emptyset$ as $0 \in \mathcal{A}(x, q)$. Likewise, for every $(y, r) \in \mathcal{L}$, $\mathcal{Y}(y, r) \neq \emptyset$ as $0 \in \mathcal{Y}(y, r)$. Moreover, for every $(x, q) \in \mathring{\mathcal{K}}$, each of the sets $\mathcal{A}(x, q)$ and $\bigcup_{(y,r) \in \mathcal{L}: xy + \int_0^T r_s q_s dK_s \leq 1} \mathcal{Y}(y, r)$ contain a strictly positive element; see Lemma 3.14 below.

The proof of Proposition 3.4 will be given via several lemmas. Let

\mathcal{M}' be the set of equivalent local martingale measures, under which

X'' (from Lemma 3.2, item (v)) is a uniformly integrable martingale.

Note that by [10], Theorem 5.2, \mathcal{M}' is a nonempty, convex subset of \mathcal{M} , which is also dense in \mathcal{M} in the total variation norm.

REMARK 3.6. Even though the results in [10] are obtained under the condition that S is a locally bounded process, they also hold without local boundedness assumption; see the discussion in [20], Remark 3.4.

Let \mathcal{Z}' denote the set of the corresponding càdlàg densities, that is,

$$(3.5) \quad \mathcal{Z}' := \left\{ \text{càdlàg } Z : Z_t = \mathbb{E} \left[\frac{d\mathbb{Q}}{d\mathbb{P}} \middle| \mathcal{F}_t \right], t \in [0, T], \mathbb{Q} \in \mathcal{M}' \right\}.$$

We also set

$$(3.6) \quad \Upsilon := \{ 1_{[0, \tau]}(t), t \in [0, T] : \tau \in \Theta \}.$$

LEMMA 3.7. Let the conditions of Proposition 3.4 hold, $\mathbb{Q} \in \mathcal{M}'$, $Z = Z^\mathbb{Q}$ be the corresponding element of \mathcal{Z}' , and $\Lambda \in \Upsilon$. Then there exists $r \in \mathbb{L}^1(dK)$, uniquely defined by

$$(3.7) \quad \int_0^t r_s dK_s = \mathbb{E} \left[\int_0^t \Lambda_s Z_s e_s d\kappa_s \right], \quad t \in [0, T],$$

such that $(1, r) \in \mathcal{L}$ and $Z\Lambda = (Z_t \Lambda_t)_{t \in [0, T]} \in \mathcal{Y}(1, r)$.

PROOF. Let $(x, q) \in \mathcal{K}$ and $c \in \mathcal{A}(x, q)$. Then there exists $X \in \mathcal{X}(x, q)$, such that

$$(3.8) \quad \int_0^\tau c_s d\kappa_s \leq X_\tau + \int_0^\tau q_s e_s d\kappa_s \quad \text{for every } \tau \in \Theta.$$

In particular, (3.8) holds for the particular τ , such that $\Lambda_t = 1_{[0, \tau]}(t)$, $t \in [0, T]$. Let $G_t := \int_0^t (q_s e_s)^+ d\kappa_s$, $t \in [0, T]$, where $(\cdot)^+$ denotes the positive part. By [33], Theorem III.29, page 128, $\int_0^t G_s - dZ_s$, $t \in [0, T]$, is a local martingale, so let $(\sigma_n)_{n \in \mathbb{N}}$ be its localizing sequence. Then by the monotone convergence theorem, integration by parts formula, and the

optional sampling theorem, we get

$$\begin{aligned}
 & \mathbb{E} \left[\int_0^T Z_s (q_s e_s)^+ \Lambda_s d\kappa_s \right] \\
 (3.9) \quad &= \lim_{n \rightarrow \infty} \mathbb{E} \left[\int_0^{\sigma_n \wedge \tau} Z_s dG \right] \\
 &= \lim_{n \rightarrow \infty} \mathbb{E} [Z_{\sigma_n \wedge \tau} G_{\sigma_n \wedge \tau}] - \lim_{n \rightarrow \infty} \mathbb{E} \left[\int_0^{\sigma_n \wedge \tau} G_s - dZ_s \right] \\
 &= \lim_{n \rightarrow \infty} \mathbb{E} [Z_{\sigma_n \wedge \tau} G_{\sigma_n \wedge \tau}] = \lim_{n \rightarrow \infty} \mathbb{E}^{\mathbb{Q}} [G_{\sigma_n \wedge \tau}] = \mathbb{E}^{\mathbb{Q}} [G_\tau],
 \end{aligned}$$

where in the last equality we used the monotone convergence theorem again. Here, finiteness of $\mathbb{E}^{\mathbb{Q}} [G_\tau] = \mathbb{E}^{\mathbb{Q}} [\int_0^\tau (q_s e_s)^+ d\kappa_s]$ follows from Assumption 2.5 via Lemma 3.2, part (iii). In a similar manner, we can show that

$$\begin{aligned}
 (3.10) \quad & \mathbb{E} \left[\int_0^T (q_s e_s)^- Z_s \Lambda_s d\kappa_s \right] = \mathbb{E}^{\mathbb{Q}} \left[\int_0^\tau (q_s e_s)^- d\kappa_s \right] < \infty, \\
 & \mathbb{E} \left[\int_0^T c_s Z_s \Lambda_s d\kappa_s \right] = \mathbb{E}^{\mathbb{Q}} \left[\int_0^\tau c_s d\kappa_s \right] < \infty.
 \end{aligned}$$

[19], Lemma 4, applies here and asserts that X in (3.8) is a supermartingale under \mathbb{Q} . Therefore, from (3.8), using (3.9) and (3.10), and taking expectation under \mathbb{Q} , we get

$$(3.11) \quad \mathbb{E} \left[\int_0^T c_s \Lambda_s Z_s d\kappa_s \right] \leq x + \mathbb{E} \left[\int_0^T q_s e_s Z_s \Lambda_s d\kappa_s \right].$$

Let us define

$$R_t := \mathbb{E} \left[\int_0^t \Lambda_s Z_s e_s d\kappa_s \right], \quad t \in [0, T].$$

Using the monotone class theorem, we obtain

$$(3.12) \quad \mathbb{E} \left[\int_0^T \tilde{q}_s e_s Z_s \Lambda_s d\kappa_s \right] = \int_0^T \tilde{q}_s dR_s \quad \text{for every } \tilde{q} \in \mathbb{L}^\infty(dK).$$

We claim that dR is absolutely continuous with respect to dK . First, using the $\pi - \lambda$ theorem, one can show that for every Borel-measurable subset A of $[0, T]$, we have

$$K(A) = \mathbb{E} \left[\int_0^T 1_A(t) d\kappa_t \right] \quad \text{and} \quad R(A) = \mathbb{E} \left[\int_0^T 1_A(t) \Lambda_t Z_t e_t d\kappa_t \right].$$

Thus, if for some A , $K(A) = 0$, then $\int_0^T 1_A(t) d\kappa_t = 0$ a.s. and $\kappa_t^A := \int_0^t 1_A(s) d\kappa_s$, $t \in [0, T]$, satisfies $\kappa_T^A = 0$ a.s. and $\int_0^T \Lambda_t Z_t e_t d\kappa_t^A = \int_0^T \Lambda_t Z_t e_t 1_A(t) d\kappa_t = 0$ a.s.

As dR is absolutely continuous with respect to dK , there exists a unique $r \in \mathbb{L}^1(dK)$, such that (3.7) holds. Since the left-hand side in (3.11) is nonnegative and since (x, q) is an arbitrary element of \mathcal{K} , we deduce from the definition of \mathcal{L} , (2.6), that $(1, r) \in \mathcal{L}$. Finally, it follows from (3.11) and (3.12) that $Z\Lambda \in \mathcal{Y}(1, r)$. This completes the proof of the lemma. □

REMARK 3.8. The natural convexification of the set Υ consists of nonnegative left-continuous decreasing and adapted processes D such that $D_{\theta_0} = 1$. In the context of utility-maximization constraints (and for $\theta_0 = 0$), this follows from [18] and [16]. We investigate convexification of the constraints in the later Sections 5 and 6, where we will extend Lemma 3.7 to a more general set of decreasing processes than Λ that drives the dual domain and that allows for the multiplicative decomposition of the dual minimizer.

COROLLARY 3.9. *Let the conditions of Proposition 3.4 hold, $\mathbb{Q} \in \mathcal{M}'$, Z be the càdlàg modification of the density process $\mathbb{E}[\frac{d\mathbb{Q}}{d\mathbb{P}}|\mathcal{F}_t]$, $t \in [0, T]$. Then there exists $r \in \mathbb{L}^1(dK)$, such that $Z \in \mathcal{Y}(1, r)$, where*

$$\int_0^t r_s dK_s = \mathbb{E} \left[\int_0^t Z_s e_s d\kappa_s \right], \quad t \in [0, T],$$

and $(1, r) \in \mathcal{L}$.

PROOF OF LEMMA 2.2. Let us fix $\mathbb{Q} \in \mathcal{M}'$ and let $Z \in \mathcal{Z}'$ be the corresponding density process. As in Lemma 3.7, we can show that there exists $r \in \mathbb{L}^1(dK)$, such that for every bounded and Borel measurable function q on $[0, T]$ we have

$$(3.13) \quad \int_0^t q_s r_s dK_s = \mathbb{E} \left[\int_0^t q_s Z_s |e_s| d\kappa_s \right], \quad t \in [0, T].$$

Let $\bar{q} := |q^1 - q^2|$, then as $\bar{q} = 0$, dK -a.e., we get

$$\mathbb{E} \left[\int_0^T \bar{q}_s |e_s| Z_s d\kappa_s \right] = \int_0^T \bar{q}_s r_s dK = 0.$$

Therefore, using integration by parts and via (3.13), we obtain

$$0 = \int_0^T \bar{q}_s r_s dK = \mathbb{E} \left[\int_0^T q_s Z_s |e_s| d\kappa_s \right] = \mathbb{E}^{\mathbb{Q}} \left[\int_0^T q_s |e_s| d\kappa_s \right].$$

Consequently, $\int_0^T q_s |e_s| d\kappa_s = 0$, \mathbb{Q} -a.s., and by the equivalence of \mathbb{Q} and \mathbb{P} , also \mathbb{P} -a.s. As, by construction $\int_0^t q_s |e_s| d\kappa_s = 0$, $t \in [0, T]$, is a nonnegative and nondecreasing process, whose terminal value is 0, \mathbb{P} -a.s.; we conclude that it is indistinguishable from the 0-valued process. The assertions of the lemma follows. \square

LEMMA 3.10. *Let the conditions of Proposition 3.4 hold, $(x, q) \in \mathcal{K}$, and c is a nonnegative optional process. Then $c \in \mathcal{A}(x, q)$ if and only if*

$$(3.14) \quad \mathbb{E} \left[\int_0^T c_s \Lambda_s Z_s d\kappa_s \right] \leq x + \mathbb{E} \left[\int_0^T q_s e_s \Lambda_s Z_s d\kappa_s \right] = x + \int_0^T q_s r_s dK_s,$$

for every $Z \in \mathcal{Z}'$ and $\Lambda \in \Upsilon$, where r is given by (3.7).

PROOF. Let $c \in \mathcal{A}(x, q)$. Then for every $Z \in \mathcal{Z}'$ and $\Lambda \in \Upsilon$, the validity of (3.14) follows from the definition of $\mathcal{A}(x, q)$, integration by parts formula and supermartingale property of every $X \in \mathcal{X}(x, q)$ under every $\mathbb{Q} \in \mathcal{M}'$, which in turn follows from [19], Lemma 4.

Conversely, let (3.14) holds for every $Z \in \mathcal{Z}'$ and $\Lambda \in \Upsilon$. Then we have

$$(3.15) \quad \mathbb{E} \left[\int_0^T (c_s - q_s e_s) Z_s \Lambda_s d\kappa_s \right] \leq x,$$

which, in view of the definition of Υ in (3.6), localization and integration by parts, implies that

$$\sup_{\mathbb{Q} \in \mathcal{M}', \tau \in \Theta} \mathbb{E}^{\mathbb{Q}} \left[\int_0^\tau (c_s - q_s e_s) d\kappa_s \right] \leq x.$$

For X'' given by Lemma 3.2, item (v), let us denote

$$(3.16) \quad f_t := \|q\|_{\mathbb{L}^\infty(dK)} X_t'' + \int_0^t (c_s - q_s e_s) d\kappa_s, \quad t \in [0, T].$$

It follows from Assumption 2.5 and item (v) of Lemma 3.2 that f is a nonnegative process. We observe that the proof of [27], Proposition 4.3, goes through, if we only take stopping times in Θ and measures in \mathcal{M}' . This proposition allows to conclude that there exists a non-negative càdlàg process V , such that

$$(3.17) \quad V_t = \operatorname{ess\,sup}_{\tau \in \Theta: \tau \geq t, \mathbb{Q} \in \mathcal{M}'} \mathbb{E}^{\mathbb{Q}}[f_{\tau} | \mathcal{F}_t], \quad t \in [0, T],$$

which is a supermartingale for every $\mathbb{Q} \in \mathcal{M}'$. Therefore, by the density of \mathcal{M}' in \mathcal{M} in the norm topology of $\mathbb{L}^1(\mathbb{P})$ and Fatou’s lemma, V is a supermartingale under every $\mathbb{Q} \in \mathcal{M}$. Moreover, V_0 satisfies

$$V_0 \leq x + \|q\|_{\mathbb{L}^\infty(dK)} X_0'',$$

by (3.17), (3.15), and by following the argument in the proof of Lemma 3.7.

We would like to apply the optional decomposition theorem of Følmer and Kramkov, [17], Theorem 3.1. For this, we need to show that V is a local supermartingale under every \mathbb{Q} , such that every $X \in \mathcal{X}(1)$ is a \mathbb{Q} -local supermartingale. However, \mathcal{M} is dense in the set of such measures in the norm topology of $\mathbb{L}^1(\mathbb{P})$, by the results of Delbaen and Schachermayer; see [11], Proposition 4.7. Therefore, the supermartingale property of V under every such \mathbb{Q} follows from Fatou’s lemma and supermartingale property of V under every $\mathbb{Q} \in \mathcal{M}$ established above. Therefore, by [17], Theorem 3.1, we get

$$V_t = V_0 + H \cdot S_t - A_t, \quad t \in [0, T],$$

where A is a nonnegative increasing process that starts at 0. Subtracting the constant $\|q\|_{\mathbb{L}^\infty(dK)} X_0''$ from both sides of (3.16), we get

$$\begin{aligned} \int_0^\tau (c_s - q_s e_s) d\kappa_s &= f_\tau - \|q\|_{\mathbb{L}^\infty(dK)} X_0'' \leq V_\tau - \|q\|_{\mathbb{L}^\infty(dK)} X_0'' \\ &= V_0 - \|q\|_{\mathbb{L}^\infty(dK)} X_0'' + H \cdot S_\tau - A_\tau \leq x + H \cdot S_\tau, \quad \tau \in \Theta, \end{aligned}$$

where $x + H \cdot S$ is acceptable, since V is nonnegative. Consequently, $c \in \mathcal{A}(x, q)$. This completes the proof of the lemma. \square

PROOF OF PROPOSITION 3.4. The assertions of item (i) follow from Lemma 3.10. It remains to show that the affirmations of item (ii) hold. Fix a $(y, r) \in \mathcal{L}$. If $Y \in \mathcal{Y}(y, r)$, (ii) follows from the definition of $\mathcal{Y}(y, r)$. Conversely, if (ii) holds for a nonnegative process Y , then since $(x, 0) \in \mathcal{K}$ for every $x > 0$, we have

$$\mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq 1 \quad \text{for every } c \in \mathcal{A} \left(\frac{1}{y}, 0 \right).$$

Via [31], Proposition 4.4, we deduce that $Y \in \mathcal{Y}(y)$ and is such that (ii) holds. Therefore, $Y \in \mathcal{Y}(y, r)$.

We have $\mathring{\mathcal{K}} \neq \emptyset$, where the interior is taken with respect to the norm-topology. According to Proposition 3.4(i), we have also $\mathcal{K} = (-\mathcal{L})^o$. The set \mathcal{K} , as the polar of \mathcal{L} , is convex and closed both in (strong) $\mathbb{R} \times \mathbb{L}^\infty(dK)$ and $\sigma(\mathbb{R} \times \mathbb{L}^\infty(dK), \mathbb{R} \times \mathbb{L}^1(dK))$ topologies. Having nonempty strong interior, we obtain $\mathcal{K} = \operatorname{cl} \mathring{\mathcal{K}}$ where the closure is in the strong-topology. Since \mathcal{K} is also closed in the weaker $\sigma(\mathbb{R} \times \mathbb{L}^\infty(dK), \mathbb{R} \times \mathbb{L}^1(dK))$ topology, we obtain that

$$\mathcal{K} = (-\mathcal{L})^o = \operatorname{cl} \mathring{\mathcal{K}},$$

where the closure is taken in both topologies. \square

3.2. *Preliminary properties of the value functions, in particular finiteness.* Let \mathbf{L}_+^0 be the positive orthant of \mathbf{L}^0 . The polar of a set $A \subseteq \mathbf{L}_+^0$ is defined as

$$A^o := \left\{ c \in \mathbf{L}_+^0 : \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq 1, \text{ for every } Y \in A \right\}.$$

We recall that the sets \mathcal{Z} and \mathcal{Z}' are defined in (2.7) and (3.5), respectively.

LEMMA 3.11. *Under the conditions of Theorem 3.1, we have*

$$(3.18) \quad (\mathcal{Z}')^{oo} = \mathcal{Y}(1)$$

and

$$(3.19) \quad \tilde{w}(y) = \inf_{Y \in \mathcal{Z}'} \mathbb{E} \left[\int_0^T V(t, yY_s) d\kappa_s \right] < \infty, \quad y > 0.$$

PROOF. It follows from [10], Theorem 5.2, that \mathcal{Z}' is dense in \mathcal{Z} in \mathbf{L}^0 . It follows from Fatou’s lemma that

$$(\mathcal{Z}')^o = \mathcal{Z}^o.$$

Therefore, [31], Lemma 4.2 and Proposition 4.4, imply (3.18).

One can show that \mathcal{Z}' is closed under countable convex combinations, where the martingale property follows from the monotone convergence theorem and the càdlàg structure of the limit is guaranteed by [12], Theorem VI.18; see also [26], Proposition 5.1, for more details in similar settings. Now, (3.19) follows (up to a notational change) from [31], Theorem 3.3. This completes the proof of the lemma. □

LEMMA 3.12. *Under the conditions of Theorem 3.1, for every $(x, q) \in \mathcal{K}$ and $(y, r) \in \mathcal{L}$, we have*

$$u(x, q) \leq v(y, r) + xy + \int_0^T r_s q_s dK_s.$$

PROOF. Fix an arbitrary $(x, q) \in \text{cl } \mathcal{K}$, $c \in \mathcal{A}(x, q)$ as well as $(y, r) \in \mathcal{L}$, $Y \in \mathcal{Y}(y, r)$. Using Proposition 3.4 and (2.5), we get

$$\begin{aligned} \mathbb{E} \left[\int_0^T U(t, c_s) d\kappa_s \right] &\leq \mathbb{E} \left[\int_0^T U(t, c_s) d\kappa_s \right] + xy + \int_0^T r_s q_s dK_s \\ &\quad - \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \\ &\leq \mathbb{E} \left[\int_0^T V(t, Y_s) d\kappa_s \right] + xy + \int_0^T r_s q_s dK_s. \end{aligned}$$

This implies the assertion of the lemma. □

For every (x, q) in \mathcal{K} , we define

$$(3.20) \quad \begin{aligned} \mathcal{B}(x, q) &:= \left\{ (y, r) \in \mathcal{L} : xy + \int_0^T r_s q_s dK_s \leq 1 \right\}, \\ \mathcal{D}(x, q) &:= \bigcup_{(y,r) \in \mathcal{B}(x,q)} \mathcal{Y}(y, r). \end{aligned}$$

The subsequent lemma established boundedness of $\mathcal{B}(x, q)$ for (x, q) in $\mathring{\mathcal{K}}$ in $\mathbb{R} \times \mathbb{L}^1(dK)$.

LEMMA 3.13. *Under the conditions of Theorem 3.1, for every $(x, q) \in \mathring{\mathcal{K}}$, $\mathcal{B}(x, q)$ is bounded in $\mathbb{R} \times \mathbb{L}^1(dK)$.*

PROOF. Fix an $(x, q) \in \mathring{\mathcal{K}}$. Then there exists $\varepsilon > 0$, such that for every

$$(3.21) \quad |x'| \leq \varepsilon \quad \text{and} \quad \|q'\|_{\mathbb{L}^\infty} \leq \varepsilon,$$

we have $(x + x', q + q') \in \mathring{\mathcal{K}}$. Let us fix an arbitrary $(y, r) \in \mathcal{B}(x, q)$. Then for every (x', q') satisfying (3.21), by the definitions of \mathcal{L} and $\mathcal{B}(x, q)$, respectively, we get

$$\begin{aligned} xy + \int_0^T q_s r_s dK_s + x'y + \int_0^T q'_s r_s dK_s &\geq 0, \\ xy + \int_0^T q_s r_s dK_s &\leq 1, \end{aligned}$$

which implies that

$$(3.22) \quad -x'y - \int_0^T q'_s r_s dK_s \leq xy + \int_0^T q_s r_s dK_s \leq 1.$$

Taking

$$x' = -\varepsilon \quad \text{and} \quad q' \equiv 0,$$

we deduce from (3.22) that $y \leq \frac{1}{\varepsilon}$. Also, by the definition of \mathcal{L} and Lemma 3.2, item (i), $y \geq 0$. In turn, setting

$$x' = 0 \quad \text{and} \quad q' = -\varepsilon 1_{\{r \geq 0\}} + \varepsilon 1_{\{r < 0\}},$$

we obtain from (3.22) that $\|r\|_{\mathbb{L}^1} \leq \frac{1}{\varepsilon}$. This completes the proof of the lemma. \square

LEMMA 3.14. *Let the conditions of Theorem 3.1 hold and (x, q) be an arbitrary element of $\mathring{\mathcal{K}}$. Then we have:*

- (i) $\mathcal{A}(x, q)$ contains a strictly positive process.
- (ii) The constant $\bar{y}(x, q)$ given by

$$(3.23) \quad \bar{y}(x, q) := \frac{1}{|x| + \|q\|_{\mathbb{L}^\infty(dK)} \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}[\int_0^T |e_s| d\kappa_s]},$$

takes values in $(0, \infty)$ and satisfies

$$(3.24) \quad \bar{y}(x, q) \mathcal{Z}' \subseteq \mathcal{D}(x, q).$$

In particular, for every $z > 0$, $z\mathcal{D}(x, q)$ contains a strictly positive process Y such that

$$(3.25) \quad \mathbb{E}\left[\int_0^T V(s, Y_s) d\kappa_s\right] < \infty.$$

PROOF. In order to show (i), we observe that the existence of a positive process in $\mathcal{A}(x, q)$ follows from the fact that $(x - \delta, q) \in \mathcal{K}$ for a sufficiently small δ . Now the constant-valued consumption $\delta/A > 0$, where A is the constant that dominates the terminal value of the stochastic clock κ in (2.1), is in $\mathcal{A}(x, q)$.

In order to prove (ii), let us consider $\bar{y}(x, q)$ given by (3.23). It follows from Lemma 3.2, item (iii), that $\bar{y}(x, q) \in (0, \infty)$. For this $\bar{y}(x, q)$, using Corollary 3.9, one can show (3.24). This and Lemma 3.11 (note that finiteness of \tilde{w} follows directly from (3.1)) imply that for every $z > 0$, there exists a positive $Y \in z\mathcal{D}(x, q)$, such that (3.25) holds. \square

LEMMA 3.15. *Under the conditions of Theorem 3.1, for every $(x, q) \in \mathring{\mathcal{K}}$ we have*

$$-\infty < u(x, q) < \infty.$$

and $u < \infty$ on $\mathbb{R} \times \mathbb{L}^\infty(dK)$.

PROOF. Let us fix an arbitrary $(x, q) \in \mathring{\mathcal{K}}$. Since $\mathring{\mathcal{K}}$ is an open convex cone, there exists $\lambda \in (0, 1)$, $(x_1, q_1) \in \mathring{\mathcal{K}}$, and $x_2 > 0$, such that

$$(x, q) = \lambda(x_1, q_1) + (1 - \lambda)(x_2, 0).$$

Note that $(x_2, 0) \in \mathring{\mathcal{K}}$ by Lemma 3.2. By (3.1), there $c \in \mathcal{A}(x_2, 0)$, such that

$$(3.26) \quad \mathbb{E} \left[\int_0^T U(t, (1 - \lambda)c_t) d\kappa_t \right] > -\infty.$$

As $\mathcal{A}(x_1, q_1) \neq \emptyset$ (see Remark 3.5), there exists $\tilde{c} \in \mathcal{A}(x_1, q_1)$. As $U(t, \cdot)$ is nondecreasing, we get

$$\begin{aligned} u(x, q) &\geq \mathbb{E} \left[\int_0^T U(t, \lambda\tilde{c}_t + (1 - \lambda)c_t) d\kappa_t \right] \\ &\geq \mathbb{E} \left[\int_0^T U(t, (1 - \lambda)c_t) d\kappa_t \right] > -\infty, \end{aligned}$$

where the last inequality follows from (3.26). This implies finiteness of u on $\mathring{\mathcal{K}}$ from below.

In order to show finiteness from above, let us fix a process $c \in \mathcal{A}(x, q)$, such that

$$\mathbb{E} \left[\int_0^T U(t, c_t) d\kappa_t \right] > -\infty.$$

By Lemma 3.11, there exists $Y \in \mathcal{Z}'$, such that

$$(3.27) \quad \mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right] < \infty.$$

It follows from Lemma 3.7 that $Y \in \mathcal{Y}(1, \rho)$ for some $(1, \rho) \in \mathcal{L}$. Therefore, by Proposition 3.4, we get

$$\begin{aligned} (3.28) \quad \mathbb{E} \left[\int_0^T U(s, c_s) d\kappa_s \right] &\leq \mathbb{E} \left[\int_0^T U(s, c_s) d\kappa_s \right] + x + \int_0^T \rho_s q_s dK_s \\ &\quad - \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \\ &\leq \mathbb{E} \left[\int_0^T V(s, Y_s) d\kappa_s \right] + x + \int_0^T \rho_s q_s dK_s. \end{aligned}$$

As Y satisfies (3.27), we conclude that $u(x, q) < \infty$. Moreover, for $(x, q) \in \mathcal{K}$, as $\mathcal{A}(x, q) \neq \emptyset$ by Remark 3.5, every $c \in \mathcal{A}(x, q)$ satisfies (3.28) (with the same Y). This implies that $u < \infty$ on \mathcal{K} and, therefore, by (2.11), on $\mathbb{R} \times \mathbb{L}^\infty(dK)$. This completes the proof of the lemma. \square

We recall that, for every $(x, q) \in \mathcal{K}$, $\mathcal{D}(x, q)$ is defined in (3.20). Let $cl\mathcal{D}(x, q)$ denote the closure of $\mathcal{D}(x, q)$ in $\mathbb{L}^0(d\kappa \times \mathbb{P})$. The following lemma proves a delicate point that, for $(x, q) \in \mathring{\mathcal{K}}$, by passing from $\mathcal{D}(x, q)$ to $cl\mathcal{D}(x, q)$, we do not change the auxiliary dual value function.

LEMMA 3.16. *Let the conditions of Theorem 3.1 hold and $(x, q) \in \mathring{\mathcal{K}}$. Then, for every $z > 0$, we have*

$$\begin{aligned}
 (3.29) \quad & -\infty < \inf_{Y \in \text{cl } \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(s, zY_s) d\kappa_s \right] \\
 & = \inf_{Y \in \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(s, zY_s) d\kappa_s \right] < \infty.
 \end{aligned}$$

PROOF. Finiteness from above follows from Lemma 3.14. To show finiteness of both infima in (3.29) from below, by Lemma 3.15 we deduce the existence of $c \in \mathcal{A}(x, q)$, such that

$$(3.30) \quad \mathbb{E} \left[\int_0^T U(s, c_s) d\kappa_s \right] > -\infty.$$

Let $Y \in \text{cl } \mathcal{D}(x, q)$ and let $Y^n \in \mathcal{Y}(y^n, r^n)$, $n \geq 1$, be a sequence in $\mathcal{D}(x, q)$ that converges to Y in $\mathbb{L}^0(d\kappa \times \mathbb{P})$. By Fatou’s lemma, Proposition 3.4 and the definition of the set $\mathcal{B}(x, q)$ in (3.20), we get

$$\begin{aligned}
 \mathbb{E} \left[\int_0^T Y_s c_s d\kappa_s \right] & \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T Y_s^n c_s d\kappa_s \right] \\
 & \leq \sup_{n \geq 1} \left(x y^n + \int_0^T r_s^n q_s dK_s \right) \leq 1.
 \end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
 \mathbb{E} \left[\int_0^T U(s, c_s) d\kappa_s \right] & \leq \mathbb{E} \left[\int_0^T U(s, c_s) d\kappa_s \right] + 1 - \mathbb{E} \left[\int_0^T Y_s c_s d\kappa_s \right] \\
 & \leq \mathbb{E} \left[\int_0^T V(s, Y_s) d\kappa_s \right] + 1,
 \end{aligned}$$

which together with (3.30) implies finiteness of both infima in (3.29) from below.

Let us show equality of two infima in (3.29). It follows from Lemma 3.14 that for every $z > 0$ there exists a process $Y \in z\mathcal{D}(x, q)$, such that

$$\mathbb{E} \left[\int_0^T V(s, Y_s) d\kappa_s \right] < \infty.$$

Let us fix $z > 0$ and let $\bar{Y} \in \text{cl } \mathcal{D}(x, q)$. Also, let $(Y^n)_{n \in \mathbb{N}}$ be a sequence in $\mathcal{D}(x, q)$ that converges to \bar{Y} ($d\kappa \times \mathbb{P}$)-a.e. Let us fix $\delta > 0$, then by Lemma 3.11, there exists $Z' \in \mathcal{Z}'$, such that

$$\mathbb{E} \left[\int_0^T V(t, \delta \bar{y}(x, q) Z'_t) d\kappa_t \right] < \infty,$$

where $\bar{y}(x, q)$ is defined in (3.23). Note that $\bar{y}(x, q) Z' \in \mathcal{D}(x, q)$ by Lemma 3.14 (see (3.24)). Therefore, using Fatou’s lemma and monotonicity of V in the spatial variable, we obtain

$$\begin{aligned}
 & \inf_{Y \in \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, (z + \delta) Y_s) d\kappa_s \right] \\
 & \leq \limsup_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T V(t, z Y_s^n + \delta \bar{y}(x, q) Z'_s) d\kappa_s \right] \\
 & \leq \mathbb{E} \left[\int_0^T V(t, z \bar{Y}_s + \delta \bar{y}(x, q) Z'_s) d\kappa_s \right] \\
 & \leq \mathbb{E} \left[\int_0^T V(t, z \bar{Y}_s) d\kappa_s \right].
 \end{aligned}$$

Taking the infimum over $Y \in \text{cl } \mathcal{D}(x, y)$, we deduce that

$$(3.31) \quad \inf_{Y \in \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, (z + \delta)Y_s) d\kappa_s \right] \leq \inf_{Y \in \text{cl } \mathcal{D}(x, y)} \mathbb{E} \left[\int_0^T V(t, zY_s) d\kappa_s \right].$$

Let us consider

$$\phi(z) := \inf_{Y \in \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, zY_s) d\kappa_s \right], \quad z > 0.$$

By the first part of the proof (finiteness of both infima), ϕ is finite-valued on $(0, \infty)$. Convexity of V in the spatial variable implies that ϕ is also convex. Therefore, ϕ is continuous. As (3.31) holds for every $\delta > 0$, by taking the limit as $\delta \downarrow 0$ in (3.31), we conclude that both infima in (3.29) are equal. This completes the proof of the lemma. \square

Let us define

$$(3.32) \quad \mathcal{E} := \{(y, r) \in \mathcal{L} : v(y, r) < \infty\}.$$

LEMMA 3.17. *Under the conditions of Theorem 3.1, for every $(y, r) \in \mathcal{L}$, we have*

$$v(y, r) > -\infty.$$

Therefore, $v > -\infty$ on $\mathbb{R} \times \mathbb{L}^1(dK)$. The set \mathcal{E} is a nonempty convex subset of \mathcal{L} , whose closure in $\mathbb{R} \times \mathbb{L}^1(dK)$ equals to \mathcal{L} , and such that

$$(3.33) \quad \mathcal{E} = \bigcup_{\lambda \geq 1} \lambda \mathcal{E}.$$

PROOF. Let us fix $(y, r) \in \mathcal{L}$, then finiteness of $v(y, r)$ from below follows from (3.1) and Lemma 3.12. To establish the properties of \mathcal{E} , we observe that the convexity of \mathcal{E} and (3.33) follow from convexity and monotonicity of V , respectively.

It remains to show that the closure of \mathcal{E} in $\mathbb{R} \times \mathbb{L}^1(dK)$ contains the origin. In (3.20), let us consider $(x, q) = (1, 0) \in \mathcal{K}$. In this case, we have

$$\mathcal{D}(1, 0) = \bigcup_{(y, r) \in \mathcal{L}: y \leq 1} \mathcal{Y}(y, r) \subseteq \mathcal{Y}(1),$$

where the last inclusion follows from the very definition of $\mathcal{Y}(y, r)$'s in (2.9). As, by (2.8), $\mathcal{Y}(1)$ is closed in $\mathbb{L}^0(d\kappa \times \mathbb{P})$ and $\mathcal{D}(1, 0) \subseteq \mathcal{Y}(1)$, we deduce that

$$(3.34) \quad \text{cl } \mathcal{D}(1, 0) \subseteq \mathcal{Y}(1).$$

By Lemma 3.14, $\mathcal{Z}' \subset \mathcal{D}(1, 0)$, as $\bar{y}(1, 0) = 1$. Therefore, by the bipolar theorem of Brannath and Schachermayer, [3], Theorem 1.3, we get

$$(3.35) \quad (\mathcal{Z}')^{oo} \subseteq \text{cl } \mathcal{D}(1, 0).$$

On the other hand, Lemma 3.11 asserts that

$$(3.36) \quad (\mathcal{Z}')^{oo} = \mathcal{Y}(1).$$

Combining (3.34), (3.35) and (3.36), we conclude

$$\text{cl } \mathcal{D}(1, 0) = \mathcal{Y}(1).$$

Therefore, the sets $\text{cl } \mathcal{D}(1, 0) = \mathcal{Y}(1)$ and $\mathcal{A}(1, 0)$ satisfy the precise technical assumptions of [31], Theorem 3.2, which, for every $x > 0$, grants the existence of $\hat{c}(x) \in \mathcal{A}(x, 0)$, the unique maximizer to $w(x)$, where w is defined in (2.12). For every $x > 0$, we set

$$Y(x) := U'(\cdot, \hat{c}(x)), \quad (d\kappa \times \mathbb{P})\text{-a.e.}$$

By [31], Theorem 3.2, for every $x > 0$, $Y(x)$ satisfies

$$Y(x) \in w'(x) \text{ cl } \mathcal{D}(1, 0) \quad \text{and} \quad \mathbb{E} \left[\int_0^T V(t, Y_t(x)) d\kappa_t \right] < \infty, \quad x > 0,$$

where by [31], Theorem 3.2, w is a strictly concave, differentiable function on $(0, \infty)$ that satisfies the Inada conditions. Therefore, as $w'(x)$ can be arbitrary close to 0 (by taking x large enough and by using the Inada conditions) and by Lemmas 3.14 and 3.16, we conclude that the closure of \mathcal{E} in $\mathbb{R} \times \mathbb{L}^1(dK)$ contains origin.

In order to prove that the closure of \mathcal{E} in $\mathbb{R} \times \mathbb{L}^1(dK)$ equals to \mathcal{L} , let $(y, r) \in \mathcal{L} \setminus (0, 0)$ be fixed. Let us take $\varepsilon > 0$. We want to find (\tilde{y}, \tilde{r}) , such that

$$(3.37) \quad |\tilde{y} - y| + \|\tilde{r} - r\|_{\mathbb{L}^1(dK)} < \varepsilon,$$

and

$$(3.38) \quad (\tilde{y}, \tilde{r}) \in \mathcal{E}.$$

As the closure of \mathcal{E} in $\mathbb{R} \times \mathbb{L}^1(dK)$ contains origin, we can pick $(y^0, r^0) \in \mathcal{E}$, such that

$$(3.39) \quad |y^0| + \|r^0\|_{\mathbb{L}^1(dK)} \leq \varepsilon/3$$

and $Y \in \mathcal{Y}(y^0, r^0)$, such that

$$(3.40) \quad \mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right] < \infty.$$

Let us fix $\alpha > 1$, such that

$$(3.41) \quad \frac{|y| + \|r\|_{\mathbb{L}^1(dK)}}{\alpha} \leq \varepsilon/3$$

and set $\varepsilon' := \frac{1}{\alpha} \in (0, 1)$. By (3.33), $(\alpha y^0, \alpha r^0) \in \mathcal{E}$. Let

$$\tilde{y} := (1 - \varepsilon')y + \varepsilon'\alpha y^0, \quad \tilde{r} := (1 - \varepsilon')r + \varepsilon'\alpha r^0.$$

Then

$$\begin{aligned} |y - \tilde{y}| + \|r - \tilde{r}\|_{\mathbb{L}^1(dK)} &= \varepsilon'\alpha \left| \frac{y}{\alpha} - y^0 \right| + \varepsilon'\alpha \left\| \frac{r}{\alpha} - r^0 \right\|_{\mathbb{L}^1(dK)} \\ &\leq \frac{|y| + \|r\|_{\mathbb{L}^1(dK)}}{\alpha} + |y^0| + \|r^0\|_{\mathbb{L}^1(dK)} \\ &\leq \frac{2\varepsilon}{3}, \end{aligned}$$

where in the last inequality we have used (3.39) and (3.41). Thus (\tilde{y}, \tilde{r}) satisfies (3.37). Further, as $0 \in \mathcal{Y}(y, r)$, by convexity of \mathcal{L} and using Proposition 3.4, we get

$$Y = (1 - \varepsilon')0 + \varepsilon'\alpha Y \in \mathcal{Y}(\tilde{y}, \tilde{r}),$$

which by (3.40) implies (3.38). This completes the proof of the lemma. \square

3.3. Existence and uniqueness of solutions to (2.4) and (2.10); semicontinuity and biconjugacy of u and v .

LEMMA 3.18. *Under the conditions of Theorem 3.1, the value function v is convex, proper and lower semicontinuous with respect to the topology of $\mathbb{R} \times \mathbb{L}^1(dK)$. For every $(y, r) \in \mathcal{E}$, there exists a unique solution to (2.10). Likewise, u is concave, proper and upper semicontinuous with respect to the strong topology of $\mathbb{R} \times \mathbb{L}^\infty(dK)$. For every $(x, q) \in \{u > -\infty\}$, there exists a unique solution to (2.4).*

PROOF. Let $(y^n, r^n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{L} that converges to (y, r) in $\mathbb{R} \times \mathbb{L}^1(dK)$. Passing if necessary to a subsequence, we will assume that

$$(3.42) \quad \lim_{n \rightarrow \infty} v(y^n, r^n) = \liminf_{n \rightarrow \infty} v(y^n, r^n).$$

Let $Y^n \in \mathcal{Y}(y^n, r^n)$, $n \in \mathbb{N}$, be such that

$$(3.43) \quad \mathbb{E} \left[\int_0^T V(t, Y_t^n) d\kappa_t \right] \leq v(y^n, r^n) + \frac{1}{n}, \quad n \in \mathbb{N}.$$

By passing to convex combinations and applying Komlos'-type lemma (see, e.g., [9], Lemma A1.1), we may suppose that $\tilde{Y}^n \in \text{conv}(Y^n, Y^{n+1}, \dots)$, $n \in \mathbb{N}$, converges ($d\kappa \times \mathbb{P}$)-a.e. to some \hat{Y} .

For every $(x, q) \in \mathcal{K}$ and $c \in \mathcal{A}(x, q)$, by Fatou's lemma, we have

$$\mathbb{E} \left[\int_0^T c_t \hat{Y}_t d\kappa_t \right] \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T c_t \tilde{Y}_t^n d\kappa_t \right] \leq xy + \int_0^T q_s r_s dK_s.$$

Therefore, by Proposition 3.4, $\hat{Y} \in \mathcal{Y}(y, r)$. With $\bar{y} := \sup_{n \geq 1} y^n$, we have $(\tilde{Y}^n)_{n \in \mathbb{N}} \subseteq \mathcal{Y}(\bar{y})$. Therefore, by [31], Lemma 3.5, we deduce that $V^-(t, \tilde{Y}_t^n)$, $n \in \mathbb{N}$, is a uniformly integrable sequence. Combining uniform integrability with the convexity of V in the spatial variable, we get

$$(3.44) \quad \begin{aligned} v(y, r) &\leq \mathbb{E} \left[\int_0^T V(t, \hat{Y}_t) d\kappa_t \right] \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T V(t, \tilde{Y}_t^n) d\kappa_t \right] \\ &\leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T V(t, Y_t^n) d\kappa_t \right] = \liminf_{n \rightarrow \infty} v(y^n, r^n), \end{aligned}$$

where in the last equality we have used (3.42) and (3.43). Since (y^n, q^n) was an arbitrary sequence that converges to (y, r) , lower semicontinuity of v in strong topology of $\mathbb{R} \times \mathbb{L}^1(dK)$ follows. Since \mathcal{L} is closed and $v = \infty$ outside of \mathcal{L} , we deduce that v is lower semicontinuous on $\mathbb{R} \times \mathbb{L}^1(dK)$. The function v is proper by Lemma 3.17. Note that (3.44) also implies that \mathcal{E} defined in (3.32) is $\mathbb{R} \times \mathbb{L}^1(dK)$ -norm closed. For $(y, r) \in \mathcal{E}$, by taking $(y^n, r^n) = (y, r)$, $n \in \mathbb{N}$, we deduce the existence of a minimizer to (2.10). Strict convexity of V results in the uniqueness of the minimizer to (2.10). Convexity of v follows. Upper semicontinuity of u with respect to the norm-topology of $\mathbb{R} \times \mathbb{L}^\infty(dK)$ can be proven similarly, first proving semicontinuity on \mathcal{K} by a Fatou-type argument, then using the closedness of \mathcal{K} and the definition of u outside it. \square

COROLLARY 3.19. *Under the conditions of Theorem 3.1, $-u$ and v are also lower semicontinuous with respect to the weak topologies $\sigma(\mathbb{R} \times \mathbb{L}^\infty(dK), (\mathbb{R} \times \mathbb{L}^\infty)^*(dK))$ and $\sigma(\mathbb{R} \times \mathbb{L}^1(dK), \mathbb{R} \times \mathbb{L}^\infty(dK))$, respectively.*

PROOF. The assertions of the corollary is a consequence of [2], Proposition 2.2.10; see also [15], Corollary I.2.2. \square

We recall that $\text{cl } \mathcal{D}(x, q)$ denotes the closure of $\mathcal{D}(x, q)$ in $\mathbb{L}^0(d\kappa \times \mathbb{P})$.

LEMMA 3.20. *Under the conditions of Theorem 3.1, for every (x, q) in $\mathring{\mathcal{K}}$, and a non-negative optional process c , we have*

$$\begin{aligned} c \in \mathcal{A}(x, q) \quad &\text{if and only if} \\ \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] &\leq 1 \quad \text{for every } Y \in \text{cl } \mathcal{D}(x, q). \end{aligned}$$

PROOF. Let (x, q) in $\mathring{\mathcal{K}}$, c is a nonnegative optional process such that

$$(3.45) \quad \mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq 1 \quad \text{for every } Y \in \text{cl } \mathcal{D}(x, q).$$

Consider arbitrary $Z \in \mathcal{Z}'$ and $\Lambda \in \Upsilon$. Let the corresponding r be given by (3.7) and we set

$$y' := x + \int_0^T q_s r_s dK_s.$$

If $y' = 0$, then $y\Lambda Z \in \mathcal{D}(x, q)$ for every $y > 0$. Thus, by (3.45), we obtain that $\mathbb{E}[\int_0^T c_s y \Lambda_s Z_s d\kappa_s] \leq 1$. Taking the limit as $y \rightarrow \infty$, we get

$$(3.46) \quad \begin{aligned} \mathbb{E} \left[\int_0^T c_s \Lambda_s Z_s d\kappa_s \right] &= 0 = x + \int_0^T q_s r_s dK_s \\ &= x + \mathbb{E} \left[\int_0^T q_s e_s \Lambda_s Z_s d\kappa_s \right], \end{aligned}$$

where in the last equality we have used (3.7).

If $y' > 0$, then $\frac{1}{y'} \Lambda Z \in \mathcal{D}(x, q)$ and thus by (3.45), we obtain

$$\begin{aligned} \mathbb{E} \left[\int_0^T c_s \frac{1}{y'} \Lambda_s Z_s d\kappa_s \right] &\leq 1 = \frac{x + \int_0^T q_s r_s dK_s}{y'} \\ &= \frac{1}{y'} \left(x + \mathbb{E} \left[\int_0^T q_s e_s \Lambda_s Z_s d\kappa_s \right] \right), \end{aligned}$$

where in the last equality, we have used (3.7) again. Consequently, we deduce

$$\mathbb{E} \left[\int_0^T c_s \Lambda_s Z_s d\kappa_s \right] \leq x + \mathbb{E} \left[\int_0^T q_s e_s \Lambda_s Z_s d\kappa_s \right],$$

which together with (3.46), by Lemma 3.10, imply that $c \in \mathcal{A}(x, q)$.

Conversely, let $(x, q) \in \mathring{\mathcal{K}}$, $c \in \mathcal{A}(x, q)$ and $Y \in \text{cl } \mathcal{D}(x, q)$. Then there exists a sequence $Y^n \in \mathcal{Y}(y^n, r^n)$ convergent to Y , $(d\kappa \times \mathbb{P})$ -a.e., where $(y^n, r^n) \in \mathcal{B}(x, q)$. As

$$\mathbb{E} \left[\int_0^T c_s Y_s^n d\kappa_s \right] \leq 1, \quad n \in \mathbb{N},$$

by Fatou's lemma, we get

$$\mathbb{E} \left[\int_0^T c_s Y_s d\kappa_s \right] \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T c_s Y_s^n d\kappa_s \right] \leq 1.$$

This completes the proof of the lemma. \square

LEMMA 3.21. Under the conditions of Theorem 3.1, for every (x, q) in $\mathring{\mathcal{K}}$, we have

$$(3.47) \quad u(x, q) = \inf_{(y,r) \in \mathcal{L}} \left(v(y, r) + xy + \int_0^T r_s q_s dK_s \right).$$

PROOF. Let us fix $(x, q) \in \mathring{\mathcal{K}}$. By Lemma 3.14, $\mathcal{A}(x, q)$ and $\text{cl } \mathcal{D}(x, q)$ contain strictly positive elements. Therefore, using Lemma 3.20 we deduce that the sets $\mathcal{A}(x, q)$ and $\text{cl } \mathcal{D}(x, q)$ satisfy the assumptions of [31], Theorem 3.2. From this theorem, Lemma 3.16

and the definition of the set $\mathcal{B}(x, q)$, we get

$$\begin{aligned} u(x, q) &= \inf_{z>0} \left(\inf_{Y \in \text{cl } \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, zY_s) d\kappa_s \right] + z \right) \\ &= \inf_{z>0} \left(\inf_{Y \in \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, zY_s) d\kappa_s \right] + z \right) \\ &= \inf_{z>0} \left(\inf_{(y, r) \in z\mathcal{B}(x, q)} v(y, r) + z \right) \\ &\geq \inf_{(y, r) \in \mathcal{L}} \left(v(y, r) + xy + \int_0^T q_s r_s dK_s \right). \end{aligned}$$

Combining this with the conclusion of Lemma 3.12, we deduce that (3.47) holds for every $(x, q) \in \mathring{\mathcal{K}}$. \square

Before proving the biconjugacy relations of item (iii), Theorem 3.1, we need a preliminary lemma. Essentially, following the notation in [15], we define

$$\begin{aligned} (3.48) \quad v^*(x, q) &:= \inf_{(y, r) \in \mathbb{R} \times \mathbb{L}^1(dK)} \left(v(y, r) + xy + \int_0^T q_s r_s dK_s \right), \\ &(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK), \end{aligned}$$

$$\begin{aligned} (3.49) \quad v^{**}(y, r) &:= \sup_{(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK)} \left(v^*(x, q) - xy - \int_0^T q_s r_s dK_s \right), \\ &(y, r) \in \mathbb{R} \times \mathbb{L}^1(dK). \end{aligned}$$

REMARK 3.22. In [15], conjugate convex functions are considered on general spaces V and V^* supplied with $\sigma(V, V^*)$ and $\sigma(V^*, V)$ topologies, which in our case are $V = \mathbb{R} \times \mathbb{L}^1(dK)$, $V^* = \mathbb{R} \times \mathbb{L}^\infty(dK)$. Thus, the starting point of our analysis is v , not u . We remind the reader we have already proved that the dual value function v is convex, proper and lower-semicontinuous on the space $V = \mathbb{R} \times \mathbb{L}^1(dK)$.

LEMMA 3.23. *Under the conditions of Theorem 3.1, we have*

$$(3.50) \quad v^{**} = v,$$

$$(3.51) \quad v^*(x, q) = -\infty \quad \text{for every } (x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK) \setminus \mathcal{K}.$$

PROOF. To show (3.50), we observe that by Lemma 3.18, v is lower semicontinuous in the $\mathbb{R} \times \mathbb{L}^1(dK)$ -norm topology (and, therefore, by [15], Corollary I.2.2, also in the weak topology $\sigma(\mathbb{R} \times \mathbb{L}^1(dK), \mathbb{R} \times \mathbb{L}^\infty(dK))$). As a result, by [15], Proposition I.4.1, we get (3.50).

The proof of (3.51) will be done in several steps.

Step 1. Let $(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK) \setminus \mathcal{K}$. According to Proposition 3.4, there exists $(y, r) \in \mathcal{L}$, such that

$$C := xy + \int_0^T q_s r_s dK_s < 0.$$

Therefore, as \mathcal{L} is a cone, for every $a > 0$, $(ay, ar) \in \mathcal{L}$, and we have

$$(3.52) \quad xay + \int_0^T q_s ar_s dK_s = aC < 0.$$

Note that $0 \in \mathcal{Y}(ay, ar)$, $a > 0$.

Step 2. Let us consider $Z \in \mathcal{Z}'$, such that

$$(3.53) \quad \mathbb{E} \left[\int_0^T V \left(t, \frac{1}{2} Z_t \right) d\kappa_t \right] < \infty.$$

The existence of such a Z is granted by Lemma 3.11. Further, by Corollary 3.9, there exists $\rho \in \mathbb{L}^1(dK)$, such that $(1, \rho) \in \mathcal{L}$, $Z \in \mathcal{Y}(1, \rho)$, and

$$\int_0^t \rho_s dK_s = \mathbb{E} \left[\int_0^t Z_s e_s d\kappa_s \right], \quad t \in [0, T].$$

Let us set

$$D := x + \int_0^T q_s \rho_s dK_s \in \mathbb{R}.$$

Step 3. In (3.52), let us pick

$$a = \frac{|D| + 1}{-C} > 0.$$

Then we have

$$(3.54) \quad aC + D = -|D| + D - 1 < 0.$$

Step 4. Let us define

$$(3.55) \quad Y := \frac{1}{2} Z, \quad y' := \frac{1}{2} ay + \frac{1}{2} \quad \text{and} \quad r' := \frac{1}{2} ar + \frac{1}{2} \rho.$$

Then by (3.53), we obtain

$$(3.56) \quad \mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right] < \infty.$$

As $Z \in \mathcal{Y}(1, \rho)$ and $0 \in \mathcal{Y}(ay, ar)$, by convexity of \mathcal{L} and Proposition 3.4, we have

$$(3.57) \quad Y \in \mathcal{Y}(y', r'),$$

where y' and r' are defined in (3.55). Now, it follows from (3.56) and (3.57) that $(y', r') \in \mathcal{E}$ (where \mathcal{E} is defined in (3.32)). Therefore, we obtain

$$\begin{aligned} 2 \left(xy' + \int_0^T q_s r'_s dK_s \right) &= (ay + 1)x + \int_0^T (ar_s + \rho_s) q_s dK_s \\ &= a \left(xy + \int_0^T q_s r_s dK_s \right) + x + \int_0^T q_s \rho_s dK_s \\ &= aC + D < 0, \end{aligned}$$

where the last inequality follows from (3.54). To recapitulate, we have shown the existence of (y', r') , such that

$$(3.58) \quad (y', r') \in \mathcal{E} \quad \text{and} \quad xy' + \int_0^T q_s r'_s dK_s < 0.$$

Step 5. For y' and r' defined in (3.55), as $v(y', r') < \infty$ and $xy' + \int_0^T q_s r'_s dK_s < 0$ by (3.58), from the monotonicity of V , we get

$$\infty > v(y', r') \geq v(\lambda y', \lambda r'), \quad \lambda \geq 1.$$

As $\bigcup_{\lambda \geq 1} (\lambda y', \lambda r') \subset \mathcal{L}$, we conclude via (3.58) that

$$v^*(x, q) \leq \lim_{\lambda \rightarrow \infty} \left(v(\lambda y', \lambda r') + \lambda \left(xy' + \int_0^T q_s r'_s dK_s \right) \right) = -\infty.$$

Therefore, (3.51) holds. This completes the proof of the lemma. \square

LEMMA 3.24. *Under the conditions of Theorem 3.1, we have*

$$(3.59) \quad v(y, r) = \sup_{(x, q) \in \mathcal{K}} \left(u(x, q) - xy - \int_0^T r_s q_s dK_s \right), \quad (y, r) \in \mathcal{L},$$

$$(3.60) \quad u(x, q) = \inf_{(y, r) \in \mathcal{L}} \left(v(y, r) + xy + \int_0^T q_s r_s dK_s \right), \quad (x, q) \in \mathcal{K}.$$

PROOF. Lemma 3.21 and (2.11) imply that on $\mathring{\mathcal{K}}$, for v^* defined in (3.48), we have

$$(3.61) \quad v^* = u.$$

By Lemma 3.23, $v^* = -\infty$ on $\mathbb{R} \times \mathbb{L}^\infty(dK) \setminus \mathcal{K}$. From [15], Definition I.4.1, and Lemma 3.18, respectively, we deduce that both v^* and u are upper semicontinuous in the topology of $\mathbb{R} \times \mathbb{L}^\infty(dK)$. Consequently, from (3.61), using [15], Corollary I.2.1, we get

$$(3.62) \quad v^* = u \quad \text{on } \mathbb{R} \times \mathbb{L}^\infty(dK).$$

As a result, with v^{**} being defined in (3.49), for every $(y, r) \in \mathbb{R} \times \mathbb{L}^1(dK)$, we obtain

$$(3.63) \quad u^*(y, r) := \sup_{(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK)} \left(u(x, q) - xy - \int_0^T q_s r_s dK_s \right) = v^{**}(y, r).$$

Therefore, from (3.50) in Lemma 3.23 and (3.63), we get

$$v = u^* \quad \text{on } \mathbb{R} \times \mathbb{L}^1(dK).$$

As a result, applying Lemma 3.23 again and since $u = -\infty$ outside of \mathcal{K} by (2.11), we deduce

$$\begin{aligned} v(y, r) &= \sup_{(x, q) \in \mathbb{R} \times \mathbb{L}^\infty(dK)} \left(u(x, q) - xy - \int_0^T r_s q_s dK_s \right) \\ &= \sup_{(x, q) \in \mathcal{K}} \left(u(x, q) - xy - \int_0^T r_s q_s dK_s \right), \quad (y, r) \in \mathcal{L}. \end{aligned}$$

Thus, (3.59) holds.

In turn, from (3.62) using (2.11), we conclude that

$$\begin{aligned} u(x, q) &= \inf_{(y, r) \in \mathbb{R} \times \mathbb{L}^1(dK)} \left(v(y, r) + xy + \int_0^T q_s r_s dK_s \right), \\ &= \inf_{(y, r) \in \mathcal{L}} \left(v(y, r) + xy + \int_0^T q_s r_s dK_s \right), \quad (x, q) \in \mathcal{K}, \end{aligned}$$

which proves (3.60) and extends the assertion of Lemma 3.21 to the boundary of $\mathring{\mathcal{K}}$. \square

PROOF OF THEOREM 3.1. The assertions of item (i) follow from Lemmas 3.15 and 3.17, item (ii) results from Lemma 3.18, whereas the validity of item (iii) come from Lemma 3.24. This completes the proof of the theorem. \square

4. Subdifferentiability of u . In order to establish subdifferentiability of u , we need to strengthen Assumption 2.5 and to impose the following condition.

ASSUMPTION 4.1. There exists an a.s. bounded away from 0 and ∞ process φ , such that

$$d\kappa(\omega) = \varphi dK \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

Let

$$(4.1) \quad \mathcal{P} := \{\rho : (1, \rho) \in \mathcal{L}\},$$

REMARK 4.2. \mathcal{P} defined in (4.1) needs to be uniformly integrable with respect to the measure dK in order for the proof of subdifferentiability of u to go through. Assumption 2.5 through Lemma 3.2 only implies that \mathcal{P} is \mathbb{L}^1 bounded. A stronger condition on the stochastic clock and income stream is therefore needed to obtain uniform integrability.

The following theorem characterizes subdifferentiability of u over $\mathring{\mathcal{K}}$, where we are looking for an $\mathbb{R} \times \mathbb{L}^1(dK)$ -valued subgradient. Under our assumptions, we can find elements of the subgradient which both belong to the effective domain of v , \mathcal{E} , and are bounded, that is, in $\mathbb{R} \times \mathbb{L}^\infty(dK)$.

THEOREM 4.3. *Let the conditions of Theorem 3.1 and Assumption 4.1 hold. Then for every $(x, q) \in \mathring{\mathcal{K}}$, the subdifferential of u at (x, q) is a nonempty and contains an element of \mathcal{E} , that is,*

$$(4.2) \quad \partial u(x, q) \cap \mathcal{E} \neq \emptyset.$$

Moreover, for $(x, q) \in \mathring{\mathcal{K}}$ and $(y, r) \in \mathcal{L}$, $(y, r) \in \partial u(x, q)$ if and only if the following conditions hold:

$$(4.3) \quad |v(y, r)| < \infty,$$

thus, $(y, r) \in \mathcal{E}$,

$$(4.4) \quad \mathbb{E}\left[\int_0^T \widehat{Y}_t(y, r) \widehat{c}_t(x, q) d\kappa_t\right] = xy + \int_0^T q_s r_s dK_s,$$

$$(4.5) \quad \widehat{Y}_t(y, r) = U'(t, \widehat{c}_t(x, q)), \quad (d\kappa \times \mathbb{P})\text{-a.e.},$$

where $\widehat{c}(x, q)$ and $\widehat{Y}(y, r)$ are the unique optimizers to (2.4) and (2.10), respectively.

4.1. Uniform integrability of \mathcal{P} .

LEMMA 4.4. *Let the conditions of Theorem 4.3 hold. Then \mathcal{P} is $\mathbb{L}^\infty(dK)$ -bounded and, therefore, a uniformly integrable family.*

PROOF. *Step 1.* For an arbitrary $q : [0, T] \rightarrow [0, 1]$, let us define

$$\beta(q) := \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}\left[\int_0^T q_s |e_s| d\kappa_s\right] = \sup_{\mathbb{Q} \in \mathcal{M}, \tau \in \Theta} \mathbb{E}^{\mathbb{Q}}\left[\int_0^\tau q_s |e_s| d\kappa_s\right].$$

Note that by Assumption 2.5, we have

$$\begin{aligned} \beta(q) &= \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}\left[\int_0^T q_s |e_s| d\kappa_s\right] \\ &\leq \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}\left[\int_0^T q_s X'_s \varphi dK_s\right] \\ &\leq \int_0^T q_s \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}}[CX'_s] dK_s \\ &\leq CX'_0 \int_0^T q_s dK_s, \end{aligned}$$

where $C \in \mathbb{R}$ is such that $|\varphi| \leq C$. It follows from [27], Proposition 4.3, and [17], Theorem 3.1, that there exists a nonnegative càdlàg process $V = \beta(q) + H \cdot S$, such that

$$V_t \geq \operatorname{ess\,sup}_{\mathbb{Q} \in \mathcal{M}, \tau \in \Theta, \tau \geq t} \mathbb{E}^{\mathbb{Q}} \left[\int_0^\tau q_s |e_s| d\kappa_s \mid \mathcal{F}_t \right], \quad t \in [0, T].$$

Consequently, V satisfies

$$V_\tau \geq \int_0^\tau q_s |e_s| d\kappa_s \geq \int_0^\tau q_s e_s d\kappa_s, \quad \mathbb{P}\text{-a.s.}, \tau \in \Theta,$$

and thus, $(\beta(q), -q) \in \mathcal{K}$. As a result, from the definitions of \mathcal{L} and \mathcal{P} , we get

$$\beta(q) \geq \sup_{\rho \in \mathcal{P}} \int_0^T q_s \rho_s dK_s.$$

One can see that for every $\rho \in \mathcal{P}$, we have $\rho \leq f := CX'_0, dK$ -a.e.

Step 2. For an arbitrary $q : [0, T] \rightarrow [-1, 0]$, let us set

$$\tilde{\beta}(q) := \sup_{\mathbb{Q} \in \mathcal{M}} \mathbb{E}^{\mathbb{Q}} \left[\int_0^T q_s (-|e_s|) d\kappa_s \right].$$

As in Step 1, we can construct a càdlàg process $\tilde{V} = \tilde{\beta}(q) + H \cdot S$, s.t.

$$\begin{aligned} \tilde{V}_t &\geq \operatorname{ess\,sup}_{\mathbb{Q} \in \mathcal{M}, \tau \in \Theta, \tau \geq t} \mathbb{E}^{\mathbb{Q}} \left[\int_0^T q_s (-|e_s|) d\kappa_s \mid \mathcal{F}_t \right] \\ &\geq \operatorname{ess\,sup}_{\mathbb{Q} \in \mathcal{M}, \tau \in \Theta, \tau \geq t} \mathbb{E}^{\mathbb{Q}} \left[\int_0^T q_s e_s d\kappa_s \mid \mathcal{F}_t \right]. \end{aligned}$$

This implies that $(\tilde{\beta}(q), -q) \in \operatorname{cl} \mathcal{K}$. Therefore,

$$\beta(q) \geq \int_0^T q_s \rho_s dK_s \quad \text{for every } \rho \in \mathcal{P}.$$

Similar to Step 1, one can see that $\rho \geq -f, dK$ -a.e. for every $\rho \in \mathcal{P}$.

Step 3. In view of Steps 1 and 2, uniform integrability of \mathcal{P} under dK follows from the integrability of f under dK . \square

LEMMA 4.5. *Let the assumption of Theorem 4.3 hold and $(x, q) \in \mathring{\mathcal{K}}$. Then*

$$\{r : (y, r) \in \mathcal{B}(x, q)\}$$

is \mathbb{L}^∞ -bounded, so a uniformly integrable subset of $\mathbb{L}^1(dK)$.

PROOF. By Lemma 3.13, we deduce the existence of a constant $M > 0$, such that

$$y \leq M \quad \text{for every } (y, r) \in \mathcal{B}(x, q).$$

We conclude that

$$\{r : (y, r) \in \mathcal{B}(x, q)\} \subseteq \bigcup_{0 \leq \lambda \leq M} \lambda \mathcal{P},$$

and thus by Lemma 4.4, $\{r : (y, r) \in \mathcal{B}(x, q)\}$ is a uniformly integrable family. \square

4.2. *Closedness of $\mathcal{D}(x, q)$ for every $(x, q) \in \mathring{\mathcal{K}}$.*

LEMMA 4.6. *Under the conditions of Theorem 4.3, for every $(x, q) \in \mathring{\mathcal{K}}$, the set $\mathcal{D}(x, q)$ is closed in $\mathbb{L}^0(d\kappa \times \mathbb{P})$.*

PROOF. Let $(x, q) \in \mathring{\mathcal{K}}$ and Y be an arbitrary element of $\text{cl } \mathcal{D}(x, q)$. We claim that there exists $(y, r) \in \mathcal{B}(x, q)$, such that $Y \in \mathcal{Y}(y, r)$. Let $Y^n \in \mathcal{Y}(y^n, r^n)$, $n \geq 1$, be a sequence in $\mathcal{D}(x, q)$, such that $\lim_{n \rightarrow \infty} Y^n = Y$, $(d\kappa \times \mathbb{P})$ -a.e. Since $(y^n, r^n)_{n \geq 1} \subset \mathcal{B}(x, q)$, which is bounded in the sense of Lemma 3.13, Komlos' lemma implies the existence of a subsequence of convex combinations $(\tilde{y}^n, \tilde{r}^n) \in \text{conv}((y^n, r^n), (y^{n+1}, r^{n+1}), \dots)$, $n \geq 1$, such that $(\tilde{y}^n)_{n \geq 1}$ converges to y and $(\tilde{r}^n)_{n \geq 1}$ converges to r , dK -a.e. Lemma 4.5 implies that $(\tilde{r}^n)_{n \geq 1}$ is uniformly integrable. Therefore, $(\tilde{r}^n)_{n \geq 1}$ converges to r in $\mathbb{L}^1(dK)$. Note that the corresponding sequence of convex combinations of $(Y^n)_{n \geq 1}$, $(\tilde{Y}^n)_{n \geq 1}$ converges to Y , $(d\kappa \times \mathbb{P})$ -a.e. Then we have

$$1 \geq \lim_{n \rightarrow \infty} \left(x \tilde{y}^n + \int_0^T q_s \tilde{r}_s^n dK_s \right) = xy + \int_0^T q_s r_s dK_s.$$

Therefore, $(y, r) \in \mathcal{B}(x, q)$.

Let us fix an arbitrary $(x', q') \in \mathcal{K}$ and $c \in \mathcal{A}(x', q')$. Using Proposition 3.4, Fatou's lemma and Lemma 4.5, we obtain

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\int_0^T Y_s c_s ds \right] \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T \tilde{Y}_s^n c_s ds \right] \\ &\leq \lim_{n \rightarrow \infty} \left(x' \tilde{y}^n + \int_0^T q'_s \tilde{r}_s^n dK_s \right) = x'y + \int_0^T q'_s r_s dK_s, \end{aligned}$$

where the uniform integrability of $\mathcal{B}(x, q)$ is needed once again in the last equality. From Proposition 3.4, we conclude that that $Y \in \mathcal{Y}(y, r)$. \square

PROOF OF THEOREM 4.3. Let $(x, q) \in \mathcal{K}$, $\hat{c}(x, q)$ be the minimizer to (2.4), whose existence and uniqueness are established in Lemma 3.18. Let us also set

$$\begin{aligned} \hat{Y}_t &:= U'(t, \hat{c}_t(x, q)), \quad (t, \omega) \in [0, T] \times \Omega \quad \text{and} \\ (4.6) \quad z &:= \mathbb{E} \left[\int_0^T \hat{c}_s(x, q) \hat{Y}_s d\kappa_s \right]. \end{aligned}$$

Note that the sets $\mathcal{A}(x, q)$ and $\text{cl } \mathcal{D}(x, q)$ satisfy the conditions of [31], Theorem 3.2, which implies that $\hat{Y} \in \text{zcl } \mathcal{D}(x, q)$ is the unique solution to the optimization problem

$$\inf_{Y \in \text{zcl } \mathcal{D}(x, q)} \mathbb{E} \left[\int_0^T V(t, Y_s) d\kappa_s \right] = \mathbb{E} \left[\int_0^T V(t, \hat{Y}_s) d\kappa_s \right] \in \mathbb{R},$$

where finiteness follows from Lemma 3.16. Note that by Lemma 4.6, $\hat{Y} \in \mathcal{Y}(zy, zr)$ for some $(y, r) \in \mathcal{B}(x, q)$. It follows from the definition of \hat{Y} in (4.6) that for $\hat{c}(x, q)$ and \hat{Y} we have the following relation:

$$U(t, \hat{c}_t(x, q)) = V(t, \hat{Y}_t) + \hat{c}_t(x, q) \hat{Y}_t, \quad (t, \omega) \in [0, T] \times \Omega,$$

which together with Lemma 3.21 implies that

$$\begin{aligned} (4.7) \quad u(x, q) &= v(zy, zr) + z \left(xy + \int_0^T q_s r_s dK_s \right), \\ \hat{Y}(zy, zr) &= \hat{Y}, \quad (d\kappa \times \mathbb{P})\text{-a.e.}, \end{aligned}$$

where $\widehat{Y}(zy, zr)$ is the unique minimizer to the dual problem (2.10). By (4.7), $(zy, zr) \in \mathcal{E}$. Equation (4.7) and [15], Proposition I.5.1, assert that $(zy, zr) \in \partial u(x, q)$. In particular, we get

$$\partial u(x, q) \cap \mathcal{E} \neq \emptyset,$$

that is, (4.2). Note that even though, for example, [2], Corollary 2.2.38 and Corollary 2.2.44, imply that $\partial u(x, q) \neq \emptyset$, over the interior of the effective domain, its elements are in $\mathbb{R} \times (\mathbb{L}^\infty)^*(dK)$. Relation (4.2) shows that $\partial u(x, q)$ contains at least a bounded element of $\mathcal{E} \subseteq \mathcal{L} \subset \mathbb{R} \times \mathbb{L}^1(dK)$.

Let $(x, q) \in \mathcal{K}$ and $(y, r) \in \mathcal{L}$. Suppose that (4.3), (4.4) and (4.5) hold. Then by conjugacy of U and V , we get

$$(4.8) \quad 0 = v(y, r) - u(x, q) + xy + \int_0^T q_s r_s dK_s.$$

Lemma 3.24 and [15], Proposition I.5.1, imply that $(y, r) \in \partial u(x, q) \cap \mathcal{E}$. Conversely, let $(x, q) \in \mathcal{K}$ and $(y, r) \in \mathcal{L} \cap \partial u(x, q)$. Then by [15], Proposition I.5.1, and Lemma 3.24, we deduce that (4.8) holds. Lemma 3.15 implies the finiteness of $u(x, q)$, which together with (4.8) results in the finiteness of $v(y, r)$, thus (4.3) holds and $(y, r) \in \mathcal{E}$. By Lemma 3.18, there exists a unique optimizer $\widehat{c}(x, q)$, for (2.4), and $\widehat{Y}(y, r)$, for (2.10), respectively. Therefore, from conjugacy of U and V , Proposition 3.4, and (4.8), we obtain

$$\begin{aligned} 0 &\leq \mathbb{E} \left[\int_0^T (V(t, \widehat{Y}_t(y, r)) - U(t, \widehat{c}_t(x, q)) + \widehat{Y}_t(y, r) \widehat{c}_t(x, q)) dk_t \right] \\ &\leq v(y, r) - u(x, q) + xy + \int_0^T q_s r_s dK_s = 0. \end{aligned}$$

This implies (4.4) and (4.5). This completes the proof of the theorem. \square

5. Structure of the dual feasible set. By Assumption 4.1, there exists at most countable subset $(s_k)_{k \in \mathbb{N}}$ of $[0, T]$, where κ has jumps. We define \mathcal{D}' the set of nonincreasing, left-continuous and adapted processes D that start at 1 and with the property that $D_{\theta_0} = 1, D_T \geq 0$ and that there exists some $n \in \mathbb{N}$ such that D is constant off the discrete grid $\mathcal{T}_n := \bigcup_{j=1}^{2^n} \{s_j\} \cup \{\frac{k}{2^n}T, k = 0, \dots, 2^n\}$.

LEMMA 5.1. *Let $\mathcal{G}' := \{ZD = (Z_t D_t)_{t \in [0, T]} : Z \in \mathcal{Z}', D \in \mathcal{D}'\}$. Assume the conditions of Proposition 3.4 hold. Then \mathcal{G}' is convex.*

PROOF. Let $Z^1 D^1$ and $Z^2 D^2$ be the elements of \mathcal{G}' and let $\lambda \in (0, 1)$ We need to show that $\lambda Z^1 D^1 + (1 - \lambda) Z^2 D^2 = ZD$ for some $Z \in \mathcal{Z}'$ and $D \in \mathcal{D}'$. There exists $n \in \mathbb{N}$, such that D^1 and D^2 decrease at most on \mathcal{T}_n . Let t_k 's be the elements of \mathcal{T}_n arranged in an increasing order. Let us define $Z_0 = D_0 = 1$ and for every $k \in \{0, \dots, 2^n - 1\}$, with

$$\begin{aligned} A_t &:= \lambda Z_{t_k}^1 D_t^1 + (1 - \lambda) Z_{t_k}^2 D_t^2, \\ \alpha_t &:= \frac{\lambda Z_{t_k}^1 D_t^1}{A_t} 1_{\{A_t \neq 0\}} + \frac{\lambda Z_{t_k}^1}{\lambda Z_{t_k}^1 + (1 - \lambda) Z_{t_k}^2} 1_{\{A_t = 0\}} \end{aligned}$$

(note that $A_t = 0$ if and only if both $D_t^1 = 0$ and $D_t^2 = 0$) we set

$$(5.1) \quad \begin{aligned} Z_t &:= Z_{t_k} \left(\alpha_{t_k+} \frac{Z_t^1}{Z_{t_k}^1} + (1 - \alpha_{t_k+}) \frac{Z_t^2}{Z_{t_k}^2} \right) \quad \text{for } t \in (t_k, t_{k+1}], \\ D_t &:= \frac{A_{t_k+}}{Z_{t_k}} \quad \text{for } t \in (t_k, t_{k+1}]. \end{aligned}$$

One can see that $ZD = \lambda Z^1 D^1 + (1 - \lambda)Z^2 D^2$ and that $Z \in \mathcal{Z}'$; see, for example, [17, 25, 35] and [5] for discussions of the sets of processes with similar convexity-type properties to the one given in (5.1). To show that $D \in \mathcal{D}'$, for $k \geq 1$, we observe that

$$\begin{aligned} D_{t_k+} &= \frac{A_{t_k+}}{Z_{t_k}} \\ &= \frac{\lambda Z_{t_k}^1 D_{t_k+}^1 + (1 - \lambda)Z_{t_k}^2 D_{t_k+}^2}{Z_{t_{k-1}}(\alpha_{t_{k-1}+} \frac{Z_{t_k}^1}{Z_{t_{k-1}}^1} + (1 - \alpha_{t_{k-1}+}) \frac{Z_{t_k}^2}{Z_{t_{k-1}}^2})} \\ &\leq \frac{\lambda Z_{t_k}^1 D_{t_k}^1 + (1 - \lambda)Z_{t_k}^2 D_{t_k}^2}{Z_{t_{k-1}}(\alpha_{t_{k-1}+} \frac{Z_{t_k}^1}{Z_{t_{k-1}}^1} + (1 - \alpha_{t_{k-1}+}) \frac{Z_{t_k}^2}{Z_{t_{k-1}}^2})} \\ &= \frac{\lambda Z_{t_k}^1 D_{t_k}^1 + (1 - \lambda)Z_{t_k}^2 D_{t_k}^2}{Z_{t_{k-1}}(\frac{\lambda Z_{t_{k-1}}^1 D_{t_k}^1}{A_{t_{k-1}+} Z_{t_{k-1}}^1} + \frac{(1-\lambda)Z_{t_{k-1}}^2 D_{t_k}^2}{A_{t_{k-1}+} Z_{t_{k-1}}^2})} \mathbf{1}_{\{A_{t_{k-1}+}>0\}} \\ &\quad + 0 \cdot \mathbf{1}_{\{A_{t_{k-1}+}=0\}} \\ &= \frac{A_{t_{k-1}+} \lambda Z_{t_k}^1 D_{t_k}^1 + (1 - \lambda)Z_{t_k}^2 D_{t_k}^2}{Z_{t_{k-1}} \lambda D_{t_k}^1 Z_{t_k}^1 + (1 - \lambda)D_{t_k}^2 Z_{t_k}^2} \mathbf{1}_{\{A_{t_{k-1}+}>0\}} \\ &= \frac{A_{t_{k-1}+}}{Z_{t_{k-1}}} \mathbf{1}_{\{A_{t_{k-1}+}>0\}} \\ &= D_{t_k}. \end{aligned}$$

Above, the inequality follows from the monotonicity of D^1 and D^2 . Therefore, D is nonincreasing. Also, clearly D is nonnegative. Thus $ZD \in \mathcal{G}'$. \square

The following lemma is an extension of Lemma 3.10 and amounts to a first layer of convexification of the set Υ , that is, of the budget constraints.

LEMMA 5.2. *Let the conditions of Proposition 3.4 hold, $(x, q) \in \mathcal{K}$, and c is a nonnegative optional process. Then $c \in \mathcal{A}(x, q)$ if and only if*

$$(5.2) \quad \mathbb{E} \left[\int_0^T c_s D_s Z_s d\kappa_s \right] \leq x + \mathbb{E} \left[\int_0^T q_s e_s D_s Z_s d\kappa_s \right]$$

for every $Z \in \mathcal{Z}'$ and $D \in \mathcal{D}'$.

PROOF. The idea is to use the assertion of Lemma 3.10 and to approximate a given $D \in \mathcal{D}'$ by (finite) linear combinations of the elements of Υ , where Υ is defined in (3.6).

For a stopping time τ , let us denote

$$\Lambda^\tau := \mathbf{1}_{[0, \tau]} \in \Upsilon,$$

and fix $D \in \mathcal{D}'$. Then there exists $l \in \mathbb{N}$, such that D has jumps at most on $\{t_0, t_1, \dots, t_l\}$ for some increasing t_i 's. For every $j \in \{0, \dots, l\}$, $k \in \{0, \dots, 2^n\}$, and $n \in \mathbb{N}$, let us set

$$A_{k,n,j} := \left\{ \omega : D_{t_j}(\omega) > 0 \text{ and } \frac{D_{t_j+}(\omega)}{D_{t_j}(\omega)} \in \left(\frac{k-1}{2^n}, \frac{k}{2^n} \right] \right\},$$

$$\tau^{k,n,j} := T \mathbf{1}_{A_{k,n,j}} + t_j \mathbf{1}_{A_{k,n,j}^c}.$$

Note that $D_0 = 1$ by definition of \mathcal{D}' , $A_{k,n,j} \in \mathcal{F}_{t_j}$, and

$$\frac{k-1}{2^n} \Lambda_{t_j^+}^{\tau^{k,n,j}} = \begin{cases} \frac{k-1}{2^n} & \text{on } A_{k,n,j}, \\ 0 & \text{on } A_{k,n,j}^c. \end{cases}$$

By construction, we have

$$D_{t_j^+} = D_{t_j} \lim_{n \rightarrow \infty} \sum_{k=1}^{2^n} \frac{k-1}{2^n} \Lambda_{t_j^+}^{\tau^{k,n,j}},$$

where the sequence

$$K_{t_j^+}^n := \sum_{k=1}^{2^n} \frac{k-1}{2^n} \Lambda_{t_j^+}^{k,n,j}, \quad n \in \mathbb{N},$$

is increasing on $\{D_{t_j} > 0\}$, that is,

$$K_{t_j^+}^n 1_{\{D_{t_j} > 0\}} \uparrow \frac{D_{t_j^+}}{D_{t_j}} 1_{\{D_{t_j} > 0\}}.$$

Thus, for an arbitrary $j \in \{0, \dots, l\}$, we have constructed a sequence of elements of Υ , whose finite linear combinations monotonically increase to $\frac{D_{t_j^+}}{D_{t_j}}$ on $\{D_{t_j} > 0\}$ (i.e., if $j < l$, we have approximated D on the interval (t_j, t_{j+1})).

In order to construct a sequence that approximates D at every point of its potential jumps, we first observe that for two stopping times τ and σ , we have

$$\Lambda^\tau \Lambda^\sigma = 1_{[0,\tau]} 1_{[0,\sigma]} = 1_{[0,\tau \wedge \sigma]} = \Lambda^{\tau \wedge \sigma}.$$

Therefore, for every $n \in \mathbb{N}$,

$$(5.3) \quad K_{t_j^+}^n K_{t_{j+1}^+}^n = \left(\sum_{k=1}^{2^n} \frac{k-1}{2^n} \Lambda_{t_j^+}^{k,n,j} \right) \left(\sum_{k=1}^{2^n} \frac{k-1}{2^n} \Lambda_{t_{j+1}^+}^{k,n,j+1} \right) = \sum_{i=1}^{4^n} \lambda^{n,i} \Lambda_{t_{j+1}^+}^{\sigma_{n,i}},$$

for some finite sequences of stopping times $(\sigma_{n,i})_{i=1}^{4^n}$ and $[0, 1]$ -valued constants $(\lambda^{n,i})_{i=1}^{4^n}$. Here, $\Lambda^{\sigma_{n,i}}$ are such that for both $t = t_j$ and $t = t_{j+1}$ on $\{D_{t_{j+1}} > 0\}$, we have

$$\lim_{n \rightarrow \infty} \sum_{i=1}^{4^n} \lambda^{n,i} \Lambda_{t^+}^{\sigma_{n,i}} = \frac{D_{t^+}}{D_t}.$$

Similarly, with $r(t) := \max\{i : t_i < t\}$, let us define

$$D_t^n := \prod_{j=0}^{r(t)} K_{t_j^+}^n 1_{\{D_{t_j} > 0\}}, \quad t \in [0, T], n \in \mathbb{N}.$$

As in (5.3), for every $n \in \mathbb{N}$, D^n can be written as a finite linear combination of Λ 's, such that $D_{t^+}^n \uparrow D_{t^+}$ for every $t \in \{t_0, \dots, t_l\}$.

Finally, (5.2) can be obtained from Lemma 3.10 by the approximation of D by D^n 's as above and via the monotone convergence theorem (applied separately to $(e_t)^+$ and $(e_t)^-$). \square

COROLLARY 5.3. *Let the conditions of Proposition 3.4 hold. Then, for every pair $Z \in \mathcal{Z}'$ and $D \in \mathcal{D}'$, there exists $r^{ZD} \in \mathbb{L}^1(dK)$, such that $ZD \in \mathcal{Y}(1, r^{ZD})$, where $(1, r^{ZD}) \in \mathcal{L}$.*

PROOF. The existence of r^{ZD} , such that $ZD \in \mathcal{Y}(1, r^{ZD})$ follows from Lemma 5.2 (equation (5.2)) and the approximation procedure in Lemma 5.2 (again, applied separately to $(e_t)^+$ and $(e_t)^-$) combined with the monotone convergence theorem. As, the left-hand side in (5.2) is nonnegative, $(1, r^{ZD}) \in \mathcal{L}$. \square

For a given $Z \in \mathcal{Z}'$ and $D \in \mathcal{D}'$, let us recall that r^{ZD} is given in Corollary 5.3. We set

$$\begin{aligned} \mathcal{B}'(x, q) &:= \{(y, yr^{ZD}) \in \mathcal{B}(x, q) : y > 0, Z \in \mathcal{Z}', D \in \mathcal{D}'\}, \\ \mathcal{G}(x, q) &:= \{yZD \in \mathcal{D}(x, q) : (y, yr^{ZD}) \in \mathcal{B}'(x, q)\}, \quad (x, q) \in \mathring{\mathcal{K}}, \end{aligned}$$

LEMMA 5.4. *Let the conditions of Proposition 3.4 hold, then for every $(x, q) \in \mathring{\mathcal{K}}$, the closure of the convex, solid hull of $\mathcal{G}(x, q)$ in \mathbb{L}^0 coincides with $\text{cl } \mathcal{D}(x, q)$.*

PROOF. Let $(x, q) \in \mathring{\mathcal{K}}$ be fixed. Along the lines of the proof of Lemma 3.20, one can show that

$$(\mathcal{G}(x, q))^o = \mathcal{A}(x, q).$$

Therefore,

$$(\mathcal{G}(x, q))^{oo} = (\mathcal{A}(x, q))^o = \text{cl } \mathcal{D}(x, q),$$

where in the last equality we have used the conclusion of Lemma 3.20. As $\mathcal{G}(x, q) \subset \text{cl } \mathcal{D}(x, q)$, the assertion of the lemma follows from the bipolar theorem of Brannath and Schermayer, [3], Theorem 1.3. \square

COROLLARY 5.5. *Let the conditions of Proposition 3.4 hold and $(x, q) \in \mathring{\mathcal{K}}$. Then for every maximal element Y of $\text{cl } \mathcal{D}(x, q)$ there exists $y^n \geq 0, Z^n \in \mathcal{Z}', D^n \in \mathcal{D}', n \in \mathbb{N}$, such that $(y^n Z^n D^n)_{n \in \mathbb{N}} \subset \mathcal{G}(x, q)$ and*

$$Y = \lim_{n \rightarrow \infty} y^n Z^n D^n, \quad (d\kappa \times \mathbb{P})\text{-a.e. and on } \bigcup_{n \in \mathbb{N}} \mathcal{T}_n.$$

PROOF. The $(d\kappa \times \mathbb{P})$ -a.e. convergence follows from Lemma 5.4. By passing to subsequences of convex combinations, we also deduce the convergence on $\bigcup_{n \in \mathbb{N}} \mathcal{T}_n$. \square

REMARK 5.6. It follows from Corollary 5.5 and Fatou’s lemma that the maximal elements of $\text{cl } \mathcal{D}(x, q)$ are strong supermartingales. Moreover, every maximal element of $\text{cl } \mathcal{D}(x, q)$ is an optional strong supermartingale deflator, which is optional strong supermartingale Y , such that XY is an optional strong supermartingale for every $X \in \mathcal{X}(1)$. We refer to [12], Appendix 1, for a general characterization and to [8] for results on strong, optional supermartingales as limits of martingales. The following section gives a more refined characterization of the dual minimizer.

6. Complementary slackness. For better readability of this section, we recall some notation and results that will be used below. Throughout this section, $(x, q) \in \mathring{\mathcal{K}}$ will be fixed, $\hat{c} = \hat{c}(x, q)$ is the optimizer to (2.4), \hat{V} is the corresponding wealth process, that is,

$$(6.1) \quad \hat{V} = x + \int_0^\cdot \hat{H}_s dS_s - \int_0^\cdot \hat{c}_s d\kappa_s + \int_0^\cdot q_s e_s d\kappa_s,$$

where \hat{H} is some S -integrable process, \hat{Y} be such that $\hat{Y}_t = U'(t, \hat{c}_t)$, $(d\kappa \times \mathbb{P})$ -a.e., that is. We recall that \hat{Y} is the optimizer to (2.10) for some $(y, r) \in \mathcal{E} \cap \partial u(x, q)$, that is, $\hat{Y} = \hat{Y}(y, r) \in \mathcal{Y}(y, r)$. We also denote, similar to the proof of equation (4.3) by

$$z = \mathbb{E} \left[\int_0^T \hat{c}_t \hat{Y}_t d\kappa_t \right].$$

From Corollary 5.5, we know that \widehat{Y} can be approximated by a sequence $y^n D^n Z^n$, $n \in \mathbb{N}$, where y^n is a nonnegative constant, $D^n \in \mathcal{D}'$, and $Z^n \in \mathcal{Z}'$, $n \in \mathbb{N}$, such that

$$(6.2) \quad y^n D^n Z^n \in \mathcal{Y}(y^n, r^n), \quad xy^n + \int_0^T q_s r_s^n dK_s \leq z.$$

Above, r^n is given by Corollary 5.3 (which has a similar interpretation to Lemma 3.7) by

$$\int_0^t r_s^n dK_s = y^n \mathbb{E} \left[\int_0^t Z_s^n D_s^n e_s d\kappa_s \right], \quad t \in [0, T].$$

We would like to point out that, unlike in Lemma 3.7 or Corollary 5.3, here, the value of y^n has to be taken into account. In what follows, for any right-continuous increasing process A satisfying

$$A_t = 0, \quad 0- \leq t \leq \theta_0-, \quad A_T = 1,$$

that is, for any probability measure dA on the closed interval $[\theta_0, T]$ (extended to $[0, T]$) we will associate a process D which is left continuous and decreasing

$$D_t := 1 - A_{t-} = dA([t, T]), \quad 0 \leq t \leq T.$$

One can also think that $D_{T+} = 0$, although this is not necessary. It is clear that such $A \leftrightarrow D$ are in bijective correspondence. Below, all processes A 's and D 's (with indexes) will be in such bijective correspondence, except for the case of the limiting process \widehat{A} (which is right continuous) and the limiting process \widehat{D} (that may be not left continuous). This will be in a similar but more subtle correspondence. More precisely, we have the following.

THEOREM 6.1. *Let the conditions of Theorem 4.3 hold and fix $(x, q) \in \mathring{\mathcal{K}}$. Let \widehat{V} be the optimal wealth process, \widehat{c} the optimal consumption and $\widehat{Y}_t = U'(t, \widehat{c}_t)$, $(d\kappa \times \mathbb{P})$ -a.e. Then there exists an $(y, r) \in \partial u(x, q)$ such that $\widehat{Y} = \widehat{Y}(y, r)$, that is, $\widehat{Y} \in \mathcal{Y}(y, r)$,*

$$\mathbb{E} \left[\int_0^T V(t, \widehat{Y}_t) d\kappa_t \right] = \inf_{Y \in \mathcal{Y}(y, r)} \mathbb{E} \left[\int_0^T V(t, Y_t) d\kappa_t \right],$$

such that \widehat{Y} can be decomposed as

$$\widehat{Y} = y \widehat{Z} \widehat{D},$$

where \widehat{Z} is a strong supermartingale $\widehat{Z} = \lim_{n \rightarrow \infty} Z^n$ (for $Z_n \in \mathcal{D}'$, where the limit is in the sense of [8]) and a right-continuous increasing process \widehat{A} with

$$\widehat{A}_t = 0 \quad \forall 0- \leq t \leq \theta_0-, \quad \widehat{A}_T = 1,$$

and a decreasing process \widehat{D} with $\widehat{D}_t = 1$, $0 \leq t \leq \theta_0$ and satisfying the complementary slackness condition

$$(6.3) \quad \mathbb{P}(\widehat{D}_t \in [1 - \widehat{A}_t, 1 - \widehat{A}_{t-}], \forall \theta_0 \leq t \leq T) = 1, \\ \mathbb{P} \left(\int_{[\theta_0, T]} 1_{\{\widehat{V}_{t-} \neq 0, \widehat{V}_t \neq 0\}} d\widehat{A}_t \right) = 0.$$

The proof of the Theorem 6.1 is aligned in several results.

LEMMA 6.2. *Let the conditions of Theorem 4.3 hold. With*

$$z := \mathbb{E} \left[\int_0^T \widehat{c}_s \widehat{Y}_s d\kappa_s \right],$$

there exist $y^n > 0$, $Z^n \in \mathcal{Z}'$ and $D^n \in \mathcal{D}'$, such that $y^n Z^n D^n \in z\mathcal{G}(x, q)$, $n \in \mathbb{N}$ and

$$(6.4) \quad \begin{aligned} \widehat{Y} &= \lim_{n \rightarrow \infty} y^n Z^n D^n, \quad (d\kappa \times \mathbb{P})\text{-a.e.}, \\ (y^n, r^n) &\rightarrow (y, r) \in \mathcal{L}, \quad y > 0, \end{aligned}$$

where the last convergence is weak in L^1 and

$$z = xy + \int_0^T q_s r_s dK_s, \quad (y, r) \in \partial u(x, q), \quad \widehat{Y} = \widehat{Y}(y, r).$$

For A^n , $n \in \mathbb{N}$, being in relation to D^n exactly as described before Theorem 6.1 we have

$$(6.5) \quad \mathbb{E} \left[\int_{\theta_0}^T \widehat{V}_t Z_t^n dA_t^n \right] = \mathbb{E} \left[Z_T^n \int_{\theta_0}^T \widehat{V}_t dA_t^n \right] \rightarrow 0.$$

PROOF. Optimality of \widehat{Y} and Corollary 5.5 imply the first convergence in (6.4). According to Lemma 4.5, the sequence $(r^n)_{n \in \mathbb{N}}$ is uniformly integrable, so it is relatively compact in the weak topology of $L^1(dK)$, by the Dunford–Pettis theorem. According to the Eberlein–Smulian theorem, it has a further convergent subsequence. As an immediate consequence of the same Lemma 4.5 and the very definition of $\mathcal{B}(x, q)$, we also obtain that the sequence $(y^n)_{n \in \mathbb{N}}$ is bounded. Altogether, we have a further subsequence such that $(y^n, r^n) \rightarrow (y, r)$ where the convergence in the second component is in the weak L^1 -sense. Now, using almost identical arguments as the proof of Theorem 4.3 and Lemma 4.6 from (6.2) we obtain the whole conclusion of the lemma, except for the relation (6.5) that we prove below. By (4.4) and Fatou’s lemma, we get

$$(6.6) \quad z = \mathbb{E} \left[\int_0^T \widehat{Y}_s \widehat{c}_s d\kappa_s \right] \leq \liminf_{n \rightarrow \infty} y^n \mathbb{E} \left[\int_0^T Z_s^n D_s^n \widehat{c}_s d\kappa_s \right].$$

Let us fix $n \in \mathbb{N}$ and consider $\mathbb{E}[\int_0^T Z_s^n D_s^n \widehat{c}_s d\kappa_s]$. Using localization and integration by parts (along the lines of the proof of Lemma 3.7), we have

$$\begin{aligned} \mathbb{E} \left[\int_0^T Z_s^n D_s^n \widehat{c}_s d\kappa_s \right] &= \mathbb{E} \left[Z_T^n \int_0^T D_s^n \widehat{c}_s d\kappa_s \right] \\ &= \mathbb{E} \left[Z_T^n \int_0^T (1 - A_{s-}^n) \widehat{c}_s d\kappa_s \right] \\ &= \mathbb{E} \left[Z_T^n \left(\int_0^T \widehat{c}_s d\kappa_s - \int_0^T A_{s-}^n \widehat{c}_s d\kappa_s \right) \right] \\ &= \mathbb{E} \left[Z_T^n \left(\int_0^T \widehat{c}_s d\kappa_s - A_T^n \int_0^T \widehat{c}_s d\kappa_s \right. \right. \\ &\quad \left. \left. + \int_0^T \left(\int_0^t \widehat{c}_s d\kappa_s \right) dA_t^n \right) \right] \\ &= \mathbb{E} \left[Z_T^n \int_0^T \left(\int_0^t \widehat{c}_s d\kappa_s \right) dA_t^n \right]. \end{aligned}$$

The latter expression, using (6.1) and with $\bar{X} := \|q\|_{\mathbb{L}^\infty(dK)} X''$ (where in turn X'' is given by the assertion (v) of Lemma 3.2), we can rewrite as

$$(6.7) \quad \mathbb{E} \left[Z_T^n \int_0^T \left(\left(x + \int_0^t \widehat{H}_s dS_s + \bar{X}_t \right) + \left(\int_0^t q_s e_s d\kappa_s - \bar{X}_t \right) - \widehat{V}_t \right) dA_t^n \right].$$

Let us denote

$$T_1 := \mathbb{E} \left[Z_T^n \int_0^T \left(x + \int_0^t \widehat{H}_s dS_s + \bar{X}_t \right) dA_t^n \right],$$

$$T_2 := \mathbb{E} \left[Z_T^n \int_0^T \left(\int_0^t q_s e_s d\kappa_s - \bar{X}_t \right) dA_t^n \right].$$

It follows from nonnegativity of \widehat{V} in (6.1), Lemma 3.2 and nonnegativity of \widehat{c} that

$$(6.8) \quad \begin{aligned} x + \int_0^t \widehat{H}_s dS_s + \bar{X}_t &\geq \int_0^t \widehat{c}_s d\kappa_s + \bar{X}_t - \int_0^t q_s e_s d\kappa_s \\ &\geq \bar{X}_t - \int_0^t q_s e_s d\kappa_s \geq 0, \quad t \in [0, T], \end{aligned}$$

that is, the integrand in T_1 is nonnegative. Let \mathbb{Q}^n be the probability measure, whose density process with respect to \mathbb{P} is Z^n . As $(x + \int_0^\cdot \widehat{H}_s dS_s + \bar{X})$ and \bar{X} are local martingales under \mathbb{Q}^n , by [33], Theorem III.27, page 128, $A_-^n \cdot (x + \int_0^\cdot \widehat{H}_s dS_s + \bar{X})$ and $A_-^n \cdot \bar{X}$ are local martingales. Let $(\tau_k)_{k \in \mathbb{N}}$ be a localizing sequence for both $A_-^n \cdot (x + \int_0^\cdot \widehat{H}_s dS_s + \bar{X})$ and $A_-^n \cdot \bar{X}$. By the monotone convergence theorem and integration by parts, we get

$$(6.9) \quad \begin{aligned} T_1 &= \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} \left(x + \int_0^t \widehat{H}_s dS_s + \bar{X}_t \right) dA_t^n \right] \\ &= \lim_{k \rightarrow \infty} \left(\mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} (-A_{t-}^n) d \left(x + \int_0^t \widehat{H}_s dS_s + \bar{X}_t \right) \right. \right. \\ &\quad \left. \left. + A_{\tau_k}^n \left(x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k} \right) \right] \right) \\ &= \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n} \left[A_{\tau_k}^n \left(x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k} \right) \right]. \end{aligned}$$

Let us consider T_2 . With $\mathcal{E}^q := \int_0^\cdot q_s e_s d\kappa_s$, Lemma 3.2 implies positivity of $\mathcal{E}_t^q - \bar{X}_t$, $t \in [0, T]$, which in turn allows to invoke the monotone convergence theorem, and we obtain

$$\begin{aligned} T_2 &= \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^T (\mathcal{E}_t^q - \bar{X}_t) dA_t^n \right] \\ &= \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} \mathcal{E}_t^q dA_t^n - \int_0^{\tau_k} \bar{X}_t dA_t^n \right]. \end{aligned}$$

By positivity of \bar{X} and the monotone convergence theorem, we have

$$\lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} \bar{X}_t dA_t^n \right] = \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^T \bar{X}_t dA_t^n \right].$$

Therefore, $\lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} \mathcal{E}_t^q dA_t^n \right] = \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^T \mathcal{E}_t^q dA_t^n \right]$. We deduce that

$$\mathbb{E}^{\mathbb{Q}^n} \left[\int_0^{\tau_k} \bar{X}_t dA_t^n \right] = \mathbb{E}^{\mathbb{Q}^n} \left[-(A_-^n \cdot \bar{X})_{\tau_k} + \bar{X}_{\tau_k} A_{\tau_k}^n \right] = \mathbb{E}^{\mathbb{Q}^n} \left[\bar{X}_{\tau_k} A_{\tau_k}^n \right].$$

Whereas, in the other term in T_2 , we get

$$\begin{aligned} \mathbb{E}^{\mathbb{Q}^n} \left[\int_0^T \mathcal{E}_t^q dA_t^n \right] &= \mathbb{E}^{\mathbb{Q}^n} \left[\mathcal{E}_T^q A_T^n - (A_-^n \cdot \mathcal{E}^q)_T \right] \\ &= \mathbb{E}^{\mathbb{Q}^n} \left[\mathcal{E}_T^q - (A_-^n \cdot \mathcal{E}^q)_T \right] \\ &= \mathbb{E}^{\mathbb{Q}^n} \left[((1 - A_-^n) \cdot \mathcal{E}^q)_T \right] \\ &= \mathbb{E}^{\mathbb{Q}^n} \left[(D^n \cdot \mathcal{E}^q)_T \right]. \end{aligned}$$

Using integration by parts and localization, as in the proof of Lemma 3.7, we can rewrite the latter expression as

$$\mathbb{E}\left[\int_0^T D_s^n Z_s^n q_s e_s d\kappa_s\right].$$

We conclude that

$$T_2 = -\lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[\bar{X}_{\tau_k} A_{\tau_k}^n] + \mathbb{E}\left[\int_0^T D_s^n Z_s^n q_s e_s d\kappa_s\right].$$

Combining this with (6.9), we obtain

$$\begin{aligned} T_1 + T_2 &= \mathbb{E}\left[\int_0^T D_s^n Z_s^n q_s e_s d\kappa_s\right] \\ &\quad + \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}\left[A_{\tau_k}^n\left(x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k}\right)\right] \\ &\quad - \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[A_{\tau_k}^n \bar{X}_{\tau_k}]. \end{aligned}$$

As both limits in the right-hand side exist and by positivity of $(x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k})$, established in (6.8), we can bound the difference of the limits as

$$\begin{aligned} &\lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}\left[A_{\tau_k}^n\left(x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k}\right)\right] - \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[A_{\tau_k}^n \bar{X}_{\tau_k}] \\ &\leq \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}\left[x + \int_0^{\tau_k} \widehat{H}_s dS_s + \bar{X}_{\tau_k}\right] - \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[A_{\tau_k}^n \bar{X}_{\tau_k}] \\ &\leq \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}\left[x + \int_0^{\tau_k} \widehat{H}_s dS_s\right] + \lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[(1 - A_{\tau_k}^n)\bar{X}_{\tau_k}]. \end{aligned}$$

By definition of \mathcal{M}' , \bar{X} is a uniformly integrable martingale under \mathbb{Q}^n . Therefore, as $(1 - A_{\tau_k}^n)$ is bounded, we can pass the limit inside of the expectation to obtain $\lim_{k \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^n}[(1 - A_{\tau_k}^n)\bar{X}_{\tau_k}] = 0$. In turn $x + \int_0^{\tau_k} \widehat{H}_s dS_s$ is a supermartingale under \mathbb{Q}^n (see the argument in the proof of Lemma 3.7). We conclude that

$$T_1 + T_2 \leq x + \mathbb{E}\left[\int_0^T D_s^n Z_s^n q_s e_s d\kappa_s\right] = x + \int_0^T q_s \rho_s^n dK_s,$$

for some ρ^n , which is well defined by Corollary 5.3, and such that $y^n(1, \rho^n) \in z\mathcal{B}(x, q)$, since $y^n Z^n D^n \in z\mathcal{G}(x, q)$. Thus, from (6.6) and (6.7), we get

$$z \leq \left(x + \int_0^T q_s \rho_s^n dK_s\right) y^n - \lim_{n \rightarrow \infty} y^n \mathbb{E}\left[Z_T^n \int_0^T \widehat{V}_t dA_t^n\right].$$

By optimality of \widehat{Y} , $z \geq (x + \int_0^T q_s \rho_s^n dK_s) y^n \geq 0$. Therefore, by nonnegativity of $\mathbb{E}[Z_T^n \times \int_0^T \widehat{V}_t dA_t^n]$, and since y^n converges to a strictly positive limit, we conclude that

$$\lim_{n \rightarrow \infty} \mathbb{E}\left[Z_T^n \int_0^T \widehat{V}_t dA_t^n\right] = 0.$$

Applying integration by parts and localization, we deduce the assertion of the lemma. \square

REMARK 6.3. We emphasize again that dA^n are probability measures on $[\theta_0, T]$, which can have mass at the endpoints, and $D_t^n = 1 - A_{t-}^n$, $\theta_0 \leq t \leq T$, $D_{T+}^n = 0$.

In the subsequent part, we will follow the notation of Lemma 6.2 and we will work with a further subsequence, still denoted by n , such that $y_n \rightarrow y > 0$,

$$(6.10) \quad Z^n D^n \rightarrow \frac{\widehat{Y}}{y}, \quad (d\kappa \times \mathbb{P})\text{-a.e.},$$

and

$$(6.11) \quad \sum_{k=n}^{\infty} \mathbb{E} \left[Z_T^n \int_0^T \widehat{V}_t dA_t^n \right] \leq \frac{2^{-n}}{n},$$

where the existence of a subsequence satisfying (6.11) follows from (6.5). The next result is a Komlos-type lemma, largely based on the results in [8], applied to the *double-sequence* of processes $(Z^n, (D^n)^{-1})$. We observe that the process $(D^n)^{-1}$ takes values in $[1, \infty]$ is increasing, left-continuous and satisfies

$$(D_t^n)^{-1} = 1, \quad 0 \leq t \leq \theta_0.$$

LEMMA 6.4. *Let the conditions of Theorem 4.3 hold. In the notation of Lemma 6.2, for each n , there exist a finite index $N(n)$ and convex weights*

$$\alpha_{n,k} \geq 0, \quad k = n, \dots, N(n), \quad \sum_{k=n}^{N(n)} \alpha_{n,k} = 1,$$

and there exists a strong optional supermartingale \widehat{Z} and a nondecreasing (not necessarily left-continuous) process \widehat{D} with $\widehat{D}_t = 0, \forall 0 \leq t \leq \theta_0$, such that, simultaneously:

1. $\widetilde{Z}^n := \sum_{k=n}^{N(n)} \alpha_{n,k} Z^k \rightarrow \widehat{Z}$ in the sense of [8], that is, for any stopping time $0 \leq \tau \leq T$ we have

$$\widetilde{Z}_\tau^n \rightarrow \mathbb{P} \widehat{Z}_\tau,$$

and
2.

$$\mathbb{P} \left(\sum_{k=n}^{N(n)} \alpha_{n,k} (D_t^k)^{-1} \rightarrow (\widehat{D}_t)^{-1}, \forall 0 \leq t \leq T+ \right) = 1$$

We have set $D_{T+} = D_{T+}^k = 0$.

Furthermore, with the notation

$$(6.12) \quad \widehat{A}_t = 1 - \widehat{D}_{t+}, \quad 0 \leq t \leq T, \quad \widehat{A}_{0-} = 0,$$

we have the probability measure $d\widehat{A}$ on $[\theta_0, T]$ such that

$$\mathbb{P}(\widehat{D}_t \in [1 - \widehat{A}_t, 1 - \widehat{A}_{t-}], \forall \theta_0 \leq t \leq T) = 1.$$

Denoting by

$$(6.13) \quad \widetilde{A}_t^n := 1 - \left(\sum_{k=n}^{N(n)} \alpha_{n,k} (1 - A_t^k)^{-1} \right)^{-1} = 1 - \left(\sum_{k=n}^{N(n)} \alpha_{n,k} (D_{t+}^k)^{-1} \right)^{-1},$$

$$0 \leq t \leq T, \quad \widetilde{A}_{0-}^n = 0,$$

the pointwise convergence in time (at all times where there is continuity) additionally implies

$$d\widetilde{A}^n \rightarrow d\widehat{A}, \quad \mathbb{P}\text{-a.e.}$$

in the sense of weak convergence of probability measures on $[\theta_0, T]$.

PROOF. The proof reduces to applying the Komlos-type results in [8] to the sequence Z^n and [4], Proposition 3.4, to the sequence $(D^n)^{-1}$, simultaneously. We observe that:

1. First, no bounds are needed for D^{-1} 's since we can apply the unbounded Komlos lemma in [9], Lemma A1.1 (and Remark 1 following it) to the proof from [4], Propositions 3.4, and this works even for infinite values (according to Remark 1 after [9], Lemma A1.1). Also, predictability can be replaced by optionality, without any change to the proof.

2. The Komlos arguments can be applied to both sequences Z^n and D^n simultaneously, with the same convex weights. In order to do this, we first apply Komlos to one sequence, then replace both original sequences by their convex combinations with the weights just obtained, and then apply Komlos again for the other sequence, and update the convex weight to both sequences. \square

COROLLARY 6.5. *Let the conditions of Theorem 4.3 hold. In the notation of Lemma 6.4, we have the representation*

$$\widehat{Y} = y\widehat{Z}\widehat{D}, \quad (d\kappa \times \mathbb{P})\text{-a.e.}$$

PROOF. Consider an observation that, for nonnegative numbers $a_k, b_k, k = n, \dots, N(n)$, we have

$$\min_{k=n, \dots, N(n)} \left(\frac{a_k}{b_k} \right) \leq \frac{\sum_{k=n}^{N(n)} \alpha_{n,k} a^k}{\sum_{k=n}^{N(n)} \alpha_{n,k} b_k} \leq \max_{k=n, \dots, N(n)} \left(\frac{a_k}{b_k} \right).$$

We apply this to $a_k = Z^k, b_k = (D^k)^{-1}$, to obtain

$$\min_{k=n, \dots, N(n)} (Z^k D^k) \leq \frac{\sum_{k=n}^{N(n)} \alpha_{n,k} Z^k}{\sum_{k=n}^{N(n)} \alpha_{n,k} (D^k)^{-1}} = \tilde{Z}^n \tilde{D}^n \leq \max_{k=n, \dots, N(n)} (Z^k D^k),$$

pointwise a.e. in the product space, where $\tilde{D}^n := \frac{1}{\sum_{k=n}^{N(n)} \alpha_{n,k} (D^k)^{-1}}$. Since

$$(Z^n D^n)_{n \in \mathbb{N}} \rightarrow \widehat{Y}/y, \quad (d\kappa \times \mathbb{P})\text{-a.e.},$$

we conclude that both

$$\left(\min_{k=n, \dots, N(n)} (Z^k D^k) \right)_{n \in \mathbb{N}} \quad \text{and} \quad \left(\max_{k=n, \dots, N(n)} (Z^k D^k) \right)_{n \in \mathbb{N}}$$

converge to $\frac{\widehat{Y}}{y}$, $(d\kappa \times \mathbb{P})$ -a.e., therefore,

$$(6.14) \quad \tilde{Z}^n \tilde{D}^n \rightarrow \frac{\widehat{Y}}{y}, \quad (d\kappa \times \mathbb{P})\text{-a.e.}$$

Using (6.10) above, if we did not plan to identify the limit \widehat{Z} as a strong-supermartingale, but only as a $(d\kappa \times \mathbb{P})$ -a.e. limit in the product space, we could only apply Komlos arguments to the single sequence D^{-1} , to conclude convergence of the other convex combination \tilde{Z}^n to $\frac{\widehat{Y}}{y\widehat{D}}$, defined up to a.e. equality in the product space.

With our (stronger) double Komlos argument, we have that, in addition to (6.10) we have

$$(6.15) \quad \widehat{Z}_\tau \widehat{D}_\tau = \mathbb{P}\text{-}\lim_n \tilde{Z}_\tau^n \tilde{D}_\tau^n \quad \text{for every } [0, T]\text{-valued stopping time } \tau$$

(where \widehat{Z} is a strong supermartingale, and \widehat{D} is well defined at all times).

It remains to prove that $\widehat{Y}/y = \widehat{Z}\widehat{D}$, $(d\kappa \times \mathbb{P})$ -a.e. We point out that the convergence (6.15) is topological. Let us consider an arbitrary optional set $O \subset \Omega \times [0, T]$ and fix an upper bound M . It follows from (6.14) that

$$1_O \min\{\tilde{Z}^n \tilde{D}^n, M\} \rightarrow 1_O \min\left\{\frac{\widehat{Y}}{y}, M\right\}, \quad (d\kappa \times \mathbb{P})\text{-a.e.}$$

Therefore, we get

$$\begin{aligned} \mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\{\tilde{Z}_t^n \tilde{D}_t^n, M\} d\kappa_t\right] \\ \rightarrow \mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\left\{\frac{\widehat{Y}_t}{y}, M\right\} d\kappa_t\right]. \end{aligned} \tag{6.16}$$

Recall that the stochastic clock has a density $d\kappa_t = \varphi_t dK_t$ with respect to the deterministic clock dK_t . For each fixed t , from (6.15) we have

$$1_O(t, \cdot) \min\{\tilde{Z}_t^n \tilde{D}_t^n, M\} \varphi_t \rightarrow 1_O(t, \cdot) \min\{\widehat{Z}_t \widehat{D}_t, M\} \varphi_t \quad \text{in } -\mathbb{P}.$$

Recalling that $\mathbb{E}[\varphi_t] < \infty$ we have, for fixed t , that

$$M \times \mathbb{E}[\varphi_t] \geq \mathbb{E}[1_O(t, \cdot) \min\{\tilde{Z}_t^n \tilde{D}_t^n, M\} \varphi_t] \rightarrow \mathbb{E}[1_O(t, \cdot) \min\{\widehat{Z}_t \widehat{D}_t, M\} \varphi_t].$$

We integrate the above with respect to the deterministic clock dK_t to obtain

$$\begin{aligned} \mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\{\tilde{Z}_t^n \tilde{D}_t^n, M\} d\kappa_t\right] \\ = \int_0^T \mathbb{E}[1_O(t, \cdot) \min\{\tilde{Z}_t^n \tilde{D}_t^n, M\} \varphi_t] dK_t \\ \rightarrow \int_0^T \mathbb{E}[1_O(t, \cdot) \min\{\widehat{Z}_t \widehat{D}_t, M\} \varphi_t] dK_t \\ = \mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\{\widehat{Z}_t \widehat{D}_t, M\} d\kappa_t\right] \end{aligned} \tag{6.17}$$

Together with (6.16), we have

$$\mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\left\{\frac{\widehat{Y}_t}{y}, M\right\} d\kappa_t\right] = \mathbb{E}\left[\int_0^T 1_O(t, \cdot) \min\{\widehat{Z}_t \widehat{D}_t, M\} d\kappa_t\right],$$

which holds for any optional set O and any bound M , therefore, $\widehat{Y} = y\widehat{Z}\widehat{D}$, $(d\kappa \times \mathbb{P})$ -a.e. □

PROOF OF THE THEOREM 6.1. Since

$$\sum_{k=n}^{N(n)} \alpha_{n,k} (D_t^k)^{-1} \rightarrow (\widehat{D}_t)^{-1}, \quad t \in [0, T],$$

recalling that \widehat{A} was defined from \widehat{D} and (6.13), as the processes A^n , therefore, \tilde{A}^n only increase by jumps. Therefore, we get

$$\begin{aligned} \Delta \tilde{A}_s^n &= \frac{1}{\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_s^k}} - \frac{1}{\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_{s+}^k}} \\ &= \sum_{k=n}^{N(n)} \underbrace{\frac{\alpha_{n,k} \frac{1}{D_{s+}^k}}{\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_{s+}^k}}}_{\leq 1} \frac{\Delta A_s^k}{D_s^k} \underbrace{\frac{1}{\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_s^k}}}_{\leq 1}. \end{aligned} \tag{6.18}$$

As $\frac{\alpha_{n,k} \frac{1}{D_s^k}}{(\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_s^k})} \leq 1$ and $\sum_{k=n}^{N(n)} \alpha_{n,k} \frac{1}{D_s^k} \geq 1$, we can bound the latter term in (6.18) by $\sum_{k=n}^{N(n)} \frac{\Delta A_s^k}{D_s^k}$ for $s \in [0, T]$. We deduce that

$$\Delta \tilde{A}_s^n \leq \sum_{k=n}^{N(n)} \frac{\Delta A_s^k}{D_s^k}, \quad s \in [0, T].$$

Therefore, we have

$$\int_0^t \widehat{V}_u d\tilde{A}_u^n \leq \sum_{k=n}^{N(n)} (D_t^k)^{-1} \int_0^t \widehat{V}_u dA_u^k, \quad t \in [0, T],$$

and thus, we obtain

$$\begin{aligned} (6.19) \quad \left(\min_{k=n, \dots, N(n)} (Z_t^k D_t^k) \right) \int_0^t \widehat{V}_u d\tilde{A}_u^n &\leq \sum_{k=n}^{N(n)} (Z_t^k D_t^k) (D_t^k)^{-1} \int_0^t \widehat{V}_u dA_u^k \\ &= \sum_{k=n}^{N(n)} Z_t^k \int_0^t \widehat{V}_u dA_u^k, \quad t \in [0, T]. \end{aligned}$$

Since the process

$$L_t^n := \sum_{k=n}^{N(n)} Z_t^k \int_0^t \widehat{V}_u dA_u^k, \quad 0 \leq t \leq T$$

is a nonnegative right-continuous submartingale, the maximal inequality and (6.11) together imply

$$\mathbb{P} \left(\sup_{0 \leq t \leq T} L_t^n \geq \frac{1}{n} \right) \leq n \mathbb{E}[L_T^n] \leq 2^{-n},$$

so

$$(6.20) \quad \sup_{0 \leq t \leq T} L_t^n \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

Since $Z^n D^n \rightarrow \frac{1}{y} \widehat{Y} > 0$, $(d\kappa \times \mathbb{P})$ -a.e., consequently

$$\min_{k=n, \dots, N(n)} (Z^k D^k) \rightarrow \frac{1}{y} \widehat{Y} > 0, \quad (d\kappa \times \mathbb{P})\text{-a.e.},$$

we obtain from (6.19) and (6.20) that the increasing RC process

$$\tilde{L}_t^n := \int_0^t \widehat{V}_u d\tilde{A}_u^n, \quad 0 \leq t \leq T,$$

converges to zero in the product space. Denoting by $\mathcal{O} \subset \Omega \times [0, T]$ the exceptional set where convergence to zero does not take place, and taking into account that \tilde{L}^n are increasing, we have that for

$$T^{\mathcal{O}}(\omega) = \inf\{0 \leq t \leq T : (\omega, t) \in \mathcal{O}\},$$

we have

$$(T^{\mathcal{O}}, T] \subset \mathcal{O}.$$

Now

$$(dk \times \mathbb{P})(\mathcal{O}) = 0,$$

implies that $T^{\mathcal{O}} \geq T$, \mathbb{P} -a.s. (here used an assumption that T is the minimal time horizon in the sense that the deterministic clock K is such that $K_t < K_T$, for every $t \in [0, T)$, this assumption does not restrict generality). Thus, there exists a set Ω^* of full probability $\mathbb{P}(\Omega^*) = 1$ such that, for each $\omega \in \Omega^*$ and $t < T$ we have

$$\int_0^t \widehat{V}_u(\omega) d\tilde{A}_t^n(\omega) \rightarrow 0.$$

Let us fix a $\omega \in \Omega^*$ and such that, for this ω , the probability measure $d\tilde{A}^n(\omega)$ converges weakly to $d\widehat{A}(\omega)$ over the interval $[\theta_0(\omega), T]$. The set of such ω 's still has probability 1. The Skorokhod representation theorem asserts that there exists a new probability space Ω^ω and a sequence of random times $(t^n(\omega))_{n \in \mathbb{N}}$ as well as $\widehat{t}(\omega)$ such that the distribution of $t^n(\omega)$ is $d\tilde{A}^n$, the distribution of $\widehat{t}(\omega)$ is $d\widehat{A}(\omega)$ and

$$t^n(\omega) \rightarrow \widehat{t}(\omega), \quad \mathbb{P}^\omega\text{-a.s.}$$

on the new, artificial, probability space. Fix $t < T$. We have

$$\mathbb{E}^\omega[\widehat{V}_{t^n(\omega)}(\omega) 1_{\{t^n(\omega) \leq t\}}] = \int_{\theta_0}^t \widehat{V}_u(\omega) d\tilde{A}_u^n(\omega) \rightarrow 0.$$

One can see that

$$\xi(\omega) := \liminf_n \widehat{V}_{t^n(\omega)}(\omega) \in \{\widehat{V}_{\widehat{t}(\omega)-}(\omega), \widehat{V}_{\widehat{t}(\omega)}(\omega)\}, \quad \mathbb{P}^\omega\text{-a.s.}$$

and

$$1_{\{t^n(\omega) \leq t\}} \rightarrow 1_{\{\widehat{t}(\omega) \leq t\}} \text{ on } \{\widehat{t}(\omega) < t\}, \quad \mathbb{P}^\omega\text{-a.s.}$$

Therefore, applying Fatou's lemma on Ω^ω , we obtain

$$\mathbb{E}^\omega[\xi(\omega) 1_{\{\widehat{t}(\omega) < t\}}] = 0.$$

We recall that both $\widehat{V}(\omega)$ and $\widehat{V}_-(\omega)$ are nonnegative, and consequently we have

$$0 \leq \min\{\widehat{V}_{\widehat{t}(\omega)-}(\omega), \widehat{V}_{\widehat{t}(\omega)}(\omega)\} \leq \xi(\omega),$$

and, therefore, we get

$$\mathbb{E}^\omega[\min\{\widehat{V}_{\widehat{t}(\omega)-}(\omega), \widehat{V}_{\widehat{t}(\omega)}(\omega)\} 1_{\{\widehat{t}(\omega) < t\}}] = 0.$$

This means that the distribution of $\widehat{t}(\omega)$ over the interval $[\theta_0(\omega), t)$ only charges the complement of the set of times

$$\{u < t : \widehat{V}_{u-}(\omega) = 0 \text{ or } \widehat{V}_u(\omega) = 0\}.$$

By taking $t \rightarrow T$, one completes the proof. \square

Acknowledgments. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

The first author was supported by NSF Grants DMS-1600307 (2015–2019) and DMS-1848339 (2019–2024).

The second author was supported by NSF Grants DMS-1517664 (2015–2019).

REFERENCES

- [1] BANK, P. and KAUPPILA, H. (2017). Convex duality for stochastic singular control problems. *Ann. Appl. Probab.* **27** 485–516. MR3619793 <https://doi.org/10.1214/16-AAP1209>
- [2] BARBU, V. and PRECUPANU, T. (2012). *Convexity and Optimization in Banach Spaces*, 4th ed. *Springer Monographs in Mathematics*. Springer, Dordrecht. MR3025420 <https://doi.org/10.1007/978-94-007-2247-7>
- [3] BRANNATH, W. and SCHACHERMAYER, W. (1999). A bipolar theorem for $L_+^0(\Omega, \mathcal{F}, \mathbb{P})$. In *Séminaire de Probabilités, XXXIII. Lecture Notes in Math.* **1709** 349–354. Springer, Berlin. MR1768009 <https://doi.org/10.1007/BFb0096525>
- [4] CAMPI, L. and SCHACHERMAYER, W. (2006). A super-replication theorem in Kabanov’s model of transaction costs. *Finance Stoch.* **10** 579–596. MR2276320 <https://doi.org/10.1007/s00780-006-0022-4>
- [5] CUCHIERO, C. and TEICHMANN, J. (2015). A convergence result for the Emery topology and a variant of the proof of the fundamental theorem of asset pricing. *Finance Stoch.* **19** 743–761. MR3413934 <https://doi.org/10.1007/s00780-015-0276-9>
- [6] CUOCO, D. (1997). Optimal consumption and equilibrium prices with portfolio constraints and stochastic income. *J. Econom. Theory* **72** 33–73. MR1436126 <https://doi.org/10.1006/jeth.1996.2207>
- [7] CVITANIĆ, J., SCHACHERMAYER, W. and WANG, H. (2001). Utility maximization in incomplete markets with random endowment. *Finance Stoch.* **5** 259–272. MR1841719 <https://doi.org/10.1007/PL00013534>
- [8] CZICHOWSKY, C. and SCHACHERMAYER, W. (2016). Strong supermartingales and limits of nonnegative martingales. *Ann. Probab.* **44** 171–205. MR3456335 <https://doi.org/10.1214/14-AOP970>
- [9] DELBAEN, F. and SCHACHERMAYER, W. (1994). A general version of the fundamental theorem of asset pricing. *Math. Ann.* **300** 463–520. MR1304434 <https://doi.org/10.1007/BF01450498>
- [10] DELBAEN, F. and SCHACHERMAYER, W. (1997). The Banach space of workable contingent claims in arbitrage theory. *Ann. Inst. Henri Poincaré Probab. Stat.* **33** 113–144. MR1440258 [https://doi.org/10.1016/S0246-0203\(97\)80118-5](https://doi.org/10.1016/S0246-0203(97)80118-5)
- [11] DELBAEN, F. and SCHACHERMAYER, W. (1998). The fundamental theorem of asset pricing for unbounded stochastic processes. *Math. Ann.* **312** 215–250. MR1671792 <https://doi.org/10.1007/s002080050220>
- [12] DELLACHERIE, C. and MEYER, P.-A. (1982). *Probabilities and Potential. B: Theory of Martingales*. North-Holland Mathematics Studies **72**. North-Holland, Amsterdam. MR0745449
- [13] DUFFIE, D., FLEMING, W., SONER, H. M. and ZARIPHOPOULOU, T. (1997). Hedging in incomplete markets with HARA utility. *J. Econom. Dynam. Control* **21** 753–782. MR1455755 [https://doi.org/10.1016/S0165-1889\(97\)00002-X](https://doi.org/10.1016/S0165-1889(97)00002-X)
- [14] DUFFIE, D. and ZARIPHOPOULOU, T. (1993). Optimal investment with undiversifiable income risk. *Math. Finance* **3** 135–148.
- [15] EKELAND, I. and TEMAM, R. (1976). *Convex Analysis and Variational Problems*. North-Holland, Amsterdam. MR0463994
- [16] EL KAROUI, N. and JEANBLANC-PICQUE, M. (1998). Optimization of consumption with labor income. *Finance Stoch.* **2** 409–440.
- [17] FÖLLMER, H. and KRAMKOV, D. (1997). Optional decompositions under constraints. *Probab. Theory Related Fields* **109** 1–25. MR1469917 <https://doi.org/10.1007/s004400050122>
- [18] HE, H. and PAGÈS, H. F. (1993). Labor income, borrowing constraints, and equilibrium asset prices. *Econom. Theory* **3** 663–696. MR1306776 <https://doi.org/10.1007/BF01210265>
- [19] HUGONNIER, J. and KRAMKOV, D. (2004). Optimal investment with random endowments in incomplete markets. *Ann. Appl. Probab.* **14** 845–864. MR2052905 <https://doi.org/10.1214/105051604000000134>
- [20] HUGONNIER, J., KRAMKOV, D. and SCHACHERMAYER, W. (2005). On utility-based pricing of contingent claims in incomplete markets. *Math. Finance* **15** 203–212. MR2132189 <https://doi.org/10.1111/j.0960-1627.2005.00217.x>
- [21] JACOD, J. and SHIRYAEV, A. N. (2003). *Limit Theorems for Stochastic Processes*, 2nd ed. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **288**. Springer, Berlin. MR1943877 <https://doi.org/10.1007/978-3-662-05265-5>
- [22] KARATZAS, I. (1989). Optimization problems in the theory of continuous trading. *SIAM J. Control Optim.* **27** 1221–1259. MR1022426 <https://doi.org/10.1137/0327063>
- [23] KARATZAS, I. and KARDARAS, C. (2007). The numéraire portfolio in semimartingale financial models. *Finance Stoch.* **11** 447–493. MR2335830 <https://doi.org/10.1007/s00780-007-0047-3>
- [24] KARATZAS, I. and ŽITKOVIĆ, G. (2003). Optimal consumption from investment and random endowment in incomplete semimartingale markets. *Ann. Probab.* **31** 1821–1858. MR2016601 <https://doi.org/10.1214/aop/1068646367>

- [25] KARDARAS, C. (2013). On the closure in the Emery topology of semimartingale wealth-process sets. *Ann. Appl. Probab.* **23** 1355–1376. MR3098435 <https://doi.org/10.1214/12-aap872>
- [26] KLEIN, I., LÉPINETTE, E. and PEREZ-OSTAFE, L. (2014). Asymptotic arbitrage with small transaction costs. *Finance Stoch.* **18** 917–939. MR3255756 <https://doi.org/10.1007/s00780-014-0242-y>
- [27] KRAMKOV, D. O. (1996). Optional decomposition of supermartingales and hedging contingent claims in incomplete security markets. *Probab. Theory Related Fields* **105** 459–479. MR1402653 <https://doi.org/10.1007/BF01191909>
- [28] MALAMUD, S. and TRUBOWITZ, E. (2007). The structure of optimal consumption streams in general incomplete markets. *Math. Financ. Econ.* **1** 129–161. MR2365686 <https://doi.org/10.1007/s11579-007-0006-y>
- [29] MATOUSSI, A. and XING, H. (2018). Convex duality for Epstein–Zin stochastic differential utility. *Math. Finance* **28** 991–1019. MR3858077 <https://doi.org/10.1111/mafi.12168>
- [30] MNIF, M. and PHAM, H. (2001). Stochastic optimization under constraints. *Stochastic Process. Appl.* **93** 149–180. MR1819488 [https://doi.org/10.1016/S0304-4149\(00\)00089-2](https://doi.org/10.1016/S0304-4149(00)00089-2)
- [31] MOSTOVYI, O. (2015). Necessary and sufficient conditions in the problem of optimal investment with intermediate consumption. *Finance Stoch.* **19** 135–159. MR3292127 <https://doi.org/10.1007/s00780-014-0248-5>
- [32] MOSTOVYI, O. (2017). Optimal investment with intermediate consumption and random endowment. *Math. Finance* **27** 96–114. MR3609722 <https://doi.org/10.1111/mafi.12089>
- [33] PROTTER, P. E. (2004). *Stochastic Integration and Differential Equations: Stochastic Modelling and Applied Probability*, 2nd ed. *Applications of Mathematics (New York)* **21**. Springer, Berlin. MR2020294
- [34] RÁSONYI, M. (2018). On utility maximization without passing by the dual problem. *Stochastics* **90** 955–971. MR3854522 <https://doi.org/10.1080/17442508.2018.1457677>
- [35] ROKHLIN, D. B. (2010). On the existence of an equivalent supermartingale density for a fork-convex family of random processes. *Mat. Zametki* **87** 594–603. MR2762746 <https://doi.org/10.1134/S0001434610030338>
- [36] YU, X. (2015). Utility maximization with addictive consumption habit formation in incomplete semimartingale markets. *Ann. Appl. Probab.* **25** 1383–1419. MR3325277 <https://doi.org/10.1214/14-AAP1026>
- [37] ŽITKOVIĆ, G. (2005). Utility maximization with a stochastic clock and an unbounded random endowment. *Ann. Appl. Probab.* **15** 748–777. MR2114989 <https://doi.org/10.1214/105051604000000738>