

Renewal theorems and mixing for non Markov flows with infinite measure

Ian Melbourne^a and Dalia Terhesiu^b

^aMathematics Institute, University of Warwick, Coventry, CV4 7AL, UK. E-mail: I.Melbourne@warwick.ac.uk

^bDepartment of Mathematics, University of Exeter, Exeter EX4 4QF, UK. E-mail: d.terhesiu@exeter.ac.uk

Received 26 June 2018; revised 20 December 2018; accepted 13 February 2019

Abstract. We obtain results on mixing for a large class of (not necessarily Markov) infinite measure semiflows and flows. Erickson proved, amongst other things, a strong renewal theorem in the corresponding i.i.d. setting. Using operator renewal theory, we extend Erickson's methods to the deterministic (i.e. non-i.i.d.) continuous time setting and obtain results on mixing as a consequence.

Our results apply to intermittent semiflows and flows of Pomeau–Manneville type (both Markov and nonMarkov), and to semiflows and flows over Collet–Eckmann maps with nonintegrable roof function.

Résumé. Nous obtenons des résultats de mélange pour une large classe de flots et de semi-flots préservant une mesure de masse infinie (et qui ne sont pas nécessairement markoviens). Erickson a prouvé, entre autres choses, un théorème de renouvellement fort dans le contexte de variables aléatoires indépendantes et identiquement distribuées. En utilisant la théorie des opérateurs de renouvellement, nous étendons les méthodes d'Erickson au cas du temps continu déterministe (et donc on i.i.d.) et en déduisons des résultats sur le mélange.

Nos résultats s'appliquent à des semi-flots et flots intermittents de type Pomeau–Manneville (à la fois de type markoviens ou de type non-markoviens) ainsi qu'à des suspensions au-dessus de transformations de Collet–Eckmann pour lesquelles la fonction toit est non-intégrable.

MSC: 37A25; 37A40; 37A50; 60K05

Keywords: Flows; Renewal theorems; Operator renewal theory; Krickeberg mixing

1. Introduction

Recently, there has been increasing interest in the investigation of mixing properties for infinite measure-preserving dynamical systems [2,13,24,27,28,31–35,37,40,41,43]. Most of these results are for discrete time noninvertible systems.

For results on semiflows preserving an infinite measure, we refer to [37] (the Markov case) and [13] (which does not assume a Markov structure). The setting is that $F : Y \rightarrow Y$ is a mixing uniformly expanding map defined on a probability space (Y, μ) and $\tau : Y \rightarrow \mathbb{R}^+$ is a nonintegrable roof function with regularly varying tails:

$$\mu(y \in Y : \tau(y) > t) = \ell(t)t^{-\beta} \quad \text{for various ranges of } \beta \in [0, 1]. \quad (1.1)$$

Here, $\ell : [0, \infty) \rightarrow [0, \infty)$ is a measurable slowly varying function (so $\lim_{t \rightarrow \infty} \ell(\lambda t)/\ell(t) = 1$ for all $\lambda > 0$). Consider the suspension (Y^τ, μ^τ) and suspension semiflow $F_t : Y^\tau \rightarrow Y^\tau$ (the standard definitions are recalled in Section 3). The aim is to prove a mixing result of the form

$$\lim_{t \rightarrow \infty} a_t \int_{Y^\tau} vw \circ F_t d\mu^\tau = \int_{Y^\tau} v d\mu^\tau \int_{Y^\tau} w d\mu^\tau,$$

for a suitable normalisation $a_t \rightarrow \infty$ and suitable classes of observables $v, w : Y^\tau \rightarrow \mathbb{R}$.

Under certain hypotheses, [13,37] obtained results on mixing and rates of mixing for such semiflows. The hypotheses were of two types: (i) assumptions on “renewal operators” associated to the transfer operator of F and the roof function τ , and (ii) Dolgopyat-type assumptions of the type used to obtain mixing rates for finite measure (semi)flows [17].

As pointed out to us by Dima Dolgopyat, Péter Nándori and Doma Szász, mixing for indicator functions can be regarded as a local limit theorem and hence hypotheses of type (ii) should not be necessary.

In this paper, we show that operator renewal-theoretic assumptions (i) are indeed sufficient for obtaining the mixing results in [13,37]. The abstract framework in [13] turns out again to be flexible enough to cover nonMarkov situations. Moreover, our main results extend to flows and we are able to treat large classes of observables v, w . (Conditions of type (i) alone are not sufficient for obtaining rates of mixing; the best results remain those in [13].)

The analogous probabilistic results go back to Erickson [20] who obtained *strong renewal theorems* in an i.i.d. continuous time framework under the assumption $\beta \in (\frac{1}{2}, 1]$. (In the discrete time setting, see [22] for the i.i.d. case and [35] for the deterministic case.) Our results on mixing when $\beta \in (\frac{1}{2}, 1]$ for semiflows (Corollary 3.1 and the extensions in Section 9) and for flows (Theorem 10.5), are proved by adapting Erickson's methods to the deterministic setting.

For $\beta \leq \frac{1}{2}$, additional hypotheses are needed on the tail of τ to obtain a strong renewal theorem (and hence mixing) even for discrete time; see [15,19,22] for i.i.d. results and [24] for deterministic results (see also [41] for higher order theory in both the i.i.d. and deterministic settings). For the continuous time case, Dolgopyat & Nándori [18] obtain strong renewal theorems for a class of Markov semiflows including the range $\beta \leq \frac{1}{2}$ (again under extra hypotheses on the tail $\mu(\tau > t)$), though our main examples seem beyond their framework. In the absence of additional tail hypotheses, [20] showed how to obtain a partial result in the probabilistic setting with limit replaced by \liminf . In Corollary 3.5, we obtain such a \liminf result for semiflows with $\beta \in (0, \frac{1}{2}]$.

We now describe two families of examples to which our results apply. For definiteness, we restrict to our main mixing result Corollary 3.1 which applies when $\beta \in (\frac{1}{2}, 1]$. (Corollary 3.3 and Corollary 3.5 hold for all $\beta \in (0, 1]$.)

Example 1.1 (NonMarkovian intermittent semiflows and flows). Consider the map $f : [0, 1] \rightarrow [0, 1]$ given by

$$f(x) = x(1 + c_1 x^{1/\beta}) \bmod 1 \quad \text{where } \beta \in \left(\frac{1}{2}, 1\right], c_1 > 0.$$

This is an example of an AFN map [45], namely a nonuniformly expanding one-dimensional map with at most countably (in this case finitely) many branches with finite images and satisfying Adler's distortion condition $\sup |f''|/|f'|^2 < \infty$. Up to scaling, there is a unique absolutely continuous invariant measure μ_0 . The measure μ_0 is infinite and the density has a singularity at the neutral fixed point 0.

Let $\tau_0 : [0, 1] \rightarrow [1, \infty)$ be a roof function of bounded variation and Hölder continuous, and let f_t denote the suspension semiflow on $[0, 1]^{\tau_0}$ with invariant measure $\mu_0^{\tau_0} = \mu_0 \times \text{Lebesgue}$. Note that there is now a neutral periodic orbit of period $\tau_0(0)$.

In [13], under a Dolgopyat-type condition on τ_0 and for sufficiently regular observables v and w supported away from the neutral periodic orbit, we proved a mixing result with rates and higher order asymptotics. Here we obtain the mixing result without requiring the Dolgopyat-type condition or high regularity for the observables. It suffices that f_t has two periodic orbits (other than the neutral periodic orbit) whose periods have irrational ratio. Define

$$m(t) = \begin{cases} \log t & \beta = 1, \\ t^{1-\beta} & \beta \in (\frac{1}{2}, 1). \end{cases}$$

We show that

$$\lim_{t \rightarrow \infty} m(t) \int v w \circ f_t d\mu_0^{\tau_0} = \text{const.} \int v d\mu_0^{\tau_0} \int w d\mu_0^{\tau_0}, \quad (1.2)$$

where the constant depends only on f and τ_0 . Here, v is any continuous function supported away from the neutral periodic orbit and w is any integrable function.

Remark 1.2. If c_1 is a positive integer, then f is Markov and is a special case of the class of maps considered by [42]. In this case, it suffices that τ_0 is Hölder continuous. Moreover, it follows from [18] that the mixing result (1.2) holds for all $\beta \leq 1$. When c_1 is not an integer, f is not Markov and [18] does not apply, as far as we can tell, regardless of the value of β .

As in [33,34], we can also consider solenoidal flows with a neutral periodic orbit. Our results on mixing apply equally to such flows, see Remark 11.3.

Example 1.3 (Suspensions over unimodal maps). We consider a class of examples studied in [13, Example 1.2]. Under a Dolgopyat-type condition on τ_0 and for sufficiently regular observables v and w , we proved a mixing result with rates

and higher order asymptotics. Again, the emphasis is now on mixing rather than mixing rates, with significantly relaxed hypotheses on the roof function and the observables.

Let $f : [0, 1] \rightarrow [0, 1]$ be a C^2 unimodal map with unique non-flat critical point $x_0 \in (0, 1)$. We suppose further that f is *Collet–Eckmann* [16]: there are constants $C > 0$, $\lambda_{\text{CE}} > 1$ such that $|(f^n)'(f x_0)| \geq C \lambda_{\text{CE}}^n$ for all $n \geq 1$. It follows [26] that there is a unique acip μ_0 that is mixing up to a finite cycle. We restrict to the case when μ_0 is mixing. Finally, we suppose that x_0 satisfies *slow recurrence* in the sense that $\lim_{n \rightarrow \infty} n^{-1} \log |f^n x_0 - x_0| = 0$.

Consider a roof function $\tau_0 : [0, 1] \rightarrow \mathbb{R}^+$ of the form $\tau_0(x) = g(x)|x - x_0|^{-1/\beta}$ where $\beta \in (\frac{1}{2}, 1)$ and $g : [0, 1] \rightarrow (1, \infty)$ is differentiable, and form the suspension semiflow $f_t : [0, 1]^{\tau_0} \rightarrow [0, 1]^{\tau_0}$. Suppose that f_t has two periodic orbits whose periods have irrational ratio. We obtain the mixing property (1.2) for any continuous function v supported in $[0, 1] \times [0, 1]$ and any integrable w .

The remainder of this paper is organised as follows. In Section 2, we describe the operator renewal-theoretic hypotheses required in this paper and we state a strong renewal theorem for $\beta \in (\frac{1}{2}, 1]$ as well as related results for $\beta \leq \frac{1}{2}$. In Section 3, we show how these results lead to mixing properties for semiflows. Sections 4 and 6 are devoted to the proof of the strong renewal theorem, while Sections 7 and 8 contain the proofs of the remaining results in Section 2. Section 5 contains prerequisites from operator renewal theory.

Corollary 3.1 (mixing for semiflows) is stated for observables that are certain indicator functions. This restriction is relaxed considerably in Section 9. The corresponding result for flows is stated and proved in Section 10.

Finally, in Section 11 we return to Examples 1.1 and 1.3.

Notation. We use “big O” and \ll notation interchangeably, writing $a_n = O(b_n)$ or $a_n \ll b_n$ if there is a constant $C > 0$ such that $a_n \leq C b_n$ for all $n \geq 1$. Also, we write $a_n \sim b_n$ if $\lim_{n \rightarrow \infty} a_n/b_n = 1$.

2. Strong renewal theorem for continuous time deterministic systems

Let (Y, μ) be a probability space and let $F : Y \rightarrow Y$ be an ergodic and mixing measure-preserving transformation. Let $\tau : Y \rightarrow \mathbb{R}^+$ be a measurable nonintegrable function bounded away from zero. For convenience, we suppose that $\text{ess inf } \tau > 1$. Throughout we assume the regularly varying tail condition (1.1).

Let $\tau_n = \sum_{j=0}^{n-1} \tau \circ F^j$. Given measurable sets $A, B \subset Y$, define the renewal measure

$$U_{A,B}(I) = \sum_{n=0}^{\infty} \mu(y \in A \cap F^{-n} B : \tau_n(y) \in I), \quad (2.1)$$

for intervals $I \subset \mathbb{R}$. We write $U_{A,B}(x) = U_{A,B}([0, x])$ for $x > 0$. Our aim is to generalise [20, Theorems 1 and 2] to this set up. That is, we want to obtain the asymptotics of $U_{A,B}(t+h) - U_{A,B}(t)$ for any $h > 0$.

With the same notation as in [13], let $\overline{\mathbb{H}} = \{\text{Re } s \geq 0\}$. Given $\delta > 0$ and $L > 0$, let $\mathbb{H}_{\delta,L} = (\overline{\mathbb{H}} \cap B_{\delta}(0)) \cup \{ib : |b| \leq L\}$. Define the family of operators for $s \in \mathbb{H}$,

$$\hat{R}(s) : L^1(Y) \rightarrow L^1(Y), \quad \hat{R}(s)v = R(e^{-s\tau} v).$$

Here $R : L^1(Y) \rightarrow L^1(Y)$ is the transfer operator for F (so $\int_Y Rvw d\mu = \int_Y vw \circ F d\mu$ for all $v \in L^1(Y)$, $w \in L^\infty(Y)$).

We assume that there exists $p_0 \geq 1$, and for each $p \in (p_0, \infty)$, $\gamma \in (0, \beta)$ and $L > 0$ there exists a Banach space $\mathcal{B} = \mathcal{B}(Y)$ containing constant functions, with norm $\|\cdot\|_{\mathcal{B}}$, and constants $\delta \in (0, L)$, $\alpha_0 \in (0, 1)$ and $C > 0$ such that

- (H) (i) \mathcal{B} is compactly embedded in L^p .
(ii) $\|\hat{R}(s)^n v\|_{\mathcal{B}} \leq C(|v|_p + \alpha_0^n \|v\|_{\mathcal{B}})$ for all $s \in \mathbb{H}_{\delta,L}$, $v \in \mathcal{B}$, $n \geq 1$.
(iii) $|R(\tau^\gamma v)|_p \leq C\|v\|_{\mathcal{B}}$ for all $v \in \mathcal{B}$.

Also, most of our results require one of the following conditions:

- (S) (i) For all $b \in [-L, L]$, $b \neq 0$, the spectrum of $\hat{R}(ib) : \mathcal{B} \rightarrow \mathcal{B}$ does not contain 1.
(ii) For all $b \in [-L, L]$, $b \neq 0$, the spectral radius of $\hat{R}(ib) : \mathcal{B} \rightarrow \mathcal{B}$ is less than 1.

Hypothesis (H) is similar to [13, hypothesis (H1)]. The hypotheses in (S) are significant weakenings of [13, hypothesis (H4)] and the diophantine ratio assumption used in [37] (Dolgopyat-type condition). The remaining hypotheses in [13], namely (H2) and (H3) (re-inducing), are not required.

Remark 2.1.

(a) For ease of exposition, hypothesis (H) is stated on the half-plane $\overline{\mathbb{H}}$, though we only use s real and s imaginary in this paper. For Theorems 2.3 and 2.4, we can take $s = ib$, $b \in [-L, L]$ in (H)(ii). For Theorem 2.6, we can take $s = a$, $a \in [0, \delta)$ in (H)(ii).

(b) For our main results Theorem 2.3 and Corollary 3.1, it suffices that $\gamma > 1 - \beta$ (this is possible since $\beta > \frac{1}{2}$ in those results). For our other results which include $\beta \leq \frac{1}{2}$, it suffices that $\gamma > 0$.

In addition, as in [13], there exists $p_0 \geq 1$ depending only on β and γ such that (H) is required to hold only for one value of $p > p_0$.

Remark 2.2. In the simplest setting, studied in [37], where the map $F : Y \rightarrow Y$ is Gibbs–Markov [1,3], hypothesis (H) is satisfied with \mathcal{B} a symbolic Hölder space and $p = \infty$. See [13, Remark 2.4] and [37, Proposition 3.5] for further details. This includes the case of Markovian intermittent semiflows.

As explained in Section 11.1, this situation generalizes to the case when F is an AFU map (i.e. an AFN map as defined in Example 1.1 but uniformly expanding instead of nonuniformly expanding), with \mathcal{B} consisting of bounded variation functions, enabling us to treat the nonMarkovian intermittent semiflows in Example 1.1.

Define

$$d_\beta = \begin{cases} \frac{1}{\pi} \sin \beta\pi & \beta < 1, \\ 1 & \beta = 1, \end{cases} \quad m(t) = \begin{cases} \ell(t)t^{1-\beta} & \beta < 1, \\ \int_1^t \ell(s)s^{-1} ds & \beta = 1. \end{cases}$$

Throughout we suppose that $A, B \subset Y$ are measurable and that $1_A \in \mathcal{B}$.

Our main result generalizes [20, Theorem 1] to the present non i.i.d. set up:

Theorem 2.3 (Strong renewal theorem). Assume $\mu(\tau > t) = \ell(t)t^{-\beta}$ where $\beta \in (\frac{1}{2}, 1]$. Suppose that (H) and (S)(i) holds. Then for any $h > 0$,

$$\lim_{t \rightarrow \infty} m(t)(U_{A,B}(t+h) - U_{A,B}(t)) = d_\beta \mu(A)\mu(B)h.$$

As discussed in the Introduction, additional hypotheses are needed to obtain a strong renewal theorem when $\beta \leq \frac{1}{2}$. However, generalizing [20, Theorem 2] to the present non i.i.d. set up, we still obtain a lim inf result:

Theorem 2.4. Assume $\mu(\tau > t) = \ell(t)t^{-\beta}$ where $\beta \in (0, 1)$. Suppose that (H) and (S)(ii) holds. Then for any $h > 0$,

$$\liminf_{t \rightarrow \infty} m(t)(U_{A,B}(t+h) - U_{A,B}(t)) = d_\beta \mu(A)\mu(B)h.$$

Remark 2.5. In the i.i.d. setting, results of this type are first due to [22] for discrete time and $\beta < 1$. The results of [20] extended [22] to continuous time and incorporated the case $\beta = 1$.

For the proof of Theorem 2.4, we will need the following result which gives the asymptotics of $U_{A,B}$ for the entire range $\beta \in [0, 1]$. This implies a property for the semiflow F_t known as *weak rational ergodicity* [1,4] (see Corollary 3.3 below) and thus is of interest in its own right.

Theorem 2.6. Assume $\mu(\tau > t) = \ell(t)t^{-\beta}$ where $\beta \in [0, 1]$. Suppose that (H) holds. Then

$$\lim_{t \rightarrow \infty} t^{-1} m(t) U_{A,B}(t) = D_\beta \mu(A)\mu(B),$$

where $D_\beta = \{\Gamma(1 - \beta)\Gamma(1 + \beta)\}^{-1}$ if $\beta \in (0, 1)$ and $D_0 = D_1 = 1$.

For the proof of Theorem 2.4, we will also require the following local limit theorem with error term which may also be of interest in its own right. Let

$$q_\beta(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ibt} e^{-c_\beta |b|^\beta} db, \quad c_\beta = i \int_0^\infty e^{-i\sigma} \sigma^{-\beta} d\sigma.$$

Theorem 2.7 (LLT). Assume the setting of Theorem 2.4 with $\beta \in (0, 1)$. Let $d_n > 0$ be an increasing sequence with $d_n \rightarrow \infty$ such that $n\mu(\tau > d_n) = n\ell(d_n)d_n^{-\beta} \rightarrow 1$, as $n \rightarrow \infty$. Then for any $h > 0$ there exists $e_n > 0$ with $\lim_{n \rightarrow \infty} e_n = 0$ such that for all $t > 0, n \geq 1$,

$$\left| \mu(y \in A \cap F^{-n}B : \tau_n(y) \in [t, t + h]) - \frac{h}{d_n} q_\beta(t/d_n) \mu(A) \mu(B) \right| \leq \frac{e_n}{d_n}.$$

Alternative hypotheses. In certain examples, such as those where $F : Y \rightarrow Y$ is modelled by a Young tower with exponential tails [44], hypothesis (H)(iii) is problematic. In such cases, it is necessary as in [13] to consider alternative hypotheses.

We assume that for every (sufficiently large) $p \in (1, \infty)$, there exists a Banach space \mathcal{B} containing constant functions, with norm $\|\cdot\|_{\mathcal{B}}$, and constants $\delta > 0, \alpha_0 \in (0, 1)$ and $C > 0$ such that

(A)(i) \mathcal{B} is compactly embedded in L^p .

(ii) $\|\hat{R}(s)^n v\|_{\mathcal{B}} \leq C(|v|_{L^1} + \alpha_0^n \|v\|_{\mathcal{B}})$ for all $s \in \overline{\mathbb{H}}_{\delta, L}, v \in \mathcal{B}, n \geq 1$.

It follows from these assumptions (see Lemma 5.1(d) below), that (after possibly shrinking δ) there is a continuous family of simple eigenvalues $\lambda(s)$ for $\hat{R}(s) : \mathcal{B} \rightarrow \mathcal{B}, s \in \overline{\mathbb{H}} \cap B_\delta(0)$, with $\lambda(0) = 1$. Let $\zeta(s) \in \mathcal{B}$ be the corresponding family of eigenfunctions normalized so that $\int_Y \zeta(s) d\mu = 1$. We assume further that there exists $\beta_+ \in (\beta, 1)$ such that

(A)(iii) $|\int_Y (e^{-s\tau} - 1)(\zeta(s) - 1) d\mu| \leq C|s|^{\beta_+}$ for all $s \in \overline{\mathbb{H}} \cap B_\delta(0)$.

Theorem 2.8. Suppose that hypothesis (H) is replaced by hypothesis (A). Then Theorems 2.4, 2.6 and 2.7 remain valid. If in addition $\mu(\tau > t) = ct^{-\beta} + O(t^{-q})$ where $c > 0, \beta \in (\frac{1}{2}, 1), q > 1$, then Theorem 2.3 remains valid.

3. Mixing for infinite measure semiflows

In this section, we obtain various mixing results for semiflows as consequences of the results in Section 2.

Let $F : Y \rightarrow Y$ and $\tau : Y \rightarrow \mathbb{R}^+$ be as in Section 2. Define the suspension $Y^\tau = \{(y, u) \in Y \times \mathbb{R} : 0 \leq u \leq \tau(y)\} / \sim$ where $(y, \tau(y)) \sim (Fy, 0)$. The suspension semiflow $F_t : Y^\tau \rightarrow Y^\tau$ is given by $F_t(y, u) = (y, u + t)$, computed modulo identifications. The measure $\mu^\tau = \mu \times \text{Lebesgue}$ is ergodic, F_t -invariant and σ -finite. Since τ is nonintegrable, μ^τ is an infinite measure.

Throughout this section, we suppose that $A_1 = A \times [a_1, a_2], B_1 = B \times [b_1, b_2]$ are measurable subsets of $\{(y, u) \in Y \times \mathbb{R} : 0 \leq u \leq \tau(y)\}$ (so $0 \leq a_1 < a_2 \leq \text{ess inf}_A \tau, 0 \leq b_1 < b_2 \leq \text{ess inf}_B \tau$), and that $1_A \in \mathcal{B}$. Also, we continue to suppose that $\mu(\tau > t) = \ell(t)t^{-\beta}$ for various ranges of $\beta \in [0, 1]$.

Corollary 3.1. Assume the setting of Theorem 2.3 (alternatively Theorem 2.8), so in particular $\beta \in (\frac{1}{2}, 1]$. Then $\lim_{t \rightarrow \infty} m(t) \mu^\tau(A_1 \cap F_t^{-1} B_1) = d_\beta \mu^\tau(A_1) \mu^\tau(B_1)$.

Proof. Recall that $\text{ess inf } \tau > 1$. Let $h \in (0, 1)$ and note using (2.1) that

$$\begin{aligned} U_{A,B}(t+h) - U_{A,B}(t) &= \mu(y \in A : F^n y \in B \text{ and } \tau_n(y) \in [t, t+h] \text{ for some } n \geq 0) \\ &= \mu(y \in A : F_{t+h}(y, 0) \in B \times [0, h]). \end{aligned}$$

After dividing rectangles into smaller subrectangles, we can suppose without loss that $b_2 - b_1 < 1$. Set $h = b_2 - b_1$. Then

$$\begin{aligned} \mu^\tau(A_1 \cap F_t^{-1} B_1) &= \mu^\tau\{(y, u) \in A \times [a_1, a_2] : F_t(y, u) \in B \times [b_1, b_2]\} \\ &= \mu^\tau\{(y, u) \in A \times [a_1, a_2] : F_{t+u-b_1}(y, 0) \in B \times [0, h]\} \\ &= \int_{a_1}^{a_2} \mu\{y \in A : F_{t+u-b_1}(y, 0) \in B \times [0, h]\} du \\ &= \int_{a_1}^{a_2} (U_{A,B}(t+u-b_1) - U_{A,B}(t+u-b_1-h)) du. \end{aligned} \tag{3.1}$$

Hence

$$\begin{aligned} m(t)\mu^\tau(A_1 \cap F_t^{-1}B_1) &= \int_{a_1}^{a_2} m(t)(U_{A,B}(t+u-b_1) - U_{A,B}(t+u-b_1-h)) du \\ &= \int_{a_1}^{a_2} \frac{m(t)}{m(t+u-b_1-h)} \chi(t+u-b_1-h) du, \end{aligned}$$

where $\chi(t) = m(t)(U_{A,B}(t+h) - U_{A,B}(t))$ is bounded by Theorem 2.3. Also $m(t)/m(t+u-b_1-h)$ is bounded by Potter's bounds (see for instance [11]). Since $m(t)$ is regularly varying, we have $\lim_{t \rightarrow \infty} m(t)/m(t+u-b_1-h) = 1$ for each $u \in [0, 1]$. By Theorem 2.3, $\lim_{t \rightarrow \infty} \chi(t+u-b_1-h) = d_\beta \mu(A)\mu(B)h = d_\beta \mu(A)\mu^\tau(B_1)$ for each $u \in [0, 1]$. Hence the result follows from the bounded convergence theorem. \square

Remark 3.2. The result also holds for all sets of the form $F_r^{-1}A_1$ and $F_s^{-1}B_1$ for fixed $r, s > 0$. Indeed, by Corollary 3.1, using that $m(t) \sim m(t+s-r)$,

$$\begin{aligned} m(t)\mu^\tau(F_r^{-1}A_1 \cap F_{t+s}^{-1}B_1) &= m(t)\mu^\tau(A_1 \cap F_{t+s-r}^{-1}B_1) \\ &\rightarrow \mu^\tau(A_1)\mu^\tau(A_2) = \mu^\tau(F_r^{-1}A_1)\mu^\tau(F_s^{-1}A_2). \end{aligned}$$

Corollary 3.3 (Weak rational ergodicity). Assume the setting of Theorem 2.6 (alternatively Theorem 2.8), with $\beta \in [0, 1]$. Then

$$\lim_{t \rightarrow \infty} t^{-1}m(t) \int_0^t \mu^\tau(A_1 \cap F_x^{-1}B_1) dx = D_\beta \mu^\tau(A_1)\mu^\tau(B_1).$$

Proof. Continuing from (3.1) (with $h = b_2 - b_1$),

$$\begin{aligned} \int_0^t \mu^\tau(A_1 \cap F_x^{-1}B_1) dx &= \int_{a_1}^{a_2} \int_0^t (U_{A,B}(x+u-b_1) - U_{A,B}(x+u-b_1-h)) dx du \\ &= \int_{a_1}^{a_2} \int_0^t U_{A,B}(x+u-b_1) dx du - \int_{a_1}^{a_2} \int_{-h}^{t-h} U_{A,B}(x+u-b_1) dx du \\ &= \int_{a_1}^{a_2} \int_{t-h}^t U_{A,B}(x+u-b_1) dx du - \int_{a_1}^{a_2} \int_{-h}^0 U_{A,B}(x+u-b_1) dx du = I_1 + I_2. \end{aligned}$$

Now

$$t^{-1}m(t)I_1 = t^{-1}m(t)U_{A,B}(t) \int_{a_1}^{a_2} \int_{-h}^0 \frac{U_{A,B}(x+t+u-b_1)}{U_{A,B}(t)} dx du.$$

By Theorem 2.6, $U_{A,B}(t)$ is regularly varying so the integrand $U_{A,B}(x+t+u-b_1)/U_{A,B}(t)$ is bounded for x, u bounded and converges pointwise to 1 as $t \rightarrow \infty$. Hence

$$\lim_{t \rightarrow \infty} \int_{a_1}^{a_2} \int_{-h}^0 \frac{U_{A,B}(x+t+u-b_1)}{U_{A,B}(t)} dx du = (a_2 - a_1)h = (a_2 - a_1)(b_2 - b_1).$$

By Theorem 2.6, $t^{-1}m(t)U_{A,B}(t) = D_\beta \mu(A)\mu(B)(1+o(1))$. Hence, $\lim_{t \rightarrow \infty} t^{-1}m(t)I_1 = D_\beta \mu(A)\mu(B)(a_2 - a_1)(b_2 - b_1) = \mu^\tau(A_1)\mu^\tau(B_1)$. A simpler argument shows that $t^{-1}m(t)I_2 = o(1)$. \square

Proposition 3.4. Let $f : [0, \infty) \rightarrow \mathbb{R}$ be bounded and integrable on compact sets, and let $K \in \mathbb{R}$. Suppose that $\beta \in (0, 1)$, that $\ell(t)$ is slowly varying, and that

- (a) $\liminf_{t \rightarrow \infty} \ell(t)t^{1-\beta} f(t) \geq K$,
- (b) $\lim_{t \rightarrow \infty} \ell(t)t^{-\beta} \int_0^t f(x) dx = \beta^{-1}K$.

Then there exists a set $E \subset [0, \infty)$ of density zero such that $\lim_{t \rightarrow \infty, t \notin E} \ell(t)t^{1-\beta} f(t) = K$.

In particular, $\liminf_{t \rightarrow \infty} \ell(t)t^{1-\beta} f(t) = K$.

Proof. This is the continuous time analogue of [35, Proposition 8.2] (which is itself a version of [38, p. 65, Lemma 6.2]). We list the main steps which are proved exactly as in [35].

Step 1. Without loss of generality, $K = 0$ and $\ell(t)t^{1-\beta}$ is increasing.

Step 2. Define the nested sequence of sets $E_q = \{t > 0 : \ell(t)t^{1-\beta} f(t) > 1/q\}$, $q = 1, 2, \dots$. Then E_q has density zero for each q , i.e. $\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t 1_{E_q}(x) dx = 0$.

Step 3. By Step 2, we can choose $0 = i_0 < i_1 < i_2 < \dots$ such that $\frac{1}{t} \int_0^t 1_{E_q}(x) dx < 1/q$ for $t \geq i_{q-1}$, $q \geq 2$. Define $E = \bigcup_{q=1}^{\infty} E_q \cap (i_{q-1}, i_q)$. Then E has density zero and $\lim_{t \rightarrow \infty, t \notin E} \ell(t)t^{1-\beta} f(t) = 0$. \square

Corollary 3.5. *Assume the setting of Theorem 2.4 (alternatively Theorem 2.8), with $\beta \in (0, 1)$. Then*

- (i) $\liminf_{t \rightarrow \infty} m(t)\mu^\tau(A_1 \cap F_t^{-1}B_1) = d_\beta \mu^\tau(A_1)\mu^\tau(B_1)$, and
- (ii) *There exists a set $E \subset [0, \infty)$ of density zero such that $\lim_{t \rightarrow \infty, t \notin E} m(t)\mu^\tau(A_1 \cap F_t^{-1}B_1) = d_\beta \mu^\tau(A_1)\mu^\tau(B_1)$.*

Proof. We start from the conclusion of Theorem 2.4. Arguing as in the proof of Corollary 3.1, but with \lim replaced by \liminf and using Fatou's lemma instead of the bounded convergence theorem, we obtain

$$\liminf_{t \rightarrow \infty} \ell(t)t^{1-\beta} \mu^\tau(A_1 \cap F_t^{-1}B_1) \geq d_\beta \mu^\tau(A_1)\mu^\tau(B_1).$$

This is condition (a) in Proposition 3.4, and Corollary 3.3 is condition (b). Hence the result follows from Proposition 3.4. \square

4. Main results used in the proof of Theorem 2.3

The first result needed in the proof of the strong renewal theorem, Theorem 2.3, is an inversion formula for the symmetric measure

$$V_{A,B}(I) = \frac{1}{2}(U_{A,B}(I) + U_{A,B}(-I)).$$

Here, $U(-I) = U(\{x : -x \in I\})$ (with $U(-I) = 0$ if $I \subset [0, \infty]$). We find it convenient to adapt the formulation in [20, Section 4], but such an inversion formula goes back to [21] (see also [12, Chapter 10]).¹

By (H) and (S)(i), $\hat{T}(s) = (I - \hat{R}(s))^{-1}$ is a bounded operator on \mathcal{B} for all $s \in \mathbb{H} \setminus \{0\}$. Let $A, B \subset Y$ be measurable with $1_A \in \mathcal{B}$.

Proposition 4.1 (Analogue of [20, Inversion formula, Section 4]). *Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous compactly supported function with Fourier transform $\hat{g}(x) = \int_{-\infty}^{\infty} e^{ixb} g(b) db$ satisfying $\hat{g}(x) = O(x^{-2})$ as $x \rightarrow \infty$. Then for all $\lambda, t \in \mathbb{R}$,*

$$\int_{-\infty}^{\infty} e^{-i\lambda(x-t)} \hat{g}(x-t) dV_{A,B}(x) = \int_{-\infty}^{\infty} e^{-itb} g(b+\lambda) \operatorname{Re} \int_B \hat{T}(ib) 1_A d\mu db.$$

The second result required in the proof of Theorem 2.3 comes directly from [20] and does not require any modification in our set up. To state this result, for each $a > 0$ we let $\hat{g}_a(0) = 1$ and for $x \neq 0$, define

$$\hat{g}_a(x) = \frac{2(1 - \cos ax)}{a^2 x^2}.$$

Proposition 4.2 ([20, Lemma 8]). *Let $\{\mu_t, t > 0\}$ be a family of measures such that $\mu_t(I) < \infty$ for every compact set I and all t . Suppose that for some constant C ,*

$$\lim_{t \rightarrow \infty} \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) d\mu_t(x) = C \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) dx,$$

for all $a > 0, \lambda \in \mathbb{R}$. Then $\mu_t(I) \rightarrow C|I|$ for every bounded interval I , where $|I|$ denotes the length of I .

¹The result does not require any regular variation assumptions on $\mu(\tau > t)$, but we use the extra structure for simplicity.

Next, note that \hat{g}_a is the Fourier transform of

$$g_a(b) = \begin{cases} a^{-1}(1 - |b|/a) & |b| \leq a, \\ 0 & |b| > a. \end{cases}$$

The final result required in the proof of Theorem 2.3 is as follows.

Proposition 4.3. For all $a > 0$ and $\lambda \in \mathbb{R}$,

$$\lim_{t \rightarrow \infty} m(t) \int_{-\infty}^{\infty} e^{-itb} g_a(b + \lambda) \operatorname{Re} \int_B \hat{T}(ib) 1_A d\mu db = \pi d_\beta g_a(\lambda) \mu(A) \mu(B).$$

Proof of Theorem 2.3. With the convention $I + t = \{x : x - t \in I\}$, let

$$\mu_t(I) = 2m(t)V_{A,B}(I + t) = m(t)(U_{A,B}(I + t) + U_{A,B}(-I - t))$$

and note that for $I = [0, h]$ with $h > 0$,

$$m(t)(U_{A,B}(t + h) - U_{A,B}(t)) = \mu_t(I).$$

Now,

$$\begin{aligned} m(t) \int_{-\infty}^{\infty} e^{-i\lambda(x-t)} \hat{g}_a(x-t) dV_{A,B}(x) &= m(t) \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) dV_{A,B}(x+t) \\ &= \frac{1}{2} \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) d\mu_t(x). \end{aligned}$$

Since \hat{g}_a satisfies the assumptions of Proposition 4.1,

$$\int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) d\mu_t(x) = 2m(t) \int_{-\infty}^{\infty} e^{-itb} g_a(b + \lambda) \operatorname{Re} \int_B \hat{T}(ib) 1_A d\mu db.$$

By Proposition 4.3 together with the Fourier inversion formula $\int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) dx = 2\pi g_a(\lambda)$,

$$\lim_{t \rightarrow \infty} \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) d\mu_t(x) = 2\pi d_\beta g_a(\lambda) \mu(A) \mu(B) = d_\beta \int_{-\infty}^{\infty} e^{-i\lambda x} \hat{g}_a(x) dx \mu(A) \mu(B).$$

Hence, we have shown that the hypothesis of Proposition 4.2 holds with $C = d_\beta \mu(A) \mu(B)$. It now follows from Proposition 4.2 with $I = [0, h]$ that

$$m(t)(U_{A,B}(t + h) - U_{A,B}(t)) = \mu_t([0, h]) \rightarrow d_\beta \mu(A) \mu(B) h,$$

as $t \rightarrow \infty$. □

The proof of Propositions 4.1 and 4.3 are given in Section 6.

5. Prerequisites from operator renewal theory

In this section, we establish some estimates for $\hat{T} = (I - \hat{R})^{-1}$. The arguments closely follow [13, Section 4] (which was restricted to the case $\ell(t) = c + o(1)$ for some constant $c > 0$ and did not include the case $\beta = 1$).

The estimates are carried out under hypotheses (H) and (S)(i) in Section 5.1. The analogous results required under hypotheses (A) and (S)(i) are obtained in Section 5.2.

5.1. Estimates under hypotheses (H) and (S)(i)

Throughout this subsection, $\beta \in (0, 1]$ and $L > 0$ are fixed. We begin with $\gamma \in (0, \beta)$, $\delta \in (0, L)$ and $p > 1$ as in (H). During the subsection, the values of γ , δ and p change finitely many times; the changes in γ are arbitrarily small. Also $C > 0$ is a constant whose value changes finitely many times.

For $r \in [0, 1]$, let $\hat{T}_r(s) = (I - r\hat{R}(s))^{-1}$. Define

$$\tilde{\ell}(t) = \begin{cases} \ell(t) & \beta < 1, \\ \int_1^t \ell(s)s^{-1} ds & \beta = 1, \end{cases} \quad c_\beta = \begin{cases} i \int_0^\infty e^{-i\sigma} \sigma^{-\beta} d\sigma & \beta < 1, \\ 1 & \beta = 1. \end{cases}$$

Lemma 5.1.

- (a) $\|\hat{R}(s_1) - \hat{R}(s_2)\|_{\mathcal{B} \rightarrow L^p} \leq C|s_1 - s_2|^\gamma$ for all $s_1, s_2 \in \overline{\mathbb{H}}$.
 (b) There exists $r_0 < 1$ such that $\|\hat{T}_r(ib)\|_{\mathcal{B}} \leq C$ for all $|b| \in [\delta, L]$, $r \in [r_0, 1]$.
 (c) For all $\delta \leq b < b + h < L$,

$$\|\hat{T}(i(b+h)) - \hat{T}(ib)\|_{\mathcal{B} \rightarrow L^p} \leq Ch^\gamma.$$

(d) There exists a continuous family $\lambda(s)$, $s \in \overline{\mathbb{H}} \cap B_\delta(0)$, of simple eigenvalues for $\hat{R}(s) : \mathcal{B} \rightarrow \mathcal{B}$ with $\lambda(0) = 1$. In addition, the corresponding family of spectral projections $P(s)$ are bounded linear operators on \mathcal{B} for all $s \in \overline{\mathbb{H}} \cap B_\delta(0)$ and $\sup_{s \in \overline{\mathbb{H}} \cap B_\delta(0)} \|P(s)\|_{\mathcal{B}} < \infty$. Moreover,

$$\|P(s_1) - P(s_2)\|_{\mathcal{B} \rightarrow L^p} \leq C|s_1 - s_2|^\gamma \quad \text{for all } s_1, s_2 \in \overline{\mathbb{H}} \cap B_\delta(0).$$

- (e) Define the complementary projections $Q(s) = I - P(s)$. Then

$$\|(I - r\hat{R}(ib))^{-1}Q(ib)\|_{\mathcal{B}} \leq C \quad \text{for all } |b| < \delta, r \in [0, 1].$$

Proof. (a) Recall that $\hat{R}(s)v = R(e^{-s\tau}v)$. Since R is a positive operator,

$$|(\hat{R}(s_1) - \hat{R}(s_2))v| \leq R(|e^{-s_1\tau} - e^{-s_2\tau}| |v|) \leq 2|s_1 - s_2|^\gamma R(\tau^\gamma |v|).$$

By (H)(iii), $|(\hat{R}(s_1) - \hat{R}(s_2))v|_p \leq 2|s_1 - s_2|^\gamma |R(\tau^\gamma |v|)|_p \ll |s_1 - s_2|^\gamma \|v\|_{\mathcal{B}}$.

(b,c) Fix $b > 0$. It is immediate from hypothesis (S)(i) that $\|\hat{T}(ib)\|_{\mathcal{B}} < \infty$. Using also part (a), it follows from (H)(i,ii) and [29, Theorem 1] that there exists $h_0 > 0$, $r_0 < 1$ and $C > 0$ such that $\|\hat{T}_r(i(b+h))\|_{\mathcal{B}} \leq C$ and $\|\hat{T}(i(b+h)) - \hat{T}(ib)\|_{\mathcal{B} \rightarrow L^p} \leq C|h|^\gamma$ for all $|h| < h_0$, $r \in (r_0, 1]$. The desired estimates follow from compactness of $[\delta, L]$.

(d) This follows from (H)(i,ii) by [29, Corollary 1] exactly as in [13, Lemma 4.4] (with $\beta - \epsilon$ replaced by γ).

(e) By (H)(i,ii) and [29, Corollary 2], for $\delta > 0$ sufficiently small there exists $\rho \in (0, 1)$ such that $\|(rR(ib)Q(ib))^n\|_{\mathcal{B}} \leq \|(R(ib)Q(ib))^n\|_{\mathcal{B}} \leq C\rho^n$ for all $|b| < \delta$, $n \geq 0$. \square

Let $\zeta(s)$ denote the corresponding family of eigenfunctions normalized so that $\int_Y \zeta(s) d\mu = 1$. We have $\zeta(0) \equiv 1$ and $P(0)v = \int_Y v d\mu$ for all $v \in \mathcal{B}$. It is immediate that $\zeta(s)$ inherits the estimates obtained for $P(s)$. In particular, there is a constant $C > 0$ such that $|\zeta(s) - \zeta(0)|_p \leq C|s|^\gamma$ for all $s \in \overline{\mathbb{H}} \cap B_\delta(0)$.

Following [23] (see [13, Equation (4.2)]),

$$\lambda(s) = \int_Y e^{-s\tau} d\mu + \chi(s) \quad \text{where } \chi(s) = \int_Y (e^{-s\tau} - 1)(\zeta(s) - \zeta(0)) d\mu. \quad (5.1)$$

From now on, we fix $\delta \in (0, 1)$ so that all conclusions of Lemma 5.1 hold.

Proposition 5.2. Write $s = a + ib \in \overline{\mathbb{H}}$.

- (a) $1 - \int_Y e^{-s\tau} d\mu \sim c_\beta \tilde{\ell}(1/|s|)s^\beta$ as $s \rightarrow 0$.
 (b) When $\beta = 1$, $\text{Re}(1 - \int_Y e^{-ib\tau} d\mu) \sim \frac{\pi}{2} \ell(1/|b|)|b|$ as $b \rightarrow 0$.
 (c) $|\int_Y (e^{-i(b+h)\tau} - e^{-ib\tau}) d\mu| \leq C\tilde{\ell}(1/h)h^\beta$ for $0 < h < b < \delta$.

Proof. Part (a) is proved as in [36, Lemma 2.4] for $\beta < 1$. Suppose that $\beta = 1$ and let $G(x) = \mu(\tau > x)$. Then $1 - \int_Y e^{-s\tau} d\mu = s \int_0^\infty e^{-sx} (1 - G(x)) dx = sI_C(s) - isI_S(s)$, where

$$I_C(s) = \int_0^\infty e^{-ax} \cos bx (1 - G(x)) dx, \quad I_S(s) = \int_0^\infty e^{-ax} \sin bx (1 - G(x)) dx.$$

By [35, Proposition 6.2], we have for $a \geq |b|$ that

$$I_C(s) = \tilde{\ell}(1/a)(1 + o(1)) + O(|b|a^{-1}\ell(1/a)) = \tilde{\ell}(1/|s|)(1 + o(1)) + O(\ell(1/|s|)) \sim \tilde{\ell}(1/|s|).$$

Similarly, for $a \leq |b|$, we have $I_C(s) = \tilde{\ell}(1/|b|)(1 + o(1)) + O(a|b|^{-1}\ell(1/|b|)) \sim \tilde{\ell}(1/|s|)$. Hence $I_C(s) \sim \tilde{\ell}(1/|s|)$ as $s \rightarrow 0$. In the same way, it follows from [35, Proposition 6.2] that $|I_S(s)| \ll \ell(1/|s|)$. Part (a) for $\beta = 1$ follows immediately from these estimates. Moreover, $I_S(ib) \sim \frac{\pi}{2}\ell(1/|b|)\operatorname{sgn} b$ as $b \rightarrow 0$ by the proof of [35, Lemma 6.8]. Since $\operatorname{Re}(1 - \int_Y e^{-ib\tau} d\mu) = bI_S(ib)$, part (b) follows.

Finally, part (c) follows by the argument used in the proof of [22, Lemma 3.3.2]. □

Proposition 5.3.

- (a) $|\chi(s)| \leq C|s|^{\beta+\gamma}$ for $s \in \overline{\mathbb{H}} \cap B_\delta(0)$.
- (b) When $\beta > \frac{1}{2}$, $|\chi(i(b+h)) - \chi(ib)| \leq Cb^\beta h^\gamma$ for $0 < h < b < \delta$.

Proof. Choose $\epsilon > 0$ arbitrarily small and $r > 1$ such that $(\beta - \epsilon)r < \beta$ with conjugate exponent r' . Then $\tau^{(\beta-\epsilon)r} \in L^1$ and it follows from Hölder's inequality that

$$|\chi(s)| \leq 2|s|^{\beta-\epsilon} |\tau^{\beta-\epsilon}(\zeta(s) - 1)|_1 \leq 2|s|^{\beta-\epsilon} |\tau^{\beta-\epsilon}|_r |\zeta(s) - 1|_{r'} \ll |s|^{\beta-\epsilon+\gamma},$$

yielding part (a). Here we used that $|\zeta(s) - 1|_p = O(|s|^\gamma)$ for p as large as desired. Similarly,

$$\begin{aligned} |\chi(i(b+h)) - \chi(ib)| &\leq |(e^{i(b+h)\tau} - 1)(\zeta(i(b+h)) - \zeta(ib))|_1 + |(e^{ih\tau} - 1)(\zeta(ib) - 1)|_1 \\ &\ll (b+h)^{\beta-\epsilon} h^\gamma + h^{\beta-\epsilon} b^\gamma \ll b^\beta h^{\gamma-\epsilon}. \end{aligned}$$

(Note that $h^{\beta-\epsilon} b^\gamma = h^{\gamma-\epsilon} h^{\beta-\gamma} b^\gamma \leq h^{\gamma-\epsilon} b^\beta$ since $\gamma < \beta$ and $h < b$.) This proves part (b). □

Corollary 5.4. Write $s = a + ib \in \overline{\mathbb{H}}$.

- (a) $1 - \lambda(s) \sim c_\beta \tilde{\ell}(1/|s|)s^\beta$ as $s \rightarrow 0$.
- (b) When $\beta = 1$, $\operatorname{Re}(1 - \lambda(ib)) \sim \frac{\pi}{2}\ell(1/|b|)|b|$ as $b \rightarrow 0$.
- (c) When $\beta > \frac{1}{2}$, $|\lambda(i(b+h)) - \lambda(ib)| \leq C(\tilde{\ell}(1/h)h^\beta + b^\beta h^\gamma)$, for $0 < h < b < \delta$.
- (d) $|1 - r\lambda(ib)|^{-1} \leq C\tilde{\ell}(1/|b|)^{-1}|b|^{-\beta}$ for all $|b| < \delta$, $r \in [\frac{1}{2}, 1]$.
- (e) When $\beta = 1$, $|\operatorname{Re}(1 - r\lambda(ib))|^{-1} \leq C\ell(1/|b|)\tilde{\ell}(1/|b|)^{-2}|b|^{-1}$ for all $|b| < \delta$, $r \in [\frac{1}{2}, 1]$.

Proof. Parts (a) and (b) are immediate from (5.1) and Propositions 5.2(a,b) and 5.3(a). Part (c) follows from (5.1) and Propositions 5.2(c) and 5.3(b). By part (a),

$$|1 - r\lambda(ib)| \geq |\operatorname{Im}(1 - r\lambda(ib))| \geq \frac{1}{2}|\operatorname{Im}\lambda(ib)| \sim \frac{1}{2}|\operatorname{Im}(i^\beta c_\beta)|\tilde{\ell}(1/|b|)|b|^\beta,$$

yielding part (d). Using also that $\operatorname{Re}\lambda(ib) \in [0, 1]$ for $|b| < \delta$, we compute for $\beta = 1$ that

$$\begin{aligned} |\operatorname{Re}(1 - r\lambda(ib)^{-1})| &= \operatorname{Re}(1 - r\lambda(ib))|1 - r\lambda(ib)|^{-2} \\ &\leq \operatorname{Re}(1 - \lambda(ib))|r \operatorname{Im}(\lambda(ib))|^{-2} \leq 4\operatorname{Re}(1 - \lambda(ib))|r \operatorname{Im}(\lambda(ib))|^{-2}, \end{aligned}$$

so part (e) follows from parts (a) and (b). □

Lemma 5.5. $\hat{T}(s) = c_\beta^{-1}\tilde{\ell}(1/|s|)^{-1}s^{-\beta}(P(0) + E(s))$ for $s \in \overline{\mathbb{H}} \cap B_\delta(0)$, where $E(s)$ is a family of operators satisfying $\lim_{s \rightarrow 0} \|E(s)\|_{\mathcal{B} \rightarrow L^1} = 0$.

Proof. By Corollary 5.4(a), $(1 - \lambda(s))^{-1} \sim c_\beta^{-1} \tilde{\ell}(1/|s|)^{-1} s^{-\beta}$ as $s \rightarrow 0$. Also,

$$\hat{T}(s) = (1 - \lambda(s))^{-1} P(s) + (I - \hat{R}(s))^{-1} Q(s) = (1 - \lambda(s))^{-1} (P(0) + E(s)),$$

where

$$E(s) = P(s) - P(0) + (1 - \lambda(s))(I - \hat{R}(s))^{-1} Q(s). \tag{5.2}$$

By (H), $\|(I - \hat{R}(s))^{-1} Q(s)\|_{\mathcal{B}} = O(1)$. By Lemma 5.1(d), $\|P(s) - P(0)\|_{\mathcal{B} \rightarrow L^1} = O(|s|^\gamma)$. Hence $\|E(s)\|_{\mathcal{B} \rightarrow L^1} \ll |s|^\gamma + |s|^{\beta-\epsilon}$. \square

Lemma 5.6. *Let $\beta = 1$. Then $\operatorname{Re} \hat{T}(ib) = \frac{\pi}{2} \ell(1/|b|) \tilde{\ell}(1/|b|)^{-2} |b|^{-1} (P(0) + E(b))$ for $b \in \mathbb{R}$, $0 < |b| < \delta$, where $\lim_{b \rightarrow 0} \|E(b)\|_{\mathcal{B} \rightarrow L^1} = 0$.*

Proof. By Corollary 5.4(a,b),

$$\operatorname{Re}((1 - \lambda(ib))^{-1}) = \operatorname{Re}(1 - \lambda(ib)) |1 - \lambda(ib)|^{-2} \sim \frac{\pi}{2} \ell(1/|b|) \tilde{\ell}(1/|b|)^{-2} |b|^{-1}.$$

As in the proof of Lemma 5.5, $\operatorname{Re} \hat{T}(ib) = \{\operatorname{Re}((1 - \lambda(ib))^{-1})\} (P(0) + E(b))$ where

$$E(b) = \operatorname{Re}(1 - \lambda(ib)) \operatorname{Re}\{(1 - \lambda(ib))^{-1} (P(ib) - P(0)) + (I - R(ib))^{-1} Q(ib)\},$$

and

$$\|E(b)\|_{\mathcal{B} \rightarrow L^1} \ll \|P(ib) - P(0)\|_{\mathcal{B} \rightarrow L^1} + |1 - \lambda(ib)| \|(I - R(ib))^{-1} Q(ib)\|_{\mathcal{B} \rightarrow L^1} \ll |b|^{1-\epsilon},$$

completing the proof. \square

Corollary 5.7. *Let $\beta \leq 1$, $L > 0$. There are constants $r_0 < 1$ and $C > 0$ such that*

$$\|\operatorname{Re} \hat{T}_r(ib)\|_{\mathcal{B} \rightarrow L^1} \leq C \psi_\beta(|b|) \quad \text{for } 0 < |b| \leq L, r_0 \leq r \leq 1,$$

where

$$\psi_\beta(x) = \begin{cases} \ell(1/x)^{-1} x^{-\beta} & \beta < 1, \\ \ell(1/x) \tilde{\ell}(1/x)^{-2} x^{-1} & \beta = 1. \end{cases}$$

Proof. By Lemma 5.1(b), we can restrict to the range $|b| < \delta$ on which

$$\hat{T}_r(ib) = (I - r \hat{R}(ib))^{-1} = (1 - r \lambda(ib))^{-1} P(ib) + (I - r \hat{R}(ib))^{-1} Q(ib).$$

The result follows from the estimates for P , $(I - r \hat{R})^{-1} Q$ and $(1 - r \lambda)^{-1}$ obtained in Lemma 5.1(d,e) and Corollary 5.4(d,e). \square

Remark 5.8.

(a) Note that ψ_β is integrable on $[0, L]$ for all $\beta \leq 1$. This is clear for $\beta < 1$ while $\tilde{\ell}(1/x)^{-1}$ is an antiderivative for ψ_1 . In particular, $\sup_r |\operatorname{Re} \hat{T}_r(ib) 1_A|_1 \leq C \psi_\beta(b) \|1_A\|_{\mathcal{B}}$ which is integrable.

(b) By Karamata's theorem on integration of regularly varying sequences [11], $\tilde{\ell}$ is slowly varying and $\ell(x) = o(\tilde{\ell}(x))$ as $x \rightarrow \infty$ when $\beta = 1$. In particular, $\psi_\beta(b) \ll \tilde{\ell}(1/|b|)^{-1} |b|^{-\beta}$ for all $\beta \leq 1$.

Lemma 5.9. *Let $\beta \in (\frac{1}{2}, 1]$. For $0 < h < b < \delta$,*

$$\|\hat{T}(i(b+h)) - \hat{T}(ib)\|_{\mathcal{B} \rightarrow L^1} \leq C \{ \tilde{\ell}(1/b)^{-2} b^{-2\beta} \tilde{\ell}(1/h) h^\beta + b^{-\beta} h^\gamma \}.$$

Proof. Recall as in Lemma 5.5 that $\hat{T}(ib) = A_1(b) + A_2(b)$, where

$$A_1(b) = (1 - \lambda(ib))^{-1} P(ib), \quad A_2(b) = (I - \hat{R}(ib))^{-1} Q(ib).$$

Using Lemma 5.1(d) and Corollary 5.4(a,c),

$$\begin{aligned} \|A_1(b+h) - A_1(b)\|_{\mathcal{B} \rightarrow L^1} &\ll |1 - \lambda(i(b+h))|^{-1} \|P(i(b+h)) - P(ib)\|_{\mathcal{B} \rightarrow L^1} \\ &\quad + |1 - \lambda(ib)|^{-1} |1 - \lambda(i(b+h))|^{-1} |\lambda(i(b+h)) - \lambda(ib)| \|P(ib)\|_{\mathcal{B} \rightarrow L^1} \\ &\ll \tilde{\ell}(1/b) b^{-\beta} h^\gamma + \tilde{\ell}(1/b)^{-2} b^{-2\beta} (\tilde{\ell}(1/h) h^\beta + b^\beta h^\gamma) \\ &\ll \tilde{\ell}(1/b)^{-2} b^{-2\beta} \tilde{\ell}(1/h) h^\beta + b^{-\beta} h^{\gamma-\epsilon}. \end{aligned}$$

An argument from [33, Proposition 3.8] shows that $\|A_2(b+h) - A_2(b)\|_{\mathcal{B} \rightarrow L^1} \ll h^{\gamma-\epsilon}$, completing the proof. \square

5.2. Estimates under hypotheses (A) and (S)(i)

Let $\epsilon \in (0, \beta)$. Since $R : L^1 \rightarrow L^1$ is a contraction,

$$|(\hat{R}(s_1) - \hat{R}(s_2))v|_1 \leq |(e^{-s_1\tau} - e^{-s_2\tau})v|_1 \leq 2|s_1 - s_2|^{\beta-\epsilon} |\tau^{\beta-\epsilon} v|_1.$$

Choose $r > 1$ such that $(\beta - \epsilon)r < \beta$ with conjugate exponent r' . By Hölder's inequality and (A)(i), $|\tau^{\beta-\epsilon} v|_1 \leq |\tau^{\beta-\epsilon}|_r |v|_{r'} \ll \|v\|_{\mathcal{B}}$. Hence $\|\hat{R}(s_1) - \hat{R}(s_2)\|_{\mathcal{B} \rightarrow L^1} \ll |s_1 - s_2|^{\beta-\epsilon}$ for all $s_1, s_2 \in \overline{\mathbb{H}}$.

Using [29] as before, we deduce that the conclusions of Lemma 5.1 hold with L^p replaced by L^1 and γ replaced by $\beta - \epsilon$.

Proposition 5.10. *The conclusions of Lemmas 5.5 and 5.6 and Corollary 5.7 are unchanged under hypotheses (A) and (S)(i).*

Proof. It is immediate from hypothesis (A)(iii) that $|\chi(s)| \ll |s|^{\beta_+}$ where $\beta_+ > \beta$, and hence the proofs are unchanged. \square

Lemma 5.9 becomes:

Lemma 5.11. $\|\hat{T}(i(b+h)) - \hat{T}(ib)\|_{\mathcal{B} \rightarrow L^1} \leq C b^{-2\beta} h^{\beta-\epsilon}$ for all $0 < h < b < \delta$.

Proof. Since $\|\zeta(s)\|_{\mathcal{B}}$ is bounded, it follows again from Hölder's inequality that

$$\begin{aligned} |\chi(i(b+h)) - \chi(ib)| &\leq |(e^{i(b+h)\tau} - 1)(\zeta(i(b+h)) - \zeta(ib))|_1 + |(e^{ih\tau} - 1)(\zeta(ib) - 1)|_1 \\ &\leq 2|\zeta(i(b+h)) - \zeta(ib)|_1 + 2h^{\beta-\epsilon} |\tau^{\beta-\epsilon}|_r |(\zeta(ib) - 1)|_{r'} \ll h^{\beta-\epsilon}. \end{aligned}$$

Hence by (5.1) and Proposition 5.2(c), $|\lambda(i(b+h)) - \lambda(ib)| \ll h^{\beta-\epsilon}$. Now proceed as in the proof of Lemma 5.9. \square

The presence of the ϵ in Lemma 5.11 necessitates some alterations to the strategy in [20]. As in [13], we make use of the following refinement of Lemma 5.5.

Lemma 5.12. *Assume that $\mu(\tau > t) = ct^{-\beta} + O(t^{-q})$ where $c > 0$, $\beta \in (\frac{1}{2}, 1)$, $q > 1$. Then $c\hat{T}(ib) = c_\beta^{-1} b^{-\beta} P(0) + \tilde{E}(b)$ for $b \in [0, \delta)$, where $\|\tilde{E}(b)\|_{\mathcal{B} \rightarrow L^1} \leq C b^{-(2\beta-\beta_+)}$.*

Proof. A calculation using only the expression for $\mu(\tau > t)$ shows that $1 - \int_Y e^{-s\tau} d\mu = cc_\beta b^\beta + O(b)$ (see [13, Eq. (4.4)]). By (5.1) and the estimate $|\chi(s)| \ll |s|^{\beta_+}$ where $\beta_+ \in (\beta, 1)$, we obtain $1 - \lambda(ib) = cc_\beta b^\beta (1 + O(b^{\beta_+-\beta}))$. Hence

$$c(1 - \lambda(ib))^{-1} = c_\beta^{-1} b^{-\beta} (1 + O(b^{-(2\beta-\beta_+)})).$$

By (5.2), $c\hat{T}(ib) = c(1 - \lambda(ib))^{-1} P(0) + c(1 - \lambda(ib))^{-1} E(ib) = c_\beta^{-1} b^{-\beta} P(0) + c\tilde{E}(b)$ where $\tilde{E}(b) = (1 - \lambda(ib))^{-1} \times E(ib) + O(b^{-(2\beta-\beta_+)})$ and

$$E(ib) = P(ib) - P(0) + (1 - \lambda(ib))(I - \hat{R}(ib))^{-1} Q(ib) = O(b^{\beta-\epsilon}).$$

Hence $\tilde{E}(b) \ll b^{-\epsilon} + b^{-(2\beta-\beta_+)}$. Recall that $2\beta - \beta_+ > 2\beta - 1 > 0$, so we can choose $\epsilon \in (0, 2\beta - \beta_+)$ completing the proof. \square

6. Completion of the proof of Theorem 2.3

In this section, we give the proof of Propositions 4.1 and 4.3, thereby completing the proof of Theorem 2.3. In Sections 6.1 and 6.2, we assume hypotheses (H) and (S)(i). In Section 6.3, we show that the results remain true under hypotheses (A) and (S)(i).

6.1. Proof of Proposition 4.1

Fix $\beta \leq 1$. Throughout, we write U and V instead of $U_{A,B}$ and $V_{A,B}$. Following [12, Chapter 10] (see also [20, Section 4]), we define for $r \in (0, 1)$,

$$U_r(I) = \sum_{n=0}^{\infty} r^n \mu(y \in A \cap F^{-n}B : \tau_n(y) \in I),$$

$$V_r(I) = \frac{1}{2}(U_r(I) + U_r(-I)).$$

For $n \geq 0$, the Fourier transform of the distribution $G_n(x) = \mu(\tau_n(y) \leq x, y \in A \cap F^{-n}B)$ is given by $\int_Y 1_A 1_B \circ F^n e^{ib\tau_n} d\mu = \int_B \hat{R}(-ib)^n 1_A d\mu$. Hence

$$\begin{aligned} \int_{-\infty}^{\infty} e^{ibx} dV_r(x) &= \operatorname{Re} \int_0^{\infty} e^{ibx} dU_r(x) \\ &= \sum_{n=0}^{\infty} r^n \operatorname{Re} \int_B \hat{R}(ib)^n 1_A d\mu = \operatorname{Re} \int_B \hat{T}_r(ib) 1_A d\mu, \end{aligned}$$

where $\hat{T}_r(s) = (I - r\hat{R}(s))^{-1}$.

Let \hat{g} and g be as in the statement of Proposition 4.1. Note that dV_r is a finite measure and g is compactly supported, so $e^{ibx}g(b)$ lies in $L^1(dV_r \times db)$. Hence it follows from Fubini's theorem that for $r \in (0, 1)$,

$$\begin{aligned} \int_{-\infty}^{\infty} \hat{g}(x) dV_r(x) &= \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} e^{ibx} g(b) db \right\} dV_r(x) \\ &= \int_{-\infty}^{\infty} g(b) \left\{ \int_{-\infty}^{\infty} e^{ibx} dV_r(x) \right\} db = \int_{-\infty}^{\infty} g(b) \operatorname{Re} \int_B \hat{T}_r(ib) 1_A d\mu db. \end{aligned}$$

Replacing $g(b)$ by $g_1(b) = e^{-ibt}g(b + \lambda)$ and $\hat{g}(x)$ by $\hat{g}_1(x) = \int_{-\infty}^{\infty} e^{ibx} g_1(b) db = e^{-i\lambda(x-t)}\hat{g}(x-t)$, we obtain

$$\int_{-\infty}^{\infty} e^{-i\lambda(x-t)}\hat{g}(x-t) dV_r(x) = \int_{-\infty}^{\infty} e^{-ibt}g(b + \lambda) \operatorname{Re} \int_B \hat{T}_r(ib) 1_A d\mu db. \quad (6.1)$$

It remains to justify passing to the limit $r \rightarrow 1_-$ on both sides of (6.1).

First, we consider the left-hand side of (6.1). Since $\tau \geq 1$, we have $U(x) = U([0, x]) \leq \sum_{n=0}^{\infty} \mu(\tau_n \leq x) \leq x + 1$ for all x . Integrating by parts,

$$\int_1^{\infty} x^{-2} dU(x) = -U(1) + 2 \int_1^{\infty} U(x)x^{-3} dx < \infty.$$

Hence $\int_{|x| \geq 1} x^{-2} dV(x) < \infty$. Since $\int_{-1}^1 |\hat{g}(x-t)| dV(x) < \infty$ and $\hat{g}(x-t) = O(x^{-2})$ for each fixed t , it follows that $f(x) = e^{-i\lambda(x-t)}\hat{g}(x-t)$ is integrable with respect to $dV(x)$. But $V_r(I) \nearrow V(I)$ as $r \rightarrow 1_-$ for every measurable I , so $\lim_{r \rightarrow 1_-} \int_{-\infty}^{\infty} f(x) dV_r(x) = \int_{-\infty}^{\infty} f(x) dV(x)$ which is the required result for the left-hand side.

Finally, we consider the right-hand side of (6.1). Choose $L > 0$ such that $\operatorname{supp} g \in [-L, L]$. By Remark 5.8(a), $|\operatorname{Re} \hat{T}_r(ib) 1_A| \ll \psi_\beta(b) \|1_A\|_B$ for $|b| \leq L + |\lambda|$, where ψ_β is integrable. Hence the desired limit as $r \rightarrow 1_-$ follows from the dominated convergence theorem.

6.2. Proof of Proposition 4.3

Fix $\beta \in (\frac{1}{2}, 1]$. We follow the proof of [20, Theorem 1] (an adaptation of the argument in [22]). Let $W(b) = \text{Re} \int_B \hat{T}(ib) 1_A d\mu$.

Fix $\omega > 1$ and write $\int_{-\infty}^{\infty} e^{-itb} g_a(b + \lambda) \text{Re} \int_B \hat{T}(ib) 1_A d\mu db = I_1(t, \omega) + I_2(t, \omega)$ where

$$I_1(t, \omega) = \int_{-\omega/t}^{\omega/t} e^{-itb} g_a(b + \lambda) W(b) db, \quad I_2(t, \omega) = \int_{|b| > \omega/t} e^{-itb} g_a(b + \lambda) W(b) db.$$

Proposition 4.3 follows immediately from the estimates for $I_1(t, \omega)$ and $I_2(t, \omega)$ below.

Lemma 6.1. $\lim_{\omega \rightarrow \infty} \lim_{t \rightarrow \infty} m(t) I_1(t, \omega) = \pi d_{\beta} g_a(\lambda) \mu(A) \mu(B)$.

Proof. It follows from the definition of g_a that $|g_a(b_1) - g_a(b_2)| \leq a^{-2} |b_1 - b_2|$. Hence

$$\begin{aligned} \left| I_1(t, \omega) - g_a(\lambda) \int_{-\omega/t}^{\omega/t} e^{-itb} W(b) db \right| &\leq \int_{-\omega/t}^{\omega/t} |g_a(b + \lambda) - g_a(\lambda)| |W(b)| db \\ &\leq 2a^{-2} \omega t^{-1} \int_0^{\omega/t} |W(b)| db. \end{aligned}$$

By Remark 5.8(a), $\int_0^{\omega/t} |W(b)| db \ll \|1_A\|$ for $t > \omega/\delta$. Hence

$$\lim_{t \rightarrow \infty} m(t) I_1(t, \omega) = 2g_a(\lambda) \lim_{t \rightarrow \infty} m(t) \int_0^{\omega/t} W(b) \cos tb db.$$

For $\beta < 1$, define $\xi(b) = \mu(A)\mu(B) + \int_B E(ib) 1_A d\mu$ where E is as in Lemma 5.5. In particular, $|\xi(b)| \leq \|1_A\| + |E(ib) 1_A| \ll \|1_A\|$ and $|\xi(b) - \mu(A)\mu(B)| \leq \|E(ib)\|_{\mathcal{B} \rightarrow L^1} \|1_A\| \rightarrow 0$ as $b \rightarrow 0$. Hence

$$\begin{aligned} m(t) \int_0^{\omega/t} W(b) \cos tb db &= \ell(t) t^{1-\beta} \text{Re} \left\{ c_{\beta}^{-1} \int_0^{\omega/t} \ell(1/b)^{-1} b^{-\beta} \xi(b) \cos tb db \right\} \\ &= \text{Re} \left\{ c_{\beta}^{-1} \int_0^{\omega} [\ell(t)/\ell(t/b)] b^{-\beta} \xi(b/t) \cos b db \right\}. \end{aligned}$$

By the dominated convergence theorem,

$$\lim_{t \rightarrow \infty} m(t) \int_0^{\omega/t} W(b) \cos tb db = (\text{Re } c_{\beta}^{-1}) \int_0^{\omega} b^{-\beta} \cos b db \mu(A) \mu(B),$$

and the result for $\beta < 1$ follows.

Now suppose that $\beta = 1$ and recall that $\psi_1(b) = \ell(1/b) \tilde{\ell}(1/b)^{-2} b^{-1}$. By Lemma 5.6,

$$m(t) \int_0^{\omega/t} W(b) \cos tb db = \tilde{\ell}(t) \frac{\pi}{2} \int_0^{\omega/t} \psi_1(b) \xi(b) \cos tb db,$$

where $\xi(b)$ has the same properties as before. Now

$$\begin{aligned} \tilde{\ell}(t) \int_0^{\omega/t} \psi_1(b) \xi(b) db &= \tilde{\ell}(t) \int_0^{\omega/t} \psi_1(b) (\mu(A)\mu(B) + o(1)) db \\ &= \tilde{\ell}(t) \tilde{\ell}(t/\omega)^{-1} (\mu(A)\mu(B) + o(1)) \rightarrow \mu(A)\mu(B). \end{aligned}$$

Next,

$$\tilde{\ell}(t) \int_0^{\omega/t} \psi_1(b) \xi(b) (\cos tb - 1) db = \int_0^{\omega} \frac{\tilde{\ell}(t)}{\tilde{\ell}(t/\sigma)} \frac{\ell(t/\sigma)}{\tilde{\ell}(t/\sigma)} \xi(\sigma/t) \frac{\cos \sigma - 1}{\sigma} d\sigma.$$

By Remark 5.8(b), $\tilde{\ell}$ is slowly varying and $\ell(x) = o(\tilde{\ell}(x))$ as $x \rightarrow \infty$. By Potter's bounds, the integrand is dominated by $\sigma^{1-\epsilon}$ for any $\epsilon > 0$, so the integrand converges to zero pointwise and $\tilde{\ell}(t) \int_0^{\omega/t} \psi_1(b)\xi(b)(\cos tb - 1) db \rightarrow 0$ as $t \rightarrow \infty$. Hence $\lim_{t \rightarrow \infty} m(t) \int_0^{\omega/t} W(b) \cos tb db = \frac{\pi}{2} \mu(A)\mu(B)$ yielding the result for $\beta = 1$. \square

Lemma 6.2. *Let $\beta' \in (\frac{1}{2}, \beta)$. Then $\limsup_{t \rightarrow \infty} m(t)I_2(t, \omega) = O(\omega^{-(2\beta'-1)})$.*

Proof. It follows from evenness of g_a and $W(b)$, together with the fact that $\text{supp } g_a = [-a, a]$, that

$$I_2(t, \omega) = \int_{b > \omega/t} [e^{-itb} g_a(b + \lambda) + e^{itb} g_a(b - \lambda)] W(b) db = \int_{\omega/t}^{a+|\lambda|} h(b) W(b) db,$$

where $h(b) = e^{-itb} g_a(b + \lambda) + e^{itb} g_a(b - \lambda)$. Continuing as on [20, p. 278] down as far as [20, Equation (5.14)], we obtain $m(t)|I_2(t, \omega)| \leq a^{-1} J_1(t, \omega) + \pi a^{-2} J_2(t, \omega) + a^{-1} J_3(t, \omega)$, where

$$\begin{aligned} J_1(t, \omega) &= m(t) \int_{(\omega-\pi)/t}^{\omega/t} |W(b + \pi/t)| db, & J_2(t, \omega) &= m(t)t^{-1} \int_{\omega/t}^{a+|\lambda|} |W(b)| db, \\ J_3(t, \omega) &= m(t) \int_{\omega/t}^{a+|\lambda|} |W(b + \pi/t) - W(b)| db. \end{aligned}$$

By Remark 5.8(a), W is integrable on $[0, a + |\lambda|]$ so $J_2(t, \omega) \ll \tilde{\ell}(t)t^{-\beta} \rightarrow 0$ as $t \rightarrow \infty$. By Lemma 5.5, for $\beta < 1$,

$$J_1(t, \omega) \ll \ell(t)t^{1-\beta} \int_{\omega/t}^{(\omega+\pi)/t} \ell(1/b)^{-1} b^{-\beta} db = \int_{\omega}^{\omega+\pi} (\ell(t)/\ell(t/\sigma))\sigma^{-\beta} d\sigma \ll \omega^{-(\beta-\epsilon)},$$

for any $\epsilon > 0$ by Potter's bounds. By Lemma 5.6 and Remark 5.8(b), for $\beta = 1$,

$$J_1(t, \omega) \ll \tilde{\ell}(t) \int_{\omega/t}^{(\omega+\pi)/t} \psi_1(b) db = \tilde{\ell}(t) \{ \tilde{\ell}(t/(\omega + \pi))^{-1} - \tilde{\ell}(t/\omega) \} \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

By Lemma 5.9 with $h = \pi/t$,

$$J_3(t, \omega) \ll \tilde{\ell}(t)^2 t^{1-2\beta} \int_{\omega/t}^{\infty} \tilde{\ell}(1/b)^{-2} b^{-2\beta} db + t^{1-\beta+\epsilon-\gamma} \int_0^{a+|\lambda|} b^{-\beta} db = J_{3,1} + J_{3,2}.$$

By Potter's bounds,

$$J_{3,1} = \int_{\omega}^{\infty} [\tilde{\ell}(t)/\tilde{\ell}(t/\sigma)]^2 \sigma^{-2\beta} d\sigma \ll \int_{\omega}^{\infty} \sigma^{-2\beta'} d\sigma \ll \omega^{-(2\beta'-1)}.$$

Finally, since we are in the case $\beta > \frac{1}{2}$, we can choose $\gamma \in (1 - \beta, \beta)$ in hypothesis (H). Hence $J_{3,2} \ll t^{1-\beta+\epsilon-\gamma} = o(1)$ as $t \rightarrow \infty$ for $\epsilon > 0$ sufficiently small. \square

6.3. Modified argument under hypotheses (A) and (S)(i)

Assume hypotheses (A) and (S)(i) and that $\mu(\tau > t) = ct^{-\beta} + O(t^{-q})$ where $c > 0, \beta \in (\frac{1}{2}, 1), q > 1$. Recall that $\beta_+ > \beta$.

First, we note by Proposition 5.10 that Corollary 5.7 is unchanged. Hence the proof of Proposition 4.1 is unchanged.

For Proposition 4.3, we adopt a different strategy from before. Instead of considering $\lim_{\omega \rightarrow \infty} \limsup_{t \rightarrow \infty} I_r(t, \omega)$ for $r = 1, 2$, we consider $\lim_{t \rightarrow \infty} I_r(t, t^\kappa)$ for a suitable choice of $\kappa > 0$.

Lemma 6.3. $\lim_{t \rightarrow \infty} m(t)I_1(t, t^\kappa) = \pi d_\beta g_a(\lambda)\mu(A)\mu(B)$ for all $\kappa > 0$.

Proof. Following the proof of Lemma 6.1 and using Lemma 5.5 and Proposition 5.10,

$$\left| m(t)I_1(t, \omega) - 2m(t)g_a(\lambda) \int_0^{\omega/t} W(b) \cos tb db \right| \ll \omega t^{-\beta} \int_0^{\omega/t} b^{-\beta} db \ll \omega^{2-\beta} t^{-1}.$$

By Lemma 5.12,

$$\begin{aligned} m(t) \int_0^{\omega/t} W(b) \cos tb \, db &= t^{1-\beta} \int_0^{\omega/t} (\operatorname{Re} c_\beta^{-1} b^{-\beta} \mu(A)\mu(B) + O(b^{-(2\beta-\beta_+)}) \cos tb \, db \\ &= \operatorname{Re} c_\beta^{-1} \int_0^\omega b^{-\beta} \cos b \, db \mu(A)\mu(B) + O(t^{-(\beta_+-\beta)} \omega^{1-2\beta+\beta_+}). \end{aligned}$$

Finally, a calculation (see for example [35, Proposition 9.5]) shows that $\int_0^\omega b^{-\beta} \cos b \, db = \Gamma(1-\beta) \sin(\beta\pi/2) + O(\omega^{-\beta})$. Hence the result follows with $\omega = t^\kappa$ for any $\kappa > 0$. □

Lemma 6.4. $\lim_{t \rightarrow \infty} m(t)I_2(t, t^\kappa) = 0$ for all $\kappa > 0$ sufficiently large.

Proof. We use the same decomposition $m(t)|I_2(t, \omega)| \leq a^{-1}J_1(t, \omega) + \pi a^{-2}J_2(t, \omega) + a^{-1}J_3(t, \omega)$ as in the proof of Lemma 6.2. By Proposition 5.10, we still have $J_1(t, \omega) \ll \omega^{-(\beta-\epsilon)}$ and $J_2(t, \omega) \ll t^{-\beta}$. By Lemma 5.11 with $h = \pi/t$,

$$J_3(t, \omega) \ll t^{1-\beta} t^{-(\beta-\epsilon)} \int_{\omega/t}^\infty b^{-2\beta} \, db \ll t^\epsilon \omega^{-(2\beta-1)},$$

for any choice of $\epsilon > 0$. Now take $\omega = t^\kappa$ with $\epsilon < \kappa(2\beta - 1)$. □

7. Proof of the local limit theorem with error term

In this section, we prove Theorem 2.7. The proof combines results from Section 5 with arguments from [39]. (A related argument [3, Theorem 6.3] based on [12] gives a similar conclusion but without the error term.)

For ease of exposition, we assume hypotheses (H) and (S)(ii) throughout. However, Lemma 5.9 is not required in this section, so we can just as well use hypothesis (A) instead of hypothesis (H) by Proposition 5.10. Recall that $q_\beta(t) = \frac{1}{2\pi} \int_{-\infty}^\infty e^{ibt} e^{-c_\beta|b|^\beta} \, db$ where $c_\beta = i \int_0^\infty e^{-i\sigma} \sigma^{-\beta} \, d\sigma$.

In Section 4, we made use of the family of kernels $g_a(b) = a^{-1}g(b/a)$ with Fourier transforms $\hat{g}_a(x) = \hat{g}(ax)$, where

$$g(b) = \begin{cases} 1 - |b| & |b| \leq 1, \\ 0 & |b| > 1 \end{cases} \quad \text{and} \quad \hat{g}(x) = \frac{2(1 - \cos x)}{x^2}.$$

Since the current section closely follows [39] which uses slightly different conventions, we now use $k_a(b) = g(ab)$ with transforms $\hat{k}_a(x) = \frac{1}{2\pi} a^{-1} \hat{g}(b/a)$. (In [39], \hat{k}_a is called K_a .)

Let

$$\mu_n(I) = \mu(y \in A \cap F^{-n}B : \tau_n(y) \in I),$$

and define

$$V_n(t, h, a) = \int_{-\infty}^\infty \hat{k}_a(t-t') \mu_n([d_n t', d_n(t'+h)]) \, dt'.$$

Lemma 7.1. *Let $L > 0$. Then*

$$V_n(t, h, a) = h \{ q_\beta(t) \mu(A)\mu(B) + e(n, h, a, t) \} \quad \text{for } a \geq (Ld_n)^{-1},$$

where $e(n, h, a, t) \rightarrow 0$ as $n \rightarrow \infty, h \rightarrow 0$ and $a \rightarrow 0$, uniformly in $t \in \mathbb{R}$.

Proof. In fact, we show that

$$|V_n(t, h, a) - h q_\beta(t) \mu(A)\mu(B)| \leq \text{const.} h \{ e_1(n) + e_2(h) + e_3(a) \}$$

where $\lim_{n \rightarrow \infty} e_1(n) = \lim_{h \rightarrow 0} e_2(h) = \lim_{a \rightarrow 0} e_3(a) = 0$.

As in Section 5, we write $\hat{R}(ib) = \lambda(ib)P(ib) + \tilde{Q}(b)$ for $|b| \leq \delta$, where $\tilde{Q}(b) = R(ib)Q(ib)$. Then

$$\hat{R}(ib)^n = \lambda(ib)^n P(0) + \lambda(ib)^n (P(ib) - P(0)) + \tilde{Q}(b)^n. \tag{7.1}$$

Moreover, there exist constants $C > 0$, $\gamma > 1 - \beta$, $\alpha_1 \in (0, 1)$, where

$$\|P(ib) - P(0)\|_{\mathcal{B} \rightarrow L^1} \leq C|b|^\gamma, \quad \|\tilde{Q}(b)^n\|_{\mathcal{B}} \leq C\alpha_1^n, \quad \text{for all } |b| \leq \delta, n \geq 1. \quad (7.2)$$

Also, we can choose $C > 0$, $\alpha_1 \in (0, 1)$ so that

$$\|\hat{R}(ib)^n\|_{\mathcal{B}} \leq C\alpha_1^n \quad \text{for all } b \in [\delta, L], n \geq 1. \quad (7.3)$$

(Such an estimate for fixed $b > 0$ holds by (S)(ii). The uniform estimate follows from [29, Corollary 2, part 2].)

By Corollary 5.4(a), $1 - \lambda(ib) \sim c_\beta \ell(1/|b|)b^\beta$. Hence

$$\lambda(ib) \sim e^{-c_\beta \ell(1/|b|)|b|^\beta} \quad \text{as } b \rightarrow 0, \quad \lim_{n \rightarrow \infty} \lambda(id_n^{-1}b)^n = e^{-c_\beta |b|^\beta}. \quad (7.4)$$

Let $\beta' \in (0, \beta)$. By (7.4) and Potter's bounds, for each fixed n , there exists $C_1(n), C_2(n) > 0$ such that $|\lambda(id_n^{-1}b)|^n \leq C_1(n)e^{-C_2(n)|b|^{\beta'}}$ for all $|b| \leq \delta d_n$. Also, there exists $n_0 \geq 1$ such that $|\lambda(id_n^{-1}b)|^n \leq 2e^{-c_\beta |b|^\beta}$ for all $|b| \leq \delta d_n, n \geq n_0$. Hence there exists $C_1, C_2 > 0$ such that

$$|\lambda(id_n^{-1}b)|^n \leq C_1 e^{-C_2 |b|^{\beta'}} \quad \text{for all } |b| \leq \delta d_n, n \geq 1. \quad (7.5)$$

Now $\hat{k}_a(t) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-ibt} k_a(b) db$ and hence

$$\begin{aligned} V_n(t, h, a) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-ib(t-t')} k_a(b) db \int_{A \cap F^{-n}B} \mathbf{1}_{\{\tau_n \in [d_n t', d_n(t'+h)]\}} d\mu dt' \\ &= \frac{1}{2\pi} \int_{|b| \leq a^{-1}} e^{-ibt} k_a(b) \int_{A \cap F^{-n}B} \int_{d_n^{-1}\tau_n - h}^{d_n^{-1}\tau_n} e^{ibt'} dt' d\mu db \\ &= \frac{1}{2\pi} \int_{|b| \leq a^{-1}} e^{-itb} k_a(b) (1 - e^{-ihb})(ib)^{-1} \int_{A \cap F^{-n}B} e^{id_n^{-1}b\tau_n} d\mu db \\ &= \frac{h}{2\pi} \int_{|b| \leq a^{-1}} e^{-itb} G(b, h, a) \int_B \hat{R}(id_n^{-1}b)^n \mathbf{1}_A d\mu db, \end{aligned}$$

where $G(b, h, a) = k_a(b)(1 - e^{-ihb})(ihb)^{-1}$.

Note that $|G(b, h, a)| \leq 1$. Using (7.3) and that $a \geq (Ld_n)^{-1}$,

$$\begin{aligned} \left| \int_{\delta d_n \leq |b| \leq a^{-1}} e^{-itb} G(b, h, a) \int_B \hat{R}(id_n^{-1}b)^n \mathbf{1}_A d\mu db \right| &\leq \|1_A\|_{\mathcal{B}} \int_{\delta d_n \leq |b| \leq Ld_n} \|\hat{R}(id_n^{-1}b)^n\|_{\mathcal{B}} db \\ &= \|1_A\|_{\mathcal{B}} d_n \int_{\delta \leq |b| \leq L} \|\hat{R}(ib)^n\|_{\mathcal{B}} db \leq C \|1_A\|_{\mathcal{B}} d_n \alpha_1^n. \end{aligned}$$

Hence this term can be incorporated into $e_1(n)$.

It remains to analyse

$$\frac{h}{2\pi} \int_{|b| \leq \delta d_n} e^{-itb} G(b, h, a) \int_B \hat{R}(id_n^{-1}b)^n \mathbf{1}_A d\mu db = \frac{h}{2\pi} (I_1 + I_2 + I_3),$$

where by (7.1),

$$\begin{aligned} I_1 &= \int_{|b| \leq \delta d_n} e^{-itb} G(b, h, a) \int_B \lambda(id_n^{-1}b)^n P(0) \mathbf{1}_A d\mu db, \\ I_2 &= \int_{|b| \leq \delta d_n} e^{-itb} G(b, h, a) \int_B \lambda(id_n^{-1}b)^n (P(id_n^{-1}b) - P(0)) \mathbf{1}_A d\mu db, \\ I_3 &= \int_{|b| \leq \delta d_n} e^{-itb} G(b, h, a) \int_B \tilde{Q}(d_n^{-1}b)^n \mathbf{1}_A d\mu db. \end{aligned}$$

By (7.2) and (7.5),

$$|I_2| \leq \int_{|b| \leq \delta d_n} C_1 e^{-C_2|b|^{\beta'}} C |d_n^{-1}b|^\gamma \|1_A\|_{\mathcal{B}} db \leq CC_1 \|1_A\|_{\mathcal{B}} d_n^{-\gamma} \int_{-\infty}^{\infty} |b|^\gamma e^{-C_2|b|^{\beta'}} db \ll d_n^{-\gamma},$$

and

$$|I_3| \leq d_n \int_{|b| \leq \delta} C \alpha_1^n \|1_A\|_{\mathcal{B}} db \ll d_n \alpha_1^n.$$

Again, these terms can be incorporated into $e_1(n)$.

This leaves the term $I_1 = I'_1 \mu(A)\mu(B)$ where $I'_1 = \int_{|b| \leq \delta d_n} e^{-itb} G(b, h, a) \lambda(i d_n^{-1}b)^n db$. Write $I'_1 = J_1 + J_2 + J_3$ where

$$J_1 = \int_{|b| \leq \delta d_n} e^{-itb} k_a(b) \{ (1 - e^{-ihb})(ihb)^{-1} - 1 \} \lambda(i d_n^{-1}b)^n db,$$

$$J_2 = \int_{|b| \leq \delta d_n} e^{-itb} (k_a(b) - 1) \lambda(i d_n^{-1}b)^n db,$$

$$J_3 = \int_{|b| \leq \delta d_n} e^{-itb} \lambda(i d_n^{-1}b)^n db.$$

Since $|(1 - e^{-ihb})(ihb)^{-1} - 1| \leq \frac{1}{2}h|b|$ it follows from (7.5) that

$$|J_1| \leq h \int_{-\infty}^{\infty} C_1 e^{-C_2|b|^{\beta'}} |b| db \ll h.$$

Also,

$$|J_2| \leq \int_{-\infty}^{\infty} |k_a(b) - 1| C_1 e^{-C_2|b|^{\beta'}} db,$$

which converges to zero by the dominated convergence theorem as $a \rightarrow 0$. These are the sole contributions to e_2 and e_3 respectively.

Finally,

$$|J_3 - 2\pi q_\beta(t)| \leq \int_{|b| \leq \delta d_n} |\lambda(i d_n^{-1}b)^n - e^{-c_\beta|b|^\beta}| db + \int_{|b| \geq \delta d_n} e^{-c_\beta|b|^\beta} db,$$

which converges to zero by (7.4), (7.5) and the dominated convergence theorem as $n \rightarrow \infty$. □

Lemma 7.2. *Let $\epsilon > 0$ and $L > 0$. There exists $n_0 \geq 1$ and $h_0 > 0$ such that*

$$h(q_\beta(t)\mu(A)\mu(B) - \epsilon) \leq \mu_n([d_n t, d_n(t+h)]) \leq h(q_\beta(t)\mu(A)\mu(B) + \epsilon),$$

for all $n \geq n_0, h \in [(Ld_n)^{-1}, h_0], t \in \mathbb{R}$.

Proof. Let $\tilde{q}_\beta = q_\beta \mu(A)\mu(B)$. Since q_β is the Fourier transform of an L^1 function, \tilde{q}_β is uniformly continuous and bounded. Let $q_\infty = |\tilde{q}_\beta|_\infty$ and choose $h_1 \in (0, 1)$ such that $|\tilde{q}_\beta(t) - \tilde{q}_\beta(t')| \leq \frac{1}{4}\epsilon$ whenever $|t - t'| \leq h_1$.

For $\epsilon_1 > 0$, set $\epsilon_2 = \int_{|x| > 1/\epsilon_1} \hat{k}_1(x) dx$. We choose $\epsilon_1 \in (0, \frac{1}{6})$ sufficiently small that

$$\left(q_\infty + 2\epsilon_1 q_\infty + \frac{1}{2}\epsilon \right) (1 - \epsilon_2)^{-1} - q_\infty \leq \epsilon, \quad 2\epsilon_1 q_\infty + \epsilon_2 (q_\infty + \epsilon) \leq \frac{1}{2}\epsilon. \tag{7.6}$$

By Lemma 7.1, there exists $n_0 \geq 1$ and $h_0 \in (0, h_1)$ such that for all $n \geq n_0, h \in [(Ld_n)^{-1}, h_0], t \in \mathbb{R}$,

$$\begin{aligned} V_n(t - \epsilon_1 h, h(1 + 2\epsilon_1), \epsilon_1^2 h) &\leq h(1 + 2\epsilon_1) \tilde{q}_\beta(t - \epsilon_1 h) + \frac{1}{6}\epsilon h \\ &\leq h(1 + 2\epsilon_1) \left(\tilde{q}_\beta(t) + \frac{1}{4}\epsilon \right) + \frac{1}{6}\epsilon h \leq h \left(\tilde{q}_\beta(t) + 2\epsilon_1 q_\infty + \frac{1}{2}\epsilon \right), \end{aligned} \tag{7.7}$$

where we used the constraint $\epsilon_1 \leq \frac{1}{6}$. Also, we can ensure that

$$\begin{aligned} V_n(t + \epsilon_1 h, h(1 - 2\epsilon_1), \epsilon_1^2 h) &\geq h(1 - 2\epsilon_1) \tilde{q}_\beta(t + \epsilon_1 h) - \frac{1}{4} \epsilon h \\ &\geq h(1 - 2\epsilon_1) \left(\tilde{q}_\beta(t) - \frac{1}{4} \epsilon \right) - \frac{1}{4} \epsilon h \geq h \left(\tilde{q}_\beta(t) - 2\epsilon_1 q_\infty - \frac{1}{2} \epsilon \right). \end{aligned} \quad (7.8)$$

Now, for $|t'| \leq \epsilon_1 h$,

$$\begin{aligned} \mu_n([d_n(t + \epsilon_1 h - t'), d_n(t - \epsilon_1 h - t' + h)]) &\leq \mu_n([d_n t, d_n(t + h)]) \\ &\leq \mu_n([d_n(t - \epsilon_1 h - t'), d_n(t + \epsilon_1 h - t' + h)]). \end{aligned}$$

Also $\int_{-\infty}^{\infty} \hat{k}_1 dx = 1$, so

$$1 - \epsilon_2 = \int_{|x| \leq 1/\epsilon_1} \hat{k}_1(x) dx = \epsilon_1^2 h \int_{|x| \leq 1/\epsilon_1} \hat{k}_{\epsilon_1^2 h}(\epsilon_1^2 h x) dx = \int_{|x| \leq \epsilon_1 h} \hat{k}_{\epsilon_1^2 h}(x) dx.$$

Hence

$$\begin{aligned} V_n(t - \epsilon_1 h, h(1 + 2\epsilon_1), \epsilon_1^2 h) &= \int_{-\infty}^{\infty} \hat{k}_{\epsilon_1^2 h}(t') \mu_n([d_n(t - \epsilon_1 h - t'), d_n(t + \epsilon_1 h - t' + h)]) dt' \\ &\geq \int_{|t'| \leq \epsilon_1 h} \hat{k}_{\epsilon_1^2 h}(t') \mu_n([d_n(t - \epsilon_1 h - t'), d_n(t + \epsilon_1 h - t' + h)]) dt' \\ &\geq \int_{|t'| \leq \epsilon_1 h} \hat{k}_{\epsilon_1^2 h}(t') \mu_n([d_n t, d_n(t + h)]) dt' = (1 - \epsilon_2) \mu_n([d_n t, d_n(t + h)]). \end{aligned}$$

By (7.6) and (7.7),

$$\begin{aligned} \mu_n([d_n t, d_n(t + h)]) &\leq (1 - \epsilon_2)^{-1} V_n(t - \epsilon_1 h, h(1 + 2\epsilon_1), \epsilon_1^2 h) \\ &\leq h \left(\tilde{q}_\beta(t) + 2\epsilon_1 q_\infty + \frac{1}{2} \epsilon \right) (1 - \epsilon_2)^{-1} \leq h(\tilde{q}_\beta(t) + \epsilon). \end{aligned}$$

Arguing similarly, and exploiting the last estimate for $\mu_n([d_n t, d_n(t + h)])$,

$$\begin{aligned} V_n(t + \epsilon_1 h, h(1 - 2\epsilon_1), \epsilon_1^2 h) &\leq \int_{|t'| \leq \epsilon_1 h} \hat{k}_{\epsilon_1^2 h}(t') \mu_n([d_n(t + \epsilon_1 h - t'), d_n(t - \epsilon_1 h - t' + h)]) dt' \\ &\quad + \int_{|t'| \geq \epsilon_1 h} \hat{k}_{\epsilon_1^2 h}(t') h(q_\infty + \epsilon) dt' \\ &\leq \mu_n([d_n t, d_n(t + h)]) + \epsilon_2 h(q_\infty + \epsilon). \end{aligned}$$

By (7.6) and (7.8),

$$\begin{aligned} \mu_n([d_n t, d_n(t + h)]) &\geq V_n(t + \epsilon_1 h, h(1 - 2\epsilon_1), \epsilon_1^2 h) - \epsilon_2 h(q_\infty + \epsilon) \\ &\geq h \left(\tilde{q}_\beta(t) - 2\epsilon_1 q_\infty - \frac{1}{2} \epsilon - \epsilon_2(q_\infty + \epsilon) \right) \geq h(\tilde{q}_\beta(t) - \epsilon). \end{aligned}$$

This completes the proof. □

Proof of Theorem 2.7. After a change of variables, Lemma 7.2 reads as follows:

Let $\epsilon > 0$ and $L > 0$. There exists $n_0 \geq 1$ and $h_0 > 0$ such that

$$\sup_{t \in \mathbb{R}} d_n \left| \mu_n([t, t+h]) - \frac{h}{d_n} q_\beta(d_n^{-1}t) \mu(A) \mu(B) \right| \leq h\epsilon, \tag{7.9}$$

for all $n \geq n_0, h \in [L^{-1}, d_n h_0]$.

Fix $h > 0$ and define $e_n = \sup_{t \in \mathbb{R}} d_n |\mu_n([t, t+h]) - \frac{h}{d_n} q_\beta(d_n^{-1}t) \mu(A) \mu(B)|$. We must show that $\lim_{n \rightarrow \infty} e_n = 0$.

Let $L = 1/h$. By (7.9), for any $\epsilon > 0$ there exists $n_0 \geq 1, h_0 > 0$, such that $e_n \leq h\epsilon$ for all $n \geq n_0$ subject to the constraint $d_n h_0 \geq h$. Since $d_n \rightarrow \infty$, there exists $n_1 \geq n_0$ such that $d_n h_0 \geq h$ for all $n \geq n_1$. Hence $e_n \leq h\epsilon$ for all $n \geq n_1$ as required. \square

8. Proofs of Theorems 2.4 and 2.6

In this section, we prove Theorem 2.4 by establishing separately an upper bound (Corollary 8.3) and a lower bound (Corollary 8.4). In the process of obtaining the upper bound, we prove Theorem 2.6.

For ease of exposition, we assume hypothesis (H) throughout. Again, Lemma 5.9 is not required in this section, so we can just as well use hypothesis (A) by Proposition 5.10.

8.1. Upper bound for \liminf

In this subsection, we only require hypothesis (H) with $s \in \mathbb{R}^+$ in (H)(ii). A simplified version of the argument used in the proof of Lemma 5.5 can be used to obtain

Proposition 8.1. *Assume the setting of Theorem 2.6 with $\beta \in [0, 1]$. For $\sigma > 0$,*

$$\hat{T}(\sigma) = D_\beta' \tilde{\ell}(1/\sigma)^{-1} \sigma^{-\beta} (P(0) + E(\sigma)),$$

where $D_\beta' = \Gamma(1 - \beta)^{-1}$ for $\beta \in (0, 1)$ and $D_0' = D_1' = 1$, and $E(\sigma)$ is a family of operators satisfying $\lim_{\sigma \rightarrow 0} \|E(\sigma)\|_{\mathcal{B} \rightarrow L^1} = 0$.

We can now complete

Proof of Theorem 2.6. For $n \geq 0$, the real Laplace transform of the distribution $G_n(x) = \mu(\tau_n(y) \leq x, y \in A \cap F^{-n}B)$ is given by $\int_Y 1_A 1_B \circ F^n e^{-\sigma \tau_n} d\mu = \int_B \hat{R}(e^{-\sigma})^n 1_A d\mu$. Hence,

$$\int_{-\infty}^{\infty} e^{-\sigma t} dU_{A,B}(t) = \sum_{n=0}^{\infty} \int_B \hat{R}(e^{-\sigma})^n 1_A d\mu = \int_B \hat{T}(e^{-\sigma}) 1_A d\mu.$$

The conclusion follows from Proposition 8.1 by the continuous time version of Karamata’s Tauberian Theorem [11, Theorem 1.7.1]. \square

Lemma 8.2. *Assume the setting of Theorem 2.6 with $\beta \in (0, 1]$. Let $z : [0, \infty) \rightarrow [0, \infty)$ be integrable. Then*

$$\liminf_{t \rightarrow \infty} m(t) \int_0^t z(t-y) dU_{A,B}(y) \leq d_\beta \mu(A) \mu(B) \int_0^\infty z dx.$$

Proof. This is proved in the same way as [20, Lemma 9] using Theorem 2.6. \square

Corollary 8.3. *Assume the setting of Theorem 2.6 with $\beta \in (0, 1]$. Then for any $h > 0$,*

$$\liminf_{t \rightarrow \infty} m(t) (U_{A,B}(t+h) - U_{A,B}(t)) \leq d_\beta \mu(A) \mu(B) h.$$

Proof. Let $z = 1_{[0,h]}$. By Lemma 8.2,

$$\begin{aligned} \liminf_{t \rightarrow \infty} m(t) (U_{A,B}(t+h) - U_{A,B}(t)) &= \liminf_{t \rightarrow \infty} m(t+h) \int_0^{t+h} z(t+h-y) dU_{A,B}(y) \\ &\leq d_\beta \mu(A) \mu(B) \int_0^\infty z dx = d_\beta \mu(A) \mu(B) h, \end{aligned}$$

as required. \square

8.2. Lower bound for \liminf

Corollary 8.4. *Assume the setting of Theorem 2.4. Then for any $h > 0$,*

$$\liminf_{t \rightarrow \infty} m(t)(U_{A,B}(t+h) - U_{A,B}(t)) \geq d_\beta \mu(A) \mu(B) h.$$

Proof. Let $m \geq k \geq 0$. By (2.1) and Theorem 2.7,

$$\begin{aligned} U_{A,B}(t+h) - U_{A,B}(t) &\geq \sum_{n=k}^m \mu(y \in A \cap F^{-n} B : \tau_n(y) \in [t, t+h]) \\ &= \sum_{n=k}^m \frac{h}{d_n} q_\beta(t/d_n) \mu(A) \mu(B) + E_{k,m}, \end{aligned}$$

where $E_{k,m} = \sum_{n=k}^m e_n/d_n$.

Let $\kappa \in (1, 1/\beta)$. Then $d_n^{-1} = O(n^{-\kappa})$ and $E_{k,m} = O(\sup_{n \geq k} |e_n|) \rightarrow 0$ as $k \rightarrow \infty$.

Choosing $k = [C_1 t^\beta / \ell(t)]$ and $m = [C_2 t^\beta / \ell(t)]$, for fixed $C_2 > C_1 > 0$ and arguing word for word as in [20, Proof of eq. (7.2)], we obtain

$$\liminf_{t \rightarrow \infty} m(t)(U_{A,B}(t+h) - U_{A,B}(t)) \geq \mu(A) \mu(B) \int_{C_1}^{C_2} x^{-1/\beta} q_\beta(x^{-1/\beta}) dx.$$

Now let $C_1 \rightarrow 0$ and $C_2 \rightarrow \infty$ and use that $\int_0^\infty x^{-1/\beta} q_\beta(x^{-1/\beta}) dx = d_\beta$. □

9. General class of observables

In this section, we extend mixing for semiflows, Corollary 3.1, to cover more general classes of observables. As well as being of interest in its own right, this is useful for the extension to flows in Section 10.

Throughout, we suppose that we are in the setting of Corollary 3.1; in particular $\beta \in (\frac{1}{2}, 1]$ and hypotheses (H) and (S)(i) hold. We also suppose from now on that Y is a metric space with inner regular² Borel probability measure μ and that F and τ are almost everywhere continuous. It is well-known that mixing for infinite measure system is not a measure-theoretic property [25,30] and that care needs to be taken with the class of observables. Here we follow Krickeberg [30]. As a special case of the general theory, we prove the following result:

Theorem 9.1. *Define $H_n = \{(y, u) \in Y \times [0, \infty) : \tau(y) - n \leq u \leq \tau(y)\}$, $n \geq 1$. Then*

$$\lim_{t \rightarrow \infty} m(t) \int_{Y^\tau} v w \circ F_t d\mu^\tau = d_\beta \int_{Y^\tau} v d\mu^\tau \int_{Y^\tau} w d\mu^\tau \quad (9.1)$$

for all bounded and almost everywhere continuous functions $v : Y^\tau \rightarrow \mathbb{R}$ supported in H_n for some n , and all $w \in L^1(Y^\tau)$.

Note that this includes all bounded almost everywhere continuous observables v supported in a set of the form $A \times [a_1, a_2] \subset Y^\tau$ where $A \subset Y$, $0 < a_1 < a_2 \leq \inf_A \tau$ and $\sup_A \tau < \infty$. For the results on flows in Section 10 we require the more general class of observables in Theorem 9.1.

In the remainder of this section, we prove a more general result along the lines of [30] and use this to prove Theorem 9.1.

Let \mathcal{C} be a collection of measurable subsets $A \subset Y$ with $1_A \in \mathcal{B}$ such that

- (i) $\mu(\partial A) = 0$ for all $A \in \mathcal{C}$,
- (ii) $A_1 \cap A_2 \in \mathcal{C}$ for all $A_1, A_2 \in \mathcal{C}$,
- (iii) \mathcal{C} is a basis for the topology on Y .

In practice, we can often take \mathcal{C} to consist of all measurable sets $A \subset Y$ with $1_A \in \mathcal{B}$ and $\mu(\partial A) = 0$. This is the case for the examples in Section 11.

² μ is inner regular if $\mu(A) = \sup \mu(K : K \subset A, A \text{ compact})$ for all open sets $A \subset Y$.

Proposition 9.2. Let $\mathcal{C}' = \{A \times [a_1, a_2] \subset Y^\tau : A \in \mathcal{C}\}$. Let \mathcal{D} be the ring generated by \mathcal{C}' and let $H \in \mathcal{D}$. Then (9.1) holds for all bounded and almost everywhere continuous functions $v : Y^\tau \rightarrow \mathbb{R}$ supported in H , and all $w \in L^1(Y^\tau)$.

Proof. It is immediate that conditions (i)–(iii) for \mathcal{C} are inherited by the collection \mathcal{C}' of subsets of Y^τ (with μ replaced by μ^τ).

Write $q(t) = d_\beta^{-1}m(t)$. By Corollary 3.1,

$$\lim_{t \rightarrow \infty} q(t) \mu^\tau(A \cap F_t^{-1}B) = \mu^\tau(A) \mu^\tau(B), \quad (9.2)$$

for all $A \in \mathcal{C}'$ and all measurable rectangles $B \subset Y^\tau$. The argument now proceeds as in [30, Section 2]. We provide the details for completeness.

Step 1: Let $B \subset Y^\tau$ be a measurable rectangle. Then (9.2) holds for all $A \in \mathcal{C}'$ and hence (using condition (ii)) for all finite unions and differences of elements of \mathcal{C}' . In other words, (9.2) holds for all $A \in \mathcal{D}$.

Step 2: Let $B \subset Y^\tau$ be a measurable rectangle. Recall that $H \in \mathcal{D}$ and let $A \subset H$ such that $\mu^\tau(\partial A) = 0$. Suppose that $K \subset \text{Int } A$ is compact. Since \mathcal{C}' is a basis and \mathcal{D} is stable under finite unions, there exists $D \in \mathcal{D}$ such that $K \subset D \subset \text{Int } A$. Using also the inner regularity of μ^τ ,

$$\begin{aligned} \mu^\tau(A) &= \mu^\tau(\text{Int } A) = \sup\{\mu^\tau(K) : K \subset \text{Int } A, K \text{ compact}\} \\ &= \sup\{\mu^\tau(D) : D \subset A, D \in \mathcal{D}\}. \end{aligned}$$

Similarly,

$$\mu^\tau(H \setminus A) = \sup\{\mu^\tau(D) : D \subset H \setminus A, D \in \mathcal{D}\} = \sup\{\mu^\tau(H \setminus D) : D \supset A, D \in \mathcal{D}\},$$

so $\mu^\tau(A) = \inf\{\mu^\tau(D) : D \supset A, D \in \mathcal{D}\}$. Hence for any $\epsilon > 0$, there exist $D_1, D_2 \in \mathcal{D}$ such that $D_1 \subset A \subset D_2$ and $\mu^\tau(D_2) - \mu^\tau(D_1) < \epsilon$. Since (9.2) holds for D_1 and D_2 and $\mu^\tau(D_1 \cap F_t^{-1}B) \leq \mu^\tau(A \cap F_t^{-1}B) \leq \mu^\tau(D_2 \cap F_t^{-1}B)$,

$$\begin{aligned} (\mu^\tau(A) - \epsilon) \mu^\tau(B) &\leq \mu^\tau(D_1) \mu^\tau(B) \leq \liminf_{t \rightarrow \infty} q(t) \mu^\tau(A \cap F_t^{-1}B) \\ &\leq \limsup_{t \rightarrow \infty} q(t) \mu^\tau(A \cap F_t^{-1}B) \leq \mu^\tau(D_2) \mu^\tau(B) \leq (\mu^\tau(A) + \epsilon) \mu^\tau(B). \end{aligned}$$

As ϵ is arbitrary, we have verified that (9.2) holds for all $A \subset H$ with $\mu^\tau(\partial A) = 0$. In other words, $\lim_{t \rightarrow \infty} q(t) \int_{Y^\tau} v \times 1_B \circ F_t d\mu^\tau = \int_{Y^\tau} v d\mu^\tau \mu^\tau(B)$ where $v = 1_A$. This extends to all finite linear combinations $v = \sum c_j 1_{A_j}$ by linearity. We will refer to such functions v as step functions.

Step 3: Let $B \subset Y^H$ be a measurable rectangle and suppose that v is as in the statement of the proposition. We claim that for any $\epsilon > 0$ there exist step functions v_1 and v_2 such that $v_1 \leq v \leq v_2$ and $\int_{Y^\tau} v_2 d\mu^\tau - \int_{Y^\tau} v_1 d\mu^\tau < \epsilon$. Then

$$\begin{aligned} \left(\int_{Y^\tau} v d\mu^\tau - \epsilon \right) \mu^\tau(B) &\leq \int_{Y^\tau} v_1 d\mu^\tau \mu^\tau(B) \leq \liminf_{t \rightarrow \infty} q(t) \int_{Y^\tau} v 1_B \circ F_t d\mu^\tau \\ &\leq \limsup_{t \rightarrow \infty} q(t) \int_{Y^\tau} v 1_B \circ F_t d\mu^\tau \leq \int_{Y^\tau} v_2 d\mu^\tau \mu^\tau(B) \leq \left(\int_{Y^\tau} v d\mu^\tau + \epsilon \right) \mu^\tau(B). \end{aligned}$$

Hence (9.1) holds for all v of the desired form and all indicator functions $w = 1_B$ where B is a measurable rectangle.

To prove the claim, let $\delta > 0$ such that $\delta(\mu^\tau(Y) + 2|v|_\infty) < \epsilon/2$ and let I be a closed interval covering the image of v . We can write I as a finite union of closed intervals I_1, \dots, I_N with $\text{diam } I_j < \delta$ intersecting only at endpoints.

Let $A_j = v^{-1}(I_j)$ and define Z to be the set of discontinuity points of v . Then $\partial A_j \subset Z \cup v^{-1}(\partial I_j)$ for all j . Hence $\mu^\tau(\partial A_j) \leq \mu^\tau(v^{-1}(\partial I_j))$.

Also, there are at most countably many $x_k \in \mathbb{R}$ such that $\mu^\tau(v^{-1}(x_k)) > 0$. We can modify the intervals I_j slightly so that $x_k \notin \partial I_j$ for all j, k . This ensures that $\mu^\tau(\partial A_j) = 0$ for all j .

As in Step 2, it follows from inner regularity of μ^τ that for each j there exists $D_j \in \mathcal{D}$ with $D_j \subset A_j$ such that $\mu^\tau(A_j \setminus D_j) < \delta/N$. Now define

$$v_1 = \sum_{D_j} \inf_{D_j} v 1_{D_j} + \inf_Y v 1_{H \setminus \cup D_j}, \quad v_2 = \sum_{D_j} \sup_{D_j} v 1_{D_j} + \sup_Y v 1_{H \setminus \cup D_j}.$$

Then $v_1 \leq v \leq v_2$. Also,

$$\begin{aligned} \int_{Y^\tau} v_2 d\mu^\tau - \int_{Y^\tau} v d\mu^\tau &\leq \sum \mu^\tau(D_j) \left(\sup_{D_j} v - \inf_{D_j} v \right) + 2\mu^\tau \left(H \setminus \bigcup D_j \right) |v|_\infty \\ &\leq \mu^\tau(Y)\delta + 2|v|_\infty \sum \mu^\tau(A_j \setminus D_j) < \delta(\mu^\tau(Y) + 2|v|_\infty) < \epsilon/2. \end{aligned}$$

Similarly, $\int_{Y^\tau} v d\mu^\tau - \int_{Y^\tau} v_1 d\mu^\tau < \epsilon/2$ verifying the claim.

Step 4: To prove the general result, suppose without loss that $v \geq 0$ and let $w \in L^1(Y^\tau)$. By a more standard approximation argument than the one in Step 3, there exist simple functions w_1 and w_2 such that $w_1 \leq w \leq w_2$ and $\int_{Y^\tau} w_2 d\mu^\tau - \int_{Y^\tau} w_1 d\mu^\tau < \epsilon$. The result follows. \square

Proof of Theorem 9.1. Let $\mathcal{C}'' = \mathcal{C}' \cup \{E_n, n \geq 1\}$ where \mathcal{C}' is the collection of rectangles in Proposition 9.2 and $E_n = \bigcup_{j=1}^n F_j^{-1}(Y \times [0, 1])$. Let $\mathcal{I} = \{C \cap E_n : C \in \mathcal{C}', n \geq 1\}$ and define $\mathcal{C}''' = \mathcal{C}'' \cup \mathcal{I}$. Then \mathcal{C}''' is closed under finite intersections, and hence conditions (i)–(iii) are satisfied by the collection \mathcal{C}''' . We claim that property (9.2) holds for all $A \in \mathcal{C}'''$. Certainly, the sets E_n lie in the ring generated by \mathcal{C}''' , and $H_n \subset E_n$, so the conclusion follows from the approximate argument used to prove Proposition 9.2.

It remains to verify the claim. By Corollary 3.1, property (9.2) holds for all $A \in \mathcal{C}'$. By Remark 3.2, this holds also for the sets E_n . Finally, if $I \in \mathcal{I}$, then I is contained in one of the rectangles in \mathcal{C}' and $\mu^\tau(\partial I) = 0$. Hence 1_I is a bounded and almost everywhere continuous function supported in a rectangle in \mathcal{C}' . The claim follows from Proposition 9.2. \square

10. Mixing for infinite measure flows

In this section, we show how mixing for semiflows extends to mixing for flows.

10.1. Assumptions and disintegration

We suppose throughout that $F_t : Y^\tau \rightarrow Y^\tau$ is a suspension semiflow over a map $F : Y \rightarrow Y$ with nonintegrable almost everywhere continuous roof function $\tau : Y \rightarrow \mathbb{R}^+$ satisfying $\text{ess inf } \tau > 1$ and $\mu(\tau > t) = \ell(t)t^{-\beta}$, $\beta \in (\frac{1}{2}, 1]$, and we assume that hypotheses (H) and (S)(i) hold.

Let $X = Y \times N$ where Y and N are bounded metric space. Let $f(y, z) = (Fy, G(y, z))$ where $F : Y \rightarrow Y$ and $G : Y \times N \rightarrow N$ are continuous almost everywhere. The projection $\pi : X \rightarrow Y$, $\pi(y, z) = y$, defines a semiconjugacy between f and F . There exists a unique f -invariant ergodic probability measure μ_X on X such that $\pi_*\mu_X = \mu$, see for instance [9, Section 6].

Define $\tau : X \rightarrow \mathbb{R}^+$ by setting $\tau(y, z) = \tau(y)$ and define the suspension $X^\tau = \{(x, u) \in X \times \mathbb{R} : 0 \leq u \leq \tau(x)\} / \sim$ where $(x, \tau(x)) \sim (fx, 0)$. The suspension flow $f_t : X^\tau \rightarrow X^\tau$ is given by $f_t(x, u) = (x, u + t)$ computed modulo identifications, with ergodic invariant measure $\mu_X^\tau = \mu_X \times \text{Lebesgue}$.

Under two additional assumptions (F1) and (F2) below, we show in Theorem 10.5 that Corollary 3.1 for the semiflow F_t applies equally to the flow f_t .

First, we assume contractivity along N :

(F1) $\lim_{n \rightarrow \infty} d(f^n(y, z), f^n(y, z')) = 0$ for all $z, z' \in N$ uniformly in $y \in Y$.

Recall that R denotes the transfer operator for $F : Y \rightarrow Y$.

Proposition 10.1. *Fix $z_0 \in N$. Suppose $v \in C^0(X)$. Then the limit*

$$\eta_y(v) = \lim_{n \rightarrow \infty} (R^n v_n)(y), \quad v_n(y) = v \circ f^n(y, z_0),$$

exists for almost every $y \in Y$ and defines a probability measure supported on $\pi^{-1}(y)$. Moreover $y \mapsto \eta_y(v) = \int_{\pi^{-1}(y)} v d\eta_y$ is integrable and $\int_X v d\mu_X = \int_Y \int_{\pi^{-1}(y)} v d\eta_y d\mu(y)$.

Proof. See for instance [14, Proposition 3]. \square

Remark 10.2. The proof of [14, Proposition 3] shows that the sequence $R^n v_n$ is Cauchy in $L^\infty(Y)$. If the metric on Y can be chosen so that $R^n v_n$ is continuous for each n , then $\bar{v} \in C^0(Y)$. (In fact, it can often be shown that \bar{v} is Hölder when v is Hölder [14].)

Note that $X^\tau = Y^\tau \times N$. Given $v \in C^0(X^\tau)$, define

$$\bar{v} : Y^\tau \rightarrow \mathbb{R}, \quad \bar{v}(y, u) = \int_{x \in \pi^{-1}(y)} v(x, u) d\eta_y(x).$$

Then

$$\int_{X^\tau} v d\mu_X^\tau = \int_{Y^\tau} \bar{v}(y, u) d\mu^\tau(y, u).$$

We require the additional assumption:

(F2) The function $\bar{v} : Y^\tau \rightarrow \mathbb{R}$ is almost everywhere continuous.

Remark 10.3. If v is uniformly continuous, then for any $\epsilon > 0$ there exists $\delta < 0$ such that $|\bar{v}(y, u) - \bar{v}(y, u')| < \epsilon$ for all $(y, u), (y, u') \in Y^\tau$ with $|u - u'| < \delta$. This combined with Remark 10.2 shows that condition (F2) is easily satisfied in practice for a large class of observables $v \in C^0(X^\tau)$.

Remark 10.4. The set up in this section (skew product $X = Y \times N$, roof function τ constant in the N direction) is not very restrictive. Suppose that $T_t : M \rightarrow M$ is a smooth flow defined on a Riemannian manifold M and that Λ is a partially hyperbolic attractor, so there exists a continuous DT_t -invariant splitting $T_\Lambda M = E^s \oplus E^{cu}$ where E^s is uniformly contracting and dominates E^{cu} . By [7, Proposition 3.2, Theorem 4.2], the stable bundle E^s extends to a neighbourhood U of Λ and integrates to a T_t -invariant collection \mathcal{W}^s of stable leaves that topologically foliate U .

This means that we can choose a topological submanifold $X \subset M$ that is a cross-section to the flow T_t formed as a union of stable leaves, and automatically the roof function τ is constant along stable leaves. (This construction has been widely used recently [5,6,8,10].) Assuming for convenience the existence of a global chart for \mathcal{W}^s , we obtain a Poincaré map $f : X \rightarrow X$ where $X = Y \times N$ with N playing the role of the stable direction. Moreover, f has the desired skew product form $f(y, z) = (Fy, G(y, z))$, where $F : Y \rightarrow Y$ is defined by quotienting along the stable leaves, and condition (F1) is automatically satisfied. Also (F2) holds by Remark 10.2. Hence our set up holds in its entirety provided $F : Y \rightarrow Y$ and $\tau : Y \rightarrow \mathbb{Z}^+$ satisfy the required properties.

10.2. The mixing result

Choose a subset H of Y^τ as in Proposition 9.2.

Theorem 10.5. Suppose that $\mu(\tau > n) = \ell(n)n^{-\beta}$ where $\beta \in (\frac{1}{2}, 1]$. Let $v \in C^0(X^\tau)$ be supported in $C \times N$ where C is a closed subset of $\text{Int } H$. Let $w \in C^0(X^\tau)$ be uniformly continuous and supported on a set of finite measure. Assume that (H), (S1), (F1) and (F2) hold. Then

$$\lim_{t \rightarrow \infty} m(t) \int_{X^\tau} vw \circ f_t d\mu_X^\tau = d_\beta \int_{X^\tau} v d\mu_X^\tau \int_{X^\tau} w d\mu_X^\tau.$$

Proof. Following [10], we define $w_s : Y^\tau \rightarrow \mathbb{R}, s > 0$, by setting

$$w_s(y, u) = \overline{w \circ f_s} = \int_{x \in \pi^{-1}(y)} w \circ f_s(x, u) d\eta_y(x).$$

Note that $\int_{Y^\tau} |w_s| d\mu^\tau \leq \int_{X^\tau} |w| \circ f_s d\mu_X^\tau = \int_{X^\tau} |w| d\mu_X^\tau$ so $w_s \in L^1(Y^\tau)$ for all s .

The semiconjugacy $\pi : X \rightarrow Y$ extends to a measure-preserving semiconjugacy $\pi^\tau : X^\tau \rightarrow Y^\tau, \pi^\tau(x, u) = (\pi x, u)$. Write $m(t) \int_{X^\tau} vw \circ f_t d\mu_X^\tau = I_1(s, t) + I_2(s, t)$ where

$$I_1(s, t) = m(t) \int_{X^\tau} vw_s \circ \pi^\tau \circ f_{t-s} d\mu_X^\tau,$$

$$I_2(s, t) = m(t) \int_{X^\tau} v(w \circ f_s - w_s \circ \pi^\tau) \circ f_{t-s} d\mu_X^\tau.$$

For $t > s$,

$$I_1(s, t) = m(t) \int_{X^\tau} vw_s \circ F_{t-s} \circ \pi^\tau d\mu_X^\tau = m(t) \int_{Y^\tau} \bar{v}w_s \circ F_{t-s} d\mu^\tau.$$

Since \bar{v} is bounded and almost everywhere continuous, supported in H , and $w_s \in L^1(Y^\tau)$, it follows from Proposition 9.2 that for all $s > 0$,

$$\lim_{t \rightarrow \infty} I_1(s, t) = d_\beta \int_{Y^\tau} \bar{v} d\mu^\tau \int_{Y^\tau} w_s d\mu^\tau = d_\beta \int_{X^\tau} v d\mu_X^\tau \int_{X^\tau} w d\mu_X^\tau.$$

Choose $\psi : Y^\tau \rightarrow [0, 1]$ continuous such that $\text{supp } v \subset \text{supp } \psi \times N \subset H \times N$. Define

$$D_s : Y^\tau \rightarrow \mathbb{R}, \quad D_s(y, u) = \text{diam } w \circ f_s((\pi^\tau)^{-1}(y, u)).$$

Note that $|D_s| \leq 2|w|_\infty$ and $\mu^\tau(\text{supp } D_s) \leq \mu_X^\tau(f_s^{-1} \text{supp } w) = \mu_X^\tau(\text{supp } w) < \infty$, so $D_s \in L^1(Y^\tau)$. Also, $|w \circ f_s(x, u) - w_s \circ \pi^\tau(x, u)| \leq D_s \circ \pi^\tau(x, u)$. Hence for $t > s$,

$$|I_2(s, t)| \leq |v|_\infty m(t) \int_{Y^\tau} \psi \circ \pi^\tau D_s \circ \pi^\tau \circ f_{t-s} d\mu_X^\tau = |v|_\infty m(t) \int_{Y^\tau} \psi D_s \circ F_{t-s} d\mu_Y^\tau.$$

Since $\psi \in C^0(Y^\tau)$ is supported in H and $D_s \in L^1(Y^\tau)$, it again follows from Proposition 9.2 that for all $s > 0$,

$$\limsup_{t \rightarrow \infty} I_2(s, t) \leq |v|_\infty d_\beta \int_{Y^\tau} \psi d\mu^\tau \int_{Y^\tau} D_s d\mu^\tau.$$

By uniform continuity of w and (F1), $\lim_{s \rightarrow \infty} |D_s|_\infty = 0$. Hence $|D_s|_1 \leq |D_s|_\infty \mu^\tau(\text{supp } D_s) \leq |D_s|_\infty \mu_X^\tau(\text{supp } w) \rightarrow 0$ as $s \rightarrow \infty$. This combined with the estimates for I_1 and I_2 yields the desired result. \square

11. Examples

In this section, we demonstrate how the methods in this paper apply to the examples described in the Introduction.

11.1. NonMarkovian intermittent semiflows and flows

Let $f_t : [0, 1]^{\tau_0} \rightarrow [0, 1]^{\tau_0}$ be an intermittent semiflow as in Example 1.1. The first step is to pass from the original suspension semiflow on $[0, 1]^{\tau_0}$ to a suspension of the form Y^τ where (Y, μ) is a probability space and τ is a nonintegrable roof function.

We take $Y \subset [0, 1]$ to be the interval of domain of the rightmost branch of the AFN map $f : [0, 1] \rightarrow [0, 1]$. Define the first return map $F = f^\sigma : Y \rightarrow Y$ where $\sigma = \min\{n \geq 1 : f^n y \in Y\}$. Then $\mu = (\mu_0|_Y)/\mu_0(Y)$ is an absolutely continuous invariant probability measure for F . Define the induced roof function $\tau : Y \rightarrow \mathbb{R}^+$ given by $\tau(y) = \sum_{\ell=0}^{\sigma(y)-1} \tau_0(f^\ell y)$. Let $F_t : Y^\tau \rightarrow Y^\tau$ be the corresponding suspension semiflow with infinite invariant measure μ^τ .

Since τ_0 is Hölder, it is standard that $\mu(\tau > t) \sim ct^{-\beta}$ for some $c > 0$ (see for example [13, Proposition 9.1]).

Proposition 11.1. *Suppose that f_t has two periodic orbits (other than the neutral one) whose periods have irrational ratio. Then hypotheses (H) and (S)(i) hold with $\mathcal{B} = \text{BV}$ being the space of bounded variation functions on Y , with norm $\|v\|_{\text{BV}} = |v|_1 + \text{Var } v$.*

Proof. Hypotheses (H)(i,iii) are verified in [13, Proposition 9.2]. Also, hypothesis (H)(ii) is verified in [13, Proposition 9.2] for $s \in \mathbb{H} \cap B_\delta(0)$.

To complete the verification of (H)(ii), we proceed as follows. Since the density $d\mu/d \text{Leb}$ lies in BV and is bounded above and below, it suffices to work with the non-normalised transfer operator $\hat{P}(ib)v = P(e^{ib\tau} v)$ where $\int_Y P v w d \text{Leb} = \int_Y v w \circ F d \text{Leb}$.

Let $\lambda = \inf g|_Y > 1$. Fix $L > 0$. It suffices to show that there exists a constant C' such that

$$\|\hat{P}(ib)^n v\|_{\text{BV}} \leq C' n |v|_1 + C' n \lambda^{-n} \text{Var } v,$$

for all $|b| \leq L$, $n \geq 1$, $v \in \text{BV}$.

Let $n \geq 1$ and let $\{I\}$ be the partition of domains of branches for F^n . There is a constant C_0 independent of n such that $\text{supp } 1/(F^n)' \leq C_0 \text{diam } I$ for all I . Also $F' \geq \lambda$, so $|1/(F^n)'| \leq 1/\lambda^n$ for all n .

Write

$$\hat{P}(ib)^n v = \sum_I \{\zeta_n e^{ib\tau_n} v\} \circ \psi_I 1_{F^n I},$$

where $\zeta_n = 1/(F^n)'$, ψ_I is the inverse branch $(F^n|_I)^{-1}$, and $\tau_n = \sum_{j=0}^{n-1} \tau \circ F^j$ (not to be confused with τ_0). We have the standard estimate

$$\begin{aligned} |\hat{P}(ib)^n v|_1 &\leq |\hat{P}(ib)^n v|_\infty \leq \sum_I \sup_I (\zeta_n |v|) \leq \sum_I \sup_I \zeta_n (\inf_I |v| + \text{Var}_I v) \\ &\leq \sum_I \sup_I \zeta_n (\text{diam } I)^{-1} \int_I |v| + \sum_I \lambda^{-n} \text{Var}_I v \leq C_0 |v|_1 + \lambda^{-n} \text{Var } v. \end{aligned}$$

Next,

$$\begin{aligned} \text{Var}(\hat{P}(ib)^n v) &\leq \sum_I \text{Var}_I (\zeta_n e^{ib\tau_n} v) + 2 \sum_I \sup_I (\zeta_n |v|) \\ &\leq \sum_I \text{Var}_I (\zeta_n v) + \sum_I \sup_I (\zeta_n |v|) \text{Var}_I e^{ib\tau_n} + 2C_0 |v|_1 + 2\lambda^{-n} \text{Var } v. \end{aligned}$$

A standard argument shows that

$$\sum_I \text{Var}_I (\zeta_n v) \leq C_1 |v|_1 + \lambda^{-n} \text{Var } v,$$

where $C_1 = \sup_n |(F^n)''/[(F^n)']^2|$. Also,

$$\text{Var}_I e^{ib\tau_n} \leq |b| \text{Var}_I \tau_n \leq L \sum_{j=0}^{n-1} \text{Var}_I (\tau \circ F^j) = L \sum_{j=0}^{n-1} \text{Var}_{F^j I} \tau.$$

Let a be the domain of a branch for F . Then $\tau|_a = \sum_{\ell=0}^{\sigma(a)-1} \tau_0 \circ f^\ell$. Since the images $f^\ell a$ are disjoint for $\ell < \sigma(a)$, it follows that $\text{Var}_a \tau \leq \text{Var } \tau_0$. But $F^j I$ lies in such a domain a , so $\text{Var}_{F^j I} \tau \leq \text{Var } \tau_0$ and it follows that $\text{Var}_I e^{ib\tau_n} \leq \text{Ln Var } \tau_0$. Hence

$$\sum_I \sup_I (\zeta_n |v|) \text{Var}_I e^{ib\tau_n} \leq \text{Ln Var } \tau_0 \sum_I \sup_I (\zeta_n |v|) \leq \text{Ln Var } \tau_0 (C_0 |v|_1 + \lambda^{-n} \text{Var } v).$$

Combining these estimates we have shown that $\|\hat{P}(ib)^n v\|_{\text{BV}} \leq (3C_0 + C_1 + C_0 L \text{Var } \tau_0)n |v|_1 + (4 + L \text{Var } \tau_0)n \lambda^{-n} \text{Var } v$ as required.

Passing to the L^2 adjoint of $\hat{R}(ib)$, to verify (S)(i) it is equivalent to rule out the possibility that there exists $b \neq 0$ and a BV eigenfunction $v : Y \rightarrow S^1$ such that $e^{ib\tau} v \circ F = v$. Suppose that $y \in Y$ is a periodic point of period k for F . Now, BV functions have one-sided limits, and F is orientation preserving, so $v(y+) = v(F^k(y+))$. Substituting into the equation $e^{ib\tau_k} v \circ F^k = v$ we obtain $e^{ibq} = 1$ where $q = \tau_k(y+)$ is the period of the corresponding periodic orbit for f_I . This is impossible under the periodic orbit assumption, so the BV eigenfunction v cannot exist. \square

It follows from Theorem 9.1 that mixing for F_t holds for all bounded almost everywhere continuous \hat{v} supported in $H_n = \{(y, u) \in Y \times [0, \infty) : \tau(y) - n \leq u \leq \tau(y)\}$ for some $n \geq 1$, and all $\hat{w} \in L^1(Y^\tau)$.

Let $v, w : [0, 1]^{\tau_0} \rightarrow \mathbb{R}$ be observables where v is bounded and almost everywhere continuous and w is integrable. The projection $\pi : Y^\tau \rightarrow [0, 1]^{\tau_0}$, $\pi(y, u) = f_u(y, 0)$, defines a measure-preserving semiconjugacy from $F_t : Y^\tau \rightarrow Y^\tau$ to $f_t : [0, 1]^{\tau_0} \rightarrow [0, 1]^{\tau_0}$. Define the lifted observables $\hat{v} = v \circ \pi$, $\hat{w} = w \circ \pi : Y^\tau \rightarrow \mathbb{R}$. Then mixing for F_t holds provided \hat{v} is supported in an H_n and hence the desired mixing result (1.2) holds for f_t and the observables v and w . This includes all (finite linear combinations of) observables v supported in $A \times [0, \inf_A \tau_0]$ where $A \subset \{\sigma \leq j\}$ for some $j \geq 1$. (For such an observable v , we have $\text{supp } \hat{v} \subset H_n$ for $n \geq j|\tau_0|_\infty$.)

We can enlarge the class of observables v to include all bounded almost everywhere continuous functions that vanish on a neighborhood of the neutral fixed point. First, by adjoining preimages of Y we can enlarge Y so that it contains $[\epsilon, 1]$ for any prescribed $\epsilon > 0$. Hence we can suppose without loss that $\text{supp } v \subset \{(x, u) \in [0, 1]^{\tau_0} : x \in Y\}$. Since Y is the first return for F , it follows that $\text{supp } \hat{v} \subset \{(y, u) \in Y^\tau : u \leq \tau_0(y)\}$. Let $Y_j = \{y \in Y : \sigma(y) = j\}$. Define $C' = C \times [0, |\tau_0|_\infty]$ where $C = \bigcup_{j \geq 1} \{Y_j : |\tau_0|_\infty < \inf_{Y_j} \tau\}$. For the remaining Y_j , we have $j \leq \inf_{Y_j} \tau \leq |\tau_0|_\infty$ so $\sup_{Y_j} \tau \leq j|\tau_0|_\infty \leq |\tau_0|_\infty^2$. Hence $\text{supp } \hat{v} \subset C' \cup H_n$ for $n \geq |\tau_0|_\infty^2$. Such observables are covered by Section 9: Take \mathcal{C} to be the collection of finite unions of intervals in Y and define C' as in Proposition 9.2. Certainly $C' \in \mathcal{C}$. Define C'' as in the proof of Theorem 9.1. Then $C', H_n \in C''$, so $C' \cup H_n$ lies in the ring generated by C'' . In particular, mixing holds for observables such as \hat{v} supported in $C' \cup H_n$.

Remark 11.2. To verify hypothesis (S)(ii) it suffices to rule out the possibility that there exists $b \neq 0$, $\lambda \in S^1$ and a BV eigenfunction $v : Y \rightarrow S^1$ such that $e^{ib\tau} v \circ F = \lambda v$. But then every period $q = \tau(y)$ corresponding to a fixed point y for F satisfies $e^{ibq} = \lambda$. Hence hypothesis (S)(ii) holds provided this set of periods is not contained in a lattice of the form $a_1 + a_2\mathbb{Z}$ for some $a_1, a_2 > 0$.

Remark 11.3. Combining this example with Remark 10.4 leads to examples of partially hyperbolic intermittent flows preserving an infinite measure. See [33,34] for similar examples in the discrete time invertible setting. In addition to extending to continuous time, our examples are an improvement over those in [33,34] as far as mixing is concerned, since we require no assumptions on smoothness of foliations (in contrast to [33]) or Markov structure (in contrast to [34]).

11.2. Suspensions over unimodal maps

Let $f_t : [0, 1]^{\tau_0} \rightarrow [0, 1]^{\tau_0}$ be a suspension over a unimodal map $f : [0, 1] \rightarrow [0, 1]$ as described in Example 1.3. We sketch the main ingredients following [13, Section 10].

By [13, Lemma 10.2(a)], $\mu_0(\tau_0 > t) = ct^{-\beta} + O(t^{-2\beta})$ where the constant $c > 0$ is given explicitly. By [44], $f : [0, 1] \rightarrow [0, 1]$ is modelled by a Young tower $F : Y \rightarrow Y$ where Y is a tower with exponential tails over a suitable inducing set $Z \subset [0, 1]$. The roof function τ_0 lifts to a roof function $\tau : Y \rightarrow \mathbb{R}^+$ satisfying $\mu(\tau > t) = ct^{-\beta} + O(t^{-2\beta})$ where μ is the SRB measure on Y .

To prove (1.2), it remains to verify hypotheses (A) and (S)(i). In [13, Section 8.1], a new function space \mathcal{B} is defined for Young towers with exponential tails, and hypothesis (A)(i,ii) are verified. This relies on a technical condition called (H3) in [13] which is verified in [13, Lemma 10.3]. (The Lasota–Yorke inequality (A)(ii) is proved in [13, Theorem B.2] for $s \in \mathbb{H} \cap B_1(0)$ but holds equally for $s \in \mathbb{H} \cap B_L(0)$ for any $L > 0$.) By [13, Proposition 8.6 and Lemma 10.4], hypothesis (A)(iii) is satisfied. Finally, hypothesis (S)(i) is immediate from the quasicompactness assumptions (A)(i,ii) and the assumption about periodic orbits for f_t .

Acknowledgements

This research began as a result of discussions with Jon Aaronson, Henk Bruin, Dima Dolgopyat, Péter Nándori, Françoise Pène and Doma Szász at the thematic program *Mixing Flows and Averaging Methods* at the Erwin Schrödinger Institute (ESI), Vienna, April/May 2016. We are particularly grateful to Bruin, Dolgopyat, Nándori and Pène for continued discussions on this topic and to the referee for several helpful suggestions.

The research of DT and IM was supported in part by funding from ESI. The research of IM was supported in part by a European Advanced Grant *StochExtHomog* (ERC AdG 320977).

References

- [1] J. Aaronson. *An Introduction to Infinite Ergodic Theory. Math. Surveys and Monographs* **50**. Amer. Math. Soc., Providence, 1997. MR1450400 <https://doi.org/10.1090/surv/050>
- [2] J. Aaronson. Rational weak mixing in infinite measure spaces. *Ergodic Theory Dynam. Systems* **33** (2013) 1611–1643. MR3122146 <https://doi.org/10.1017/etds.2012.102>
- [3] J. Aaronson and M. Denker. Local limit theorems for partial sums of stationary sequences generated by Gibbs–Markov maps. *Stoch. Dyn.* **1** (2001) 193–237. MR1840194 <https://doi.org/10.1142/S0219493701000114>
- [4] T. M. Adams and C. E. Silva. Weak rational ergodicity does not imply rational ergodicity. *Israel J. Math.* **214** (2016) 491–506. MR3540624 <https://doi.org/10.1007/s11856-016-1371-0>
- [5] V. Araújo, O. Butterley and P. Varandas. Open sets of axiom A flows with exponentially mixing attractors. *Proc. Amer. Math. Soc.* **144** (2016) 2971–2984. MR3487229 <https://doi.org/10.1090/proc/13055>
- [6] V. Araújo and I. Melbourne. Exponential decay of correlations for nonuniformly hyperbolic flows with a $C^{1+\alpha}$ stable foliation, including the classical Lorenz attractor. *Ann. Henri Poincaré* **17** (2016) 2975–3004. MR3556513 <https://doi.org/10.1007/s00023-016-0482-9>
- [7] V. Araújo and I. Melbourne. Existence and smoothness of the stable foliation for sectional hyperbolic attractors. *Bull. Lond. Math. Soc.* **49** (2017) 351–367. MR3656303 <https://doi.org/10.1112/blms.12037>
- [8] V. Araújo, I. Melbourne and P. Varandas. Rapid mixing for the Lorenz attractor and statistical limit laws for their time-1 maps. *Comm. Math. Phys.* **340** (2015) 901–938. MR3406637 <https://doi.org/10.1007/s00220-015-2471-0>
- [9] V. Araujo, M. J. Pacifico, E. R. Pujals and M. Viana. Singular-hyperbolic attractors are chaotic. *Trans. Amer. Math. Soc.* **361** (2009) 2431–2485. MR2471925 <https://doi.org/10.1090/S0002-9947-08-04595-9>
- [10] A. Avila, S. Gouëzel and J. Yoccoz. Exponential mixing for the Teichmüller flow. *Publ. Math. Inst. Hautes Études Sci.* **104** (2006) 143–211. MR2264836 <https://doi.org/10.1007/s10240-006-0001-5>
- [11] N. H. Bingham, C. M. Goldie and J. L. Teugels. *Regular Variation. Encyclopedia of Mathematics and Its Applications* **27**. Cambridge University Press, Cambridge, 1987. MR1015093
- [12] L. Breiman. *Probability*. Addison-Wesley Publishing Company, Reading, MA, 1968. MR0229267

- [13] H. Bruin, I. Melbourne and D. Terhesiu. Rates of mixing for non-Markov infinite measure semiflows. *Trans. Amer. Math. Soc.* **371** (2019) 7343–7386. MR3939580 <https://doi.org/10.1090/tran/7582>
- [14] O. Butterley and I. Melbourne. Disintegration of invariant measures for hyperbolic skew products. *Israel J. Math.* **219** (2017) 171–188. MR3642019 <https://doi.org/10.1007/s11856-017-1477-z>
- [15] F. Caravenna and R. A. Doney. Local large deviations and the strong renewal theorem. Preprint, 2016. Available at arXiv:1612.07635.
- [16] P. Collet and J.-P. Eckmann. Positive Liapunov exponents and absolute continuity for maps of the interval. *Ergodic Theory Dynam. Systems* **3** (1983) 13–46. MR0743027 <https://doi.org/10.1017/S0143385700001802>
- [17] D. Dolgopyat. Prevalence of rapid mixing in hyperbolic flows. *Ergodic Theory Dynam. Systems* **18** (1998) 1097–1114. MR1653299 <https://doi.org/10.1017/S0143385798117431>
- [18] D. Dolgopyat and P. Nándori. Infinite measure renewal theorem and related results. *Bull. Lond. Math. Soc.* To appear. Available at arXiv:1709.04074.
- [19] R. A. Doney. One-sided local large deviation and renewal theorems in the case of infinite mean. *Probab. Theory Related Fields* **107** (1997) 451–465. MR1440141 <https://doi.org/10.1007/s004400050093>
- [20] K. B. Erickson. Strong renewal theorems with infinite mean. *Trans. Amer. Math. Soc.* **151** (1970) 263–291. MR0268976 <https://doi.org/10.2307/1995628>
- [21] W. Feller and S. Orey. A renewal theorem. *J. Math. Mech.* **10** (1961) 619–624. MR0130721
- [22] A. Garsia and J. Lamperti. A discrete renewal theorem with infinite mean. *Comment. Math. Helv.* **37** (1962/1963) 221–234. MR0148121 <https://doi.org/10.1007/BF02566974>
- [23] S. Gouëzel. Characterization of weak convergence of Birkhoff sums for Gibbs–Markov maps. *Israel J. Math.* **180** (2010) 1–41. MR2735054 <https://doi.org/10.1007/s11856-010-0092-z>
- [24] S. Gouëzel. Correlation asymptotics from large deviations in dynamical systems with infinite measure. *Colloq. Math.* **125** (2011) 193–212. MR2871313 <https://doi.org/10.4064/cm125-2-5>
- [25] A. B. Hajian and S. Kakutani. Weakly wandering sets and invariant measures. *Trans. Amer. Math. Soc.* **110** (1964) 136–151. MR0154961 <https://doi.org/10.2307/1993640>
- [26] M. Jakobson. Absolutely continuous invariant measures for one-parameter families of one-dimensional maps. *Comm. Math. Phys.* **81** (1981) 39–88. MR0630331
- [27] J. Kautzsch, M. Kesseböhmer and T. Samuel. On the convergence to equilibrium of unbounded observables under a family of intermittent interval maps. *Ann. Henri Poincaré* **17** (2016) 2585–2621. MR3535873 <https://doi.org/10.1007/s00023-015-0451-8>
- [28] J. Kautzsch, M. Kesseböhmer, T. Samuel and B. O. Stratmann. On the asymptotics of the α -Farey transfer operator. *Nonlinearity* **28** (2015) 143–166. MR3297130 <https://doi.org/10.1088/0951-7715/28/1/143>
- [29] G. Keller and C. Liverani. Stability of the spectrum for transfer operators. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **XXVIII** (1999) 141–152. MR1679080
- [30] K. Krickeberg. Strong mixing properties of Markov chains with infinite invariant measure. In *Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 2* 431–446. Univ. California Press, Berkeley, Calif, 1967. MR0219140
- [31] M. Lenci. On infinite-volume mixing. *Comm. Math. Phys.* **298** (2010) 485–514. MR2669446 <https://doi.org/10.1007/s00220-010-1043-6>
- [32] M. Lenci. Uniformly expanding Markov maps of the real line: Exactness and infinite mixing. *Discrete Contin. Dyn. Syst.* **37** (2017) 3867–3903. MR3639443 <https://doi.org/10.3934/dcds.2017163>
- [33] C. Liverani and D. Terhesiu. Mixing for some non-uniformly hyperbolic systems. *Ann. Henri Poincaré* **17** (2016) 179–226. MR3437828 <https://doi.org/10.1007/s00023-015-0399-8>
- [34] I. Melbourne. Mixing for invertible dynamical systems with infinite measure. *Stoch. Dyn.* **15** (2015), 1550012. MR3332272 <https://doi.org/10.1142/S0219493715500124>
- [35] I. Melbourne and D. Terhesiu. Operator renewal theory and mixing rates for dynamical systems with infinite measure. *Invent. Math.* **189** (2012) 61–110. MR2929083 <https://doi.org/10.1007/s00222-011-0361-4>
- [36] I. Melbourne and D. Terhesiu. First and higher order uniform dual ergodic theorems for dynamical systems with infinite measure. *Israel J. Math.* **194** (2013) 793–830. MR3047092 <https://doi.org/10.1007/s11856-012-0154-5>
- [37] I. Melbourne and D. Terhesiu. Operator renewal theory for continuous time dynamical systems with finite and infinite measure. *Monatsh. Math.* **182** (2017) 377–431. MR3600409 <https://doi.org/10.1007/s00605-016-0922-0>
- [38] K. Petersen. *Ergodic Theory. Cambridge Studies in Adv. Math.* **2**. Cambridge Univ. Press, Cambridge, 1983. MR0833286 <https://doi.org/10.1017/CBO9780511608728>
- [39] C. Stone. A local limit theorem for nonlattice multi-dimensional distribution functions. *Ann. Math. Stat.* **36** (1965) 546–551. MR0175166 <https://doi.org/10.1214/aoms/1177700165>
- [40] D. Terhesiu. Improved mixing rates for infinite measure preserving systems. *Ergodic Theory Dynam. Systems* **35** (2015) 585–614. MR3316926 <https://doi.org/10.1017/etds.2013.59>
- [41] D. Terhesiu. Mixing rates for intermittent maps of high exponent. *Probab. Theory Related Fields* **166** (2016) 1025–1060. MR3568045 <https://doi.org/10.1007/s00440-015-0690-0>
- [42] M. Thaler. Transformations on $[0, 1]$ with infinite invariant measures. *Israel J. Math.* **46** (1983) 67–96. MR0727023 <https://doi.org/10.1007/BF02760623>
- [43] M. Thaler. The asymptotics of the Perron–Frobenius operator of a class of interval maps preserving infinite measures. *Studia Math.* **143** (2000) 103–119. MR1813362 <https://doi.org/10.4064/sm-143-2-103-119>
- [44] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math.* **147** (1998) 585–650. MR1637655 <https://doi.org/10.2307/120960>
- [45] R. Zweimüller. Ergodic structure and invariant densities of non-Markovian interval maps with indifferent fixed points. *Nonlinearity* **11** (1998) 1263–1276. MR1644385 <https://doi.org/10.1088/0951-7715/11/5/005>