

CRAMÉR THEOREM FOR GAMMA RANDOM VARIABLES

SOLESNE BOURGUIN

SAMM, EA 4543, Université Paris 1 Panthéon Sorbonne, 90 Rue de Tolbiac, 75634 Paris, France
email: solesne.bourguin@univ-paris1.fr

CIPRIAN A. TUDOR¹

Laboratoire Paul Painlevé, Université Lille 1, 59655 Villeneuve d'Ascq, France
Associate member of the team SAMM, Université Paris 1 Panthéon-Sorbonne
email: tudor@math.univ-lille1.fr

Submitted January 12, 2011, accepted in final form June 16, 2011

AMS 2000 Subject classification: 60F05, 60H05, 91G70

Keywords: Cramér's theorem, Gamma distribution, multiple stochastic integrals, limit theorems, Malliavin calculus

Abstract

In this paper we discuss the following problem: given a random variable $Z = X + Y$ with Gamma law such that X and Y are independent, we want to understand if then X and Y each follow a Gamma law. This is related to Cramér's theorem which states that if X and Y are independent then $Z = X + Y$ follows a Gaussian law if and only if X and Y follow a Gaussian law. We prove that Cramér's theorem is true in the case of the Gamma distribution for random variables living in a Wiener chaos of fixed order but the result is not true in general. We also give an asymptotic variant of our result.

1 Introduction

Cramér's theorem (see [1]) says that the sum of two independent random variables is Gaussian if and only if each summand is Gaussian. One direction is elementary to prove, that is, given two independent random variables with Gaussian distribution, then their sum follows a Gaussian distribution. The second direction is less trivial and its proof requires powerful results from complex analysis (see [1]).

In this paper, we treat the same problem for Gamma distributed random variables. A Gamma random variable, denoted usually by $\Gamma(a, \lambda)$, is a random variable with probability density function given by $f_{a,\lambda}(x) = \frac{\lambda^a}{\Gamma(a)} x^{a-1} e^{-\lambda x}$ if $x > 0$ and $f_{a,\lambda}(x) = 0$ otherwise. The parameters a and λ are strictly positive and Γ denotes the usual Gamma function.

It is well known that if $X \sim \Gamma(a, \lambda)$ and $Y \sim \Gamma(b, \lambda)$ and X is independent of Y , then $X + Y$ follows the law $\Gamma(a + b, \lambda)$. The purpose of this paper is to understand the converse implication, i.e. whether or not (or under what conditions), if X and Y are two independent random

¹PARTIALLY SUPPORTED BY THE ANR GRANT 'MASTERIE' (10 BLAN 0121 01)

variables such that $X + Y \sim \Gamma(a + b, \lambda)$ and $\mathbf{E}(X) = \mathbf{E}(\Gamma(a, \lambda))$, $\mathbf{E}(X^2) = \mathbf{E}(\Gamma(a, \lambda)^2)$ and $\mathbf{E}(Y) = \mathbf{E}(\Gamma(b, \lambda))$, $\mathbf{E}(Y^2) = \mathbf{E}(\Gamma(b, \lambda)^2)$, it holds that $X \sim \Gamma(a, \lambda)$ and $Y \sim \Gamma(b, \lambda)$. We will actually focus our attention on the so-called centered Gamma distribution $F(\nu)$. We will call ‘centered Gamma’ the random variables of the form

$$F(\nu) \stackrel{\text{Law}}{=} 2G(\nu/2) - \nu, \quad \nu > 0,$$

where $G(\nu/2) := \Gamma(\nu/2, 1)$ has a Gamma law with parameters $\nu/2, 1$. This means that $G(\nu/2)$ is a (a.s. strictly positive) random variable with density $g(x) = \frac{x^{\frac{\nu}{2}-1}e^{-x}}{\Gamma(\nu/2)} \mathbf{1}_{(0,\infty)}(x)$. The characteristic function of the law $F(\nu)$ is given by

$$\mathbf{E}\left(e^{i\lambda F(\nu)}\right) = \left(\frac{e^{-i\lambda}}{\sqrt{1-2i\lambda}}\right)^\nu, \quad \lambda \in \mathbb{R}. \quad (1)$$

We will find the following answer: if X and Y are two independent random variables, each living in a Wiener chaos of fixed order (and these orders are allowed to be different) then the fact that the sum $X + Y$ follows a centered Gamma distribution implies that X and Y each follow a Gamma distribution. On the other hand, for random variables having an infinite Wiener-Itô chaos decomposition, the result is not true even in very particular cases (for so-called strongly independent random variables). We construct a counter-example to illustrate this fact.

Our tools are based on a criterion given in [6] to characterize the random variables with Gamma distribution in terms of Malliavin calculus.

Our paper is structured as follows. Section 2 contains some notations and preliminaries. In Section 3 we prove the Cramér theorem for Gamma distributed random variables in Wiener chaos of finite orders and we also give an asymptotic version of this result. In Section 4 we show that the result does not hold in the general case.

2 Some notations and definitions

Let $(W_t)_{t \in T}$ be a classical Wiener process on a standard Wiener space $(\Omega, \mathcal{F}, \mathbf{P})$. If $f \in L^2(T^n)$ with $n \geq 1$ integer, we introduce the multiple Wiener-Itô integral of f with respect to W . The basic references are the monographs [3] or [4]. Let $f \in \mathcal{S}_n$ be an elementary function with n variables that can be written as $f = \sum_{i_1, \dots, i_n} c_{i_1, \dots, i_n} \mathbf{1}_{A_{i_1} \times \dots \times A_{i_n}}$ where the coefficients satisfy $c_{i_1, \dots, i_n} = 0$ if two indices i_k and i_l are equal and the sets $A_i \in \mathcal{B}(T)$ are pairwise disjoint. For such a step function f we define

$$I_n(f) = \sum_{i_1, \dots, i_n} c_{i_1, \dots, i_n} W(A_{i_1}) \dots W(A_{i_n})$$

where we put $W(A) = \int_0^1 \mathbf{1}_A(s) dW_s$. It can be seen that the application I_n constructed above from \mathcal{S}_n to $L^2(\Omega)$ is an isometry on \mathcal{S}_n in the sense

$$\mathbf{E}(I_n(f)I_m(g)) = n! \langle f, g \rangle_{L^2(T^n)} \text{ if } m = n \quad (2)$$

and

$$\mathbf{E}(I_n(f)I_m(g)) = 0 \text{ if } m \neq n.$$

Since the set \mathcal{S}_n is dense in $L^2(T^n)$ for every $n \geq 1$ the mapping I_n can be extended to an isometry from $L^2(T^n)$ to $L^2(\Omega)$ and the above properties hold true for this extension.

It also holds that $I_n(f) = I_n(\tilde{f})$ where \tilde{f} denotes the symmetrization of f defined by

$$\tilde{f}(x_1, \dots, x_n) = \frac{1}{n!} \sum_{\sigma} f(x_{\sigma(1)}, \dots, x_{\sigma(n)}),$$

σ running over all permutations of $\{1, \dots, n\}$. We will need the general formula for calculating products of Wiener chaos integrals of any orders m, n for any symmetric integrands $f \in L^2(T^m)$ and $g \in L^2(T^n)$, which is

$$I_m(f)I_n(g) = \sum_{\ell=0}^{m \wedge n} \ell! \binom{m}{\ell} \binom{n}{\ell} I_{m+n-2\ell}(f \otimes_{\ell} g) \tag{3}$$

where the contraction $f \otimes_{\ell} g$ is defined by

$$\begin{aligned} & (f \otimes_{\ell} g)(s_1, \dots, s_{m-\ell}, t_1, \dots, t_{n-\ell}) \\ &= \int_{T^{m+n-2\ell}} f(s_1, \dots, s_{m-\ell}, u_1, \dots, u_{\ell}) g(t_1, \dots, t_{n-\ell}, u_1, \dots, u_{\ell}) du_1 \dots du_{\ell}. \end{aligned} \tag{4}$$

Note that the contraction $(f \otimes_{\ell} g)$ is an element of $L^2(T^{m+n-2\ell})$ but it is not necessarily symmetric. We will denote its symmetrization by $(f \otimes_{\ell} g)$.

We recall that any square integrable random variable which is measurable with respect to the σ -algebra generated by W can be expanded into an orthogonal sum of multiple stochastic integrals

$$F = \sum_{n \geq 0} I_n(f_n) \tag{5}$$

where $f_n \in L^2(T^n)$ are (uniquely determined) symmetric functions and $I_0(f_0) = \mathbf{E}(F)$.

We denote by D the Malliavin derivative operator that acts on smooth functionals of the form $F = g(W(\varphi_1), \dots, W(\varphi_n))$ (here g is a smooth function with compact support and $\varphi_i \in L^2(T)$ for $i = 1, \dots, n$)

$$DF = \sum_{i=1}^n \frac{\partial g}{\partial x_i}(W(\varphi_1), \dots, W(\varphi_n)) \varphi_i.$$

We can define the i -th Malliavin derivative $D^{(i)}$ iteratively. The operator $D^{(i)}$ can be extended to the closure $\mathbb{D}^{p,2}$ of smooth functionals with respect to the norm

$$\|F\|_{p,2}^2 = \mathbf{E}(F^2) + \sum_{i=1}^p \mathbf{E}(\|D^i F\|_{L^2(T^i)}^2).$$

The adjoint of D is denoted by δ and is called the divergence (or Skorohod) integral. Its domain $\text{Dom}(\delta)$ coincides with the class of stochastic processes $u \in L^2(\Omega \times T)$ such that

$$|\mathbf{E}(\langle DF, u \rangle)| \leq c \|F\|_2$$

for all $F \in \mathbb{D}^{1,2}$ and $\delta(u)$ is the element of $L^2(\Omega)$ characterized by the duality relationship

$$\mathbf{E}(F \delta(u)) = \mathbf{E}(\langle DF, u \rangle).$$

For adapted integrands, the divergence integral coincides with the classical Itô integral.

Let L be the Ornstein-Uhlenbeck operator defined on $\text{Dom}(L) = \mathbb{D}^{2,2}$. We have

$$LF = - \sum_{n \geq 0} n I_n(f_n)$$

if F is given by (5). There exists a connection between δ, D and L in the sense that a random variable F belongs to the domain of L if and only if $F \in \mathbb{D}^{1,2}$ and $DF \in \text{Dom}(\delta)$ and then $\delta DF = -LF$. Let us consider a multiple stochastic integral $I_q(f)$ with symmetric kernel $f \in L^2(T^q)$. We denote the Malliavin derivative of $I_q(f)$ by $DI_q(f)$. We have

$$D_\theta I_q(f) = qI_{q-1}(f^{(\theta)}),$$

where $f^{(\theta)} = f(t_1, \dots, t_{q-1}, \theta)$ is the $(q-1)$ th order kernel obtained by parametrizing the q th order kernel f by one of the variables.

For any random variable $X, Y \in \mathbb{D}^{1,2}$ we use the following notations

$$G_X = \langle DX, -DL^{-1}X \rangle_{L^2(T)}$$

and

$$G_{X,Y} = \langle DX, -DL^{-1}Y \rangle_{L^2(T)}.$$

Finally, we will use the notation $X \perp Y$ to denote that two random variables X and Y are independent.

The following facts are key points in our proofs:

Fact 1: Let $X = I_{q_1}(f)$ and $Y = I_{q_2}(g)$ where $f \in L^2(T^{q_1})$ and $g \in L^2(T^{q_2})$ are symmetric functions. Then X and Y are independent if and only if (see [8])

$$f \otimes_1 g = 0 \text{ a.e. on } T^{q_1+q_2-2}.$$

Fact 2: Let $X = I_q(f)$ with $f \in L^2(T^q)$ symmetric. Assume that $\mathbf{E}(X^2) = \mathbf{E}(F(\nu)^2) = 2\nu$. Then X follows a centered Gamma law $F(\nu)$ with $\nu > 0$ if and only if (see [5])

$$\|DX\|_{L^2(T)}^2 - 2qX - 2q\nu = 0 \text{ almost surely.}$$

Fact 3: Let $(f_k)_{k \geq 1}$ be a sequence in $L^2(T^q)$ such that $\mathbf{E}(I_q(f_k)^2) \xrightarrow[k \rightarrow +\infty]{} 2\nu$. Then the sequence $X_k = I_q(f_k)$ converges in distribution, as $k \rightarrow \infty$, to a Gamma law, if and only if (see [5])

$$\|DX_k\|_{L^2(T)}^2 - 2qX_k - 2q\nu \xrightarrow[k \rightarrow +\infty]{} 0 \text{ in } L^2(\Omega).$$

Remark: In this particular paper, we will restrict ourselves to an underlying Hilbert space (to the Wiener process we will be working with in the upcoming sections) of the form $\mathfrak{H} = L^2(T)$ for the sake of simplicity. However, all the results presented in the upcoming sections remain valid on a more general separable Hilbert space as the underlying space.

3 (Asymptotic) Cramér theorem for multiple integrals

In this section, we will prove Cramér's theorem for random variables living in fixed Wiener chaoses. More precisely, our context is as follows: we assume that $X = I_{q_1}(f)$ and $Y = I_{q_2}(h)$ and X, Y are independent. We also assume that $\mathbf{E}(X^2) = \mathbf{E}(F(\nu_1)^2) = 2\nu_1$ and $\mathbf{E}(Y^2) = \mathbf{E}(F(\nu_2)^2) = 2\nu_2$. Here ν, ν_1, ν_2 denotes three strictly positive numbers such that $\nu_1 + \nu_2 = \nu$. We assume that $X + Y$ follows a Gamma law $F(\nu)$ and we will prove that $X \sim F(\nu_1)$ and $Y \sim F(\nu_2)$.

Let us first give the following two auxiliary lemmas that will be useful throughout the paper.

Lemma 1. *Let $q_1, q_2 \geq 1$ be integers, and let $X = I_{q_1}(f)$ and $Y = I_{q_2}(h)$, where $f \in L^2(T^{q_1})$ and $h \in L^2(T^{q_2})$ are symmetric functions. Assume moreover that X and Y are independent. Then, we have $DX \perp DY$, $X \perp DY$ and $Y \perp DX$.*

Proof: From Fact 1 in Section 2, $f \otimes_1 h = 0$ a.e on $T^{q_1+q_2-2}$ and by extension $f \otimes_r h = 0$ a.e on $T^{q_1+q_2-2r}$ for every $1 \leq r \leq q_1 \wedge q_2$. We will now prove that for every $\theta, \psi \in T$, we also have $f^{(\theta)} \otimes_1 h^{(\psi)} = 0$ a.e on $T^{q_1+q_2-4}$, $f^{(\theta)} \otimes_1 h = 0$ a.e on $T^{q_1+q_2-3}$ and $f \otimes_1 h^{(\psi)} = 0$ a.e. on $T^{q_1+q_2-3}$. Indeed, we have

$$\begin{aligned} (f^{(\theta)} \otimes_1 h^{(\psi)})(t_1, \dots, t_{q_1-2}, s_1, \dots, s_{q_2-2}) &= \int_T f(t_1, \dots, t_{q_1-2}, u, \theta) h(s_1, \dots, s_{q_2-2}, u, \psi) du \\ &= 0 \end{aligned}$$

as a particular case of $f \otimes_1 h = 0$ a.e.. By extension, we also have $f^{(\theta)} \otimes_r h^{(\psi)} = 0$ for $1 \leq r \leq (q_1 - 1) \wedge (q_2 - 1)$. Similarly,

$$\begin{aligned} (f^{(\theta)} \otimes_1 h)(t_1, \dots, t_{q_1-2}, s_1, \dots, s_{q_2-1}) &= \int_T f(t_1, \dots, t_{q_1-2}, u, \theta) h(s_1, \dots, s_{q_2-1}, u) du \\ &= 0. \end{aligned} \tag{6}$$

Clearly $f^{(\theta)} \otimes_r h = 0$ for $1 \leq r \leq (q_1 - 1) \wedge q_2$. Given the symmetric roles played by f and h , we also have $f \otimes_1 h^{(\psi)} = 0$ and then $f \otimes_r h^{(\psi)} = 0$ for $1 \leq r \leq q_1 \wedge (q_2 - 1)$.

Let us now prove that $DX \perp DY$. Since for every $\theta, \psi \in T$, $D_\theta X = q_1 I_{q_1-1}(f^{(\theta)})$ and $D_\psi Y = q_2 I_{q_2-1}(h^{(\psi)})$, it suffices to show that the random variables $I_{q_1-1}(f^{(\theta)})$ and $I_{q_2-1}(h^{(\psi)})$ are independent. To do this, we will use the criterion for the independence of multiple integrals given in [8]. We need to check that $f^{(\theta)} \otimes_1 h^{(\psi)} = 0$ a.e. on $T^{q_1+q_2-4}$ and this follows from above.

It remains to prove that $X \perp DY$ and $DX \perp Y$. Given the symmetric roles played by X and Y , we will only prove that $X \perp DY$. That is equivalent to the independence of the random variables $I_{q_1}(f)$ and $I_{q_2-1}(h^{(\psi)})$ for every $\theta \in T$, which follows from [8] (see Fact 1 in Section 2) and (6). Thus, we have $X \perp DY$ and $DX \perp Y$. ■

Let us recall the following definition (see [7]).

Definition 1. *Two random variables $X = \sum_{n \geq 0} I_n(f_n)$ and $Y = \sum_{m \geq 0} I_m(h_m)$ are called strongly independent if for every $m, n \geq 0$, the random variables $I_n(f_n)$ and $I_m(h_m)$ are independent.*

We have the following lemma about strongly independent random variables.

Lemma 2. *Let $X = \sum_{n \geq 0} I_n(f_n)$ and $Y = \sum_{m \geq 0} I_m(h_m)$ ($f_n \in L^2(T^n), h_m \in L^2(T^m)$) symmetric for every $n, m \geq 1$) be two centered random variables in the space $\mathbb{D}^{1,2}$. Then, if X and Y are strongly independent, we have*

$$\langle DX, -DL^{-1}Y \rangle_{L^2(T)} = \langle DY, -DL^{-1}X \rangle_{L^2(T)} = 0.$$

Proof: We have, for every $\theta \in T$,

$$D_\theta X = \sum_{n \geq 1} n I_{n-1}(f_n^{(\theta)}) \text{ and } -D_\theta L^{-1}Y = \sum_{m \geq 1} I_{m-1}(h_m^{(\theta)}).$$

Therefore, we can write

$$\begin{aligned} \langle DX, -DL^{-1}Y \rangle_{L^2(T)} &= \sum_{n,m \geq 1} n \int_T I_{n-1}(f_n(t_1, \dots, t_{n-1}, \theta)) I_{m-1}(h_m(t_1, \dots, t_{m-1}, \theta)) d\theta \\ &= \sum_{n,m \geq 1} n \int_T \sum_{r=0}^{(n-1) \wedge (m-1)} r! \binom{n-1}{r} \binom{m-1}{r} I_{n+m-2r-2}(f_n^{(\theta)} \otimes_r h_m^{(\theta)}) d\theta. \end{aligned}$$

The strong independence of X and Y gives us that $f_n^{(\theta)} \otimes_r h_m^{(\theta)} = 0$ for every $1 \leq r \leq (n-1) \wedge (m-1)$. Thus, we obtain

$$\langle DX, -DL^{-1}Y \rangle_{L^2(T)} = \sum_{n,m \geq 1} n \int_T I_{n+m-2}(f_n^{(\theta)} \otimes h_m^{(\theta)}) d\theta.$$

Using a Fubini type result, we can write

$$\begin{aligned} \langle DX, -DL^{-1}Y \rangle_{L^2(T)} &= \sum_{n,m \geq 1} n I_{n+m-2} \left(\int_T f_n^{(\theta)} \otimes h_m^{(\theta)} d\theta \right) \\ &= \sum_{n,m \geq 1} n I_{n+m-2}(f_n \otimes_1 h_m). \end{aligned}$$

Again, the strong independence of X and Y gives us that $f_n \otimes_1 h_m = 0$ a.e and we finally obtain $\langle DX, -DL^{-1}Y \rangle_{L^2(T)} = 0$, and similarly $\langle DY, -DL^{-1}X \rangle_{L^2(T)} = 0$. \blacksquare

Let us first remark that the Cramér theorem holds for random variables in the same Wiener chaos of fixed order.

Proposition 1. *Let $X = I_m(f)$ and $Y = I_m(h)$ with $m \geq 2$ fixed and f, h symmetric functions in $L^2(T^m)$. Fix $\nu_1, \nu_2, \nu > 0$ such that $\nu_1 + \nu_2 = \nu$. Assume that $X + Y$ follows the law $F(\nu)$ and X is independent of Y . Also suppose that $\mathbf{E}(X^2) = \mathbf{E}(F(\nu_1)^2) = 2\nu_1$ and $\mathbf{E}(Y^2) = \mathbf{E}(F(\nu_2)^2) = 2\nu_2$. Then $X \sim F(\nu_1)$ and $Y \sim F(\nu_2)$.*

Proof: By a result in [5] (see Fact 2 in Section 2), $X + Y$ follows the law $F(\nu)$ is equivalent to

$$\left\| |DI_m(f+h)| \right\|_{L^2(T)}^2 - 2mI_m(f+h) - 2m\nu = 0 \text{ a.s. } \quad (7)$$

On the other hand

$$\begin{aligned} &\mathbf{E} \left(\left\| |DI_m(f+h)| \right\|_{L^2(T)}^2 - 2mI_m(f+h) - 2m\nu \right)^2 \\ &= \mathbf{E} \left(\left\| |DI_m(f)| \right\|_{L^2(T)}^2 + \left\| |DI_m(h)| \right\|_{L^2(T)}^2 + 2\langle DI_m(f), DI_m(h) \rangle_{L^2(T)} \right. \\ &\quad \left. - 2mI_m(f) - 2mI_m(h) - 2m(\nu_1 + \nu_2) \right)^2 \\ &= \mathbf{E} \left(\left(\left\| |DI_m(f)| \right\|_{L^2(T)}^2 - 2mI_m(f) - 2m\nu_1 \right)^2 \right) + \mathbf{E} \left(\left(\left\| |DI_m(h)| \right\|_{L^2(T)}^2 - 2mI_m(h) - 2m\nu_2 \right)^2 \right) \\ &\quad + \mathbf{E} \left(\left(\left\| |DI_m(f)| \right\|_{L^2(T)}^2 - 2mI_m(f) - 2m\nu_1 \right) \left(\left\| |DI_m(h)| \right\|_{L^2(T)}^2 - 2mI_m(h) - 2m\nu_2 \right) \right). \quad (8) \end{aligned}$$

Above we used the fact that $\langle DI_m(f), DI_m(h) \rangle_{L^2(T)} = 0$ as a consequence of Lemma 1. It is also easy to remark that, from Lemma 1

$$\begin{aligned} &\mathbf{E} \left(\left(\left\| |DI_m(f)| \right\|_{L^2(T)}^2 - 2mI_m(f) - 2m\nu_1 \right) \left(\left\| |DI_m(h)| \right\|_{L^2(T)}^2 - 2mI_m(h) - 2m\nu_2 \right) \right) \\ &= \mathbf{E} \left(\left\| |DI_m(f)| \right\|_{L^2(T)}^2 - 2mI_m(f) - 2m\nu_1 \right) \mathbf{E} \left(\left\| |DI_m(h)| \right\|_{L^2(T)}^2 - 2mI_m(h) - 2m\nu_2 \right) = 0. \end{aligned}$$

Using this and by combining (7) and (8), we obtain that

$$\mathbf{E} \left(\left(\|DI_m(f)\|_{L^2(T)}^2 - 2mI_m(f) - 2mv_1 \right)^2 \right) + \mathbf{E} \left(\left(\|DI_m(h)\|_{L^2(T)}^2 - 2mI_m(h) - 2mv_2 \right)^2 \right) = 0.$$

The left hand side of this last equation is equal to zero and is the sum of two non negative quantities (as expectations of squares). This implies that each of the summands are equal to zero. Thus,

$$\mathbf{E} \left(\left(\|DI_m(f)\|_{L^2(T)}^2 - 2mI_m(f) - 2mv_1 \right)^2 \right) = \mathbf{E} \left(\left(\|DI_m(h)\|_{L^2(T)}^2 - 2mI_m(h) - 2mv_2 \right)^2 \right) = 0$$

and consequently $X \sim F(v_1)$ and $Y \sim F(v_2)$. ■

Remark 1. We mention that the above Proposition 1 is a particular case of Theorem 3. We prefer to state it and prove it separately because its proof is simpler and does not require the techniques used in the proof of Theorem 3. Using Fact 3 in Section 2, an asymptotic variant of the above result can be stated. We will state it here because it is a particular case of Theorem 4 proved later in our paper.

Theorem 1.2 in [5] gives a characterization of (asymptotically) centered Gamma random variable which are given by a multiple Wiener-Itô integral. There is not such a characterization for random variable living in a finite or infinite sum of Wiener chaos; only an upper bound for the distance between the law of a random variable in $\mathbb{D}^{1,2}$ and the Gamma distribution has been proven in [6], Theorem 3.11. It turns out, that for the case of a sum of independent multiple integrals, it is possible to characterize the relation between its distribution and the Gamma distribution. We will prove this fact in the following theorem.

Theorem 1. Fix $v_1, v_2, v > 0$ such that $v_1 + v_2 = v$ and let $F(v)$ be a real-valued random variable with characteristic function given by (1). Fix two even integers $q_1 \geq 2$ and $q_2 \geq 2$. For any symmetric kernels $f \in L^2(T^{q_1})$ and $h \in L^2(T^{q_2})$ such that

$$\mathbf{E} \left(I_{q_1}(f)^2 \right) = q_1! \|f\|_{L^2(T^{q_1})}^2 = 2v_1 \quad \text{and} \quad \mathbf{E} \left(I_{q_2}(h)^2 \right) = q_2! \|h\|_{L^2(T^{q_2})}^2 = 2v_2, \tag{9}$$

and such that $X = I_{q_1}(f)$ and $Y = I_{q_2}(h)$ are independent, define the random variable

$$Z = X + Y = I_{q_1}(f) + I_{q_2}(h).$$

Under those conditions, the following two conditions are equivalent:

(i) $\mathbf{E} \left(\left(2v + 2Z - \langle DZ, -DL^{-1}Z \rangle_{L^2(T)} \right)^2 \right) = 0$, where D is the Malliavin derivative operator and L is the infinitesimal generator of the Ornstein-Uhlenbeck semigroup;

(ii) $Z \stackrel{\text{Law}}{=} F(v)$;

Proof: Proof of (ii) \rightarrow (i). Suppose that $Z \sim F(v)$. We easily obtain that

$$\mathbf{E} \left(Z^3 \right) = \mathbf{E} \left(F(v)^3 \right) = 8v \quad \text{and} \quad \mathbf{E} \left(Z^4 \right) = \mathbf{E} \left(F(v)^4 \right) = 12v^2 + 48v. \tag{10}$$

Consequently,

$$\mathbf{E} \left(Z^4 \right) - 12\mathbf{E} \left(Z^3 \right) = \mathbf{E} \left(F(v)^4 \right) - 12\mathbf{E} \left(F(v)^3 \right) = 12v^2 - 48v. \tag{11}$$

Then we will use the fact that for every multiple integral $I_q(f)$

$$\mathbf{E} \left(I_q(f)^3 \right) = q!(q/2)! \binom{q}{q/2}^2 \langle f, f \tilde{\otimes}_{q/2} f \rangle_{L^2(T^q)}. \quad (12)$$

and

$$\mathbf{E} \left(I_q(f)^4 \right) = 3 \left[q! \|f\|_{L^2(T^q)}^2 \right]^2 + \frac{3}{q} \sum_{p=1}^{q-1} q^2(p-1)! \binom{q-1}{p-1}^2 p! \binom{q}{p}^2 (2q-2p)! \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q-p)})}^2. \quad (13)$$

We will now compute $\mathbf{E}(Z^3)$, $\mathbf{E}(Z^4)$ and $\mathbf{E}(Z^4) - 12\mathbf{E}(Z^3)$ by using the above two relations (12) and (13). We have $Z^2 = (I_{q_1}(f) + I_{q_2}(h))^2 = I_{q_1}(f)^2 + I_{q_2}(h)^2 + 2I_{q_1}(f)I_{q_2}(h)$ and thus, by using the independence between $I_{q_1}(f)$ and $I_{q_2}(h)$,

$$\mathbf{E}(Z^3) = \mathbf{E}(I_{q_1}(f)^3) + \mathbf{E}(I_{q_2}(h)^3).$$

Using relation (12), we can write

$$\mathbf{E}(Z^3) = q_1!(q_1/2)! \binom{q_1}{q_1/2}^2 \langle f, f \tilde{\otimes}_{q_1/2} f \rangle_{L^2(T^{q_1})} + q_2!(q_2/2)! \binom{q_2}{q_2/2}^2 \langle h, h \tilde{\otimes}_{q_2/2} h \rangle_{L^2(T^{q_2})}. \quad (14)$$

For $\mathbf{E}(Z^4)$, we combine relations (9) and (13) with the independence between $I_{q_1}(f)$ and $I_{q_2}(h)$ to obtain

$$\begin{aligned} \mathbf{E}(Z^4) &= \mathbf{E}(Z^2 Z^2) = \mathbf{E}(I_{q_1}(f)^4) + \mathbf{E}(I_{q_2}(h)^4) + 6\mathbf{E}(I_{q_1}(f)^2 I_{q_2}(h)^2) \\ &= 3 \left[q_1! \|f\|_{L^2(T^{q_1})}^2 \right]^2 + \frac{3}{q_1} \sum_{p=1}^{q_1-1} q_1^2(p-1)! \binom{q_1-1}{p-1}^2 p! \binom{q_1}{p}^2 (2q_1-2p)! \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q_1-p)})}^2 \\ &\quad + 3 \left[q_2! \|h\|_{L^2(T^{q_2})}^2 \right]^2 + \frac{3}{q_2} \sum_{p=1}^{q_2-1} q_2^2(p-1)! \binom{q_2-1}{p-1}^2 p! \binom{q_2}{p}^2 (2q_2-2p)! \|h \tilde{\otimes}_p h\|_{L^2(T^{2(q_2-p)})}^2 \\ &\quad + 24v_1 v_2. \end{aligned}$$

Using the fact that $q_1! \|f\|_{L^2(T^{q_1})}^2 = 2v_1$ and $q_2! \|h\|_{L^2(T^{q_2})}^2 = 2v_2$, we can write

$$\begin{aligned} \mathbf{E}(Z^4) - 12\mathbf{E}(Z^3) &= 12v_1^2 + 12v_2^2 - 48v_1 - 48v_2 + 24v_1 v_2 \\ &\quad + \frac{3}{q_1} \sum_{p=1, p \neq q_1/2}^{q_1-1} q_1^2(p-1)! \binom{q_1-1}{p-1}^2 p! \binom{q_1}{p}^2 (2q_1-2p)! \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q_1-p)})}^2 \\ &\quad + \frac{3}{q_2} \sum_{p=1, p \neq q_2/2}^{q_2-1} q_2^2(p-1)! \binom{q_2-1}{p-1}^2 p! \binom{q_2}{p}^2 (2q_2-2p)! \|h \tilde{\otimes}_p h\|_{L^2(T^{2(q_2-p)})}^2 \\ &\quad + 24q_1! \|f\|_{L^2(T^{q_1})}^2 + 3q_1(q_1/2-1)! \binom{q_1-1}{q_1/2-1}^2 (q_1/2)! \binom{q_1}{q_1/2}^2 q_1! \|f \tilde{\otimes}_{q_1/2} f\|_{L^2(T^{q_1})}^2 \\ &\quad + 24q_2! \|h\|_{L^2(T^{q_2})}^2 + 3q_2(q_2/2-1)! \binom{q_2-1}{q_2/2-1}^2 (q_2/2)! \binom{q_2}{q_2/2}^2 q_2! \|h \tilde{\otimes}_{q_2/2} h\|_{L^2(T^{q_2})}^2 \\ &\quad - 12q_1!(q_1/2)! \binom{q_1}{q_1/2}^2 \langle f, f \tilde{\otimes}_{q_1/2} f \rangle_{L^2(T^{q_1})} \\ &\quad - 12q_2!(q_2/2)! \binom{q_2}{q_2/2}^2 \langle h, h \tilde{\otimes}_{q_2/2} h \rangle_{L^2(T^{q_2})}. \end{aligned} \quad (15)$$

Recall that $v = v_1 + v_2$ and note that $12v_1^2 + 12v_2^2 - 48v_1 - 48v_2 + 24v_1v_2 = 12v^2 - 48v$. Also note that

$$\begin{aligned} & 24q_1! \|f\|_{L^2(T^{q_1})}^2 + 3q_1(q_1/2 - 1)! \binom{q_1 - 1}{q_1/2 - 1}^2 (q_1/2)! \binom{q_1}{q_1/2}^2 q_1! \|f \tilde{\otimes}_{q_1/2} f\|_{L^2(T^{q_1})}^2 \\ & - 12q_1!(q_1/2)! \binom{q_1}{q_1/2}^2 \langle f, f \tilde{\otimes}_{q_1/2} f \rangle_{L^2(T^{q_1})} \\ & = \frac{3}{2} \frac{(q_1!)^5}{((q_1/2)!)^6} \|f \tilde{\otimes}_{q_1/2} f - c_{q_1} f\|_{L^2(T^{q_1})}^2, \end{aligned}$$

where c_{q_1} is defined by $c_{q_1} = \frac{1}{(q_1/2)!(q_1/2-1)^2} = \frac{4}{(q_1/2)!(q_1/2)^2}$ and a similar relation holds for the function h with q_2, c_{q_2} instead of q_1, c_{q_1} respectively, where $c_{q_2} = \frac{1}{(q_2/2)!(q_2/2-1)^2} = \frac{4}{(q_2/2)!(q_2/2)^2}$.

$$\begin{aligned} \mathbf{E}(Z^4) - 12\mathbf{E}(Z^3) &= 12v^2 - 48v \\ &+ \frac{3}{q_1} \sum_{p=1, p \neq q_1/2}^{q_1-1} q_1^2(p-1)! \binom{q_1-1}{p-1}^2 p! \binom{q_1}{p}^2 (2q_1-2p)! \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q_1-p)})}^2 \\ &+ \frac{3}{2} \frac{(q_1!)^5}{((q_1/2)!)^6} \|f \tilde{\otimes}_{q_1/2} f - c_{q_1} f\|_{L^2(T^{q_1})}^2 \\ &+ \frac{3}{q_2} \sum_{p=1, p \neq q_2/2}^{q_2-1} q_2^2(p-1)! \binom{q_2-1}{p-1}^2 p! \binom{q_2}{p}^2 (2q_2-2p)! \|h \tilde{\otimes}_p h\|_{L^2(T^{2(q_2-p)})}^2 \\ &+ \frac{3}{2} \frac{(q_2!)^5}{((q_2/2)!)^6} \|h \tilde{\otimes}_{q_2/2} h - c_{q_2} h\|_{L^2(T^{q_2})}^2. \end{aligned}$$

From (ii), it follows that

$$\begin{aligned} & \frac{3}{q_1} \sum_{p=1, p \neq q_1/2}^{q_1-1} q_1^2(p-1)! \binom{q_1-1}{p-1}^2 p! \binom{q_1}{p}^2 (2q_1-2p)! \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q_1-p)})}^2 \\ &+ \frac{3}{2} \frac{(q_1!)^5}{((q_1/2)!)^6} \|f \tilde{\otimes}_{q_1/2} f - c_{q_1} f\|_{L^2(T^{q_1})}^2 \\ &+ \frac{3}{q_2} \sum_{p=1, p \neq q_2/2}^{q_2-1} q_2^2(p-1)! \binom{q_2-1}{p-1}^2 p! \binom{q_2}{p}^2 (2q_2-2p)! \|h \tilde{\otimes}_p h\|_{L^2(T^{2(q_2-p)})}^2 \\ &+ \frac{3}{2} \frac{(q_2!)^5}{((q_2/2)!)^6} \|h \tilde{\otimes}_{q_2/2} h - c_{q_2} h\|_{L^2(T^{q_2})}^2 = 0, \end{aligned}$$

which leads to the conclusion as all the summands are positive, that is

$$\begin{aligned} & \|f \tilde{\otimes}_{q_1/2} f - c_{q_1} f\|_{L^2(T^{q_1})} = \|h \tilde{\otimes}_{q_2/2} h - c_{q_2} h\|_{L^2(T^{q_2})} = 0 \text{ and} \\ & \|f \tilde{\otimes}_p f\|_{L^2(T^{2(q_1-p)})} = \|h \tilde{\otimes}_r h\|_{L^2(T^{2(q_2-p)})} = 0 \end{aligned} \tag{16}$$

for every $p = 1, \dots, q_1 - 1$ such that $p \neq q_1/2$ and for every $r = 1, \dots, q_2 - 1$ such that $r \neq q_2/2$;

This implies

$$\begin{aligned} \|f \tilde{\otimes}_{q_1/2} f - c_{q_1} f\|_{L^2(T^{q_1})} &= \|h \tilde{\otimes}_{q_2/2} h - c_{q_2} h\|_{L^2(T^{q_2})} = 0 \text{ and} \\ \|f \otimes_p f\|_{L^2(T^{2(q_1-p)})} &= \|h \otimes_r h\|_{L^2(T^{2(q_2-p)})} = 0 \end{aligned} \quad (17)$$

for every $p = 1, \dots, q_1 - 1$ such that $p \neq q_1/2$ and for every $r = 1, \dots, q_2 - 1$ such that $r \neq q_2/2$ (see [5], Theorem 1.2.).

We will compute $\mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right)$. Let us start with G_Z .

$$\begin{aligned} G_Z &= \langle DZ, -DL^{-1}Z \rangle_{L^2(T)} = \langle DI_{q_1}(f) + DI_{q_2}(h), -DL^{-1}I_{q_1}(f) - DL^{-1}I_{q_2}(h) \rangle_{L^2(T)} \\ &= \langle DI_{q_1}(f), -DL^{-1}I_{q_1}(f) \rangle_{L^2(T)} + \langle DI_{q_2}(h), -DL^{-1}I_{q_2}(h) \rangle_{L^2(T)} \\ &\quad + \langle DI_{q_1}(f), -DL^{-1}I_{q_2}(h) \rangle_{L^2(T)} + \langle DI_{q_2}(h), -DL^{-1}I_{q_1}(f) \rangle_{L^2(T)}. \end{aligned}$$

From Lemma 2, it follows that $\langle DI_{q_1}(f), -DL^{-1}I_{q_2}(h) \rangle_{L^2(T)} = \langle DI_{q_2}(h), -DL^{-1}I_{q_1}(f) \rangle_{L^2(T)} = 0$. Thus,

$$G_Z = q_1^{-1} \|DI_{q_1}(f)\|_{L^2(T)}^2 + q_2^{-1} \|DI_{q_2}(h)\|_{L^2(T)}^2.$$

It follows that

$$\begin{aligned} &\mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) \\ &= \mathbf{E} \left(\left(2\nu_1 + 2\nu_2 + 2I_{q_1}(f) + 2I_{q_2}(h) - q_1^{-1} \|DI_{q_1}(f)\|_{L^2(T)}^2 - q_2^{-1} \|DI_{q_2}(h)\|_{L^2(T)}^2 \right)^2 \right) \\ &= \mathbf{E} \left(\left(q_1^{-1} \|DI_{q_1}(f)\|_{L^2(T)}^2 - 2I_{q_1}(f) - 2\nu_1 \right)^2 \right) \\ &\quad + \mathbf{E} \left(\left(q_2^{-1} \|DI_{q_2}(h)\|_{L^2(T)}^2 - 2I_{q_2}(h) - 2\nu_2 \right)^2 \right) \\ &\quad + 2\mathbf{E} \left(\left(q_1^{-1} \|DI_{q_1}(f)\|_{L^2(T)}^2 - 2I_{q_1}(f) - 2\nu_1 \right) \left(q_2^{-1} \|DI_{q_2}(h)\|_{L^2(T)}^2 - 2I_{q_2}(h) - 2\nu_2 \right) \right). \end{aligned}$$

We use Lemma 1 to write

$$\mathbf{E} \left(\left(q_1^{-1} \|DI_{q_1}(f)\|_{L^2(T)}^2 - 2I_{q_1}(f) - 2\nu_1 \right) \left(q_2^{-1} \|DI_{q_2}(h)\|_{L^2(T)}^2 - 2I_{q_2}(h) - 2\nu_2 \right) \right) = 0.$$

Thus,

$$\begin{aligned} \mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) &= q_1^{-1} \mathbf{E} \left(\left(\|DI_{q_1}(f)\|_{L^2(T)}^2 - 2q_1 I_{q_1}(f) - 2q_1 \nu_1 \right)^2 \right) \\ &\quad + q_2^{-1} \mathbf{E} \left(\left(\|DI_{q_2}(h)\|_{L^2(T)}^2 - 2q_2 I_{q_2}(h) - 2q_2 \nu_2 \right)^2 \right). \end{aligned}$$

Relation (17) and the calculations contained in [5] imply that the above two summands vanish. It finally follows from this that

$$\mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) = 0.$$

Proof of (i) \rightarrow (ii). Suppose that (ii) holds. We have proven that

$$\mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) = 0 \Rightarrow \begin{cases} \mathbf{E} \left(\left(\|DI_{q_1}(f)\|_{L^2(T)}^2 - 2q_1 I_{q_1}(f) - 2q_1 \nu_1 \right)^2 \right) = 0 \\ \mathbf{E} \left(\left(\|DI_{q_2}(h)\|_{L^2(T)}^2 - 2q_2 I_{q_2}(h) - 2q_2 \nu_2 \right)^2 \right) = 0. \end{cases}$$

From Theorem 1.2 in [5] it follows that $I_{q_1}(f) \sim F(\nu_1)$ and $I_{q_2}(h) \sim F(\nu_2)$. $I_{q_1}(f)$ and $I_{q_2}(h)$ being independent, we use the convolution property of Gamma random variables to state that $Z = I_{q_1}(f) + I_{q_2}(h) \sim F(\nu_1 + \nu_2) \sim F(\nu)$. ■

Remark 2. The proof of the above theorem shows that the affirmations (i) and (ii) are equivalent with relations (10), (11), (16) and (17).

Following exactly the lines of the proof of Theorem 1 it is possible to characterize random variables given by a sum of independent multiple integrals that converges in law to a Gamma distribution.

Theorem 2. Fix $\nu_1, \nu_2, \nu > 0$ such that $\nu_1 + \nu_2 = \nu$ and let $F(\nu)$ be a real-valued random variable with characteristic function given by (1). Fix two even integers $q_1 \geq 2$ and $q_2 \geq 2$. For any sequence $(f_k)_{k \geq 1} \subset L^2(T^{q_1})$ and $(h_k)_{k \geq 1} \subset L^2(T^{q_2})$ (f_k and h_k are symmetric for every $k \geq 1$) such that

$$\mathbf{E} \left(I_{q_1}(f_k)^2 \right) = q_1! \|f_k\|_{L^2(T^{q_1})}^2 \xrightarrow{k \rightarrow +\infty} 2\nu_1 \quad \text{and} \quad \mathbf{E} \left(I_{q_2}(h_k)^2 \right) = q_2! \|h_k\|_{L^2(T^{q_2})}^2 \xrightarrow{k \rightarrow +\infty} 2\nu_2,$$

and such that $X_k = I_{q_1}(f_k)$ and $Y_k = I_{q_2}(h_k)$ are independent for any $k \geq 1$, define the random variable

$$Z_k = X_k + Y_k = I_{q_1}(f_k) + I_{q_2}(h_k) \quad \forall k \geq 1.$$

Under those conditions, the following two conditions are equivalent:

(i) $\mathbf{E} \left(\left(2\nu + 2Z_k - \langle DZ_k, -DL^{-1}Z_k \rangle_{L^2(T)} \right)^2 \right) \xrightarrow{k \rightarrow +\infty} 0;$

(ii) $Z_k \xrightarrow[k \rightarrow +\infty]{\text{Law}} F(\nu);$

Cramér’s theorem for Gamma random variables in the setting of multiple stochastic integrals is a corollary of Theorem 1. We have the following :

Theorem 3. Let $Z = X + Y = I_{q_1}(f) + I_{q_2}(h)$, $q_1, q_2 \geq 2$, $f \in L^2(T^{q_1}), h \in L^2(T^{q_2})$ symmetric, be such that X, Y are independent and

$$\mathbf{E} \left(Z^2 \right) = 2\nu, \mathbf{E} \left(X^2 \right) = q_1! \|f\|_{L^2(T^{q_1})}^2 = 2\nu_1, \mathbf{E} \left(Y^2 \right) = q_2! \|h\|_{L^2(T^{q_2})}^2 = 2\nu_2$$

with $\nu = \nu_1 + \nu_2$. Furthermore, let’s assume that $Z \sim F(\nu)$. Then,

$$X \sim F(\nu_1) \quad \text{and} \quad Y \sim F(\nu_2).$$

Proof: Theorem 1 states that $Z \sim F(\nu) \Leftrightarrow \mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) = 0$ and we proved that

$$\mathbf{E} \left((2\nu + 2Z - G_Z)^2 \right) = \mathbf{E} \left((2\nu_1 + 2X - G_X)^2 \right) + \mathbf{E} \left((2\nu_2 + 2Y - G_Y)^2 \right).$$

Both summands being positive, it follows that

$$\mathbf{E} \left((2\nu_1 + 2X - G_X)^2 \right) = 0 \quad \text{and} \quad \mathbf{E} \left((2\nu_2 + 2Y - G_Y)^2 \right) = 0.$$

Applying theorem 1 to X and Y separately gives us $\mathbf{E} \left((2\nu_1 + 2X - G_X)^2 \right) \Leftrightarrow X \sim F(\nu_1)$ and $\mathbf{E} \left((2\nu_2 + 2Y - G_Y)^2 \right) \Leftrightarrow Y \sim F(\nu_2)$. ■

It is immediate to give an asymptotic version of Theorem 3.

Theorem 4. Let $Z_k = X_k + Y_k = I_{q_1}(f_k) + I_{q_2}(h_k)$, $f_k \in L^2(T^{q_1})$, $h_k \in L^2(T^{q_2})$ symmetric for $k \geq 1$, $q_1, q_2 \geq 2$, be such that X_k, Y_k are independent for every $k \geq 1$ and

$$\mathbf{E}\left(Z_k^2\right) \xrightarrow{k \rightarrow +\infty} 2\nu, \mathbf{E}\left(X_k^2\right) = q_1! \|f\|_{L^2(T^{q_1})}^2 \xrightarrow{k \rightarrow +\infty} 2\nu_1, \mathbf{E}\left(Y_k^2\right) = q_2! \|h\|_{L^2(T^{q_2})}^2 \xrightarrow{k \rightarrow +\infty} 2\nu_2$$

with $\nu = \nu_1 + \nu_2$. Furthermore, let's assume that $Z_k \xrightarrow{k \rightarrow +\infty} F(\nu)$ in distribution. Then,

$$X_k \xrightarrow{k \rightarrow +\infty} F(\nu_1) \quad \text{and} \quad Y_k \xrightarrow{k \rightarrow +\infty} F(\nu_2).$$

Remark 3. i) From Corollary 4.4. in [5] it follows that actually there are no Gamma distributed random variables in a chaos of order bigger or equal than 4. (We actually conjecture that a Gamma distributed random variable given by a multiple integral can only live in the second Wiener chaos). In this sense Theorem 3 contains a limited number of examples. By contrary, the asymptotic Cramér theorem (Theorem 4) is more interesting and more general since there exists a large class of variables which are asymptotically Gamma distributed.

ii) Theorem 3 cannot be applied directly to random variables with law $\Gamma(a, \lambda)$ (as defined in the Introduction) because such random variables are not centered and then they cannot live in a finite Wiener chaos. But, it is not difficult to understand that if $X = I_{q_1} + c$ is a random variable which is independent of $Y = I_{q_2} + d$ (and assume that the first two moments of X and Y are the same as the moment of the corresponding Gamma distributions), and if $X + Y \sim \Gamma(a + b, \lambda)$ then X has the distribution $\Gamma(a, \lambda)$ and Y has the distribution $\Gamma(b, \lambda)$.

iii) Several results of the paper (Lemmas 1 and 2) holds for strongly independent random variables. Nevertheless, the key results (Theorems 1 and 2 that allows to prove Cramér's theorem and its asymptotic variant are not true for strongly independent random variables (actually the implication ii) \rightarrow i) in these results, whose proof is based on the differential equation satisfied by the characteristic function of the Gamma distribution, does not work.

4 Counterexample in the general case

We will see in this section that Theorem 3 does not hold for random variables which have a chaos decomposition into an infinite sum of multiple stochastic integrals. We construct a counterexample in this sense. What is more interesting is that the random variables defined in the below example are not only independent, they are *strongly independent* (see the definition above).

Example 1. Let $\epsilon(\lambda)$ denote the exponential distribution with parameter λ and let $b(p)$ denote the Bernoulli distribution with parameter p . Let $X = A - 1$ and $Y = 2\varpi B - 1$, where $A \sim \epsilon(1)$, $B \sim \epsilon(1)$, $\varpi \sim b(\frac{1}{2})$ and A, B and ϖ are mutually independent. This implies that X and Y are independent. We have $\mathbf{E}(X) = \mathbf{E}(Y) = 0$ as well as $\mathbf{E}(X^2) = 1$ and $\mathbf{E}(Y^2) = 3$. Consider also $Z = X + Y$. Observe that X, Y and Z match every condition of theorem 3, but X and Y are not multiple stochastic integrals in a fixed Wiener chaos (see the next proposition for more details). We have the following : $Z \sim F(2)$, but Y is not Gamma distributed.

Proof: We know that

$$\mathbf{E}\left(e^{itX}\right) = \mathbf{E}\left(e^{it(A-1)}\right) = e^{-it} \mathbf{E}\left(e^{itA}\right) = \frac{e^{-it}}{1-it}$$

and that

$$\begin{aligned}\mathbf{E}\left(e^{itY}\right) &= \mathbf{E}\left(e^{it(2\varpi B-1)}\right) = e^{-it}\mathbf{E}\left(e^{it2\varpi B}\right) = e^{-it}\left(\frac{1}{2}\mathbf{E}\left(e^{it2B}\right) + \frac{1}{2}\right) \\ &= e^{-it}\left(\frac{1}{2}\frac{1}{1-2it} + \frac{1}{2}\right) = e^{-it}\frac{1-it}{1-2it}.\end{aligned}$$

Observe at this point that the characteristic function of Y proves that Y is not Gamma distributed. Let us compute the characteristic function of Z . We have

$$\mathbf{E}\left(e^{itZ}\right) = \mathbf{E}\left(e^{it(X+Y)}\right) = \mathbf{E}\left(e^{itX}\right)\mathbf{E}\left(e^{itY}\right) = \frac{e^{-it}}{1-it}e^{-it}\frac{1-it}{1-2it} = \frac{e^{-2it}}{1-2it} = \mathbf{E}\left(e^{itF(2)}\right).$$

■

Remark 4. It is also possible to construct a similar example for the laws $\Gamma(a, \lambda), \Gamma(b, \lambda)$ instead of $F(v_1), F(v_2)$.

The following proposition shows that this counterexample accounts for independent random variables but also for strongly independent random variables.

Proposition 2. X and Y as defined in Example 1 are strongly independent.

Proof: In order to prove that X and Y are strongly independent, we need to compute their Wiener chaos expansions in order to emphasize the fact that all the components of these Wiener Chaos expansions are mutually independent. Consider a standard Brownian motion B indexed on $L^2(T) = L^2((0, T))$. Consider $h_1, \dots, h_5 \in L^2(T)$ such that $\|h_i\|_{L^2(T)} = 1$ for every $1 \leq i \leq 5$ and such that $W(h_i)$ and $W(h_j)$ are independent for every $1 \leq i, j \leq 5, i \neq j$. First notice that the random variables $A = \frac{1}{2}(W(h_1)^2 + W(h_2)^2)$ and $B = \frac{1}{2}(W(h_4)^2 + W(h_5)^2)$ are independent (this is obvious) and have the exponential distribution with parameter 1. Also, note that the random variable $\varpi = \frac{1}{2}\text{sign}(W(h_3)) + \frac{1}{2}$ has the Bernoulli distribution and is independent from A and B . As in Example 1, set $X = A - 1$ and $Y = 2\varpi B - 1$. X and Y are as defined in Example 1. Let us now compute their Wiener chaos decompositions. We have

$$A = \frac{1}{2}(W(h_1)^2 + W(h_2)^2) = \frac{1}{2}(I_1(h_1)^2 + I_1(h_2)^2) = \frac{1}{2}(2 + I_2(h_1^{\otimes 2}) + I_2(h_2^{\otimes 2})),$$

and similarly $B = \frac{1}{2}(2 + I_2(h_4^{\otimes 2}) + I_2(h_5^{\otimes 2}))$. Therefore, we have

$$X = I_2\left(\frac{h_1^{\otimes 2} + h_2^{\otimes 2}}{2}\right).$$

From [2], Lemma 3, we know that

$$\text{sign}(W(h_3)) = \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)}),$$

where $b_{2k+1} = \frac{2(-1)^k}{(2k+1)\sqrt{2\pi k!2^k}}$. It follows that $\varpi = \frac{1}{2} + \frac{1}{2} \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)})$, and

$$\begin{aligned} Y &= \left(1 + \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)})\right) \left(1 + \frac{1}{2} I_2(h_4^{\otimes 2}) + \frac{1}{2} I_2(h_5^{\otimes 2})\right) - 1 \\ &= \frac{1}{2} I_2(h_4^{\otimes 2}) + \frac{1}{2} I_2(h_5^{\otimes 2}) + \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)}) + \frac{1}{2} \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)}) I_2(h_4^{\otimes 2}) \\ &\quad + \frac{1}{2} \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)}) I_2(h_5^{\otimes 2}). \end{aligned}$$

Using the multiplication formula for multiple stochastic integrals, we obtain

$$\begin{aligned} Y &= \frac{1}{2} I_2(h_4^{\otimes 2}) + \frac{1}{2} I_2(h_5^{\otimes 2}) + \sum_{k \geq 0} b_{2k+1} I_{2k+1}(h_3^{\otimes(2k+1)}) \\ &\quad + \frac{1}{2} \sum_{k \geq 0} b_{2k+1} \sum_{r=0}^{(2k+1) \wedge 2} r! \binom{2}{r} \binom{2k+1}{r} I_{2k+3-2r}(h_3^{\otimes(2k+1)} \otimes_r h_4^{\otimes 2}) \\ &\quad + \frac{1}{2} \sum_{k \geq 0} b_{2k+1} \sum_{r=0}^{(2k+1) \wedge 2} r! \binom{2}{r} \binom{2k+1}{r} I_{2k+3-2r}(h_3^{\otimes(2k+1)} \otimes_r h_5^{\otimes 2}). \end{aligned}$$

At this point, it is clear that X and Y are strongly independent. ■

References

- [1] H. Cramér (1936): Über eine Eigenschaft der normalen Verteilungsfunktion. *Math. Z.*, 41(2), 405-414. [MR1545629](#)
- [2] Y. Hu and D. Nualart (2005): Some processes associated with fractional Bessel processes. *Journal of Theoretical Probability*, 18 no. 2, 377-397. [MR2137449](#)
- [3] P. Malliavin (1997): *Stochastic Analysis*. Springer-Verlag. [MR1450093](#)
- [4] D. Nualart (2006): *Malliavin Calculus and Related Topics. Second Edition*. Springer. [MR2200233](#)
- [5] I. Nourdin and G. Peccati (2009): Noncentral convergence of multiple integrals. *The Annals of Probability*, 37 no. 4, 1412-1426. [MR2546749](#)
- [6] I. Nourdin and G. Peccati (2007): Stein's method on Wiener chaos. *Probability Theory and Related Fields*. 145 (1-2), 75-118. [MR2520122](#)
- [7] C.A. Tudor (2008): Asymptotic Cramér's theorem and analysis on Wiener space. Preprint, to appear in *Séminaire de Probabilités, Lecture Notes in Mathematics*.
- [8] A.S. Ustunel and M. Zakai (1989): On independence and conditioning on Wiener space. *The Annals of Probability*, 17 no. 4, 1441-1453. [MR1048936](#)