



Vol. 10 (2005), Paper no. 38, pages 1260-1285.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

### Limit Theorems for Self-normalized Large Deviation <sup>1</sup>

**Qiyang Wang**

School of Mathematics and Statistics

The University of Sydney, Sydney

NSW 2006, Australia

E-mail: [qiyang@maths.usyd.edu.au](mailto:qiyang@maths.usyd.edu.au)

URL: <http://www.maths.usyd.edu.au/u/qiyang/>

**Abstract.** Let  $X, X_1, X_2, \dots$  be i.i.d. random variables with zero mean and finite variance  $\sigma^2$ . It is well known that a finite exponential moment assumption is necessary to study limit theorems for large deviation for the standardized partial sums. In this paper, limit theorems for large deviation for self-normalized sums are derived only under finite moment conditions. In particular, we show that, if  $EX^4 < \infty$ , then

$$\frac{P(S_n/V_n \geq x)}{1 - \Phi(x)} = \exp \left\{ -\frac{x^3 EX^3}{3\sqrt{n}\sigma^3} \right\} \left[ 1 + O\left(\frac{1+x}{\sqrt{n}}\right) \right],$$

for  $x \geq 0$  and  $x = O(n^{1/6})$ , where  $S_n = \sum_{i=1}^n X_i$  and  $V_n = (\sum_{i=1}^n X_i^2)^{1/2}$ .

**Key Words and Phrases:** Cramér large deviation, limit theorem, self-normalized sum.

**AMS Subject Classification (2000):** Primary 60F05, Secondary 62E20.

Submitted to EJP on September 24, 2004. Final version accepted on September 16, 2005.

---

<sup>1</sup>This research is supported in part by ARC DP0451722

# 1 Introduction and main results

Let  $X, X_1, X_2, \dots$ , be a sequence of non-degenerate independent and identically distributed (i.i.d.) random variables with zero mean. Set

$$S_n = \sum_{i=1}^n X_i, \quad V_n^2 = \sum_{i=1}^n X_i^2, \quad n \geq 1.$$

The self-normalized version of the classical central limit theorem states that, as  $n \rightarrow \infty$ ,

$$\sup_x |P(S_n \geq xV_n) - \{1 - \Phi(x)\}| \rightarrow 0,$$

if and only if the distribution of  $X$  is in the domain of attraction of the normal law, where  $\Phi(x)$  denotes the standard normal distribution function. This beautiful self-normalized central limit theorem was conjectured by Logan, Mallows, Rice and Shepp (1973), and latterly proved by Gine, Götze and Mason (1997). For a short summary of developments that have eventually led to Gine, Götze and Mason (1997), we refer to the Introduction of the latter paper.

The self-normalized central limit theorem is useful when  $x$  is not too large or when the error is well estimated. There are two approaches for estimating the error of the normal approximation. One approach is to investigate the absolute error in the self-normalized central limit theorem via Berry-Esseen bounds or Edgeworth expansions. This has been done by many researchers. For details, we refer to Slavova (1985), Hall (1988) and Bentkus and Götze (1996) for the Berry-Esseen bounds, Wang and Jing (1999) for an exponential nonuniform Berry-Esseen bound, Hall (1987) as well as Hall and Jing (1995) for Edgeworth expansions. See also van Zwet (1984), Friedrich (1989), Bentkus, Bloznelis and Götze (1996), Bentkus, Götze and van Zwet (1997), Putter and van Zwet (1998) and Wang, Jing and Zhao (2000). Another approach is to estimate the relative error  $P(S_n \geq xV_n)/(1 - \Phi(x))$ . In this direction, Jing, Shao and Wang (2003) refined Shao (1999), Wang and Jing (1999) as well as Chistyakov and Götze (2003), and obtained the following result: if  $0 < \sigma^2 = EX^2 < \infty$ , then there exists an absolute constant  $A > 0$  such that

$$\exp \left\{ -A(1+x)^2 \tilde{\Delta}_{n,x} \right\} \leq \frac{P(S_n \geq xV_n)}{1 - \Phi(x)} \leq \exp \left\{ A(1+x)^2 \tilde{\Delta}_{n,x} \right\}, \quad (1.1)$$

for all  $x \geq 0$  satisfying  $\tilde{\Delta}_{n,x} \leq 1/A$ , where

$$\tilde{\Delta}_{n,x} = \sigma^{-2} EX^2 I_{|X| \geq \sqrt{n}\sigma/(1+x)} + (1+x)\sigma^{-3} n^{-1/2} E|X|^3 I_{|X| \leq \sqrt{n}\sigma/(1+x)}.$$

Jing, Shao and Wang (2003) actually established (1.1) for independent random variables that are not necessarily identically distributed. It follows from (1.1) that if  $E|X|^3 < \infty$ , then

$$\frac{P(S_n \geq xV_n)}{1 - \Phi(x)} = 1 + O(1)(1+x)^3 n^{-1/2} \sigma^{-3} E|X|^3, \quad (1.2)$$

for  $0 \leq x \leq An^{1/6}\sigma/(E|X|^3)^{1/3}$ .

Result (1.2) is useful in statistics because it provides not only the relative error but also a Berry-Esseen type rate of convergence. Indeed, as a direct consequence of this result, it has been shown in Jing, Shao and Wang (2003) that bootstrapped studentized  $t$ -statistics possess large deviation properties in the region  $0 \leq x \leq o(n^{1/6})$  under only a finite third moment condition. However, (1.1) as well as (1.2) does not capture the term with  $n^{-1/2}$  explicitly. This short has limited further applications of the self-normalized large deviation.

In this paper we investigate the limit theorems for self-normalized large deviation. Under finite moment conditions, a leading term with  $n^{-1/2}$  in (1.1) and (1.2) is obtained explicitly.

**THEOREM 1.1.** *Assume that  $EX^4 < \infty$ . Then, for  $x \geq 0$  and  $x = O(n^{1/6})$ ,*

$$\frac{P(S_n \geq xV_n)}{1 - \Phi(x)} = \exp\left\{-\frac{x^3 EX^3}{3\sqrt{n}\sigma^3}\right\} \left[1 + O\left(\frac{1+x}{\sqrt{n}}\right)\right], \quad (1.3)$$

$$\frac{P(S_n \leq -xV_n)}{\Phi(-x)} = \exp\left\{\frac{x^3 EX^3}{3\sqrt{n}\sigma^3}\right\} \left[1 + O\left(\frac{1+x}{\sqrt{n}}\right)\right]. \quad (1.4)$$

If in addition  $EX^3 = 0$ , then, for  $|x| = O(n^{1/6})$ ,

$$P(S_n \leq xV_n) - \Phi(x) = O(n^{-1/2}e^{-x^2/2}). \quad (1.5)$$

Write  $\mathcal{L}_{n,x} = (1+x)\rho_n + \Delta_{n,x}$ , where  $\rho_n = E|X|^3/(\sqrt{n}\sigma^3)$  and

$$\Delta_{n,x} = (1+x)^3 \sigma^{-3} n^{-1/2} E|X|^3 I_{\{|X| > \sqrt{n}\sigma/(1+x)\}} + (1+x)^4 \sigma^{-4} n^{-1} EX^4 I_{\{|X| \leq \sqrt{n}\sigma/(1+x)\}}.$$

Furthermore, we obtain the following bounds which refine (1.1) under finite third moment condition in the region  $0 \leq x \leq c\rho_n^{-1/2}$ , where  $c$  is an absolute constant.

**THEOREM 1.2.** *Assume that  $E|X|^3 < \infty$ . Then, there exists a positive absolute constant  $c$  such that, for  $0 \leq x \leq c\rho_n^{-1/2}$ ,*

$$\exp\left(-\frac{x^3 EX^3}{3\sqrt{n}\sigma^3} - A\mathcal{L}_{n,x}\right) \leq \frac{P(S_n \geq xV_n)}{1 - \Phi(x)} \leq \exp\left(-\frac{x^3 EX^3}{3\sqrt{n}\sigma^3} + A\mathcal{L}_{n,x}\right). \quad (1.6)$$

where  $A$  is an absolute positive constant.

*REMARK 1.1.* Similar results to those in Theorem 1.1 hold for the standardized mean under much stronger conditions. For instance, it follows from Section 5.8 of Petrov (1995) that, for  $x \geq 0$  and  $x = O(n^{1/6})$ ,

$$\frac{P(S_n \geq x\sqrt{n}\sigma)}{1 - \Phi(x)} = \exp\left\{\frac{x^3 EX^3}{6\sqrt{n}\sigma^3}\right\} \left[1 + O\left(\frac{1+x}{\sqrt{n}}\right)\right],$$

$$\frac{P(S_n \leq -x\sqrt{n}\sigma)}{\Phi(-x)} = \exp\left\{-\frac{x^3 EX^3}{6\sqrt{n}\sigma^3}\right\} \left[1 + O\left(\frac{1+x}{\sqrt{n}}\right)\right].$$

hold only when Cramér's condition is satisfied, i.e.,  $Ee^{tX} < \infty$  for  $t$  being in a neighborhood of zero. We also notice that there are different formulae for the self-normalized and standardized cases.

*REMARK 1.2.* It is readily seen that  $\mathcal{L}_{n,x_n} = o(x_n^3/\sqrt{n})$  for  $x_n \rightarrow \infty$  and  $x_n = O(\rho_n^{-1/2})$ . Hence  $-\frac{x^3 EX^3}{3\sqrt{n}\sigma^3}$  in (1.6) provides a leading term in this case. However it remains an open problem for more refined results.

This paper is organized as follows. The proofs of main results will be given in Section 3. Next section we present two auxiliary theorems that will be used in the proofs of main results. The proofs of these auxiliary theorems will be postponed to Sections 4 and 5 respectively. Without loss of generality, throughout the paper, we assume  $\sigma^2 = EX^2 = 1$  and denote by  $A, A_1, A_2, \dots$  and  $c, C, C_1, C_2, \dots$  absolute positive constants, which may be different at each occurrence. If a constant depends on a parameter, say  $u$ , then we write  $A(u)$ . In addition to the notation for  $\mathcal{L}_{n,x}$  and  $\Delta_{n,x}$  defined in Theorem 1.2, we always let

$$\rho_n = n^{-1/2}E|X|^3 \quad \text{and} \quad \Psi_n(x) = \left[1 - \Phi(x)\right] \exp\left(-\frac{x^3 EX^3}{3\sqrt{n}}\right).$$

## 2 Two auxiliary theorems

Throughout the section we assume that  $X, X_1, X_2, \dots$ , are i.i.d. random variables satisfying  $EX = 0, EX^2 = 1$  and  $E|X|^3 < \infty$ . Two theorems in this section are established under quite general setting, which will be interesting in themselves. The proofs of these two theorems will be given in Sections 4 and 5 respectively.

**THEOREM 2.1.** Let  $h = x/B_n$ , where  $B_n$  is a sequence of positive constants with

$$|B_n^2 - n| \leq C_1 n EX^2 I_{\{|X| \geq \sqrt{n}/(1+x)\}}. \quad (2.1)$$

Suppose that  $\eta_j := \eta_n(x, X_j)$ ,  $1 \leq j \leq n$ , satisfy the conditions:

$$\left| Ee^{h\eta_j} - 1 - \frac{x^2}{2B_n^2} + \frac{x^3}{3B_n^3} EX^3 \right| \leq C_2 n^{-1} \Delta_{n,x}, \quad (2.2)$$

$$\left| E\eta_j e^{h\eta_j} - \frac{x}{B_n} \right| \leq C_3 x^2 n^{-1/2} \rho_n, \quad (2.3)$$

$$\left| E\eta_j^2 e^{h\eta_j} - 1 \right| \leq C_4 x \rho_n, \quad (2.4)$$

$$E|\eta_j|^3 e^{h\eta_j} \leq C_5 E|X|^3. \quad (2.5)$$

Then, for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small and for any  $|b| \leq x^2/4$ ,

$$P\left(\sum_{j=1}^n \eta_j \geq (x + bx^{-1})B_n\right) \leq (1 + 2|b|x^{-2})e^{-b}\Psi_n(x) \exp(A\mathcal{L}_{n,x}); \quad (2.6)$$

for  $2 \leq x \leq c_1\rho_n^{-1/2}$  with  $c_1$  sufficiently small,

$$P\left(\sum_{j=1}^n \eta_j \geq x B_n\right) \geq \Psi_n(x) \exp(-A\mathcal{L}_{n,x}). \quad (2.7)$$

**THEOREM 2.2.** Write, for  $1 \leq m \leq n$ ,

$$T_m = \frac{1}{\sqrt{n}} \sum_{j=1}^m \zeta_j \quad \text{and} \quad \Lambda_{n,m} = \frac{1}{n^2} \sum_{k=1}^{m-1} \sum_{j=k+1}^n \psi_{k,j},$$

where  $\zeta_j := \zeta_n(x, X_j)$  and  $\psi_{k,j} := \psi_n(x, X_k, X_j)$  satisfy the conditions:

$$\left| E\zeta_j^2 - 1 + \frac{x}{\sqrt{n}} EX^3 \right| \leq C_6 \Delta_{n,x}/(1+x)^2, \quad (2.8)$$

$$\left| E\zeta_j^3 - EX^3 \right| \leq C_7 \sqrt{n} \Delta_{n,x}/(1+x)^3, \quad (2.9)$$

$$E\zeta_j = 0, \quad |\zeta_j| \leq C_8 \sqrt{n}/(1+x), \quad E\zeta_j^4 \leq C_9 EX^4 I_{\{|X| \leq \sqrt{n}/(1+x)\}}, \quad (2.10)$$

$$E(\psi_{k,j}|X_k) = E(\psi_{k,j}|X_j) = 0, \quad \text{for } k \neq j, \quad (2.11)$$

$$E|\psi_{k,j}| \leq C_{10} |x|, \quad E|\psi_{k,j}|^{3/2} \leq C_{11} |x|^{3/2} (E|X|^3)^2. \quad (2.12)$$

Then, for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small, and for any constants sequence  $\lambda_n(x)$  satisfying  $|\lambda_n(x)| \leq C \Delta_{n,x}/(1+x)$ ,

$$P\left(T_n + \Lambda_{n,n} \geq x + \lambda_n(x)\right) \leq \Psi_n(x) (1 + A\mathcal{L}_{n,x}) + A_1(x\rho_n)^{3/2}. \quad (2.13)$$

### 3 Proofs of main results

*Proof of Theorem 1.1.* Note that  $\mathcal{L}_{n,x} \asymp (1+x)/\sqrt{n}$  for  $x \geq 0$  and  $x = O(n^{1/6})$ . Theorem 1.1 follows immediately from Theorem 1.2. We omit the details.

*Proof of Theorem 1.2.* Without loss of generality, assume  $x \geq 2$ . If  $0 \leq x < 2$ , the results are direct consequences of the Berry-Esseen bound (cf. Bentkus and Götze (1996))

$$\left| P(S_n \geq xV_n) - \{1 - \Phi(x)\} \right| \leq A \rho_n.$$

We first provide four lemmas. For simplicity of presentation, define  $\tau = \sqrt{n}/(1+x)$  and assume  $2 \leq x \leq c \rho_n^{-1/2}$  with  $c$  sufficiently small throughout the section except where we point out.

**LEMMA 3.1.** *We have,*

$$P\left(S_n - \frac{x}{2\sqrt{n}}(V_n^2 - n) \geq x\sqrt{n}\right) \geq \Psi_n(x) \exp(-A \mathcal{L}_{n,x}); \quad (3.1)$$

and for  $2 \leq x \leq c \rho_n^{-1}$  with  $c$  sufficiently small and for arbitrary  $|\delta| \leq x^2/4$ ,

$$P\left(S_n - \frac{x}{2\sqrt{n}}(V_n^2 - n) \geq (x + \delta x^{-1})\sqrt{n}\right) \leq (1 + 2|\delta|x^{-2})e^{-\delta} \Psi_n(x) \exp(A \mathcal{L}_{n,x}). \quad (3.2)$$

*Proof.* We first prove (3.2). Let  $h = x/\sqrt{n}$  and  $\eta_j = X_j - \frac{x}{2\sqrt{n}}(X_j^2 - 1)$ . It follows from (4.12)-(4.14) in Wang and Jing (1999) that

$$\begin{aligned} \left| E\eta_1 e^{h\eta_1} - \frac{x}{\sqrt{n}} \right| &\leq 16 x^2 n^{-1/2} \rho_n, \\ \left| E\eta_1^2 e^{h\eta_1} - 1 \right| &\leq 30 x \rho_n, \\ E|\eta_1|^3 e^{h\eta_1} &\leq 30 E|X|^3. \end{aligned}$$

Thus (3.2) follows immediately from (2.6) in Theorem 2.1 with  $B_n^2 = n$  if we prove

$$\left| Ee^{h\eta_1} - 1 - \frac{x^2}{2n} + \frac{x^3}{3n^{3/2}} EX^3 \right| \leq C n^{-1} \Delta_{n,x}. \quad (3.3)$$

Without loss of generality, assume  $x \leq \rho_n^{-1}/16$ . By noting  $E|X|^3 \geq (EX^2)^{3/2} = 1$ , we get  $h \leq 1/4$ . This implies that  $h\eta_1 = h^2/2 + hX_1 - (hX_1)^2/2 \leq 1$  and  $|h\eta_1|^k e^{h\eta_1} \leq \sup_{s \leq 1} |s|^k e^s \leq e$  for  $k = 0, 1, \dots, 4$ . Therefore, using Taylor's expansion

$$\left| e^x - \sum_{j=0}^k \frac{x^j}{j!} \right| \leq \frac{|x|^{k+1}}{(k+1)!} e^{x \vee 0}, \quad \text{for } k \geq 1,$$

we obtain that

$$\begin{aligned}
Ee^{h\eta_1} &= E\left[1 + h\eta_1 + \frac{1}{2}(h\eta_1)^2 + \frac{1}{6}(h\eta_1)^3 + \frac{\theta}{24}(h\eta_1)^4\right]I_{(|X_1|\leq\tau)} \\
&\quad + E\left(1 + h\eta_1 + \frac{\theta_1}{2}\right)I_{(|X_1|>\tau)} \\
&= 1 + \frac{1}{2}E(h\eta_1)^2I_{(|X_1|\leq\tau)} + \frac{1}{6}E(h\eta_1)^3I_{(|X_1|\leq\tau)} \\
&\quad + \frac{\theta_1}{2}P(|X_1| > \tau) + \frac{\theta}{24}E(h\eta_1)^4I_{(|X_1|\leq\tau)}, \tag{3.4}
\end{aligned}$$

where  $|\theta| \leq e$  and  $|\theta_1| \leq e$ . Since  $EX^2 = 1$ , it is readily seen that

$$\begin{aligned}
\left|E(h\eta_1)^2I_{(|X_1|\leq\tau)} - h^2 + E(hX)^3\right| &\leq 8n^{-1}\Delta_{n,x}, \\
\left|E(h\eta_1)^3I_{(|X_1|\leq\tau)} - E(hX)^3\right| &\leq 32n^{-1}\Delta_{n,x}, \\
E(h\eta_1)^4I_{(|X_1|\leq\tau)} &\leq 16n^{-1}\Delta_{n,x}.
\end{aligned}$$

Taking these estimates back into (3.4), we obtain (3.3), and hence (3.2).

The proof of (3.1) is similar by using (2.7). We omit the details. The proof of Lemma 3.1 is now complete.  $\square$

The next lemma is from Lemma 6.4 in Jing, Shao and Wang (2003).

**LEMMA 3.2.** *Let  $\{\xi_i, 1 \leq i \leq n\}$  be a sequence of independent random variables with  $E\xi_i = 0$  and  $E\xi_i^2 < \infty$ . Then*

$$P\left[\sum_{i=1}^n \xi_i \geq a\left\{4D_n + \left(\sum_{i=1}^n \xi_i^2\right)^{1/2}\right\}\right] \leq 8e^{-a^2/2} \tag{3.5}$$

where  $D_n = \left(\sum_{i=1}^n E\xi_i^2\right)^{1/2}$ .

In Lemmas 3.3-3.4, we use the notation:

$$\begin{aligned}
\bar{X}_i &= X_i I_{\{|X_i|\leq\tau\}}, & \bar{S}_n &= \sum_{i=1}^n \bar{X}_i, & \bar{V}_n^2 &= \sum_{i=1}^n \bar{X}_i^2, & B_n^2 &= \sum_{i=1}^n E\bar{X}_i^2 \\
S_n^{(i)} &= S_n - X_i, & V_n^{(i)} &= (V_n^2 - X_i^2)^{1/2}.
\end{aligned}$$

**LEMMA 3.3.** *We have*

$$P(S_n \geq xV_n) \leq A\Psi_n(x) \exp(A\mathcal{L}_{n,x}). \tag{3.6}$$

*Proof.* Let  $\Omega_n = (1 - x^{-1}/2, 1 + x^{-1}/2)$ . Recalling  $2 \leq x \leq c\rho_n^{-1/2}$  with  $c$  sufficiently small, it follows from (2.27)-(2.29) in Shao (1999) that

$$\begin{aligned} & P(S_n \geq xV_n, V_n^2/n \notin \Omega_n) \\ & \leq P(S_n \geq xV_n, V_n^2 \geq 9n) + P\{S_n \geq xV_n, n(1 + x^{-1}/2) \leq V_n^2 \leq 9n\} \\ & \quad + P\{S_n \geq xV_n, V_n^2 \leq n(1 - x^{-1}/2)\} \\ & \leq 4 \exp(-x^2/2 - x/8 + Ax^3\rho_n) \\ & \leq A\Psi_n(x). \end{aligned}$$

This, together with (3.2), implies that

$$\begin{aligned} P(S_n \geq xV_n) &= P(S_n \geq xV_n, V_n^2/n \in \Omega_n) + P(S_n \geq xV_n, V_n^2/n \notin \Omega_n) \\ &\leq P\left\{S_n - \frac{x}{2\sqrt{n}}(V_n^2 - n) \geq (x - x^{-1}/4)\sqrt{n}\right\} + P(S_n \geq xV_n, V_n^2/n \notin \Omega_n) \\ &\leq A\Psi_n(x) \exp(A\mathcal{L}_{n,x}), \end{aligned}$$

as required. The proof of Lemma 3.3 is complete.  $\square$

**LEMMA 3.4.** *We have*

$$P(S_n \geq xV_n) \leq \Psi_n(x) \exp(A\mathcal{L}_{n,x}) + A_1 e^{-3x^2}; \quad (3.7)$$

$$P(S_n \geq xV_n) \leq \Psi_n(x) \exp(A\mathcal{L}_{n,x}) + A_1 (x\rho_n)^{3/2}. \quad (3.8)$$

*Proof.* As in Wang and Jing (1999), for  $x \geq 2$ ,

$$\begin{aligned} P(S_n \geq xV_n) &\leq P(\bar{S}_n \geq x\bar{V}_n) + \sum_{i=1}^n P[S_n^{(i)} \geq (x^2 - 1)^{1/2}V_n^{(i)}]P(|X_i| > \tau). \\ &= K_n + I_n, \quad \text{say.} \end{aligned} \quad (3.9)$$

It follows easily from Lemma 3.3 that for all  $i$

$$P[S_n^{(i)} \geq (x^2 - 1)^{1/2}V_n^{(i)}] \leq A\Psi_n(x) \exp(A\mathcal{L}_{n,x}).$$

This, together with  $P(|X_i| > \tau) \leq n^{-1}\Delta_{n,x}$ , implies that

$$I_n \leq A\Delta_{n,x} \Psi_n(x) \exp(A\mathcal{L}_{n,x}) \quad (3.10)$$

In view of (3.9) and (3.10), the inequalities (3.7) and (3.8) will follow if we prove

$$K_n \leq \Psi_n(x) \exp(A \mathcal{L}_{n,x}) + A_1 e^{-3x^2}, \quad (3.11)$$

$$K_n \leq \Psi_n(x) \exp(A \mathcal{L}_{n,x}) + A_1 (x\rho_n)^{3/2}, \quad (3.12)$$

for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small.

We first prove (3.11). Let  $D_n^2 = \sum_{i=1}^n E\bar{X}_i^4$  and  $\xi_i = \bar{X}_i^2 - E\bar{X}_i^2$ . By the inequality  $(1+y)^{1/2} \geq 1+y/2-y^2$  for any  $y \geq -1$  and Lemma 3.2,

$$\begin{aligned} K_n &= P\left[\bar{S}_n \geq x\left\{B_n^2 + \sum_{i=1}^n (\bar{X}_i^2 - E\bar{X}_i^2)\right\}^{1/2}\right] \\ &\leq P\left[\bar{S}_n \geq xB_n\left\{1 + \frac{1}{2B_n^2} \sum_{i=1}^n \xi_i - \frac{1}{B_n^4} \left(\sum_{i=1}^n \xi_i\right)^2\right\}\right] \\ &\leq P\left[\left|\sum_{i=1}^n \xi_i\right| \geq \sqrt{6}x\left\{4D_n + \left(\sum_{i=1}^n \xi_i^2\right)^{1/2}\right\}\right] \\ &\quad + P\left[\bar{S}_n \geq xB_n\left\{1 + \frac{1}{2B_n^2} \sum_{i=1}^n \xi_i - \frac{1}{B_n^4} \left(\sum_{i=1}^n \xi_i\right)^2\right\},\right. \\ &\quad \left.\left|\sum_{i=1}^n \xi_i\right| \leq \sqrt{6}x\left\{4D_n + \left(\sum_{i=1}^n \xi_i^2\right)^{1/2}\right\}\right] \\ &\leq 8e^{-3x^2} + K_{n,1}, \end{aligned} \quad (3.13)$$

where, after some algebra (see, Jing, Shao and Wang (2003), for example),

$$K_{n,1} \leq P\left(\sum_{j=1}^n \eta_j \geq xB_n\right),$$

with  $\eta_i = \bar{X}_i - 2^{-1}xB_n^{-1}(\bar{X}_i^2 - E\bar{X}_i^2) + 24x^3B_n^{-3}(\bar{X}_i^4 + 16E\bar{X}_i^4)$ . As in the proof of Lemma 3.1, tedious but elementary calculations show that the inequalities (2.1)-(2.5) hold true for  $B_n^2 = \sum_{i=1}^n E\bar{X}_i^2$  and the  $\eta_i$  defined above. Therefore it follows from (3.2) in Theorem 2.1 that

$$K_{n,1} \leq \Psi_n(x) \exp(A \mathcal{L}_{n,x}), \quad (3.14)$$

for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small. Take this estimate back into (3.13), we get the desired (3.11).

We next prove (3.12). Note that

$$\begin{aligned}
 (\bar{V}_n^2 - n)^2 &= \sum_{j=1}^n (\bar{X}_j^2 - 1)^2 + \sum_{k \neq j} (\bar{X}_k^2 - 1)(\bar{X}_j^2 - 1) \\
 &= \sum_{j=1}^n (\bar{X}_j^2 - 1)^2 + \sum_{k \neq j} (\bar{X}_k^2 - E\bar{X}_k^2)(\bar{X}_j^2 - E\bar{X}_j^2) \\
 &\quad + (n-1)(E\bar{X}_j^2 - 1) \sum_{j=1}^n (2\bar{X}_j^2 - 1 - E\bar{X}_j^2).
 \end{aligned}$$

By the inequality  $(1+y)^{1/2} \geq 1 + y/2 - y^2$  for any  $y \geq -1$  again, we have

$$\begin{aligned}
 K_n &= P\left[\bar{S}_n \geq x\sqrt{n} \left\{1 + \frac{1}{n} \sum_{i=1}^n (\bar{X}_i^2 - 1)\right\}^{1/2}\right] \\
 &\leq P\left[\bar{S}_n \geq x\sqrt{n} \left\{1 + \frac{1}{2n}(\bar{V}_n^2 - n) - \frac{1}{n^2}(\bar{V}_n^2 - n)^2\right\}\right] \\
 &= P\left(\frac{1}{\sqrt{n}} \sum_{j=1}^n \zeta_j + \frac{1}{n^2} \sum_{k=1}^{n-1} \sum_{j=k+1}^n \psi_{k,j} \geq x - \lambda_n(x)\right), \tag{3.15}
 \end{aligned}$$

where

$$\begin{aligned}
 \zeta_j^* &= \bar{X}_j - \frac{x}{2\sqrt{n}}(\bar{X}_j^2 - 1) + \frac{x}{n^{3/2}}(\bar{X}_j^2 - 1)^2 \\
 &\quad + \frac{(n-1)x}{n^{3/2}}(E\bar{X}_1^2 - 1) \sum_{j=1}^n (2\bar{X}_j^2 - 1 - E\bar{X}_j^2), \\
 \zeta_j &= \zeta_j^* - E\zeta_j^* \\
 &= \bar{X}_j - E\bar{X}_j - \frac{x}{2\sqrt{n}} \left(1 + \frac{2}{n} + \frac{2(n-1)}{n} EX^2 I_{(|X| \geq \tau)}\right) (\bar{X}_j^2 - E\bar{X}_j^2) \\
 &\quad + \frac{x}{n^{3/2}} (\bar{X}_j^4 - E\bar{X}_j^4), \\
 \psi_{k,j} &= 2x(\bar{X}_k^2 - E\bar{X}_k^2)(\bar{X}_j^2 - E\bar{X}_j^2), \\
 \lambda_n(x) &= \sqrt{n}E\zeta_1^* \\
 &= \sqrt{n}EXI_{(|X| \geq \tau)} - \frac{x}{2}EX^2I_{(|X| \geq \tau)} + \frac{x}{n}E(\bar{X}_1^2 - 1)^2 - \frac{(n-1)x}{n} \left( EX^2I_{(|X| \geq \tau)} \right)^2.
 \end{aligned}$$

It is easy to see that  $\zeta_j$  satisfy the conditions (2.8)- (2.10),  $\psi_{k,j}$  satisfy the conditions (2.11)- (2.12) in Theorem 2.2 and  $|\lambda_n(x)| \leq 3xn^{-1} + 3(1+x)^{-1}\Delta_{n,x} \leq 6(1+x)^{-1}\Delta_{n,x}$ . Hence, in view of (3.15), (3.12) follows immediately Theorem 2.2. This also completes the proof of Lemma 3.4.  $\square$

After these preliminaries, we are now ready to prove Theorem 1.2.

As is well-known (see Wang and Jing (1999) for example),

$$P(S_n \geq xV_n) \geq P\left(S_n - \frac{x}{2\sqrt{n}}(V_n^2 - n) \geq x\sqrt{n}\right).$$

The left hand inequality of (1.6) follows from Lemma 3.1 immediately.

To prove the right hand inequality of (1.6), we use Lemmas 3.4. If  $\Psi_n(x) \geq (x\rho_n)^{1/2}$ , then by (3.12),

$$P(S_n \geq xV_n) \leq \Psi_n(x) \exp(A\mathcal{L}_{n,x})\left(1 + A_1x\rho_n\right) \leq \Psi_n(x) \exp\left(A_2\mathcal{L}_{n,x}\right). \quad (3.16)$$

Recall we may assume that  $2 \leq x \leq \rho_n^{-1}/16$ . It is readily seen that

$$\{1 - \Phi(x)\}^{-3} \exp\left\{\frac{x^3 EX^3}{\sqrt{n}}\right\} \leq (2\pi)^{3/2} x^3 e^{2x^2}.$$

This implies that, when  $\Psi_n(x) \leq (x\rho_n)^{1/2}$ ,

$$e^{-3x^2} \leq A\{1 - \Phi(x)\}^3 \exp\left\{-\frac{x^3 EX^3}{\sqrt{n}}\right\} \leq Ax\rho_n \Psi_n(x).$$

Therefore, by (3.7),

$$P(S_n \geq xV_n) \leq \Psi_n(x) \exp(A\mathcal{L}_{n,x})\left(1 + A_1x\rho_n\right) \leq \Psi_n(x) \exp(A\mathcal{L}_{n,x}). \quad (3.17)$$

Collecting the estimates (3.16) and (3.17), we get the right hand inequality of (1.6). The proof of Theorem 1.2 is now complete.

## 4 Proof of Theorem 2.1

The proof of Theorem 2.1 is based on the conjugate method. To employ the method, let  $\xi_1, \dots, \xi_n$  be independent random variables with  $\xi_j$  having distribution function  $V_j(u)$  defined by

$$V_j(u) = E\left\{e^{h\eta_j} I(\eta_j \leq u)\right\} / Ee^{h\eta_j}, \quad \text{for } j = 1, \dots, n,$$

Also define  $M_n^2(h) = \sum_{j=1}^n \text{Var}(\xi_j)$ ,

$$G_n(t) = P\left\{\frac{\sum_{j=1}^n (\xi_j - E\xi_j)}{M_n(h)} \leq t\right\} \quad \text{and} \quad R_n(h) = \frac{(x + bx^{-1})B_n - \sum_{j=1}^n E\xi_j}{M_n(h)}.$$

Since  $EX^2 = 1$ , we have  $E|X|^3 \geq 1$  and there exists a positive constant  $c_0$  such that  $EX^2 I_{(|X| \geq \tau)} \leq 1/(4C_1)$  for  $2 \leq x \leq c_0 \rho_n^{-1}$ . Let  $c_1 = 2^8 \max\{1, C_1, \dots, C_5\}$  and  $c_2 = \min\{c_0, c_1^{-1}\}$ . Taking account of the conditions (2.1)-(2.5), it can be shown by a elementary method that, for  $2 \leq x \leq c_2 \rho_n^{-1}$ ,

$$\begin{aligned} Ee^{h\eta_j} &= 1 + \frac{x^2}{2n} - \frac{EX^3}{3} \left(\frac{x}{\sqrt{n}}\right)^3 + O_1 n^{-1} \Delta_{n,x} \\ &= \exp \left\{ \frac{x^2}{2n} - \frac{EX^3}{3} \left(\frac{x}{\sqrt{n}}\right)^3 + O_2 n^{-1} \Delta_{n,x} \right\}, \end{aligned} \quad (4.1)$$

$$E\xi_j = E\eta_j e^{h\eta_j} / Ee^{h\eta_j} = \frac{x}{\sqrt{n}} + O_3 x^2 n^{-1/2} \rho_n, \quad (4.2)$$

$$\text{Var}(\xi_j) = E\eta_j^2 e^{h\eta_j} / Ee^{h\eta_j} - (E\xi_j)^2 = 1 + O_4 x \rho_n, \quad (4.3)$$

$$E|\xi_j|^3 \leq E|\eta_j|^3 e^{h\eta_j} / Ee^{h\eta_j} \leq 2C_5 n^{1/2} \rho_n, \quad (4.4)$$

where  $|O_j| \leq 1/(2c_2)$ , for  $j = 1, \dots, 4$ .

Using (4.3)-(4.4) and (2.1), we get, for  $2 \leq x \leq c_2 \rho_n^{-1}$ ,

$$n/2 \leq M_n^2(h) = n + O_4 n x \rho_n \leq 3n/2, \quad (4.5)$$

$$\begin{aligned} |R_n(h)| &= \left| \frac{(x + bx^{-1})B_n - \sum_{j=1}^n E\xi_j}{M_n(h)} \right| \\ &\leq (3/2)|b|x^{-1} + 2(C_1 + |O_3|) x^2 \rho_n, \end{aligned} \quad (4.6)$$

$$\begin{aligned} |hM_n(h) - x| &\leq \frac{1}{x} |h^2 M_n^2(h) - x^2| \\ &\leq x \left\{ \left| \frac{M_n^2(h)}{n} - 1 \right| + \left| \frac{1}{B_n^2} - \frac{1}{n} \right| M_n^2(h) \right\} \\ &\leq (4C_1 + |O_4|) x^2 \rho_n. \end{aligned} \quad (4.7)$$

Let  $A_n(h) = R_n(h) + hM_n(h)$ ,  $c_3 = \frac{1}{4}(6C_1 + 2|Q_3| + |O_3|)^{-1}$  and  $c_4 = \min\{c_2, c_3\}$ . It follows from (4.6)-(4.7) that, for  $2 \leq x \leq c_4 \rho_n^{-1}$ ,

$$\begin{aligned} |A_n(h) - x| &\leq |R_n(h)| + |hM_n(h) - x| \\ &\leq (3/2)|b|x^{-1} + (6C_1 + 2|O_3| + |O_4|) x^2 \rho_n \end{aligned} \quad (4.8)$$

and (recall  $|b| \leq x^2/4$ ),

$$x/2 \leq A_n(h) \leq 3x/2. \quad (4.9)$$

After these preliminaries, we next give the proof of Theorem 2.1.

Write  $\delta_n(x) = (x + bx^{-1})B_n$ . By the conjugate method,

$$\begin{aligned}
 P\left(\sum_{j=1}^n \eta_j > \delta_n(x)\right) &= \left(\prod_{j=1}^n Ee^{h\eta_j}\right) \int_{\delta_n(x)}^{\infty} e^{-hu} dP\left(\sum_{j=1}^n \xi_j \leq u\right), \\
 &= \left(\prod_{j=1}^n Ee^{h\eta_j}\right) \int_0^{\infty} e^{-h\delta_n(x) - hM_n(h)v} dG_n\{v + R_n(h)\}, \\
 &= \left(\prod_{j=1}^n Ee^{h\eta_j}\right) e^{-x^2 - b} \left( \int_0^{\infty} e^{-hM_n(h)v} d\left[G_n\{v + R_n(h)\} - \Phi\{v + R_n(h)\}\right] \right. \\
 &\quad \left. + \int_0^{\infty} e^{-hM_n(h)v} d\Phi\{v + R_n(h)\} \right) \\
 &:= I_0(h) e^{-x^2 - b} (I_1(h) + I_2(h)). \tag{4.10}
 \end{aligned}$$

It follows from (4.1) that, for  $2 \leq x \leq c_4 \rho_n^{-1}$ ,

$$\begin{aligned}
 \exp\left\{\frac{x^2}{2} - \frac{x^3}{3\sqrt{n}} EX^3 - A\Delta_{n,x}\right\} &\leq I_0(h) = \left(Ee^{h\eta_j}\right)^n \\
 &\leq \exp\left\{\frac{x^2}{2} - \frac{x^3}{3\sqrt{n}} EX^3 + A\Delta_{n,x}\right\}. \tag{4.11}
 \end{aligned}$$

Next we estimate  $I_2(h)$ . We have

$$\begin{aligned}
 I_2(h) &= \frac{1}{\sqrt{2\pi}} \int_0^{\infty} e^{-hM_n(h)v - \frac{1}{2}(v + R_n(h))^2} dv \\
 &= \frac{e^{-R_n^2(h)/2}}{\sqrt{2\pi}} \int_0^{\infty} e^{-(hM_n(h) + R_n(h))v - \frac{1}{2}v^2} dv \\
 &:= \frac{e^{-R_n^2(h)/2}}{\sqrt{2\pi}} I_3(h). \tag{4.12}
 \end{aligned}$$

Write  $\psi(x) = \{1 - \Phi(x)\}/\Phi'(x) = e^{x^2/2} \int_x^{\infty} e^{-y^2/2} dy$ . Clearly,  $\psi\{A_n(h)\} = I_3(h)$ , and for  $x \geq 2$ ,

$$\frac{1}{2x} \leq \psi(x) \leq \frac{1}{x} \quad \text{and} \quad |\psi'(x)| = |x\psi(x) - 1| \leq x^{-2}.$$

These estimates, together with (4.8) and (4.9), imply that for  $2 \leq x \leq c_4 \rho_n^{-1}$ ,

$$\begin{aligned}
 I_3(h) &= \psi(x) + \psi'(\theta) \{A_n(h) - x\}, \quad [\text{where } \theta \in (x/2, 3x/2)] \\
 &= \psi(x) [O_5 + O_6 x \rho_n],
 \end{aligned}$$

where  $|O_5 - 1| \leq 2|b|x^{-2}$  and  $|O_6| \leq A$ . Therefore, for  $2 \leq x \leq c_4 \rho_n^{-1}$ ,

$$I_2(h) = e^{x^2/2} \{1 - \Phi(x)\} e^{-R_n^2(x)/2} (O_5 + O_6 x \rho_n). \quad (4.13)$$

As for  $I_1(h)$ , by (4.4)-(4.5), integration by parts and Berry-Esseen theorem, we get

$$|I_1(h)| \leq 2 \sup_v |G_n(v) - \Phi(v)| \leq 4M_n^{-3}(h) \sum_{j=1}^n E|\xi_j|^3 \leq 16 C_5 \rho_n.$$

This implies that for  $x \geq 2$ ,

$$I_1(h) = O_7 x \rho_n e^{x^2/2} \{1 - \Phi(x)\}, \quad (4.14)$$

where  $|O_7| \leq A$ .

It follows easily from (4.10)-(4.11) and (4.13)-(4.14) that for  $2 \leq x \leq c_4 \rho_n^{-1}$ ,

$$\begin{aligned} P\left(\sum_{j=1}^n \eta_j \geq \delta_n(x)\right) &\leq (1 + 2|b|x^{-2})e^{-b}\Psi_n(x) \exp\left(A \Delta_{n,x}\right) (1 + A_1 x \rho_n) \\ &\leq (1 + 2|b|x^{-2})e^{-b}\Psi_n(x) \exp\left(A \mathcal{L}_{n,x}\right). \end{aligned}$$

This proves (2.6). Similarly, by letting  $b = 0$ , it follows from (4.10)-(4.11) and (4.13)-(4.14) that

$$\begin{aligned} &P\left(\sum_{j=1}^n \eta_j \geq x B_n\right) \\ &\geq \Psi_n(x) \exp\{-A\Delta_{n,x} - R_n^2(h)/2\} \left[1 - \left\{|O_6| + |O_7|e^{R_n^2(h)/2}\right\} x \rho_n\right] \\ &\geq \Psi_n(x) \exp\{-A\Delta_{n,x}\} \left[1 - A_1 x \rho_n\right] \\ &\geq \Psi_n(x) \exp\{-A_2 \mathcal{L}_{n,x}\}, \end{aligned}$$

for  $2 \leq x \leq c_4 \rho_n^{-1/2}$ , where we have used the fact that  $|R_n(h)| \leq 1$  by (4.6), and also

$$R_n^2(h) \leq 4(C_1 + |O_3|)^2 x^4 n^{-1} (E|X|^3)^2 \leq 8(C_1 + |O_3|)^2 \Delta_{n,x}$$

since  $(E|X|^3)^2 \leq 2\{E|X|^3 I_{(|x|>\tau)}\}^2 + 2EX^4 I_{(|x|\leq\tau)}$ . This proves (2.7) and hence complete the proof of Theorem 2.1.

## 5 Proof of Theorem 2.2

The idea for the proof of Theorem 2.2 is similar to Proposition 5.4 in Jing, Shao and Wang (2002), but we need some different details. Throughout this section, we use the following notations:  $g(t, x) = Ee^{it\zeta_1/\sqrt{n}}$  and

$$\Omega_n(x, t) = \left\{ (x, t) : 1 + x \leq \frac{1}{3}(1 + C_6)^{-1}/\rho_n, |t| \leq \frac{1}{6}(1 + C_7)^{-1}/\rho_n \right\}.$$

The proof of Theorem 2.2 is based on the following lemmas.

**LEMMA 5.1.** *If  $(x, t) \in \Omega_n(x, t)$ , then*

$$|g(t, x)| \leq e^{-t^2/6n}, \quad (5.1)$$

$$\left| g^n(t, x) - e^{-\frac{t^2}{2}} \right| \leq A(1+x)\rho_n(t^2 + |t|^3)e^{-t^2/6}, \quad (5.2)$$

$$\left| g^n(t, x) - e^{-\frac{t^2}{2}} \left[ 1 + n\{g(t, x) - 1\} + \frac{t^2}{2} \right] \right| \leq A(1+x)^2 \rho_n^2 (|t|^3 + |t|^8) e^{-t^2/6}. \quad (5.3)$$

*Proof.* It follows from Taylor's expansion of  $e^{ix}$  that

$$\left| g(t, x) - 1 + \frac{t^2 E\zeta_1^2}{2n} \right| \leq \frac{1}{6} |t|^3 n^{-3/2} E|\zeta_1|^3. \quad (5.4)$$

In view of (2.8) and (2.9), we have that

$$|E\zeta_1^2 - 1| \leq (1 + C_6)(1 + x)\rho_n \quad \text{and} \quad E|\zeta_1|^3 \leq (1 + C_7)E|X|^3. \quad (5.5)$$

Taking these estimates back into (5.4), we obtain for  $(x, t) \in \Omega_n(x, t)$ ,

$$\begin{aligned} |g(t, x)| &\leq 1 - \frac{t^2}{2n} + \frac{t^2}{2n} |E\zeta_1^2 - 1| + \frac{1}{6} |t|^3 n^{-3/2} E|\zeta_1|^3 \\ &\leq 1 - \frac{t^2}{2n} + \frac{t^2}{6n} + \frac{t^2}{6n} \leq e^{-t^2/6n}. \end{aligned}$$

This proves (5.1).

Using (5.1), we have that for  $(x, t) \in \Omega_n(x, t)$  and  $|t|^3 \geq (1 + C_7)^{-1}/\rho_n$ ,

$$\left| g^n(t, x) - e^{-\frac{t^2}{2}} \right| \leq 2e^{-t^2/6} \leq 2(1 + C_7)\rho_n |t|^3 e^{-t^2/6}. \quad (5.6)$$

On the other hand, by using (5.5), we get that for  $(x, t) \in \Omega_n(x, t)$  and  $|t|^3 \leq (1 + C_7)^{-1}/\rho_n$ ,

$$g(t, x) = 1 - r_1(t, x), \quad (5.7)$$

where

$$r_1(t, x) = \frac{t^2}{2n} E\zeta_1^2 + \eta \frac{|t|^3}{6n^{3/2}} E|\zeta_1|^3, \quad |\eta| \leq 1,$$

having  $|r_1(t, x)| \leq 1/4$  and

$$|r_1(t, x)|^2 \leq \frac{t^4}{2n^2} (E|\zeta_1|^3)^{4/3} + \frac{|t|^6}{18n^3} (E|\zeta_1|^3)^2 \leq \frac{5}{9} (1 + C_7) |t|^3 n^{-1} \rho_n. \quad (5.8)$$

Therefore, it follows from  $\ln(1 + z) = z + \eta_1 z^2$ , whenever  $|z| < 1/2$ , where  $|\eta_1| \leq 1$ , that for  $(x, t) \in \Omega_n(x, t)$  and  $|t|^3 \leq (1 + C_7)^{-1}/\rho_n$ ,

$$\ln g(t, x) = -\frac{t^2}{2n} E\zeta_1^2 + \theta |t|^3 n^{-1} \rho_n, \quad |\theta| \leq 1 + C_7, \quad (5.9)$$

$$\ln g^n(t, x) = -\frac{t^2}{2} E\zeta_1^2 + \theta |t|^3 \rho_n = -\frac{t^2}{2} + r(t, x), \quad (5.10)$$

where, by using (5.5) again,

$$|r(t, x)| \leq \frac{t^2}{2} |E\zeta_1^2 - 1| + |\theta| |t|^3 \rho_n \leq t^2/3.$$

Also, we have

$$|r(t, x)| \leq \frac{t^2}{2} |E\zeta_1^2 - 1| + |\theta| |t|^3 \rho_n \leq A(t^2 + |t|^3)(1 + x)\rho_n.$$

These estimates, together with  $|e^z - 1| \leq |z|e^{|z|}$ , imply that for  $(x, t) \in \Omega_n(x, t)$  and  $|t|^3 \leq (1 + C_7)^{-1}/\rho_n$ ,

$$\left| g^n(t, x) - e^{-\frac{t^2}{2}} \right| \leq e^{-\frac{t^2}{2}} \left| e^{r(t, x)} - 1 \right| \leq A(1 + x)\rho_n(t^2 + |t|^3)e^{-t^2/6}. \quad (5.11)$$

Now, (5.2) follows from (5.6) and (5.11).

If we instead the estimate of  $r_1(t, x)$  in (5.8) by

$$|r_1(t, x)|^2 \leq \frac{t^4}{2n^2} (E\zeta_1^2)^2 + \frac{|t|^6}{18n^3} (E|\zeta_1|^3)^2 \leq 2n^{-2} t^4,$$

we can rewrite (5.9) and (5.10) as

$$\begin{aligned} \ln g(t, x) &= g(t, x) - 1 + \theta_1 n^{-2} t^4, \quad |\theta_1| \leq 2, \\ \ln g^n(t, x) &= n\{g(t, x) - 1\} + \theta n^{-1} t^4. \end{aligned}$$

Therefore, by using (5.5) and (5.7), we obtain for  $(x, t) \in \Omega_n(x, t)$ ,

$$\begin{aligned} \left| \ln g^n(t, x) + \frac{t^2}{2} \right| &\leq \left| n\{g(t, x) - 1\} + \frac{t^2}{2} \right| + 2n^{-1}t^4 \\ &\leq \frac{t^2}{2}|E\zeta_1^2 - 1| + \frac{|t|^3}{6\sqrt{n}}E|\zeta_1|^3 + 2n^{-1}t^4 \leq t^2/3 \end{aligned}$$

(recalling  $|t| \leq \frac{1}{6}\rho_n^{-1} \leq \sqrt{n}/6$ ). Also, we have

$$\begin{aligned} \left| \ln g^n(t, x) + \frac{t^2}{2} \right| &\leq \frac{t^2}{2}|E\zeta_1^2 - 1| + \frac{|t|^3}{6\sqrt{n}}E|\zeta_1|^3 + 2n^{-1}t^4 \\ &\leq A(1+x)\rho_n(t^2 + t^4). \end{aligned}$$

Now, by using  $|e^z - 1 - z| \leq \frac{z^2}{2}e^{|z|}$ , we obtain that for  $(x, t) \in \Omega_n(x, t)$ ,

$$\begin{aligned} g^n(t, x) - e^{-\frac{t^2}{2}} &= e^{-\frac{t^2}{2}} \left\{ e^{\ln g^n(t, x) + \frac{t^2}{2}} - 1 \right\} \\ &= e^{-\frac{t^2}{2}} \left[ n\{g(t, x) - 1\} + \frac{t^2}{2} + r^*(t, x) \right], \end{aligned}$$

where

$$\begin{aligned} |r^*(t, x)| &\leq 2n^{-1}t^4 + \frac{1}{2} \left| \ln g^n(t, x) + \frac{t^2}{2} \right|^2 \exp \left\{ \left| \ln g^n(t, x) + \frac{t^2}{2} \right| \right\} \\ &\leq A(1+x)^2 \rho_n^2 (|t|^4 + |t|^8) e^{t^2/3}. \end{aligned}$$

This implies (5.3). The proof of Lemma 5.1 is now complete.  $\square$

**LEMMA 5.2.** *If  $(x, t) \in \Omega_n(x, t)$ , then*

$$\left| E\Lambda_{n,n} e^{itT_n} \right| \leq Ax\rho_n^2 t^2 e^{-t^2/6}, \quad (5.12)$$

$$\left| Ee^{it(T_n + \Lambda_{n,n})} - g^n(t, x) \right| \leq Ax\rho_n^2 t^2 e^{-t^2/6} + A_1 x^{3/2} \rho_n^2 |t|^{3/2} \quad (5.13)$$

and for any  $1 \leq m \leq n$ ,

$$\left| Ee^{it(T_n + \Lambda_{n,n})} \right| \leq (1 + Ax|t|) e^{-(m-2)t^2/(12n)} + A_1 m n^{-1} x^{3/2} \rho_n^2 |t|^{3/2}. \quad (5.14)$$

*Proof.* It follows from (2.11), (2.12), (5.5) and Holder's inequality that

$$\begin{aligned} \left| E\psi_{1,2} e^{it(\zeta_1 + \zeta_2)/\sqrt{n}} \right| &= \left| E\psi_{1,2} \left( e^{it\zeta_1/\sqrt{n}} - 1 \right) \left( e^{it\zeta_2/\sqrt{n}} - 1 \right) \right| \\ &\leq \left( \frac{|t|}{\sqrt{n}} \right)^2 E \{ |\psi_{1,2}| |\zeta_1| |\zeta_2| \} \\ &\leq \frac{t^2}{n} \{ E|\psi_{1,2}|^{3/2} \}^{2/3} \{ E|\zeta_1|^3 \}^{2/3} \leq Ax\rho_n^2 t^2. \end{aligned}$$

Therefore, it follows from independence of  $\zeta_j$  and Lemma 5.1 that

$$\left| E\Lambda_{n,n}e^{itT_n} \right| \leq \left| E\psi_{1,2}e^{it(\zeta_1+\zeta_2)/\sqrt{n}} \right| |g(t, x)|^{n-2} \leq Ax\rho_n^2 t^2 e^{-t^2/6}.$$

This proves (5.12).

To prove (5.13) and (5.14), put

$$\Lambda_{n,m}^* = \Lambda_{n,n} - \Lambda_{n,m} = \frac{1}{n^2} \sum_{k=m+1}^{n-1} \sum_{j=k+1}^n \psi_{k,j} \quad \text{and} \quad \Lambda_{n,m}^* = 0, \quad m \geq n.$$

By (2.12) and  $|e^{iz} - 1 - iz| \leq 2|z|^{3/2}$ ,

$$\begin{aligned} & \left| Ee^{it(T_n+\Lambda_{n,n})} - Ee^{it(T_n+\Lambda_{n,m}^*)} - itE\Lambda_{n,m}e^{it(T_n+\Lambda_{n,m}^*)} \right| \\ & \leq 2|t|^{3/2}E|\Lambda_{n,m}|^{3/2} \leq A|t|^{3/2}mn^{-2}E|\psi_{1,2}|^{3/2} \leq Amn^{-1}x^{3/2}\rho_n^2|t|^{3/2}. \end{aligned} \quad (5.15)$$

Therefore, (5.13) follows easily from (5.12) and (5.15) with  $m = n$ . In view of independence of  $\zeta_j$ , on the other hand, (5.15) implies that for any  $1 \leq m \leq n$ ,

$$\begin{aligned} \left| Ee^{it(T_n+\Lambda_{n,n})} \right| & \leq |g^m(t, x)| + Ax|t| \left| g^{m-2}(t, x) \right| + Amn^{-1}x^{3/2}\rho_n^2|t|^{3/2} \\ & \leq (1 + Ax|t|)e^{-(m-2)t^2/(6n)} + Amn^{-1}x^{3/2}\rho_n^2|t|^{3/2} \end{aligned}$$

where we have used the estimate (recalling (2.12)):

$$\begin{aligned} E|\Lambda_{n,m}e^{it(T_n+\Lambda_{n,m}^*)}| & = E\left| \frac{1}{n^2} \sum_{k=1}^m \sum_{j=k+1}^n \psi_{k,j} e^{it(T_n+\Lambda_{n,m}^*)} \right| \\ & \leq |g(t, x)|^{m-2} E|\psi_{1,2}| \leq Ax|g(t, x)|^{m-2}. \end{aligned}$$

This gives (5.14). The proof of Lemma 5.2 is now complete.  $\square$

**LEMMA 5.3.** *Let  $F$  be a distribution function with the characteristic function  $f$ . Then for all  $y \in R$  and  $T > 0$  it holds that*

$$\lim_{z \downarrow y} F(z) \leq \frac{1}{2} + V.P. \int_{-T}^T \exp(-iyt) \frac{1}{T} K\left(\frac{t}{T}\right) f(t) dt. \quad (5.16)$$

$$\lim_{z \uparrow y} F(z) \geq \frac{1}{2} - V.P. \int_{-T}^T \exp(-iyt) \frac{1}{T} K\left(-\frac{t}{T}\right) f(t) dt, \quad (5.17)$$

where

$$V.P. \int_{-T}^T = \lim_{h \downarrow 0} \left( \int_{-T}^{-h} + \int_h^T \right),$$

and  $2K(s) = K_1(s) + iK_2(s)/(\pi s)$ ,

$$K_1(s) = 1 - |s|, \quad K_2(s) = \pi s(1 - |s|)\cot \pi s + |s|, \quad \text{for } |s| < 1,$$

and  $K(s) \equiv 0$  for  $|s| \geq 1$ .

Proof of Lemma 5.3 can be found in Prawitz (1972).

**LEMMA 5.4.** *It holds that if  $0 \leq x \leq c\rho_n^{-1}$  with  $c \leq \frac{1}{6}(1 + C_6)^{-1}$ , then for any  $y \in R$ ,*

$$|I^+(y)|, |I^-(y)| \leq A\rho_n e^{-y^2/2} + A_1(1 + x^{3/2})\rho_n^2, \quad (5.18)$$

where  $K_1(s)$  is defined as in Lemma 5.3,

$$\begin{aligned} I^+(y) &= \frac{1}{T} \int_{-T}^T e^{-iyt} K_1\left(\frac{t}{T}\right) E e^{it(S_n + \Lambda_{n,n})} dt, \\ I^-(y) &= \frac{1}{T} \int_{-T}^T e^{-iyt} K_1\left(-\frac{t}{T}\right) E e^{it(S_n + \Lambda_{n,n})} dt, \quad T = \frac{1}{6}(1 + C_7)^{-1}/\rho_n. \end{aligned}$$

*Proof.* We only prove (5.18) for  $I^+(y)$ . Without loss of generality, we assume  $\rho_n \leq 12^{-3}$ . This assumption implies that  $1 + x \leq \frac{1}{3}(1 + C_6)^{-1}/\rho_n$  for  $0 \leq x \leq c\rho_n^{-1}$  with  $c \leq \frac{1}{6}(1 + C_6)^{-1}$ . Let  $I^+ = I^+(y)$  and  $T_1 = \rho_n^{-1/3}$ . We have that

$$I^+ = \int_{-T_1}^{T_1} \cdots + \int_{T_1 \leq |t| \leq T} \cdots := I_1 + I_2.$$

It is easy to see that  $[12nt^{-2} \log |t|] \leq n - 2$  if  $|t| \geq 12$  and  $n \geq 6$ . Hence, recalling  $|T_1| \geq 12$  and  $\sqrt{n} \geq 12/E|X|^3 \geq 12$ , by (5.14) with  $m = [12nt^{-2} \log |t|] + 2$ ,

$$|I_2| \leq \frac{1}{T} \int_{T_1 \leq |t| \leq T} |E e^{it(S_n + \Lambda_{n,n})}| dt \leq A(1 + x^{3/2})\rho_n^2. \quad (5.19)$$

Noting  $K_1(s) = 1 - |s|$ , for  $|s| < 1$ , we obtain  $|I_1| \leq |I_{11}| + |I_{12}|$ , where

$$I_{11} = \frac{1}{T} \int_{-T_1}^{T_1} e^{-iyt} E e^{it(S_n + \Lambda_{n,n})} dt, \quad I_{12} = \frac{2}{T^2} \int_0^{T_1} t |E e^{it(S_n + \Lambda_{n,n})}| dt.$$

It follows from (5.14) with  $m = [12nt^{-2} \log |t|] + 2$  again that

$$|I_{12}| \leq \frac{2}{T^2} \left\{ \int_0^{12} t dt + \int_{12}^{T_1} t |E e^{it(S_n + \Lambda_{n,n})}| dt \right\} \leq A(1 + x^{3/2})\rho_n^2. \quad (5.20)$$

On the other hand, noting that

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-iyt-t^2/2} dt = e^{-y^2/2},$$

it follows from Lemmas (5.1)-(5.2) (i.e, (5.2) and (5.13)) that

$$\begin{aligned} |I_{11}| &\leq \frac{1}{T} \left| \int_{-\infty}^{\infty} e^{-iyt-t^2/2} dt \right| + \frac{1}{T} \int_{|t| \geq T_1} e^{-t^2/2} dt + \frac{1}{T} \int_{-T_1}^{T_1} \left| Ee^{it(S_n+\Lambda_{n,n})} - e^{-t^2/2} \right| dt \\ &\leq A\rho_n e^{-y^2/2} + A_1(1+x^{3/2})\rho_n^2. \end{aligned}$$

Collecting all these estimates, we conclude the proof of Lemma 5.4.  $\square$

**LEMMA 5.5.** *The integral*

$$\begin{aligned} J^+(y) &= \frac{i}{\pi} V.P. \int_{-T}^T e^{-iyt} K_2\left(\frac{t}{T}\right) Ee^{it(S_n+\Lambda_{n,n})} \frac{dt}{t}, \\ J^-(y) &= \frac{i}{\pi} V.P. \int_{-T}^T e^{-iyt} K_2\left(-\frac{t}{T}\right) Ee^{it(S_n+\Lambda_{n,n})} \frac{dt}{t}, \quad T = \frac{1}{6}(1+C_7)^{-1}/\rho_n, \end{aligned}$$

satisfy that, for any  $y \in R$  and  $0 \leq x \leq c\rho_n^{-1}$  with  $c \leq \frac{1}{6}(1+C_6)^{-1}$ ,

$$\left| J^+(y) + 1 - 2\Phi(y) - 2\mathcal{L}_n(y) \right| \leq A(1+x^{3/2})\rho_n^{3/2}, \quad (5.21)$$

$$\left| J^-(y) + 1 - 2\Phi(y) - 2\mathcal{L}_n(y) \right| \leq A(1+x^{3/2})\rho_n^{3/2}, \quad (5.22)$$

where  $K_2(s)$  is defined as in Lemma 5.3 and

$$\mathcal{L}_n(y) = n \left\{ E\Phi\left(y - \frac{\zeta_1}{\sqrt{n}}\right) - \Phi(y) \right\} - \frac{1}{2}\Phi^{(2)}(y).$$

*Proof.* We only prove (5.21). Similar to the proof of Lemma 5.4, we assume  $\rho_n \leq 12^{-3}$ , which implies  $1+x \leq \frac{1}{3}(1+C_6)^{-1}/\rho_n$  for  $0 \leq x \leq c\rho_n^{-1}$  with  $c \leq \frac{1}{6}(1+C_6)^{-1}$ . Write  $J^+ = J^+(y)$  and  $f_n(t, x) = \left[ 1 + n\{g(t, x) - 1\} + \frac{t^2}{2} \right] e^{-t^2/2}$ . The  $J^+$  can be rewrite as

$$J^+ = J_{11} + J_{12} + J_{13} + J_2,$$

where

$$\begin{aligned} J_{11} &= \frac{i}{\pi} V.P. \int_{-T_1}^{T_1} e^{-iyt} f_n(t, x) \frac{dt}{t}, \\ J_{12} &= \frac{i}{\pi} V.P. \int_{-T_1}^{T_1} e^{-iyt} \left\{ Ee^{it(S_n+\Lambda_{n,n})} - f_n(t, x) \right\} \frac{dt}{t}, \\ J_{13} &= \frac{i}{\pi} V.P. \int_{-T_1}^{T_1} e^{-iyt} \left\{ K_2\left(\frac{t}{T}\right) - 1 \right\} Ee^{it(S_n+\Lambda_{n,n})} \frac{dt}{t}, \\ J_2 &= \frac{i}{\pi} V.P. \int_{T_1 \leq |t| \leq T} e^{-iyt} K_2\left(\frac{t}{T}\right) Ee^{it(S_n+\Lambda_{n,n})} \frac{dt}{t} \end{aligned}$$

and  $T_1 = \rho_n^{-1/3}$ . Similar to (5.19), it follows that  $|J_2| \leq A(1 + x^{3/2})\rho_n^2$ . By using (5.3) and (5.13), we have

$$\begin{aligned} |J_{12}| &\leq \int_{-T_1}^{T_1} \left| Ee^{it(S_n + \Lambda_{n,n})} - g^n(t, x) \right| \frac{dt}{|t|} + \int_{-T_1}^{T_1} \left| g^n(t, x) - f_n(t, x) \right| \frac{dt}{|t|} \\ &\leq A \left\{ (1+x)^2 \rho_n^2 + x^{3/2} \rho_n^{3/2} \right\}. \end{aligned}$$

Noting that  $|K_2(s) - 1| \leq As^2$ , for  $|s| \leq 1/2$  (cf., e.g., Lemma 2.1 in Bentkus (1994), similar to (5.20), it can be easily shown that

$$|J_{13}| \leq AT^{-2} \int_{-T_1}^{T_1} |t| \left| Ee^{it(S_n + \Lambda_{n,n})} \right| dt \leq A(1 + x^{3/2})\rho_n^2.$$

On the other hand, simple calculation shows that

$$\frac{i}{2\pi} V.P. \int_{-\infty}^{\infty} e^{-iyt} f_n(t, x) \frac{dt}{t} = -\frac{1}{2} + \Phi(y) + \mathcal{L}_n(y).$$

Therefore, it follows from all these estimates and  $x \leq c\rho_n^{-1}$  that

$$\begin{aligned} &\left| J^+ + 1 - 2\Phi(y) - 2\mathcal{L}_n(y) \right| \\ &\leq \left| J_{11} + 1 - 2\Phi(y) - 2\mathcal{L}_n(y) \right| + |J_{12}| + |J_{13}| + |J_2| \\ &\leq \int_{|t| \geq T_1} \frac{1}{|t|} \left| f_n(t, x) \right| dt + A \left\{ (1+x)^2 \rho_n^2 + x^{3/2} \rho_n^{3/2} \right\} \\ &\leq A \left\{ (1+x)^2 \rho_n^2 + x^{3/2} \rho_n^{3/2} \right\} \\ &\leq A(1 + x^{3/2}) \rho_n^{3/2}. \end{aligned}$$

This also completes the proof of Lemma 5.5. □

**LEMMA 5.6.** For any  $|y| \leq A(1+x)$ ,

$$\left| \mathcal{L}_n(y) - \frac{EX^3}{\sqrt{2\pi n}} \left( \frac{y^2}{6} - \frac{yx}{2} \right) e^{-y^2/2} \right| \leq A \left\{ \rho_n + \Delta_{n,x}/(1+x) \right\} e^{-y^2/2}, \quad (5.23)$$

where  $\mathcal{L}_n(y)$  is defined as in Lemma 5.5. For  $0 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small, and  $y_0 = x + \lambda_n(x)$ , where  $|\lambda_n(x)| \leq C \Delta_{n,x}/(1+x)$ ,

$$\left| e^{-y_0^2/2} - e^{-x^2/2} \right| \leq A \rho_n, \quad (5.24)$$

$$\left| \Phi(y_0) - \Phi(x) \right| \leq A \left\{ \rho_n + \Delta_{n,x}/(1+x) \right\} e^{-x^2/2} + A_1 \rho_n^2, \quad (5.25)$$

$$\left| \mathcal{L}_n(y_0) + \frac{EX^3}{3\sqrt{n}} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} \right| \leq A \left\{ \rho_n + \Delta_{n,x}/(1+x) \right\} e^{-x^2/2} + A_1(1+x^2)\rho_n^2. \quad (5.26)$$

*Proof.* We first note that

$$\Phi^{(2)}(y) = -\frac{y}{\sqrt{2\pi}}e^{-y^2/2}, \quad \Phi^{(3)}(y) = \frac{y^2-1}{\sqrt{2\pi}}e^{-y^2/2} \quad (5.27)$$

and

$$|\Phi^{(4)}(y)| \leq A(1+|y|^3)e^{-y^2/2}. \quad (5.28)$$

Using (5.28),  $|y| \leq A(1+x)$  and  $|\zeta_1| \leq A\sqrt{n}/(1+x)$ , it can be easily seen that for  $|\theta| \leq 1$ ,

$$\left| \Phi^{(4)}\left(y + \theta \frac{|\zeta_1|}{\sqrt{n}}\right) \right| \leq A(1+|y|^3) \exp\left\{-\frac{y^2}{2} + \frac{|y||\zeta_1|}{\sqrt{n}}\right\} \leq A(1+x)^3 e^{-y^2/2}.$$

This, together with Taylor's expansion and  $E\zeta_1^4 \leq C_9 EX^4 I_{(|X| \leq \tau)}$ , implies that

$$\begin{aligned} Q_n &:= \left| E\Phi\left(y - \frac{\zeta_1}{\sqrt{n}}\right) - \Phi(y) - \frac{E\zeta_1^2}{2n}\Phi^{(2)}(y) + \frac{E\zeta_1^3}{6n^{3/2}}\Phi^{(3)}(y) \right| \\ &\leq \frac{1}{24n^2} E\zeta_1^4 \Phi^{(4)}\left(y + \theta \frac{|\zeta_1|}{\sqrt{n}}\right) \quad (|\theta| \leq 1) \\ &\leq A EX^4 I_{(|X| \leq \tau)} (1+x)^3 n^{-2} e^{-y^2/2}. \end{aligned}$$

Therefore, taking account of (2.8)-(2.9) and (5.27), we have

$$\begin{aligned} &\left| \mathcal{L}_n(y) - \frac{EX^3}{\sqrt{2\pi n}} \left(\frac{y^2}{6} - \frac{yx}{2}\right) e^{-y^2/2} \right| \\ &\leq \left| \mathcal{L}_n(y) - \frac{xEX^3}{2\sqrt{n}}\Phi^{(2)}(y) - \frac{EX^3}{6\sqrt{n}}\Phi^{(3)}(y) \right| + \frac{\rho_n}{6\sqrt{2\pi}} e^{-y^2/2} \\ &\leq nQ_n + \frac{1}{2} |\Phi^{(2)}(y)| \left| E\zeta_1^2 - 1 - \frac{\alpha x}{\sqrt{n}} EX^3 \right| \\ &\quad + \frac{1}{6n^{1/2}} |\Phi^{(3)}(y)| \left| E\zeta_1^3 - \beta EX^3 \right| + \frac{\rho_n}{6\sqrt{2\pi}} e^{-y^2/2} \\ &\leq A\{\rho_n + \Delta_{n,x}/(1+x)\} e^{-y^2/2}. \end{aligned}$$

This concludes (5.23).

We next prove (5.24)-(5.26). It can be easily seen that  $\Delta_{n,x} \leq (1+x)^3 \rho_n^{-1}$ , and for  $x \leq c\rho_n^{-1}$  with  $c$  small enough,

$$\begin{aligned} |y_0^2 - x^2| &\leq 2x|\lambda_n(x)| + \lambda_n^2(x) \leq (2C + C^2)\Delta_{n,x} \leq x^2/3, \\ -(x - |y_0 - x|)^2 &\leq -x^2 + 2x|\lambda_n(x)| \leq -x^2 + 2C\Delta_{n,x} \leq -3x^2/4. \end{aligned}$$

These estimates imply that

$$\begin{aligned} \left| e^{-y_0^2/2} - e^{-x^2/2} \right| &\leq \frac{1}{2} |y_0^2 - x^2| e^{-x^2/2 + |y_0^2 - x^2|/2} \\ &\leq \frac{A}{2} \rho_n (1+x)^3 e^{-x^2/3} \leq A \rho_n, \end{aligned} \quad (5.29)$$

$$\begin{aligned} \left| \Phi(y_0) - \Phi(x) \right| &\leq |y_0 - x| \Phi'(x) + \frac{(y_0 - x)^2}{2} \left| \Phi^{(2)}(x + \theta|y_0 - x|) \right| \quad (|\theta| \leq 1) \\ &\leq A \frac{\Delta_{n,x}}{1+x} e^{-x^2/2} + \frac{\rho_n^2}{2} (1+x)^7 e^{-(x-|y_0-x|)^2/2} \\ &\leq A \frac{\Delta_{n,x}}{1+x} e^{-x^2/2} + A_1 \rho_n^2. \end{aligned} \quad (5.30)$$

This proves (5.24) and (5.25). On the other hand, we have

$$\left| -\frac{y_0^2}{6} + \frac{y_0 x}{2} - \frac{1}{3} x^2 \right| \leq \frac{1}{6} |y_0^2 - x^2| + \frac{x}{2} |y_0 - x| \leq A \Delta_{n,x}.$$

This, together with (5.27)-(5.24), implies that

$$\begin{aligned} \left| \mathcal{L}_n(y_0) + \frac{EX^3}{3\sqrt{n}} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} \right| &\leq A \rho_n \Delta_{n,x} e^{-y_0^2/2} + A_1 \{ \rho_n + \Delta_{n,x}/(1+x) \} e^{-y_0^2/2} \\ &\leq A \{ \rho_n + \Delta_{n,x}/(1+x) \} e^{-x^2/2} + A_1 (1+x^2) \rho_n^2. \end{aligned}$$

This gives (5.26). The proof of Lemma 5.6 is now complete.  $\square$

After these lemmas, we are now ready to prove Theorem 2.2.

By using (5.17) and Lemmas 5.4-5.5, we obtain for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small,

$$\begin{aligned} P\left(T_n + \Delta_{n,n} \geq x + \lambda_n(x)\right) &\leq \frac{1}{2} \left[ I^-(y_0) + 1 - J^-(y_0) \right] \\ &\leq 1 - \Phi(y_0) + \mathcal{L}_n(y_0) + A \rho_n e^{-y_0^2/2} + A_1 x^{3/2} \rho_n^{3/2}, \end{aligned}$$

where  $y_0 = x + \lambda_n(x)$ . Furthermore, it follows from Lemma 5.6 that

$$\begin{aligned} P\left(T_n + \Delta_{n,n} \geq x + \lambda_n(x)\right) &\leq 1 - \Phi(x) - \frac{EX^3}{3\sqrt{n}} \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} + A(\rho_n + \Delta_{n,x}/x) e^{-x^2/2} + A_1 x^{3/2} \rho_n^{3/2}. \end{aligned} \quad (5.31)$$

Using the well-known inequality

$$\frac{1}{\sqrt{2\pi}} \left( \frac{1}{x} - \frac{1}{x^3} \right) e^{-x^2/2} \leq 1 - \Phi(x) \leq \frac{1}{\sqrt{2\pi} x} e^{-x^2/2}, \quad x > 0,$$

we have for  $x \geq 2$ ,

$$\left| x^3 \{1 - \Phi(x)\} - \frac{x^2}{\sqrt{2\pi}} e^{-x^2/2} \right| \leq \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

and  $e^{-x^2/2} \leq Ax \{1 - \Phi(x)\}$ . Taking these estimates back into (5.31), we get for  $2 \leq x \leq c\rho_n^{-1}$  with  $c$  sufficiently small,

$$P\left(T_n + \Delta_{n,n} \geq x + \lambda_n(x)\right) \leq \{1 - \Phi(x)\} \left[1 - \frac{x^3 EX^3}{3\sqrt{n}} + A\mathcal{L}_{n,x}\right] + A_1 x^{3/2} \rho_n^{3/2}.$$

Now we conclude the result if we prove

$$1 - \frac{x^3 EX^3}{3\sqrt{n}} + A\mathcal{L}_{n,x} \leq \exp\left\{-\frac{x^3 EX^3}{3\sqrt{n}}\right\} \left[1 + A_1 \mathcal{L}_{n,x}\right]. \quad (5.32)$$

In fact, (5.32) is obvious for  $EX^3 < 0$ . In the case that  $EX^3 > 0$ , if  $\frac{x^3 EX^3}{3\sqrt{n}} \leq 2$ , then

$$\begin{aligned} 1 - \frac{x^3 EX^3}{3\sqrt{n}} + A\mathcal{L}_{n,x} &\leq \exp\left\{-\frac{x^3 EX^3}{3\sqrt{n}}\right\} + A\mathcal{L}_{n,x} \\ &\leq \exp\left\{-\frac{x^3 EX^3}{3\sqrt{n}}\right\} \left[1 + Ae^2 \mathcal{L}_{n,x}\right]. \end{aligned}$$

On the other hand, it follows easily from the definition of  $\Delta_{n,x}$  that for  $2 \leq x \leq c\rho_n^{-1}$ ,

$$A(x\rho_n + \Delta_{n,x}) \leq (1 + x^3 EX^3 / \sqrt{n})/9,$$

by choosing  $c$  sufficiently small. Therefore, if  $\frac{x^3 EX^3}{3\sqrt{n}} \geq 2$ , then

$$1 - \frac{x^3 EX^3}{3\sqrt{n}} + A\mathcal{L}_{n,x} \leq 0 \leq \exp\left\{-\frac{x^3 EX^3}{3\sqrt{n}}\right\} \left[1 + A\mathcal{L}_{n,x}\right].$$

Collecting all these estimates, we get the desired (5.32). This also completes the proof of Theorem 2.2.

## REFERENCES

- Bentkus, V., Bloznelis, M. and Götze, F. (1996). A Berry-Esseen bound for Student's statistic in the non-i.i.d. case. *J. Theoret. Probab.* **9**, 765-796.
- Bentkus, V. and Götze, F. (1996). The Berry-Esseen bound for Student's statistic. *Ann. Probab.* **24**, 491-503.

- Bentkus, V., Götze, F. and van Zwet, W.R. (1997). An Edgeworth expansion for symmetric statistics. *Ann. Statist.* **25**, 851-896.
- Chistyakov, G. P., Gtze, F. (2003). Moderate deviations for Student's statistic. *Theory Probab. Appl.* **47**, 415-428.
- Friedrich, K. O. (1989). A Berry-Esseen bound for functions of independent random variables, *Ann. Statist.*, **17** 170-183.
- Gine, E., Götze, F. and Mason, D.M. (1997). When is the Student  $t$ -statistic asymptotically standard normal? *Ann. Probab.* **25**, 1514-1531.
- Hall, P. (1987). Edgeworth expansion for student's  $t$ -statistic under minimal conditions. *Ann. Probab.* **15**, 920-931.
- Hall, P. (1988). On the effect of random norming on the rate of convergence in the central limit theorem. *Ann. Probab.* **16**, 1265-1280.
- Hall, P. and Jing, B.-Y. (1995). Uniform convergence bounds for confidence Intervals and Berry-Esseen Theorems for Edgeworth Expansion. *Ann. Statist.* **23**, 363-375.
- He, X. and Shao, Q.M. (2000). On parameters of increasing dimensions. *J. Multivariate Anal.* **73**, 120-135.
- Jing, B.-Y., Shao, Q.-M. and Wang, Q. (2003). Self-normalized Cramér-type large deviation for independent random variables. *Ann. Probab.* **31**, 2167-2215.
- Logan, B.F., Mallows, C.L., Rice, S.O. and Shepp, L.A. (1973). Limit distributions of self-normalized sums. *Ann. Probab.* **1**, 788-809.
- Petrov, V.V. (1995). *Limit Theorems of Probability Theory: Sequences of Independent Random Variables*. Oxford Science Publications.
- Prawitz, H. (1972). Limits for a distribution, if the characteristic function is given in a finite domain. *Skand. Aktuar. Tidskr.* 138-154.
- Putter, H. and van Zwet, W.R. (1998). Empirical Edgeworth expansions for symmetric statistics. *Ann. Stat.* **26**, 1540-1569.
- Slavova, V.V. (1985). On the Berry-Esseen bound for Student's statistic. In: *Stability Problems for Stochastic Models*, Eds. V.V. Kalashnikov and V.M. Zolotarev. *Lecture Notes in Math.* **1155**, 355-390. Springer, Berlin.
- Shao, Q. M. (1999). Cramér-type large deviation for Student's  $t$  statistic. *J. Theoret. Probab.* **12**, 387-398.

- van Zwet W. R. (1984). A Berry-Esséen Bound for symmetric statistics. *Z-wahrsch. verw. Gebiete*, **66**, 425-440.
- Wang, Q. and Jing, B.-Y. (1999). An exponential non-uniform Berry-Esseen bound for self-normalized sums. *Ann. Probab.* **27**, 2068-2088.
- Wang, Q., Jing, B.-Y. and Zhao, L. C. (2000). The Berry-Esséen bound for studentized statistics. *Ann. Probab.*, **28** 511-535.