

NOTE ON A CHARACTERIZING PROPERTY OF THE EXPONENTIAL DISTRIBUTION

BY N. KRISHNAJI

Indian Institute of Management, Calcutta

0. Summary. In this note we discuss a characteristic property of the exponential distribution which is closely related to the "lack-of-memory" characterization.

1. Introduction. We begin with a random variable (rv) X having an exponential distribution:

$$(1) \quad \Pr(X > x) = U(x) = \exp(-kx), \quad k > 0, \quad x \geq 0$$

and note that

$$(2) \quad U(y+z) - U(y)U(z) = 0 \quad y \geq 0, z \geq 0.$$

The lack-of-memory characterization is obtained by showing that the only solution of (2), where $1 - U(\cdot)$ is the cdf of some rv X , is of the form (1) (Feller [1]).

Now if Y and Z are arbitrary rv's, it follows from (2) that

$$(3) \quad \int_0^\infty \int_0^\infty [U(y+z) - U(y)U(z)] dG(y) dH(z) = 0$$

where $G(\cdot)$ and $H(\cdot)$ are the cdf's of Y and Z respectively. If we further assume that Y and Z are nonnegative, independent and independent of X , it is easy to see that (3) implies that

$$(4) \quad \Pr(X > Y+Z) = \Pr(X > Y) \Pr(X > Z).$$

In the following section we give conditions under which the validity of (4) implies that X is exponential. We can see immediately that if (4) is valid for all Y and Z in any class of rv's that includes all the nonnegative degenerates then (2) is implied and X must be exponential.

2. Characterizations. We introduce families \mathcal{F} of nonnegative rv's that have the property

(a) For each interval $I \subset [0, \infty)$ with positive Lebesgue measure $\Pr(Y \in I) > 0$ for some $Y \in \mathcal{F}$.

In what follows we write $\Pr(X > x) = U(x)$ and

$$D(y, z) = U(y+z) - U(y)U(z).$$

THEOREM. If (i) X is a nonnegative rv and $U(x)$ is continuous on $(0, \infty)$ and either (b.1) $D(y, z) \geq 0, y \geq 0, z \geq 0$ or (b.2) $D(y, z) \leq 0, y \geq 0, z \geq 0$;

(ii) \mathcal{A} and \mathcal{B} are families satisfying condition (a) such that every member of \mathcal{A} is independent of every member of \mathcal{B} and equation (4) holds whenever $Y \in \mathcal{A}$ and $Z \in \mathcal{B}$, then X is exponential.

Received October 27, 1969; revised June 1, 1970.

PROOF. Suppose (b.1) holds and $D(y_0, z_0) = \delta > 0$ for some $y_0 \geq 0, z_0 \geq 0$. Then, since D is continuous there is a square neighborhood S of (y_0, z_0) with sides of positive Lebesgue measure such that $D > \delta/2$ in S . In accordance with condition (a) we can then choose Y and Z with cdf's $G(\cdot)$ and $H(\cdot)$ respectively such that the product GXH assigns a positive measure to S ; for this choice the contribution of S to the integral in (3) is positive. But this is impossible because by hypothesis (3) holds for this pair of G and H also. The case (b.2) can also be treated in a similar way. This implies (2) and since $1 - U(x)$ is a cdf, X is exponential.

By properly choosing the families \mathcal{A} and \mathcal{B} we can easily prove the following.

COROLLARY. *If X, Y and Z are mutually independent nonnegative rv's such that X has a continuous cdf and satisfies either (b.1) or (b.2) and Y and Z assign positive measures to every interval $I \subset [0, \infty)$ of positive Lebesgue measure, then*

- (i) (4) holds iff X is exponential and
- (ii) $\Pr(X - Y > t) = \Pr(X > Y)\Pr(X > t), t \geq 0$ iff X is exponential.

Lastly we state a result which does not require the conditions (a), (b.1) and (b.2).

PROPOSITION. *If X and Y are independent and nonnegative such that $\Pr(X - Y > 0) > 0$, then*

$$(5) \quad \Pr(X - Y > x + y) = \Pr(X - Y > x)\Pr(X > y), \quad x \geq 0, y \geq 0$$

iff X is exponential.

PROOF. The "if" part is obvious from Section 1. Writing $W(x) = \Pr(X - Y > x)$ (5) implies that $W(x + y) = W(x)U(y), x \geq 0, y \geq 0$ and hence $W(s) = W(0)U(s), s \geq 0$, so that $W(0)U(x + y) = W(0)U(x)U(y)$ for $x \geq 0, y \geq 0$ which implies (2) since $W(0) > 0$.

Acknowledgment. The author is indebted to the referee for suggesting substantial improvements to an original draft of this note. Thanks are also due to an Associate Editor for useful comments.

REFERENCE

- [1] FELLER, W. (1957). *An Introduction to Probability Theory and its Applications* 1. Wiley, New York.