

REFERENCES

- [1] S. BOCHNER AND K. CHANDRASEKHARAN, "Fourier transforms," *Ann. Math. Studies*, No. 19 (1949), Princeton.
- [2] J. M. HAMMERSLEY, "The distribution of distance in a hypersphere," *Ann. Math. Stat.*, Vol. 21 (1950), pp. 447-452.
- [3] R. D. LORD, "The use of the Hankel transform in statistics. I, General theory and examples," to be published in *Biometrika*.
- [4] G. N. WATSON, *A treatise on Bessel functions*, Cambridge University Press, 1st edn., 1922, 2nd edn., 1944.

◆

EXTREME VALUES IN SAMPLES FROM m -DEPENDENT STATIONARY STOCHASTIC PROCESSES

BY G. S. WATSON

University of Melbourne, Australia

Summary. The limiting distributions for the order statistics of n successive observations in a sequence of independent and identically distributed random variables are shown to hold also when the sequence is generated by a stationary stochastic process of a certain moving average type.

A sequence of random variables $\{x_i\}$ has been called m -dependent [3] if $|i - j| > m$ implies that x_i and x_j are independent. If the variables in a strictly stationary sequence are m -dependent and have a finite upper bound to their range of variation, the largest in a sample of n successive members tends with probability one to this upper bound. This is a simple extension of Dodd's results [1] for the case of independence.

The following theorem shows that when this upper bound is infinite, the asymptotic distribution of the largest in such a sample is the same as in the case of independence.

THEOREM. *Let $\{x_i\}$ be a sequence of random variables, unbounded above and generated by an m -dependent strictly stationary stochastic process with the property that*

$$(1) \quad \lim_{c \rightarrow \infty} \frac{1}{P(x_i > c)} \max_{|i-j| \leq m} P[(x_i > c), (x_j > c)] = 0.$$

Then, if $\xi = n P[x_i > c_n(\xi)]$, for ξ fixed,

$$\lim_{n \rightarrow \infty} P[x_i \leq c_n(\xi); i = 1, \dots, n] = e^{-\xi}$$

PROOF. Using the formula for the probabilities of the joint occurrence of a set of events in terms of probabilities of occurrence of their contraries (Feller [2],

Received 8/19/52, revised 7/2/54.

p. 61), we have, for any even integer $l \leq n$ and for $i = 1, \dots, n$, that $P[x_i \leq c_n(\xi)]$ is bounded below and above, respectively, by

$$1 - \sum P(x_i > c) + \dots + (-1)^{l-1} \sum P[(x_{i_1} > c), \dots, (x_{i_{l-1}} > c)],$$

$$1 - \sum P(x_i > c) + \dots + (-1)^l \sum P[(x_{i_1} > c), \dots, (x_{i_l} > c)],$$

where, for brevity, $c = c_n(\xi)$. Clearly, $\sum P(x_i > c) = nP(x_i > c) = \xi$.

Now

$$\sum P[(x_i > c), (x_j > c)]$$

$$= \sum_{i=1}^m (n - i)P[(x_1 > c), (x_{i+1} > c)] + P(x_i > c)^2 \frac{1}{2}(n - m - 1)(n - m).$$

But

$$\sum_{i=1}^m (n - i)P[(x_1 > c), (x_{i+1} > c)]$$

$$\leq mn \left[1 - \frac{(m + 1)}{2n} \right] \max_{|i-j| \leq m} P[(x_i > c), (x_j > c)]$$

$$= m\xi \left[1 - \frac{m + 1}{2n} \right] \max_{|i-j| \leq m} \frac{P[(x_i > c), (x_j > c)]}{P(x_i > c)}.$$

Since, as $n \rightarrow \infty$, with ξ fixed, $c = c_n(\xi) \rightarrow \infty$, condition (1) shows that the last expression tends to zero. Hence

$$\lim_{n \rightarrow \infty} \sum P[(x_i > c), (x_j > c)] = \frac{1}{2}\xi^2.$$

The general sum $\sum P[(x_{i_1} > c), \dots, (x_{i_q} > c)]$ contains $\binom{n}{q}$ terms. Of these, there are order n terms in which none of the x_i appearing ever differs in its subscript by more than m from its nearest neighbours, order n^2 terms in which only one x_i differs in its subscript by more than m from its nearest neighbours, and so on, provided that n is large enough for all the cases to occur. There are $\sim n^q/q!$ terms in which each x_i is separated in its subscript by more than m from its neighbours. These terms may be said to belong to the first, second, \dots , q th class. The sum of terms of the first class will be less than a constant times $n \max_{|i-j| \leq m} P[(x_i > c), (x_j > c)]$, the sum of terms of the second class will be less than a constant times $n^2 P(x_i > c) \max_{|i-j| \leq m} P[(x_i > c), (x_j > c)]$ and so on until we reach the q th class, where the sum is $[n^q/q! + O(n^{q-1})] P(x_i > c)^q$. Thus, by (1), the only terms contributing to the sum as $n \rightarrow \infty$ are those of the q th class; these yield $\xi^q/q!$ asymptotically.

Thus for any even integer l we have shown that

$$\sum_{q=0}^{l-1} \frac{(-\xi)^q}{q!} \leq \lim_{n \rightarrow \infty} P(x_i \leq c_n(\xi)) \leq \sum_{q=0}^l \frac{(-\xi)^q}{q!}, \quad i = 1, \dots, n$$

which proves the theorem.

To show that this theorem covers a class of stochastic processes of practical interest, it is shown next that the condition (1) of the theorem is true in strictly stationary processes which are normal. For this, it suffices to show that

$$(2) \quad \frac{P[(x > c), (y > c)]}{P(x > c)} \rightarrow 0, \quad (c \rightarrow \infty),$$

where x and y have a bivariate normal distribution with means zero, variances unity and covariance ρ , with $|\rho| < 1$. Now

$$P[(x > c), (y > c)] = \frac{1}{2\pi\sqrt{1-\rho^2}} \int_c^\infty \int_c^\infty \exp\left[\frac{-1}{2(1-\rho^2)}(x^2 - 2\rho xy + y^2)\right] dx dy.$$

The substitution $x = r/c + c$ and $y = t/c + c$ leads to

$$P[(x > c), (y > c)] = \frac{\exp[-c^2/(1+\rho)]}{2\pi c^2\sqrt{1-\rho^2}} \int_0^\infty \int_0^\infty \exp\left[-\frac{r^2 - 2prt + t^2}{2c^2(1-\rho^2)}\right] \exp\left[-\frac{r+t}{1+\rho}\right] dr dt \\ \sim \frac{1}{2\pi} \exp\left(\frac{-c^2}{1+\rho}\right) \left[\frac{(1+\rho)^{3/2}}{\sqrt{1-\rho}} \frac{1}{c^2} - 0\left(\frac{1}{c^4}\right)\right], \quad c \text{ large.}$$

Since $P(x > c) \sim (1/\sqrt{2\pi}) \exp(-\frac{1}{2}c^2)$, statement (2) follows.

Acknowledgement. The author wishes to record his gratitude to a referee for suggesting a change in the assumptions made in an earlier form in this paper.

REFERENCES

- [1] E. L. DODD, "The greatest and the least variate under general laws of error," *Trans. Amer. Math. Soc.*, Vol. 25 (1923), p. 525-539.
- [2] W. FELLER, *An introduction to probability theory and its applications*, John Wiley and Sons, Inc., New York., 1950.
- [3] W. HOEFFDING AND H. ROBBINS, "The central limit theorem for dependent random variables," *Duke Math. J.*, Vol. 15 (1948), pp. 773-780.

EXPRESSION OF THE k -STATISTICS k_9 AND k_{10} IN TERMS OF POWER SUMS AND SAMPLE MOMENTS

BY M. ZIA UD-DIN

Panjab University, Lahore, Pakistan

The k statistics are of interest to workers in the theory of sampling distributions and moment statistics. They are related also to certain aspects of the theory of numbers and combinatory analysis, as indicated by Dressel [1].

The k statistics were introduced by Fisher in 1928 [2] to estimate the cumulants