

INVARIANCE PRINCIPLES FOR THE LAW OF THE ITERATED LOGARITHM FOR MARTINGALES AND PROCESSES WITH STATIONARY INCREMENTS

BY C. C. HEYDE AND D. J. SCOTT¹

Australian National University

The main result in this paper is an invariance principle for the law of the iterated logarithm for square integrable martingales subject to fairly mild regularity conditions on the increments. When specialized to the case of identically distributed increments the result contains that of Stout [16] as well as the invariance principle for independent random variables of Strassen [17]. The martingale result is also used to obtain an invariance principle for the iterated logarithm law for a wide class of stationary ergodic sequences and a corollary is given which extends recent results of Oodaira and Yoshihara [10] on ϕ -mixing processes.

1. Introduction. This paper is basically divided into two parts. In the first part a general invariance principle for the law of the iterated logarithm for martingales is obtained. This result (Theorem 1) specializes, in the case of identically distributed increments, to a result (Corollary 2) which contains the invariance principle of Strassen [17] for sums of independent and identically distributed random variables. Corollary 2 also contains Stout's [16] result giving a classical Hartman-Wintner form of iterated logarithm law for stationary ergodic martingales and its minor generalization by Heyde [6]. Our approach to Theorem 1 follows essentially the pattern of that of Theorem 3 of [17] in that it is based on the use of a form of the Skorokhod representation theorem.

In the second part of the paper we make use of Corollary 2 to establish an invariance principle for the law of the iterated logarithm for a certain class of (strictly) stationary processes (Theorem 2). The basic idea is a representation for the increments of the stationary process in terms of the increments of a stationary martingale plus other terms whose sum is negligible under suitable norming. This idea is due to Gordin [4], who has used it to prove a central limit theorem for stationary ergodic sequences and it has been further developed, in that context, by Scott [14]. As a corollary to Theorem 2 we obtain Corollary 3 which extends results of Oodaira and Yoshihara [10], who have given an invariance principle for the law of the iterated logarithm for stationary ϕ -mixing sequences.

Previous work on iterated logarithm results for stationary processes has proceeded via classical methods. Our approach via the Skorokhod representation is

Received March 27, 1972; revised October 6, 1972.

¹ Now at La Trobe University.

AMS 1970 subject classifications. Primary: 60B10, 60F15, Secondary: 60G10, 60G45.

Key words and phrases. Invariance principles, iterated logarithm law, martingales, stationary ergodic processes, ϕ -mixing.

more direct, gives ready access to the invariance principle, and avoids the necessity of obtaining detailed estimates of probabilities via, for example, rates of convergence to normality. In principle the result of Theorem 2 could be obtained by classical methods by paralleling the work of Chover [2] but a prerequisite would be estimates of rates of convergence to normality along the lines of those of Heyde [5] for sums of independent random variables.

2. Martingale results. Let $\{S_n, \mathcal{F}_n; n \geq 0\}$ be a martingale on the probability space (Ω, \mathcal{A}, p) where $\mathcal{F}_0 = \{\phi, \Omega\}$ and $\mathcal{F}_n = \sigma$ -field generated by S_1, S_2, \dots, S_n for $n > 0$. Let $S_0 = X_0 = 0$ almost surely (a.s.) and $S_n = \sum_{k=1}^n X_k$ for $n \geq 1$. Further, let $s_n^2 = ES_n^2 < \infty$.

We consider the metric space (C, ρ) of all real-valued continuous functions on $[0, 1]$ with

$$\rho(x, y) = \sup_{0 \leq t \leq 1} |x(t) - y(t)| \quad \text{for } x, y \in C.$$

Let K be the set of absolutely continuous $x \in C$ such that $x(0) = 0$ and

$$\int_0^1 [\dot{x}(t)]^2 dt \leq 1$$

where \dot{x} denotes the derivative of x determined almost everywhere with respect to Lebesgue measure. Define a real function $g(\cdot)$ on $[0, \infty)$ by

$$g(t) = \sup \{n : s_n^2 \leq t\}.$$

Let ξ be a standard Wiener process (Brownian motion) on $[0, \infty)$ and define a sequence of real random functions $\xi_n(\cdot)$ on $[0, 1]$, for $n > g(e)$ by

$$\xi_n(t) = [\phi(s_n^2)]^{-1} \xi(s_n^2 t) \quad t \in [0, 1]$$

where $\phi(\cdot)$ is a real function on $[e, \infty)$ given by

$$\phi(t) = (2t \log \log t)^{\frac{1}{2}} \quad t \in [e, \infty).$$

We also define a sequence of real random functions $\eta_n(\cdot)$ on $[0, 1]$, for $n > g(e)$, by

$$\eta_n(t) = [\phi(s_n^2)]^{-1} [S_k + (s_n^2 t - s_k^2)(s_{k+1}^2 - s_k^2)^{-1} X_{k+1}]$$

$$s_k^2 \leq s_n^2 t \leq s_{k+1}^2, \quad k = 0, 1, \dots, n - 1.$$

THEOREM 1. *If $s_n^2 \rightarrow \infty$ and*

- (1) $\sum_{n=1}^{\infty} s_n^{-4} E\{X_n^4 I(|X_n| < \delta s_n)\} < \infty$ for some $\delta > 0$,
- (2) $\sum_{n=1}^{\infty} s_n^{-1} E\{|X_n| I(|X_n| \geq \epsilon s_n)\} < \infty$ for all $\epsilon > 0$,
- (3) $s_n^{-2} \sum_{k=1}^n X_k^2 \rightarrow_{a.s.} 1$ as $n \rightarrow \infty$

hold, $I(\cdot)$ denoting the indicator function, then $\{\eta_n; n > g(e)\}$ is relatively compact in C and the set of its limit points coincides with K . (“ $\rightarrow_{a.s.}$ ” means “converges a.s. to.”)

COROLLARY 1. *If $s_n^2 \rightarrow \infty$ as $n \rightarrow \infty$,*

$$(4) \quad \sum_{n=1}^{\infty} s_n^{-4} E X_n^4 < \infty$$

and (3) holds then $\{\eta_n; n > g(e)\}$ is relatively compact in C and the set of its limit points coincides with K .

COROLLARY 2. *If the X_n are identically distributed and satisfy (3), then $\{\eta_n; n > e[E(X_1^2)]^{-1}\}$ is relatively compact in C and the set of its limit points coincides with K .*

The condition (3) is, unfortunately, often difficult to check. It holds, for example, if $\{X_n\}$ is a stationary ergodic sequence.

The result of Theorem 1 appears to be new even in the case of independent random variables. Other iterated logarithm type results for martingales have been obtained by Stout [15] and Strassen [18] but they involve random norming and do not seem to be directly comparable with the present results.

3. Martingale proofs. The first step is to define a truncated martingale sequence $\{S_n^*, \mathcal{F}_n^*; n \geq 0\}$. For $\delta > 0$, let $X_0^* = 0$ a.s., $\mathcal{F}_0^* = \{\phi, \Omega\}$ and for $k > 0$,

$$X_k^* = X_k I(|X_k| < \delta s_k) - E\{X_k I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\},$$

where $\mathcal{F}_k^* = \sigma$ -field generated by X_1^*, \dots, X_k^* . Then we let

$$S_n^* = \sum_{k=0}^n X_k^* \quad \text{for } n \geq 0,$$

and $s_n^{*2} = ES_n^{*2} \leq s_n^2 < \infty$. For $n > g(e)$ we define a sequence of real random functions $\eta_n^*(\cdot)$ on $[0, 1]$ by

$$\begin{aligned} \eta_n^*(t) &= [\phi(s_n^2)]^{-1} [S_k^* + (s_n^2 t - s_k^2)(s_{k+1}^2 - s_k^2)^{-1} X_{k+1}^*] \\ &\quad s_k^2 \leq t s_n^2 \leq s_{k+1}^2, \quad k = 0, 1, \dots, n - 1. \end{aligned}$$

We first observe that

$$(5) \quad \sup_{0 \leq t \leq 1} |\eta_n^*(t) - \eta_n(t)| \xrightarrow{\text{a.s.}} 0 \quad \text{as } n \rightarrow \infty$$

under the conditions of Theorem 1. This will of course be true if

$$(6) \quad s_n^{-1} \sum_{k=1}^n |X_k| I(|X_k| \geq \delta s_k) \xrightarrow{\text{a.s.}} 0 \quad \text{as } n \rightarrow \infty$$

and

$$(7) \quad s_n^{-1} \sum_{k=1}^n |E\{X_k I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\}| \xrightarrow{\text{a.s.}} 0 \quad \text{as } n \rightarrow \infty,$$

and these results follow straightforwardly from (2) via the Kronecker lemma and the monotone convergence theorem.

Now we introduce the Skorokhod representation. From Theorem 4.3 of Strassen [18] we may conclude that there exists a sequence $T_0 = 0, T_1, T_2, \dots$ of nonnegative random variables such that

$$(\bar{S}_1, \bar{S}_2, \dots, \bar{S}_n) \equiv (\xi(T_1), \xi(T_1 + T_2), \dots, \xi(\sum_{k=1}^n T_k))$$

is distributed as

$$(8) \quad (S_1^*, S_2^*, \dots, S_n^*)$$

for each $n \geq 1$. Further, if $\bar{S}_0 = 0$ a.s., $\bar{\mathcal{F}}_0 = \{\phi, \Omega\}$, $\bar{\mathcal{F}}_n$ for $n \geq 1$ is the σ -field generated by $\bar{S}_1, \dots, \bar{S}_n$, $\mathcal{G}_0 = \mathcal{F}_0$, \mathcal{G}_n is the σ -field generated by $\xi(t)$ for

$0 \leq t \leq \sum_{k=1}^n T_k$ for $n \geq 1$ and $\bar{S}_n = \sum_{k=0}^n \bar{X}_k$ for $n \geq 0$, then

$$(9) \quad E\{T_n^r | \mathcal{G}_{n-1}\} = E\{\bar{X}_n^{2r} | \bar{\mathcal{F}}_{n-1}\} \quad \text{a.s.} \quad \text{for } n \geq 1.$$

Also, for $r > 1$,

$$(10) \quad E\{T_n^r | \mathcal{G}_{n-1}\} \leq L_r E\{\bar{X}_n^{2r} | \bar{\mathcal{F}}_{n-1}\} \quad \text{a.s.} \quad \text{for } n \geq 1,$$

where L_r is a constant depending only on r .

Define $\tilde{\eta}_n(\cdot)$ on $[0, 1]$, for $n > g(e)$, by

$$\tilde{\eta}_n(t) = [\phi(s_n^2)]^{-1} [\bar{S}_k + (s_n^2 t - s_k^2)(s_{k+1}^2 - s_k^2)^{-1} \bar{X}_{k+1}]$$

$$s_k^2 \leq s_n^2 t \leq s_{k+1}^2, k = 0, 1, \dots, n - 1,$$

and $\eta(\cdot)$ on $[0, \infty)$ by

$$\eta(t) = \bar{S}_n + \bar{X}_{n+1}(t - s_n^2)(s_{n+1}^2 - s_n^2)^{-1} \quad s_n^2 \leq t \leq s_{n+1}^2.$$

Then

$$\eta(t) = \phi(s_n^2) \tilde{\eta}_n(s_n^{-2} t) \quad \text{for } t \in [0, s_n^2]$$

for any $n > g(e)$.

Our proof follows the work of Strassen in [17]. In fact, in view of (5) and (8) above and the results of [17], Theorem 1 will follow provided we can show that $s_{n+1}^2/s_n^2 \rightarrow 1$ and $\rho(\tilde{\eta}_n, \xi_n) \rightarrow_{\text{a.s.}} 0$ and this last condition is implied by

$$(11) \quad P(\lim_{t \rightarrow \infty} [\phi(t)]^{-1} \sup_{\tau \leq t} |\eta(\tau) - \xi(\tau)| = 0) = 1.$$

In order to prove that $s_{n+1}^2/s_n^2 \rightarrow 1$, we first note that (2) gives, via an application of the Kronecker Lemma,

$$s_n^{-1} \sup_{k \leq n} |X_k| I(|X_k| \geq \epsilon s_k) \rightarrow_{\text{a.s.}} 0, \quad \text{for any } \epsilon > 0,$$

so that upon squaring we have *a fortiori*

$$s_n^{-2} \sup_{k \leq n} X_k^2 I(|X_k| \geq \epsilon s_n) \rightarrow_{\text{a.s.}} 0.$$

Thus

$$s_n^{-2} \sup_{k \leq n} X_k^2 \leq \epsilon^2 + s_n^{-2} \sup_{k \leq n} X_k^2 I(|X_k| \geq \epsilon s_n),$$

and since ϵ may be chosen arbitrarily small,

$$(12) \quad s_n^{-2} \sup_{k \leq n} X_k^2 \rightarrow_{\text{a.s.}} 0.$$

From (3) and Theorem 1 of Pratt [12] we then deduce that

$$(13) \quad \lim_{n \rightarrow \infty} s_n^{-2} \sup_{k \leq n} E X_k^2 \leq \lim_{n \rightarrow \infty} E s_n^{-2} \sup_{k \leq n} X_k^2 = 0,$$

and $s_{n+1}^2/s_n^2 \rightarrow 1$ follows.

We next proceed to the proof of (11) for which we need to know more about the behavior of the sequence T_1, T_2, \dots .

LEMMA 1. *Under the conditions of Theorem 1,*

$$s_n^{-2} \sum_{k=1}^n E\{T_k | \mathcal{G}_{k-1}\} \rightarrow_{\text{a.s.}} 1 \quad \text{as } n \rightarrow \infty.$$

PROOF. Using (8) and (9), we need only show

$$(14) \quad s_n^{-2} \sum_{k=1}^n E\{X_k^{*2} | \mathcal{F}_{k-1}^*\} \rightarrow_{\text{a.s.}} 1 \quad \text{as } n \rightarrow \infty.$$

Now

$$(15) \quad s_n^{-2} \sum_{k=1}^n E\{X_k^{*2} | \mathcal{F}_{k-1}^*\} \\ = s_n^{-2} \sum_{k=1}^n [E\{X_k^2 I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\} - (E\{X_k I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\})^2]$$

and

$$(16) \quad s_n^{-2} \sum_{k=1}^n (E\{X_k I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\})^2 \\ \leq s_n^{-1} \delta \sum_{k=1}^n E\{|X_k| I(|X_k| \geq \delta s_k) | \mathcal{F}_{k-1}^*\} \rightarrow_{a.s.} 0 \quad \text{as } n \rightarrow \infty .$$

Also, using Proposition IV. 6.1, page 147 of Neveu [8],

$$(17) \quad s_n^{-2} \sum_{k=1}^n (X_k^2 I(|X_k| < \delta s_k) - E\{X_k^2 I(|X_k| < \delta s_k) | \mathcal{F}_{k-1}^*\}) \rightarrow_{a.s.} 0 \\ \text{as } n \rightarrow \infty$$

since

$$\sum_{n=1}^{\infty} s_n^{-4} E[X_n^2 I(|X_n| < \delta s_n) - E\{X_n^2 I(|X_n| < \delta s_n) | \mathcal{F}_{n-1}^*\}]^2 \\ \leq \sum_{n=1}^{\infty} s_n^{-4} E[X_n^4 I(|X_n| < \delta s_n)] < \infty$$

using (1). Consequently, from (15), (16) and (17), it suffices to show that

$$s_n^{-2} \sum_{k=1}^n X_k^2 I(|X_k| < \delta s_k) \rightarrow_{a.s.} 1 \quad \text{as } n \rightarrow \infty$$

or equivalently, in view of (3),

$$(18) \quad s_n^{-2} \sum_{k=1}^n X_k^2 I(|X_k| \geq \delta s_k) \rightarrow_{a.s.} 0 .$$

However,

$$s_n^{-2} \sum_{k=1}^n X_k^2 I(|X_k| \geq \delta s_k) \leq (s_n^{-1} \sup_{k \leq n} |X_k|) s_n^{-1} \sum_{k=1}^n |X_k| I(|X_k| \geq \delta s_k)$$

and (18) follows from (2) and (12).

LEMMA 2. Under the conditions of Theorem 1,

$$s_n^{-2} \sum_{k=1}^n (T_k - E\{T_k | \mathcal{G}_{k-1}\}) \rightarrow_{a.s.} 0 \quad \text{as } n \rightarrow \infty .$$

PROOF. Again using Proposition IV. 6.1, page 147 of [8], the result will follow if

$$\sum_{n=1}^{\infty} s_n^{-4} E(T_n - E\{T_n | \mathcal{G}_{n-1}\})^2 < \infty .$$

Now

$$\sum_{n=1}^{\infty} s_n^{-4} E(T_n - E\{T_n | \mathcal{G}_{n-1}\})^2 \\ \leq \sum_{n=1}^{\infty} s_n^{-4} E T_n^2 \\ \leq L_2 \sum_{n=1}^{\infty} s_n^{-4} E X_n^{*4} \quad \text{(by (10))} \\ = L_2 \sum_{n=1}^{\infty} s_n^{-4} E(X_n I(|X_n| < \delta s_n) - E\{X_n I(|X_n| < \delta s_n) | \mathcal{F}_{n-1}^*\})^4 \\ \leq L_2 \sum_{n=1}^{\infty} s_n^{-4} [E(X_n^4 I(|X_n| < \delta s_n)) \\ + 15\delta^3 s_n^3 E\{E\{X_n I(|X_n| < \delta s_n) | \mathcal{F}_{n-1}^*\}\}] \\ \leq L_2 \sum_{n=1}^{\infty} s_n^{-4} [E(X_n^4 I(|X_n| < \delta s_n)) + 15\delta^3 s_n^3 E(|X_n| I(|X_n| \geq \delta s_n))] \\ < \infty$$

by (1) and (2).

We are now in a position to complete the proof of Theorem 1. Since $EX_k^2 < \infty$ for any k ,

$$(19) \quad g(t) \rightarrow \infty \quad \text{as } t \rightarrow \infty .$$

Then also

$$(20) \quad \begin{aligned} |tS_{g(t)}^{-2} - 1| &\leq |S_{g(t)}^{-2} EX_{g(t)+1}^2| \\ &= |S_{g(t)+1}^{-2} EX_{g(t)+1}^2| |S_{g(t)+1}^2 S_{g(t)}^{-2}| \\ &\rightarrow 0 \end{aligned} \quad \text{as } t \rightarrow \infty$$

by (13) and (19). Similarly,

$$(21) \quad tS_{g(t)+1}^{-2} \rightarrow 1 \quad \text{as } t \rightarrow \infty .$$

From (19), (20), (21) and Lemmas 1 and 2 we obtain

$$t^{-1} \sum_{k=1}^{g(t)} T_k \rightarrow_{\text{a.s.}} 1 \quad \text{as } t \rightarrow \infty$$

and

$$t^{-1} \sum_{k=1}^{g(t)+1} T_k \rightarrow_{\text{a.s.}} 1 \quad \text{as } t \rightarrow \infty .$$

Finally, we note that

$$|\eta(t) - \xi(t)| \leq \max \{ |\xi(\sum_{k=1}^{g(t)} T_k) - \xi(t)|, |\xi(\sum_{k=1}^{g(t)+1} T_k) - \xi(t)| \} .$$

From this point, it is obvious that Strassen's proof on page 217 of [17] may be followed to obtain (11), which completes the proof of Theorem 1.

Corollaries 1 and 2 follow easily from Theorem 1 using standard arguments of the kind of, for example, Neveu [8], pages 153, 154.

4. Stationary process results. We consider a probability space (Ω, \mathcal{A}, P) with an ergodic automorphism T . Let $L_2(P)$ be the Hilbert space of random variables with finite second moment, and write $(EX^2)^{\frac{1}{2}}$ as $\|X\|$. Define U on $L_2(P)$ by $UX(\omega) = X(T\omega)$ for $X \in L_2(P)$, $\omega \in \Omega$. Then U is a unitary operator. If \mathcal{M}_0 is a σ -field such that $\mathcal{M}_0 \subset \mathcal{A}$ and $\mathcal{M}_0 \subset T^{-1}(\mathcal{M}_0)$, define $\mathcal{M}_k = T^{-k}(\mathcal{M}_0)$, $\mathcal{M}_{-\infty} = \bigcap_{k=-\infty}^{\infty} \mathcal{M}_k$ and $\mathcal{M}_{+\infty} = \sigma$ -field generated by $\bigcup_{k=-\infty}^{\infty} \mathcal{M}_k$.

We consider a particular random variable $X_0 \in L_2(P)$. Defining $X_k = U^k X_0$, we see that $\dots, X_{-1}, X_0, X_1, \dots$ is a doubly infinite stationary ergodic sequence. Put $X^{(0)} = 0$ and $X^{(n)} = \sum_{k=0}^{n-1} X_k$ for $n \geq 1$ and define a sequence of random functions $\theta_n(\cdot)$ on $[0, 1]$ by

$$\begin{aligned} \theta_n(t) &= [\phi(\|X^{(n)}\|^2)]^{-1} (X^{(k)} + (nt - k)X_k) \\ & \quad k \leq nt \leq k + 1, k = 0, 1, \dots, n - 1 . \end{aligned}$$

Also define

$$g = \sup \{ n : \|X^{(n)}\|^2 \leq e \} .$$

THEOREM 2. *If*

$$(22) \quad \sum_{m=0}^{\infty} (\|E\{X_0 | \mathcal{M}_{-m}\}\| + \|X_0 - E\{X_0 | \mathcal{M}_m\}\|) < \infty$$

then

$$\lim_{n \rightarrow \infty} \|X^{(n)}\|/n^{\frac{1}{2}} = \sigma$$

exists for $0 \leq \sigma < \infty$. If $\sigma > 0$ then $g < \infty$, $\{\theta_n; n > g\}$ is relatively compact and the set of its limit points coincides with K .

PROOF. The key to the proof is a representation

$$(23) \quad X_0 = Y_0 + UZ_0 - Z_0$$

where Y_0, Z_0 belong to $L_2(P)$ and, writing $Y_k = U^k Y_0$, $Y^{(0)} = 0$ and $Y^{(n)} = \sum_{k=1}^n Y_k$ for $k \geq 1$, $\{Y^{(n)}, \mathcal{M}_{n-1}; n \geq 0\}$ is a stationary ergodic zero-mean martingale. This representation, which is due to Gordin, holds under the condition (22); see Scott [14] for full details.

Setting $\|Y_0\| = \sigma$ we have $\|Y^{(n)}\| = \sigma n^{\frac{1}{2}}$. Also, from Minkowski's inequality and (23),

$$\| \|X^{(n)}\| - \sigma n^{\frac{1}{2}}/n^{\frac{1}{2}} \| \leq 2\|Z_0\|/n^{\frac{1}{2}} \rightarrow 0$$

since $Z_0 \in L_2(P)$ and thus

$$\lim_{n \rightarrow \infty} \|X^{(n)}\|/n^{\frac{1}{2}} = \sigma.$$

We henceforth assume $\sigma > 0$. Then

$$(24) \quad \lim_{n \rightarrow \infty} \|X^{(n)}\|/\sigma n^{\frac{1}{2}} = 1$$

and $g < \infty$. Also, from (24) we readily deduce that

$$(25) \quad \phi(\|X^{(n)}\|^2)/\phi(n\sigma^2) \rightarrow 1.$$

Now define a sequence of random functions $\zeta_n(\cdot)$ on $[0, 1]$ by

$$\zeta_n(t) = [\phi(n\sigma^2)]^{-1} [Y^{(k)} + (nt - k)Y_k] \\ k \leq nt \leq k + 1, k = 0, 1, \dots, n - 1.$$

Then by Corollary 2 we know that $\{\zeta_n; n > e/\sigma^2\}$ is relatively compact and the set of its limit points coincides with K . Thus, the proof will be complete if we can show that

$$(26) \quad \sup_{0 \leq t \leq 1} |\theta_n(t) - \zeta_n(t)| \rightarrow_{a.s.} 0.$$

We first consider

$$(27) \quad \sup_{0 \leq t \leq 1} |\zeta_n(t) - \phi(n\sigma^2)[\phi(\|X^{(n)}\|^2)]^{-1} \zeta_n(t)| \\ \leq |1 - \phi(n\sigma^2)[\phi(\|X^{(n)}\|^2)]^{-1}| \sup_{0 \leq t \leq 1} |\zeta_n(t)|.$$

Now

$$\limsup_{n \rightarrow \infty} \sup_{0 \leq t \leq 1} |\zeta_n(t)| = \sup_{x \in K} \sup_{0 \leq t \leq 1} |x(t)| \quad a.s. \\ \leq 1 \quad a.s.$$

From this and (25) and (27) we conclude that

$$(28) \quad \sup_{0 \leq t \leq 1} |\zeta_n(t) - \phi(n\sigma^2)[\phi(\|X^{(n)}\|^2)]^{-1} \zeta_n(t)| \rightarrow_{a.s.} 0.$$

Now, writing $Z_k = U^k Z_0$, consider

$$\sup_{0 \leq t \leq 1} |\theta_n(t) - \phi(n\sigma^2)[\phi(\|X^{(n)}\|^2)]^{-1} \zeta_n(t)| \\ = \sup_{1 \leq k \leq n} [\phi(\|X^{(n)}\|^2)]^{-1} |Z_k - Z_0| \quad \text{(by (23))} \\ \leq 2 \sup_{0 \leq k \leq n} [\phi(\|X^{(n)}\|^2)]^{-1} |Z_k|$$

which goes almost surely to zero provided

$$\sup_{0 \leq k \leq n} [\phi(n\sigma^2)]^{-1} |Z_k| = (n\sigma^2 \log \log n\sigma^2)^{-\frac{1}{2}} \sup_{0 \leq k \leq n} |Z_k| \xrightarrow{\text{a.s.}} 0$$

(by (25)) and thus if

$$(n \log \log n)^{-\frac{1}{2}} |Z_n| \xrightarrow{\text{a.s.}} 0.$$

But, this last result follows from the Borel–Cantelli Lemma since $Z_0 \in L_2(P)$. Thus

$$\sup_{0 \leq t \leq 1} |\theta_n(t) - \phi(n\sigma^2)[\phi(\|X^{(n)}\|^2)]^{-1} \zeta_n(t)| \xrightarrow{\text{a.s.}} 0$$

and taking this with (28), (26) follows and the proof is complete.

As an application of Theorem 2, we shall see how it extends some recent results of Oodaira and Yoshihara [9], [10]. The papers [9], [10] in turn improve results of Iosifescu [7] and Reznik [13] as well as the (strictly) stationary case of results of Philipp [11].

Let $\{X_j, -\infty < j < \infty\}$ be a stationary process defined on a probability space (Ω, \mathcal{A}, P) . Write \mathcal{M}_k for the σ -field generated by \dots, X_{k-1}, X_k ; \mathcal{M}_∞ for the σ -field generated by $\dots, X_{-1}, X_0, X_1, \dots$ and \mathcal{M}_k^∞ for the σ -field generated by X_k, X_{k+1}, \dots . The sequence $\{X_j\}$ is called ϕ -mixing if for each $k (-\infty < k < \infty)$ and each $n (n \geq 1)$,

$$\sup_{B \in \mathcal{M}_{k+n}^\infty} \text{ess sup } |P(B | \mathcal{M}_k) - P(B)| \leq \phi_n \downarrow 0 \quad \text{as } n \rightarrow \infty.$$

COROLLARY 3. *If $\{X_n\}$ is a stationary ϕ -mixing process with $EX_0 = 0$, $E|X_0|^{2+\delta} < \infty$ some $\delta \geq 0$ and $\sum_{n=1}^\infty [\phi_n]^{(1+\delta)/(2+\delta)} < \infty$, then*

$$\lim_{n \rightarrow \infty} \|X^{(n)}\|^2/n = EX_0^2 + 2 \sum_{i=1}^\infty EX_0 X_i = \sigma^2 \geq 0.$$

If $\sigma > 0$ then $g < \infty$ and $\{\theta_n; n > g\}$ is relatively compact and the set of its limit points coincides with K .

PROOF. First note that $\{X_n\}$ is stationary and ϕ -mixing and hence ergodic. Further, there exists an ergodic transformation T on \mathcal{M}_∞ such that $X_k(\omega) = X_0(T^k\omega)$, $\omega \in \Omega$ (e.g. Doob [3], Chapter X) and, since \mathcal{M}_0 is the σ -field generated by \dots, X_{-1}, X_0 our present situation fits within the framework of Theorem 2.

Next we need to check (22). This is a simple matter using Lemma 1, page 170 of Billingsley [1]. We have

$$\sum_{m=0}^\infty (\|E\{X_0 | \mathcal{M}_{-m}\} \| + \|X_0 - E\{X_0 | \mathcal{M}_m\} \|) = \sum_{m=0}^\infty \|E\{X_0 | \mathcal{M}_{-m}\} \|$$

and

$$\begin{aligned} E[E\{X_0 | \mathcal{M}_{-m}\}]^2 &= E[X_0 E\{X_0 | \mathcal{M}_{-m}\}] \\ &\leq 2\phi_m^{(1+\delta)/(2+\delta)} [E|E\{X_0 | \mathcal{M}_{-m}\}|^{(2+\delta)/(1+\delta)}]^{(1+\delta)/(2+\delta)} [E|X_0|^{2+\delta}]^{1/(2+\delta)} \\ &\leq 2\phi_m^{(1+\delta)/(2+\delta)} [E[E\{X_0 | \mathcal{M}_{-m}\}]^2]^{\frac{1}{2}} [E|X_0|^{2+\delta}]^{1/(2+\delta)} \end{aligned}$$

so that

$$\|E\{X_0 | \mathcal{M}_{-m}\} \| \leq 2\phi_m^{(1+\delta)/(2+\delta)} [E|X_0|^{2+\delta}]^{1/(2+\delta)}$$

and (22) holds under the conditions of the corollary.

Finally, from Lemma 3, page 172 of [1],

$$\lim_{n \rightarrow \infty} \|X^{(n)}\|^2/n = EX_0^2 + 2 \sum_{i=1}^{\infty} E(X_0 X_i)$$

and the corollary follows from Theorem 2.

In Theorem 1 of Oodaira and Yoshihara [10], an invariance principle for the law of the iterated logarithm is given under the conditions of Corollary 3 plus the requirements that $\int_{|x| \geq N} x^2 dP(X_0 \leq x) = O((\log N)^{-\delta})$ as $N \rightarrow \infty$ if $\delta = 0$ or $\phi_n = O(n^{-1-\varepsilon})$ for some $\varepsilon > (1 + \delta)^{-1}$ if $\delta > 0$.

REFERENCES

- [1] BILLINGSLEY, P. (1968). *Convergence of Probability Measures*. Wiley, New York.
- [2] CHOVER, J. (1967). On Strassen's version of the log log law. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **8** 83-90.
- [3] DOOB, J. L. (1953). *Stochastic Processes*. Wiley, New York.
- [4] GORDIN, M. I. (1969). The central limit theorem for stationary processes. *Soviet Math. Dokl.* **10** 1174-1176.
- [5] HEYDE, C. C. (1969). Some properties of metrics in a study on convergence to normality. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **11** 181-192.
- [6] HEYDE, C. C. (1971). An iterated logarithm result for martingales and its application in estimation theory for autoregressive processes. To appear in *J. Appl. Probability* **9**.
- [7] IOSIFESCU, M. (1968). The law of the iterated logarithm for a class of dependent random variables. *Theor. Probability Appl.* **13** 304-313.
- [8] NEVEU, J. (1965). *Mathematical Foundations of the Calculus of Probability*. Holden-Day, San Francisco.
- [9] OODAIRA, H. and YOSHIHARA, K. (1971a). The law of the iterated logarithm for stationary processes satisfying mixing conditions. *Kôdai Math. Sem. Rep.* **23** 311-334.
- [10] OODAIRA, H. and YOSHIHARA, K. (1971b). Note on the law of the iterated logarithm for stationary processes satisfying mixing conditions. *Kôdai Math. Sem. Rep.* **23** 335-342.
- [11] PHILIPP, W. (1969). The law of the iterated logarithm for mixing stochastic processes. *Ann. Math. Statist.* **40** 1985-1991.
- [12] PRATT, J. W. (1960). On interchanging limits and integrals. *Ann. Math. Statist.* **31** 74-77.
- [13] REZNIK, M. Kh. (1968). The law of the iterated logarithm for some classes of stationary processes. *Theor. Probability Appl.* **13** 606-621.
- [14] SCOTT, D. J. (1971). Central limit theorems for martingales and for processes with stationary increments, using a Skorokhod representation approach. To appear in *Adv. Appl. Probability*.
- [15] STOUT, W. F. (1970). A martingale analogue of Kolmogorov's law of the iterated logarithm. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **15** 279-290.
- [16] STOUT, W. F. (1970). The Hartman-Wintner law of the iterated logarithm for martingales. *Ann. Math. Statist.* **41** 2158-2160.
- [17] STRASSEN, V. (1964). An invariance principle for the law of the iterated logarithm. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **3** 211-226.
- [18] STRASSEN, V. (1967). Almost sure behavior of sums of independent random variables and martingales. *Proc. Fifth Berkeley Symp. Math. Statist. Prob.* **2** 315-343. Univ. of California Press.

DEPARTMENT OF STATISTICS
 AUSTRALIAN NATIONAL UNIVERSITY
 BOX 4, P.O. CANBERRA, A.C.T. 2600
 AUSTRALIA