

## CURRENT STATUS AND RIGHT-CENSORED DATA STRUCTURES WHEN OBSERVING A MARKER AT THE CENSORING TIME<sup>1</sup>

BY MARK J. VAN DER LAAN AND NICHOLAS P. JEWELL

*University of California, Berkeley*

We study nonparametric estimation with two types of data structures. In the first data structure  $n$  i.i.d. copies of  $(C, N(C))$  are observed, where  $N$  is a finite state counting process jumping at time-variables of interest and  $C$  a random monitoring time. In the second data structure  $n$  i.i.d. copies of  $(C \wedge T, I(T \leq C), N(C \wedge T))$  are observed, where  $N$  is a counting process with a final jump at time  $T$  (e.g., death). This data structure includes observing right-censored data on  $T$  and a marker variable at the censoring time.

In these data structures, easy to compute estimators, namely (weighted)-pool-adjacent-violator estimators for the marginal distributions of the unobservable time variables, and the Kaplan–Meier estimator for the time  $T$  till the final observable event, are available. These estimators ignore seemingly important information in the data. In this paper we prove that, at many continuous data generating distributions the ad hoc estimators yield asymptotically efficient estimators of  $\sqrt{n}$ -estimable parameters.

**1. Introduction.** In this paper we study nonparametric estimation with two types of data structures. First, we discuss these two data structures in detail. Subsequently, we provide an overview of the rest of the paper.

1.1. *Current status data on a finite counting process.* Consider a finite state counting process  $N(t) = \sum_{j=1}^k I(T_j \leq t)$ ,  $T_1 < \dots < T_k$ , where  $T_j$  is the time-variable at which a specified event occurs and where  $N$  jumps from value  $j - 1$  to  $j$  at time  $T_j$ . The number of jumps  $k$  is fixed and known. We allow that there is a positive probability that the counting process never reaches jump  $j_0$  for any particular  $j_0 \in \{1, \dots, k\}$ ; since  $T_1 < \dots < T_k$ , this implies that there is also a positive probability that  $N$  never reaches jump  $j$  for  $j = j_0, \dots, k$ : that is, we allow multivariate distributions of  $(T_1, \dots, T_k)$  with  $P(T_j = \infty) > 0$  for  $j = j_0, \dots, k$ . In this manner we allow applications in which the number of jumps of  $N$  is random on  $\{1, \dots, k\}$ .

We consider the data structure  $(C, N(C))$  for a single random monitoring time  $C$ . The only assumption is that  $C$  is independent of  $N$ : the cumulative

---

Received October 1998; revised November 2001.

<sup>1</sup>Supported by a FIRST award (GM53722) from the National Institute of General Medical Sciences and the National Institutes of Health.

AMS 2000 subject classifications. Primary 62G07; secondary 62F12.

Key words and phrases. Asymptotically linear estimator, asymptotically efficient estimator, current status data, right-censored data, isotonic regression.

distribution  $G$  of  $C$ , and the probability distribution  $F$  of  $N$  are unspecified. Note that the distribution of  $N$ , denoted by  $F$ , is not a cumulative distribution function, but a probability distribution that is identified by the multivariate cumulative distribution of  $(T_1, \dots, T_k)$ .

Such data structures occur in cross-sectional studies where each subject is monitored once. For example, in some carcinogenicity experiments, one can only determine a discretized occult tumor size at time  $t$  in a randomly sampled mouse, as measured by  $N(t)$ , by sacrificing a mouse at time  $t$ . In this example,  $T_1$  might represent time till onset of the tumor and  $T_2, \dots, T_k$  might correspond with times till increasing sizes of the tumor. Similarly,  $T_j$  might denote the age at which a child has mastered the  $j$ th skill among a set of  $k$  skills ordered in difficulty. We refer to Jewell and van der Laan (1995) for additional applications.

The distribution of  $(C, N(C))$  depends on the distribution of  $\vec{T} = (T_1, \dots, T_k)$  only through the marginal distributions  $F_j$  of  $T_j$ ,  $j = 1, \dots, k$  (see Section 2). In this problem, the NPMLE of the distribution of  $T_j$  requires an iterative algorithm. On the other hand, an ad hoc method for estimation of the distribution of  $T_j$  is directly available: reduce the observation  $(C, N(C))$  to a standard current status observation  $(C, \Delta_j = I(T_j \leq C))$  on  $T_j$ . Then one can estimate the distribution of  $T_j$  with the NPMLE based on the reduced current status observations, which we will refer to as the reduced data NPMLE (RNPMLE). This estimator provides regular and asymptotically linear estimators of pathwise differentiable functionals of  $F_j$  such as  $\mu_j = \int (1 - F_j)(u)r(u) du$ , for a given  $r$ , in the nonparametric model under certain conditions [Groeneboom and Wellner (1992)]. Previous work and examples of traditional current status data on a time variable  $T$  can be found in Diamond, McDonald and Shah (1986), Jewell and Shiboski (1990), Diamond and McDonald (1992), Keiding (1991) and Sun and Kalbfleisch (1993). In its nonparametric setting, the current status data structure is also known as case I interval censored data [Groeneboom and Wellner (1992)]. Current status data commonly arise in epidemiological investigations of the natural history of disease and in animal tumorigenicity experiments. Jewell, Malani and Vittinghoff (1994) give two examples that arise from studies of Human Immunodeficiency Virus (HIV) disease.

Note that the RNPMLE of  $F_j$  ignores the value of  $N(C)$ , beyond information on whether  $N(C) \geq j$  or not. For example, if  $N(t)$  is tumor size in a carcinogenicity experiment, then the simple current status estimator of the distribution of time,  $T_1$ , till onset of tumor would not distinguish between an observation  $(C, N(C))$  with  $N(C)$  large and an observation  $(C, N(C))$  with  $N(C)$  small but larger than 0, while the latter observation seems to suggest that onset occurred recently. Nonetheless, we establish that the RNPMLE yields efficient estimators of pathwise differentiable parameters at a large class of continuous data generating distributions of interest.

1.2. *Current status data on a finite counting process when the final event is right censored.* We also consider the data structure  $(\tilde{T}_k \equiv C \wedge T_k, N(\tilde{T}_k))$  for a finite state counting process  $N(t) = \sum_{j=1}^k I(T_j \leq t)$ , where  $T_k$  represents the final event (say death) which is right censored by the monitoring time  $C$ , and  $k$  is known. Note that this observation includes observing the failure indicator  $I(\tilde{T}_k = T_k)$ . For example, consider a carcinogenicity experiment with mice in which  $T_1$  is time till onset of colon tumor,  $T_2$  time to liver metastasis and  $T_3$  time to death from tumor, where we assume that colon tumors do not cause death except through liver failure secondary to metastasis. Here  $C$  is either a sacrificing time or time till death from any unrelated cause.

Consider another example concerning estimation of the survival function of the time  $T = J - I$  between time  $I$  at seroconversion and time  $J$  at death of a hemophiliac patient infected with HIV. For this purpose we observe  $n$  i.i.d. subjects in a fixed time-interval of 10 years. If we assume that the time  $I$  at seroconversion of the subject is observed (which is approximately true for hemophiliacs), then the subject's survival time  $T$  is right censored by  $C \equiv 10 - I$ , where  $T$  will play the role of  $T_k$ . We define  $T_j$  as the time till a given monotone "surrogate" process  $Z(t)$  achieves a particular value among a set of  $k - 1$  increasing values,  $j = 1, \dots, k - 1$ , where we assume that death  $T = T_k$  always and only occurs after the value  $Z(T_{k-1})$  has been reached. Let  $N(t) = \sum_{j=1}^k I(T_j \leq t)$  be the counting process. Here  $Z(t)$  measures the progression of the disease of the subject  $t$  years after seroconversion; for example,  $Z(t)$  might be a measure of viral load of the subject  $t$  years after seroconversion, where it may be reasonable to assume that the viral load is a nondecreasing process in the absence of treatment.

Suppose that for every subject who did not die before the end of the study  $C$  one measures the "surrogate"  $Z(C)$  at time  $C$  only. In other words, we observe failure times only for subjects who fail before end of follow up and for every subject who is alive at end of follow up we also have a marker indicating future prognosis. Note that the observed data on a subject is given by  $(\tilde{T} = T \wedge C, Z(\tilde{T}))$ . We only assume that  $C$  is independent of  $Z$ . A seemingly ad hoc estimator of  $S(t) = P(T > t)$  is the Kaplan–Meier estimator which simply ignores the marker information. In this example, a natural question is whether one can improve on the Kaplan–Meier estimator using the information in the surrogate process  $Z$ . In this paper we prove that the Kaplan–Meier estimator is asymptotically efficient at many continuous data generating distributions for which  $F_j$  have compact support.

A special case of this data structure has been treated in the literature. Consider a carcinogenicity experiment with  $N(t) = \sum_{j=1}^2 I(T_j \leq t)$ ,  $T_1$  is time till onset of tumor and  $T_2$  is time till death from tumor. Thus one observes  $(\tilde{T}_2 \equiv C \wedge T_2, N(\tilde{T}_2))$ . This data structure has been considered in Kodell, Shaw and Johnson (1982), Dinse and Lagakos (1982), Turnbull and Mitchell (1984), van der Laan, Jewell and Peterson (1997), and recently Groeneboom (1998). The NPMLE for this data structure requires an iterative algorithm: Turnbull and Mitchell (1984)

implemented the NPMLE via the EM-algorithm (using an initial distribution with point masses at each data point so that the EM-algorithm indeed converges to the NPMLE), while Groeneboom (1998) implements the NPMLE by maximizing the actual likelihood with a modern optimization algorithm. In this problem, an ad hoc estimator of the distribution of  $T_2$  is the Kaplan–Meier estimator based on the reduced data  $(\tilde{T}_2, \Delta_2 = I(\tilde{T}_2 = T_2))$ . In Dinse and Lagakos (1982), the Kaplan–Meier estimator of  $F_2$  was proposed and it was suggested that the NPMLE might be more efficient than the Kaplan–Meier estimator. In van der Laan, Jewell and Peterson (1997) it is shown that the Kaplan–Meier is efficient under a weak condition on  $(F_1, F_2)$ . Moreover, an isotonic regression estimator of  $F_1$  was provided: note that estimation of  $F_1$  is complicated by the fact that for some subjects one only observes  $T_2$  and thus that  $T_1 < T_2$ , where  $T_2$  cannot be viewed as an independent monitoring time for  $T_1$ . We note here that, in van der Laan, Jewell and Peterson (1997), a simulation study was carried out which incorrectly implements the NPMLE, so that finite sample comparisons between the Kaplan–Meier estimator and the NPMLE remain open to study [specifically the derivation of the score equations in van der Laan, Jewell and Peterson (1997) for the NPMLE was not valid since the authors incorrectly assumed that the NPMLE  $\hat{F}_1$  is strictly smaller than the NPMLE  $\hat{F}_2$ ].

1.3. *Organization and overview of results.* In Section 2 we prove, for the data structure of Section 1.1, that if the  $F_j$ 's are continuous with Lebesgue density bounded away from zero on  $[0, \tau_j]$  and zero elsewhere, and  $G$  is also continuous, then any estimator of a parameter  $\mu = \Phi(F) \in \mathbb{R}$  that is regular and asymptotically linear at  $P_{F,G}$  is also asymptotically efficient. The complexity of the NPMLE is discussed including that it is more efficient at many data generating distributions with singular pairs  $F_{j_1}, F_{j_2}$ ; for example,  $F_1$  discrete and  $F_2$  continuous.

In Section 3, we prove an analogous result for the nonparametric model with the data structure  $(C \wedge T_k, N(C \wedge T_k))$ . This shows that the Kaplan–Meier estimator of the distribution of  $T_k$ , based on the reduced data  $(\tilde{T}_k, \Delta_k \equiv I(T_k \leq C))$ , is asymptotically efficient at many continuous data generating distributions, extending the result in van der Laan, Jewell and Peterson (1997) for the case  $k = 2$ . Moreover, simple isotonic regression estimators for the distributions  $F_j$ ,  $j = 1, \dots, k - 1$ , are proposed that also yield asymptotically efficient estimators of smooth functionals by our general result.

## 2. Current status data on a counting process.

2.1. *Traditional current status data.* Traditional current status data can be viewed as current status data on a simple counting process as follows. Let  $T$  be a univariate failure time of interest and define the process  $\Delta(t) = I(T \leq t)$  as the counting process with one single jump at point  $T$ . Let  $Y = (C, \Delta(C))$  represent

current status data on  $\Delta$  at a monitoring time  $C$ . We assume that  $C$  is independent of  $T$  [i.e., of  $\Delta(\cdot)$ ]. The parameter of interest is the distribution  $F$  of  $T$ .

The properties of the NPMLE  $F_n$  of the distribution of  $T$  were established in Groeneboom and Wellner (1992). Here the NPMLE is defined as the maximum likelihood estimator over all discrete distributions with jumps at the monitoring times. Beyond proving a limit distribution result for  $F_n$ , these authors also established efficiency of smooth functionals of  $F_n$  with a closed form expression of the limit variance so that Wald-type confidence intervals are directly available. Huang and Wellner (1995) provide an alternative proof of asymptotic linearity of the NPMLE of smooth functionals of  $F$  under weak conditions.

We refer to Bickel, Klaassen, Ritov and Wellner (1993) for definitions of a *regular, asymptotically linear and efficient estimator* and *influence curve* of an estimator. The semiparametric-information bound at  $P_{F,G}$  is defined as the infimum of parametric information bounds over a specified class of parametric submodels. We choose as parametric one-dimensional submodels

$$\left\{ \varepsilon \rightarrow P_{F_\varepsilon, h_1, G_\varepsilon, h_2} : \|h_j\|_\infty < \infty, j = 1, 2, \int h_1 dF = \int h_2 dG = 0 \right\},$$

where  $dF_{\varepsilon, h_1}(\cdot) = (1 + \varepsilon h_1(\cdot)) dF(\cdot)$ ,  $dG_{\varepsilon, h_2}(\cdot) = (1 + \varepsilon h_2(\cdot)) dG(\cdot)$  and  $\varepsilon$  is the unknown parameter with parameter space  $[-\delta, \delta]$  for some small  $\delta > 0$ . The tangent space at  $P_{F,G}$  is now defined as the closure in  $L_0^2(P_{F,G})$  of the linear span of all the scores of these one-dimensional submodels, where, for a given measure  $\mu$ , we define  $L_0^2(\mu) = \{h : \int h^2 d\mu < \infty, \int h d\mu = 0\}$  as the Hilbert space endowed with inner product  $\langle h_1, h_2 \rangle_\mu = \int h_1(y)h_2(y) d\mu(y)$ . Thus the tangent space at  $P_{F,G}$  is a sub-Hilbert space of  $L_0^2(P_{F,G})$ .

In this paper it is particularly important to realize that efficiency of an estimator is a local property in the sense that a regular estimator can be efficient at a particular  $P_{F,G}$  and inefficient at another element of the model.

LEMMA 2.1. *Consider the nonparametric model for  $Y = (C, \Delta(C))$ , where  $\Delta(\cdot) \equiv I(T \leq \cdot)$ ,  $T$  has unspecified distribution  $F$  and  $C$  is independent of  $T$  with unspecified distribution  $G$ . We observe  $n$  i.i.d. observations of  $Y = (C, \Delta(C))$ . Consider the parameter  $\mu = \int (1 - F)(u)r(u) du$  for a given function  $r$ . Consider the estimator  $\mu_n = \int (1 - F_n)(u)r(u) du$ , where  $F_n$  is the NPMLE of  $F$ . We have that  $\mu_n$  is regular and asymptotically linear at any  $(F, G)$  for which  $F$  is continuous with density  $f_T > 0$  on  $[0, M]$  and zero elsewhere ( $M < \infty$ ),  $g(x) = dG/dx > 0$  on  $[0, M]$ , and  $r$  is bounded on  $[0, M]$ .*

The influence curve of  $\mu_n$  is given by

$$(1) \quad IC(Y | F, g, r) = \frac{r(C)}{g(C)}(F(C)(1 - \Delta) - (1 - F(C))\Delta).$$

The variance of IC is given by

$$VAR(IC) = \int \frac{r^2(c)}{g(c)} F(c)(1 - F(c)) dc.$$

This lemma is proved in Huang and Wellner (1995).

We can also prove the following tangent space result.

LEMMA 2.2. Consider the nonparametric model for  $Y = (C, \Delta(C))$ , where  $\Delta(\cdot) \equiv I(T \leq \cdot)$ ,  $T$  has unspecified distribution  $F$ , and  $C$  is independent of  $T$  with unspecified distribution  $G$ . We observe  $n$  i.i.d. observations of  $Y = (C, \Delta(C))$ . Suppose that:

- (1)  $F$  has a Lebesgue density  $f$  with  $f > 0$  on  $[0, \tau_F)$  and, if  $\tau_F < \infty$  ( $\tau_F = \infty$  is allowed), then  $f = 0$  on  $(\tau_F, \infty)$ , and
- (2)  $G$  has a Lebesgue density  $g$ .

We allow  $F(\{\infty\}) > 0$ . Then the tangent space at  $P_{F,G}$  equals  $L_0^2(P_{F,G})$ . This implies that an estimator of a parameter  $\mu(F)$  which is regular and asymptotically linear at  $P_{F,G}$  is also asymptotically efficient if  $F, G$  satisfy (1) and (2).

In Gill, van der Laan and Robins (1997) it is proved that if one only assumes that the conditional distribution of the observed data  $Y$ , given the full data  $T$ , satisfies “coarsening at random” (CAR), then the tangent space at  $P_{F,G}$  is saturated, that is, equals  $L_0^2(P_{F,G})$ . The tangent space generated by  $G(\cdot | T)$  under the sole assumption CAR equals  $T_{CAR} = \{v(Y) \in L_0^2(P_{F,G}) : E(v(Y) | T) = 0\}$ . Therefore, the main idea of the proof below is to show that under the independent censoring model  $G(\cdot | T) = G(\cdot)$ , the tangent space of the marginal distribution  $G$  equals  $T_{CAR}$  at a  $P_{F,G}$  satisfying (1) and (2) of Lemma 2.2. The proof below will be an ingredient of the proofs of our two main theorems.

PROOF OF LEMMA 2.2. Let  $A : L_0^2(F) \rightarrow L_0^2(P_{F,G})$ ,  $A(h)(Y) = E_F(h(T) | Y)$  be the score operator for  $F$  and let  $A^\top : L_0^2(P_{F,G}) \rightarrow L_0^2(F)$ ,  $A(V)(T) = E_G(V(Y) | T)$  be its adjoint. The closure of the range of a Hilbert space operator equals the orthogonal complement of the null-space of its adjoint; that is,  $\overline{R(A)} = N(A^\top)^\perp$ , where  $\overline{R(A)}$  is the closure of the range of the score operator and  $N(A^\top)$  is the null space of  $A^\top$ . Thus  $L_0^2(P_{F,G}) = \overline{R(A)} + N(A^\top)$ .

The data generating distribution is indexed by two locally variation-independent parameters  $F$  and  $G$ , so that the tangent space at  $P_{F,G}$  can be obtained as a sum of two tangent spaces, namely the tangent space for  $F$ , which is given by  $\overline{R(A)}$ , and the tangent space for  $G$ . For every  $h \in L_0^2(G)$  with finite supremum norm, we have that  $\varepsilon \rightarrow (1 + \varepsilon h_2) dG$  is a one-dimensional submodel through  $G$  at  $\varepsilon = 0$ . Thus the tangent space corresponding with submodels  $\varepsilon \rightarrow P_{F,G_\varepsilon}$  equals  $L_0^2(G)$ . Thus we have that the tangent space is given by  $\overline{R(A)} + L_0^2(G)$ . We conclude that it suffices to show that  $N(A^\top) = L_0^2(G)$ .

We have

$$A^\top(V)(T) = \int_0^T V(c, 0) dG(c) + \int_T^\infty V(c, 1) dG(c).$$

Thus  $\int V(c, \Delta(c)) dG(c) = 0$   $F$ -a.e. implies that

$$(2) \quad \int_0^T \{V(c, 0) - V(c, 1)\}g(c) dc = \int_0^\infty V(c, 1) dG(c) \quad \text{for } T \in [0, \tau_F).$$

Differentiation w.r.t  $T$  yields  $V(C, 0) = V(C, 1)$  on  $[0, \tau_F)$   $G$ -a.e. If  $\tau_F < \infty$  and  $c > \tau_F$ , then  $c > T$  and thus  $V(c, \Delta(c)) = V(c, 1)$ . Thus  $V(C, 0) = V(C, 1)$   $G$ -a.e. which proves  $N(A^\top) = L_0^2(G)$ .  $\square$

It is of interest to note that one can represent  $F_T(t)$  as a monotonic regression of  $\Delta$  on  $C$  since  $F(t) = E(\Delta(C) | C = t)$ . This suggests that one can estimate  $F_T$  with the estimator  $F_n(t)$  which minimizes  $\sum_{i=1}^n (\Delta(C_i) - F_T(C_i))^2$  over all distribution functions  $F_T$ .  $F_n(t)$  can be computed using the pool-adjacent-violator-algorithm [see Barlow, Bartholomew, Bremner and Brunk (1972)] which, in fact, yields the NPMLE.

2.2. *Current status data on a counting process.* Let the process of interest be a counting process  $N(t) = \sum_{j=1}^k I(T_j \leq t)$ ,  $T_1 < \dots < T_k$ , where  $T_j$  is the time-variable at which an event occurs and where  $N$  jumps from value  $j - 1$  to  $j$ . Let  $C$  be a monitoring time and consider the data structure  $Y = (C, N(C))$ . We observe  $n$  i.i.d. copies of  $Y$ . We only assume that  $C$  is independent of  $N$ .

The distribution of  $(C, N(C))$  depends on the distribution of  $\vec{T}$  only through the marginal distributions  $F_j$  of  $T_j$ ,  $j = 1, \dots, k$ . To be precise, we have (denoting  $S_j = 1 - F_j$ ), for  $j \in \{0, \dots, k\}$ ,

$$\begin{aligned} P_{F,G}(dc, N(C) = j) &= I(j = 0)S_1(c) dG(c) + I(j = k)F_k(c) dG(c) \\ &\quad + I(j = 1)\{S_2(c) - S_1(c)\} dG(c) \\ &\quad + \dots + I(j = k - 1)\{S_k(c) - S_{k-1}(c)\} dG(c). \end{aligned}$$

Thus the distribution of  $Y = (C, N(C))$  only identifies the marginal distributions of  $T_j$ ,  $j = 1, \dots, k$ .

The NPMLE does not exist in closed form and can only be computed with an iterative algorithm. For a given  $j$ , we can reduce the observation  $(C, N(C))$  to simple current status data  $(C, \Delta_j = I(T_j \leq C))$  on  $T_j$ , and estimate  $F_j$  with the RNPMLE. Under the conditions stated in Lemma 2.1, with  $F = F_j$  and  $G = G$ , this estimator provides regular and asymptotically linear estimators of smooth functionals of the type  $\mu_j = \int (1 - F_{T_j})(u)r(u) du$ , for a given  $r$  in the nonparametric model. The following theorem proves that, at a data generating distribution of  $Y$  satisfying a specified condition, any regular asymptotically linear estimator will provide asymptotically efficient estimators of smooth functionals of  $F_{T_j}$ . We decided to state a condition (3) which is easy to understand, but our proof shows that this can be weakened, for example, to allow the analogue of (3) for the case where all distributions  $G, F_1, \dots, F_k$  are discrete with a finite number of support points; that is, the support points of  $F_j$  are contained in the support

points of  $F_{j+1}$ ,  $j = 1, \dots, k - 1$ , and  $G$  is discrete with support contained in the support of  $F_k$ .

**THEOREM 2.1.** *Let  $T_1 < T_2 < \dots < T_k$  be time-variables corresponding to the chronological events of interest. Define the counting process with jumps of size 1 at these  $T_j$ 's by*

$$N(t) = \sum_{j=1}^k I(T_j \leq t).$$

*Let  $Y = (C, N(C))$ . Consider the following semiparametric model for  $Y$ : Let  $C \sim G$  be independent of  $\vec{T} \sim F$ , but leave  $G$  and  $F$  unspecified. Then, the distribution of  $Y$  only depends on the multivariate distribution  $F$  of  $\vec{T} = (T_1, \dots, T_k)$  through the marginal distributions  $F_1, \dots, F_k$  of  $T_1, \dots, T_k$ .*

*Consider a data generating distribution  $P_{F,G}$  in the model above, satisfying the following condition (3): For certain  $\tau_1 < \dots < \tau_k < \infty$ , let  $F_j$  have Lebesgue density  $f_j$  on  $[0, \tau_j]$  with*

$$(3) \quad \begin{array}{ll} f_j > 0 & \text{on } [0, \tau_j] \quad \text{and} \quad f_j = 0 \quad \text{on } (\tau_j, \infty), \quad j = 1, \dots, k, \\ F_j > F_{j+1} & \text{on } (0, \tau_j], \quad j = 1, \dots, k - 1, \\ G & \text{has Lebesgue density } g. \end{array}$$

*We allow that  $p_j \equiv P(T_j = \infty) > 0$  for  $j = j_0, \dots, k$  and  $j_0 \in \{1, \dots, k\}$ .*

*Then the tangent space at  $P_{F,G}$  equals  $L_0^2(P_{F,G})$  and is thus saturated.*

*This implies that any estimator of a real valued parameter of  $F$  that is a regular and asymptotically linear estimator at  $P_{F,G}$  is also asymptotically efficient if  $P_{F,G}$  satisfies (3). In particular, given  $j \in \{1, \dots, k\}$ , if  $P_{F,G}$  satisfies (3), and  $F_j, G$  satisfy the conditions of Lemma 2.1 for the RNPMLE of  $\mu_{F_j}$  based on  $(C, I(T_j \leq C))$  (thus with  $F = F_j$  and  $G = G$ ), then the RNPMLE of  $\mu_{F_j}$  is asymptotically efficient.*

**2.2.1. Heuristic understanding of the difference between NPMLE and RNPMLE.** To understand the difference between the NPMLE and the RNPMLE, we consider the special case  $k = 2$  in detail. In this case  $N$  can have three possible values:

$$N(C) = \begin{cases} 0, & \text{if } C < T_1, \\ 1, & \text{if } T_1 < C < T_2, \\ 2, & \text{if } C > T_2. \end{cases}$$

Let us assume that  $C$  has a Lebesgue density  $g$ . The likelihood of  $(C, N(C))$  is given by

$$P_{F_1, F_2, G}(c, N(c) = j) = S_1(c)^{I(j=0)} (S_2 - S_1)(c)^{I(j=1)} F_2(c)^{I(j=2)} g(c).$$



We note that the density  $p_{F_1, F_2, G}$  can be reparametrized as

$$p_{R, F_2, G}(c, \delta) = R(c)^{I(j=0)}(1 - R(c))^{I(j=1)}S_2(c)^{I(j \in \{0, 1\})}F_2(c)^{I(j=2)}g(c),$$

where  $R(t) \equiv S_1(t)/S_2(t)$ . Thus, if we ignore the relation between  $F_2$  and  $R$ , then the NPMLE of  $F_2$  of the likelihood corresponding with  $p_{R, F_2, G}$  would actually be equal to the reduced data NPMLE based on the reduced data  $(C, I(T_2 \leq C))$ . However,  $F_2$  and  $R$  are related since  $S_2R$  has to be a survival function. Therefore, it is not possible to determine the NPMLE by separate maximization w.r.t.  $F_2$  and  $R$ , which explains why the NPMLE and the RNPMLE of  $F_2$  differ.

Theorem 2.1 shows that this relation between  $F_2$  and  $R$  is not informative for estimation of smooth functionals of  $F_2$  at a large class of data generating distributions, since the RNPMLE, which ignores this relation, is still asymptotically efficient for estimation of  $\sqrt{n}$ -estimable parameters. Our proof of Theorem 2.1 for  $k = 2$  shows that the efficient score operator (for the definition of an efficient score operator, see the proof) of  $F_2$  equals the efficient score operator for  $F_2$  in the reduced data model based on  $(C, \Delta_2)$ . This implies that, at  $(F_1, F_2)$  satisfying (3), the efficient influence curve for any smooth functional of  $F_2$  equals the influence curve of the RNPMLE as given in Lemma 2.1. Closer inspection of the proof for  $k = 2$  also shows that, if (e.g.)  $F_2$  is continuous while  $F_1$  is discrete on  $[0, \tau_1]$ , or  $F_2$  is discrete with support not containing the support of a discrete  $F_1$ , then the efficient score operator for  $F_2$  is not the same as the efficient score operator for  $F_2$  in the reduced data model, so that, in particular, the efficient influence curves (and information bounds) differ for the two models. Thus, at such  $(F_1, F_2)$ , the RNPMLE of smooth functionals of  $F_2$  is inefficient.

Here, we provide a likelihood-based explanation of this fact. Let  $R_n$  be the NPMLE of  $R$ . The NPMLE of  $F_2$  maximizes the likelihood corresponding with  $p_{R_n, F_2}$  over all  $F_2$  for which  $S_2R_n$  is a survival function, while the RNPMLE maximizes the likelihood over all distributions  $F_2$ . Suppose now that the model consists of discrete  $F_1$ 's and continuous  $F_2$ 's. This model, though smaller than the model with  $F_1, F_2$  being unspecified, has the same semiparametric efficiency bound at a  $(F_1, F_2)$  in this smaller model as the efficiency bound in the original model. This follows from the fact that the class of one-dimensional submodels as needed to compute the tangent space can still be chosen the same. In this smaller model, an  $R = S_1/S_2$  will be discrete at the support points of  $F_1$ , and the shape of  $R$  between the support points equals the shape of  $1/S_2$ . As a consequence, since  $R$  determines the shape of  $F_2$  between the support points, knowing  $R$  in the smaller model helps enormously in estimating  $S_2$ . In particular, for a given  $R_n$ , maximizing the likelihood corresponding with  $p_{R_n, F_2}$  over  $F_2$  with  $S_2R_n$  being a survival function, is very different from maximizing this likelihood over all possible distributions  $F_2$ . This shows that the RNPMLE in the smaller model is inefficient at such  $(F_1, F_2)$ . Since the efficiency bound in the smaller model is the same as the efficiency bound in the original model, this also shows that the RNPMLE will also be inefficient at such  $(F_1, F_2)$ .

PROOF OF THEOREM 2.1. We need to prove that assumption (3) implies that the tangent space at  $P_{F,G}$  equals  $L_0^2(P_{F,G})$ , and is thus saturated. The data generating distribution  $P_{F,G}$  is indexed by  $F$  and  $G$ , where the dependence on  $F$  is only through the marginals  $F_j, j = 1, \dots, k$ . Thus, the tangent space at  $P_{F,G}$  can be obtained as a sum of two tangent spaces, namely the tangent space for  $F$  and the tangent space for  $G$ , where the latter equals  $L_0^2(G)$ . Let  $F, G$  be given and satisfy (3). We now claim that the tangent space for  $F$  is given by the closure of the sum of the  $k$  tangent spaces for  $F_j$  calculated as if the  $F_j$ 's are variation-independent parameters,  $j = 1, \dots, k$ . We will show this now. Let  $h_j \in L_0^2(F_j)$  have finite supremum norm, and let  $F_{j,\varepsilon,h_j}$  be the one-dimensional perturbation  $\varepsilon \rightarrow \int_0^\infty (1 + \varepsilon h_j) dF_j$  through  $F_j$  at  $\varepsilon = 0, j = 1, \dots, k$ . First, note that the support of  $F_{j,\varepsilon,h_j}$  equals the support of  $F_j, j = 1, \dots, k$ . Since  $F_j > F_{j+1}$  (strictly) on  $(0, \tau_j]$  we have that, given an arbitrarily small  $\delta_1 > 0$ , there exists a neighborhood  $\varepsilon \in (-\delta, \delta)$  with  $F_{j,\varepsilon,h_j} \geq F_{j+1,\varepsilon,h_{j+1}}$  on  $(\delta, \tau_j]$  for all  $j = 1, \dots, k - 1$ . Thus,  $P_{F_{j,\varepsilon,h_j}, j=1, \dots, k, G}$  satisfies the constraints  $F_j \geq F_{j+1}, j = 1, \dots, k - 1$ , of our model except on an arbitrarily small neighborhood of 0. Thus, by modifying  $h_j$  on an arbitrarily small neighborhood of 0, we can make  $\varepsilon \rightarrow P_{F_{j,\varepsilon,h_j}, j=1, \dots, k, G}$  a true one-dimensional submodel. Since a tangent space for  $F$  is obtained as the closure in  $L_0^2(F)$  of the linear span of scores of all possible one-dimensional submodels, it follows that the score of the unmodified  $\varepsilon \rightarrow P_{F_{j,\varepsilon,h_j}, j=1, \dots, k, G}$  also belongs to the tangent space. This proves our claim.

Let  $j \in \{1, \dots, k\}$  be given. For a given  $h_j \in L_0^2(F_j)$ , we consider the one-dimensional submodel  $F_{j,\varepsilon}$  given by  $\varepsilon \rightarrow (1 + \varepsilon h_j(t)) dF_j(t)$  which goes through  $F_j$  at  $\varepsilon = 0$ . For notational convenience, define the random variable  $R = N(C) + 1 \in \{1, \dots, k + 1\}$ , and let  $F_{-j}$  be the  $(k - 1)$ -dimensional vector of c.d.f.'s excluding  $F_j$ . This one-dimensional submodel  $F_{j,\varepsilon}$  implies a score for  $P_{F_{j,\varepsilon}, F_{-j}, G}$  given by

$$\begin{aligned}
 A_1(h_1) &= I(R = 1) \frac{\int_c^\infty h_1 dF_1}{S_1(c)} - I(R = 2) \frac{\int_c^\infty h_1 dF_1}{(S_2 - S_1)(c)} \quad \text{if } j = 1, \\
 A_j(h_j) &= I(R = j) \frac{\int_c^\infty h_j dF_j}{(S_j - S_{j-1})(c)} \\
 &\quad - I(R = j + 1) \frac{\int_c^\infty h_j dF_j}{(S_{j+1} - S_j)(c)} \quad \text{if } j \in \{2, \dots, k - 1\}, \\
 A_k(h_k) &= I(R = k) \frac{\int_c^\infty h_k dF_k}{(S_k - S_{k-1})(c)} - I(R = k + 1) \frac{\int_c^\infty h_k dF_k}{F_k(c)} \quad \text{if } j = k.
 \end{aligned}$$

If we define  $S_0 \equiv 0$  and  $S_{k+1} \equiv 1$ , then, for  $j = 1, \dots, k$ ,

$$A_j(h_j) = I(R = j) \frac{\int_c^\infty h_j dF_j}{(S_j - S_{j-1})(c)} - I(R = j + 1) \frac{\int_c^\infty h_j dF_j}{(S_{j+1} - S_j)(c)},$$

where we use that  $S_1 - S_0 = S_1$ , and  $S_{k+1} - S_k = F_k$ . Here  $A_j : L_0^2(F_j) \rightarrow L_0^2(P_{F,G})$  is called the score operator of  $F_j$ ,  $j = 1, \dots, k$ . The tangent space for  $F_j$  is given by the closure of the range of  $A_j$  denoted by  $\overline{R(A_j)}$ . Define  $A_F : L_0^2(F_1) \times \dots \times L_0^2(F_k) \rightarrow L_0^2(P_{F,G})$  by  $A_F(h_1, \dots, h_k) = A_1(h_1) + \dots + A_k(h_k)$ . Then, the tangent space for  $F$  equals  $\overline{R(A_F)}$  so that the tangent space at  $P_{F,G}$  is given by  $\overline{R(A_F) + L_0^2(G)}$ . Thus, to prove the theorem, it suffices to show that  $\overline{R(A_F) + L_0^2(G)} = L_0^2(P_{F,G})$  at any  $F, G$  satisfying (3).

The remaining task is to understand the range of  $A_F$ . We decompose  $A_F$  as a sum of efficient score operators  $A_j^*$ , where  $A_j^*$  is defined as  $A_j$  minus its projection, on the sum-space spanned by the ranges of the other score operators  $A_1, \dots, A_{j-1}, A_{j+1}, \dots, A_k$ ,  $j = 1, \dots, k$ . We will prove that the efficient score operator of  $F_j$  at a  $P_{F,G}$  satisfying (3) equals  $A_j^*(h_j) = E(h_j(T_j) | (C, \Delta_j = I(T_j \leq C)))$ , which is the score operator for the reduced current status data structure  $(C, \Delta_j)$ ,  $j = 1, \dots, k$ . Since the information bounds for smooth functionals of  $F_j$  are, in both models, solely expressed in terms of the efficient score operator for  $F_j$ , the latter result proves that an efficient estimator of  $\mu_j$  based on  $(C, \Delta_j)$ ,  $j = 1, \dots, k$ , like the RNPMLE, is also efficient in the model for the more informative data structure  $(C, N(C))$  [e.g., Bickel, Klaassen, Ritov and Wellner (1993)]. This proves that the RNPMLE actually yields efficient estimators. Subsequently, we show that this special structure of the efficient score operators implies that the tangent space at a  $P_{F,G}$  satisfying (3) is saturated, proving the more general statement of Theorem 2.1.

*Derivation of the efficient score operators of  $F_j$ .* Since  $E(A_l(h_l)A_m(h_m)(Y))$  is equal to 0 if  $|l - m| \geq 2$ , it will follow that the efficient score operators mainly involve projections of the type  $\Pi(A_j | \overline{R(A_{j-1})})$  and  $\Pi(A_j | \overline{R(A_{j+1})})$ . Therefore we first obtain closed form expressions, in general, for these projection operators.

If the projection  $\Pi(A_j(h_j) | \overline{R(A_{j-1})})$  is actually an element of  $R(A_{j-1})$ , then this projection is given by (compare with the formula  $X(X'X)^{-1}X'Y$  for the least squares estimator):

$$(4) \quad \Pi(A_j(h_j) | \overline{R(A_{j-1})}) = A_{j-1}(A_{j-1}^\top A_{j-1})^{-1} A_{j-1}^\top A_j(h_j),$$

where  $A_{j-1}^\top : L_0^2(P_{F,G}) \rightarrow L_0^2(F_{j-1})$  is the adjoint of  $A_{j-1} : L_0^2(F_{j-1}) \rightarrow L_0^2(P_{F,G})$ , and  $(A_{j-1}^\top A_{j-1})^{-1}$  stands for the generalized inverse of  $A_{j-1}^\top A_{j-1} : L_0^2(F_{j-1}) \rightarrow L_0^2(F_{j-1})$ . Similarly,

$$(5) \quad \Pi(A_j(h_j) | \overline{R(A_{j+1})}) = A_{j+1}(A_{j+1}^\top A_{j+1})^{-1} A_{j+1}^\top A_j(h_j).$$

The adjoint  $A_l^\top$  is defined by

$$\langle A_l(h_l), \eta \rangle_{P_F} = \langle h_l, A_l^\top(\eta) \rangle_{F_l} \quad \text{for all } h_l \in L_0^2(F_l) \text{ and } \eta \in L_0^2(P_{F,G}).$$

It is easily shown that for  $l \in \{1, \dots, k\}$ ,

$$A_l^\top(V)(T_l) = \int_0^{T_l} \{V(c, l) - V(c, l + 1)\} dG(c).$$

We have that

$$A_l^\top A_l(h_l)(T_l) = \int_0^{T_l} \phi_l(c) \int_c^\infty h_l dF_l dG(c),$$

where

$$\begin{aligned} \phi_1 &= \frac{S_2}{S_1(S_2 - S_1)}, \\ \phi_l &= \frac{S_{l+1} - S_{l-1}}{(S_{l+1} - S_l)(S_l - S_{l-1})}, \quad l = 2, \dots, k - 1, \\ \phi_k &= \frac{F_{k-1}}{(S_k - S_{k-1})F_k}, \end{aligned}$$

or, in fact, with our convention of  $S_0 = 0$  and  $S_{k+1} = 1$ ,

$$\phi_l = \frac{S_{l+1} - S_{l-1}}{(S_{l+1} - S_l)(S_l - S_{l-1})}, \quad l = 1, \dots, k.$$

Here  $\phi_l(t) \equiv 0$  if  $S_l(t) = 0$ .

If  $p_l = P(T_l = \infty) > 0$ , then we can write

$$\begin{aligned} A_l^\top A_l(h_l)(T_l) &= - \int_0^{\min(T_l, \tau_l)} \phi_l(c) \int_0^c h_l dF_l dG(c) \\ &\quad + I(T_l = \infty)h_l(\infty)p_l \int_{\tau_l}^\infty \phi_l(c) dG(c). \end{aligned}$$

Thus, given a  $K$  with  $K \ll G$ , a solution (if it exists) of  $A_l^\top A_l(h_l) = K$  has to satisfy: for  $G$ -a.e.,  $c \in [0, \tau_l]$ ,

$$(6) \quad \int_0^c h_l dF_l = - \frac{dK}{dG} \frac{1}{\phi_l(c)}, \quad l = 1, \dots, k,$$

and, if  $p_l = P(T_l = \infty) > 0$ , then the equation  $A_l^\top A_l(h_l)(\infty) = K(\infty)$  yields

$$(7) \quad h_l(\infty) = \frac{1}{p_l \int_0^\infty \phi_l(c) dG(c)} \left\{ K(\infty) - \int_0^{\tau_l} \phi_l(c) \int_c^{\tau_l} h_l dF_l dG(c) \right\}.$$

Thus, even when  $p_l > 0$ , (6) is the principal equation to solve (and will imply our conditions) since its solution  $h_l$  on  $[0, \tau_l]$  yields the complete solution  $h_l(T_l) = h_l(T_l)I_{[0, \tau_l]}(T_l) + I(T_l = \infty)h_l(\infty)$ . This two-step method for solving for  $h_l$  in  $A_l^\top A_l(h_l) = K$  first solves for  $h_l I_{[0, \tau_l]}$  and then uses that, if  $p_l > 0$ ,  $h_l(\infty)$  is a function of  $h_l I_{[0, \tau_l]}$ .

We have, for  $l \in \{1, \dots, k - 1\}$ ,

$$\begin{aligned} A_l^\top A_{l+1}(h_{l+1}) &= \int_0^{T_l} [A_{l+1}(h_{l+1})I(R = l) - A_{l+1}(h_{l+1})I(R = l + 1)] dG(c) \\ &= - \int_0^{T_l} A_{l+1}(h_{l+1})I(R = l + 1) dG(c) \\ &= - \int_0^{T_l} \frac{1}{S_{l+1} - S_l} \int_c^\infty h_{l+1} dF_{l+1} dG(c). \end{aligned}$$

We note that this element is indeed absolutely continuous w.r.t.  $G$ . Similarly, it follows that, for  $l \in \{1, \dots, k - 1\}$ ,

$$A_{l+1}^\top A_l(h_l) = - \int_0^{T_{l+1}} \frac{1}{S_{l+1} - S_l} \int_c^\infty h_l dF_l dG(c).$$

Thus,  $h_{j-1,j} \equiv (A_{j-1}^\top A_{j-1})^{-1} A_{j-1}^\top A_j(h_j)$  is the  $h$  satisfying

$$(8) \quad - \int_0^c h dF_{j-1} = - \frac{1}{S_j - S_{j-1}} \int_c^\infty h_j dF_j \frac{1}{\phi_{j-1}} \quad \text{for } G\text{-a.e., } c \in [0, \tau_{j-1}]$$

and, if  $p_{j-1} > 0$ , then  $h(\infty)$  is a simple function of  $hI_{[0, \tau_{j-1}]}$  as given above. Similarly,  $h_{j+1,j} \equiv (A_{j+1}^\top A_{j+1})^{-1} A_{j+1}^\top A_j(h_j)$  is the  $h$  satisfying

$$(9) \quad - \int_0^c h dF_{j+1} = - \frac{1}{S_{j+1} - S_j} \int_c^\infty h_j dF_j \frac{1}{\phi_{j+1}} \quad \text{for } G\text{-a.e., } c \in [0, \tau_{j+1}]$$

and, if  $p_{j+1} > 0$ , then  $h(\infty)$  is a simple function of  $hI_{[0, \tau_{j+1}]}$ . If we can take a derivative of the right-hand sides in (8) and (9) w.r.t.  $F_{j-1}$  and  $F_{j+1}$ , then, in terms of  $h$ , equations (8) and (9) have a solution. This is possible if  $F_j \ll F_l$  (i.e.,  $F_j$  is absolute continuous w.r.t.  $F_l$ ) on  $[0, \tau_l]$ ,  $l \in \{j - 1, j + 1\}$ , which holds under assumption (3) since we assumed that all  $F_j$  have positive Lebesgue density on  $[0, \tau_j]$ . The efficient score operator  $A_j^*$  also involves projections requiring existence of solutions  $h_{l-1,l}, h_{l+1,l}$  for  $l$  different from  $j$ . Therefore, the assumed condition (3) includes (via an easy to understand condition) the necessary and sufficient conditions for the existence of  $h_{l-1,l}, h_{l+1,l}$  for all possible  $l$ , as needed below.

This gives the following closed form expressions for the projections (4) and (5) by simply replacing  $\int_c^\infty h dF_l$  in  $A_l(h)$  by the expressions above. We have, for  $j = 1, \dots, k - 1$ ,

$$\begin{aligned} (10) \quad \Pi(A_j(h_j) | \overline{R(A_{j+1})}) &= A_{j+1}(h_{j+1,j}) \\ &= - \frac{\int_c^\infty h_j dF_j}{(S_{j+1} - S_j)^2 \phi_{j+1}} I(R = j + 1) \\ &\quad + \frac{\int_c^\infty h_j dF_j}{(S_{j+2} - S_{j+1})(S_{j+1} - S_j) \phi_{j+1}} I(R = j + 2) \end{aligned}$$

and, for  $j = 2, \dots, k$ ,

$$\begin{aligned}
 \Pi(A_j(h_j) \mid \overline{R(A_{j-1})}) &= A_{j-1}(h_{j-1,j}) \\
 &= -\frac{\int_c^\infty h_j dF_j}{(S_j - S_{j-1})(S_{j-1} - S_{j-2})\phi_{j-1}} I(R = j - 1) \\
 &\quad + \frac{\int_c^\infty h_j dF_j}{(S_j - S_{j-1})^2\phi_{j-1}} I(R = j).
 \end{aligned}
 \tag{11}$$

For simplicity we derive the efficient score operators for the case  $k = 3$ . (The proof generalizes to the general case.) First, define

$$A_j^l = A_j - \Pi(A_j \mid \overline{R(A_l)}).$$

The efficient score operators  $A_j^* : L_0^2(F_j) \rightarrow L_0^2(P_F)$  are given by

$$\begin{aligned}
 A_3^* &= A_3 - \Pi(A_3 \mid \overline{R(A_1 + A_2)}) = A_3 - \Pi(A_3 \mid \overline{R(A_2^l)}), \\
 A_2^* &= A_2 - \Pi(A_2 \mid \overline{R(A_1 + A_3)}) = A_2 - \Pi(A_2 \mid \overline{R(A_1)}) - \Pi(A_2 \mid \overline{R(A_3)}), \\
 A_1^* &= A_1 - \Pi(A_1 \mid \overline{R(A_2 + A_3)}) = A_1 - \Pi(A_1 \mid \overline{R(A_2^l)}).
 \end{aligned}$$

*Calculation of  $A_2^*$ .* Applying (10) and (11) with  $j = 2$  gives us

$$\begin{aligned}
 \Pi(A_2(h_2) \mid \overline{R(A_1)}) &= -\frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)(S_1 - S_0)\phi_1} I(R = 1) \\
 &\quad + \frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)^2\phi_1} I(R = 2)
 \end{aligned}$$

and

$$\begin{aligned}
 \Pi(A_2(h_2) \mid \overline{R(A_3)}) &= -\frac{\int_c^\infty h_2 dF_2}{(S_3 - S_2)^2\phi_3} I(R = 3) \\
 &\quad + \frac{\int_c^\infty h_2 dF_2}{(S_4 - S_3)(S_3 - S_2)\phi_3} I(R = 4).
 \end{aligned}$$

Thus,

$$\begin{aligned}
 A_2^*(h_2) &= \frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)\phi_1} I(R = 1) \\
 &\quad + \left\{ \frac{1}{S_2 - S_1} - \frac{1}{(S_2 - S_1)^2\phi_1} \right\} \int_c^\infty h_2 dF_2 I(R = 2) \\
 &\quad + \left\{ \frac{1}{(S_3 - S_2)^2\phi_3} - \frac{1}{S_3 - S_2} \right\} \int_c^\infty h_2 dF_2 I(R = 3) \\
 &\quad - \frac{\int_c^\infty h_2 dF_2}{(S_4 - S_3)(S_3 - S_2)\phi_3} I(R = 4).
 \end{aligned}$$

Now, notice that

$$\begin{aligned} (S_2 - S_1)S_1\phi_1 &= S_2, \\ (S_4 - S_3)(S_3 - S_2)\phi_3 &= S_4 - S_2 = F_2, \\ \frac{1}{(S_3 - S_2)^2\phi_3} - \frac{1}{S_3 - S_2} &= -\frac{1}{F_2}, \\ \frac{1}{S_2 - S_1} - \frac{1}{(S_2 - S_1)^2\phi_1} &= \frac{1}{S_2}. \end{aligned}$$

Thus (using  $\int_0^\infty h_2 dF_2 = 0$ ),

$$A_2^*(h_2) = \frac{\int_c^\infty h_2 dF_2}{S_2(c)} I(R \in \{1, 2\}) + \frac{\int_0^c h_2 dF_2}{F_2(c)} I(R \in \{3, 4\}).$$

Calculation of  $A_1^*$ . Formula (10) with  $j = 1$  gives us

$$\begin{aligned} \Pi(A_2(h_2) | \overline{R(A_3)}) &= -\frac{\int_c^\infty h_2 dF_2}{(S_3 - S_2)^2\phi_3(c)} I(R = 3) \\ &\quad + \frac{\int_c^\infty h_2 dF_2}{(S_4 - S_3)(S_3 - S_2)\phi_3(c)} I(R = 4). \end{aligned}$$

Thus,

$$\begin{aligned} A_2^3(h_2) &= A_2(h_2) - \Pi(A_2(h_2) | \overline{R(A_3)}) \\ &= \frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)(c)} I(R = 2) \\ &\quad + \left\{ \frac{1}{(S_3 - S_2)^2\phi_3(c)} - \frac{1}{(S_3 - S_2)(c)} \right\} \int_c^\infty h_2 dF_2 I(R = 3) \\ &\quad - \frac{\int_c^\infty h_2 dF_2}{(S_4 - S_3)(S_3 - S_2)\phi_3(c)} I(R = 4). \end{aligned}$$

We now note that

$$\begin{aligned} (S_4 - S_3)(S_3 - S_2)\phi_3 &= F_2, \\ \frac{1}{(S_3 - S_2)^2\phi_3(c)} - \frac{1}{(S_3 - S_2)(c)} &= -\frac{1}{F_2}. \end{aligned}$$

Thus,

$$A_2^3(h_2) = \frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)(c)} I(R = 2) - \frac{\int_c^\infty h_2 dF_2}{F_2(c)} I(R \in \{3, 4\}).$$

It is easily verified that the adjoint  $A_2^{3\top} : L_0^2(P_F) \rightarrow L_0^2(F_2)$  is given by

$$A_2^{3\top}(V) = \int_0^{T_2} \left\{ V(c, 2) - \frac{(S_3 - S_2)(c)}{F_2(c)} V(c, 3) - \frac{F_3(c)}{F_2(c)} V(c, 4) \right\} dG(c).$$

Subsequently, we can now verify that

$$A_2^{3\top} A_2^3(h_2) = \int_0^{T_2} \phi_2^3(c) \int_c^\infty h_2 dF_2 dG(c),$$

where

$$\phi_2^3 = \frac{F_1}{F_2(S_2 - S_1)}.$$

We need to find  $h_{23,1} \equiv (A_2^{3\top} A_2^3)^{-1}(K)$  with

$$K = A_2^{3\top} A_1(h_1) = - \int_0^{T_2} \frac{\int_c^\infty h_1 dF_1}{(S_2 - S_1)(c)} dG(c).$$

This solution has to satisfy on  $[0, \tau_2]$ :

$$- \int_0^c h_{23,1} dF_2 = \frac{dK}{dG}(c) \frac{1}{\phi_2^3(c)} = - \frac{F_2}{F_1}(c) \int_c^\infty h_1 dF_1$$

and, as shown previously,  $h_{23,1}(\infty)$  is a simple function of  $h_{23,1} I_{[0, \tau_2]}$ . We note that  $h_{23,1}$  exists under the assumption  $F_j \equiv F_k$  (i.e.,  $F_j \ll F_k$  and  $F_k \ll F_j$ ) on  $[0, \tau_j]$ ,  $j = 1, \dots, k - 1$ , which follows from (3). We conclude that

$$\begin{aligned} \Pi(A_1(h_1) | \overline{R(A_2^3)}) &= A_2^3(h_{23,1}) \\ &= - \frac{F_2}{F_1(S_2 - S_1)} \int_c^\infty h_1 dF_1 I(R = 2) \\ &\quad + \frac{\int_c^\infty h_1 dF_1}{F_1} I(R \in \{3, 4\}). \end{aligned}$$

Using  $F_2/(F_1(S_2 - S_1)) - 1/(S_2 - S_1) = -1/F_1$  and  $\int_c^\infty h_1 dF_1 = - \int_0^c h_1 dF_1$  yields

$$\begin{aligned} A_1^*(h_1) &= A_1(h_1) - \Pi(A_1(h_1) | \overline{R(A_2^3)}) \\ &= \frac{\int_c^\infty h_1 dF_1}{S_1(c)} I(R = 1) + \frac{\int_0^c h_1 dF_1}{F_1(c)} I(R \in \{2, 3, 4\}). \end{aligned}$$

*Calculation of  $A_3^*$ .* This calculation is very similar to the one above for  $A_1^*$  and is omitted. We have

$$A_3^*(h_3) = \frac{\int_0^c h_3 dF_3}{F_3(c)} I(R = 4) + \frac{\int_c^\infty h_3 dF_3}{S_3(c)} I(R \in \{1, 2, 3\}).$$



*Proving that the tangent space is saturated.* Given the expressions for the efficient score operators derived above, we now prove that the tangent space at a  $P_{F,G}$  satisfying (3) is saturated. Under our assumption (3), the tangent space equals  $L_0^2(G)$  (scores generated by  $G$ ) plus the closure of the range of  $A^* : L_0^2(F_1) \times \dots \times L_0^2(F_k) \rightarrow L_0^2(P_F)$  defined by

$$(h_1, \dots, h_k) \rightarrow A_1^*(h_1) + \dots + A_k^*(h_k),$$

where the marginal efficient score operators are given by  $A_j^*(h_j) = E(h_j(T_j) | (C, \Delta_j = I(T_j \leq C)))$ ,  $j = 1, \dots, k$ . The closure of the range of a Hilbert space operator equals the orthogonal complement of the null-space of its adjoint, that is,  $\overline{R(A^*)} = N(A^{*\top})^\perp$ . Thus we need to show that  $N(A^{*\top}) = L_0^2(G)$ . The adjoint  $A^{*\top} : L_0^2(P_F) \rightarrow L_0^2(F_1) \times \dots \times L_0^2(F_k)$  is given by

$$A^{*\top}(V) = (A_1^{*\top}(V), \dots, A_k^{*\top}(V)),$$

where it is easily verified that the adjoint  $A_j^{*\top} : L_0^2(P_F) \rightarrow L_0^2(F_j)$  of  $A_j^*$  is given by

$$A_j^{*\top}(V) = E(E(V(C, R) | C, \Delta_j) | T_j).$$

Consider the operator  $B_j^\top : L_0^2(C, \Delta_j) \rightarrow L_0^2(F_j)$  given by  $B_j^\top(\eta) = E(\eta(C, \Delta_j) | T_j)$ , where  $L_0^2(C, \Delta_j)$  is the space of functions of  $(C, \Delta_j)$  with finite variance and zero mean (both taken w.r.t.  $P_{F,G}$ ). Using precisely the same proof as the proof of Lemma 2.2, it follows that, if  $F_j$  has a Lebesgue density  $f_j > 0$  on  $[0, \tau_j]$ , then the null-space  $N(B_j^\top) = L_0^2(G)$ , that is, it consists of functions independent of  $\Delta_j$ . Thus, under (3),  $A_j^{*\top}(V) = 0$  implies that  $E(V(C, R) | C, \Delta_j) = E(V(C, R) | C) \equiv \phi(C)$ ,  $j = 1, \dots, k$ .

Setting  $\Delta_1 = 0$  yields  $\phi(C) = E(V(C, R) | C, \Delta_1 = 0) = V(C, 1)$ . Now, we note that

$$P(R = m | \Delta_j = 1, C = c) = I(m \geq j + 1) \frac{P(R = m | c)}{F_j(c)}, \quad j = 1, \dots, k,$$

where  $P(R = m | c) = (S_m - S_{m-1})(c)$ . Thus,  $E(V(C, R) | C, \Delta_j = 1)$  is given by

$$\sum_{m \geq j+1} V(c, m) \frac{(S_m - S_{m-1})(c)}{F_j(c)} = \phi(c), \quad j = 1, \dots, k.$$

For  $j = k$ , this equality gives  $V(c, k + 1) = \phi(c)$ . For  $j = k - 1$ , this equality gives then

$$V(c, k) \frac{(S_k - S_{k-1})(c)}{F_{k-1}(c)} = \left(1 - \frac{F_k(c)}{F_{k-1}(c)}\right) \phi(c) = \frac{(S_k - S_{k-1})(c)}{F_{k-1}(c)} \phi(c)$$

so that  $V(c, k) = \phi(c)$ . In this manner, we subsequently find  $\phi(c) = V(c, k + 1) = V(c, k) = \dots = V(c, 2)$ . This shows that  $V(C, R)$  does not depend on  $R$ . This completes the proof.  $\square$

**3. Current status data on a counting process when final event is right censored.** The following theorem proves efficiency of any regular asymptotically linear estimator at a specified rich sub-model.

**THEOREM 3.1.** *Let  $N(t)$  be a counting process  $N(t) = \sum_{j=1}^k I(T_j \leq t)$  for random variables  $T_1 < \dots < T_k$ . Let  $C$  be a random censoring time. For every subject we observe the following data structure:*

$$Y = (\tilde{T} = T_k \wedge C, \Delta = I(T_k \leq C), N(\tilde{T})).$$

*We assume that  $C$  is independent of  $(T_1, \dots, T_k)$ . The distribution of  $Y$  only depends on the multivariate distribution  $F$  of  $(T_1, \dots, T_k)$  through the marginal distributions  $F_1, \dots, F_k$  of  $(T_1, \dots, T_k)$ .*

*Consider a data generating distribution  $P_{F,G}$  in the model above satisfying the following condition (12): For certain  $\tau_1 < \dots < \tau_k < \infty$ , let  $F_j$  have Lebesgue density  $f_j$  on  $[0, \tau_j]$  with*

$$(12) \quad \begin{aligned} f_j &> 0 && \text{on } [0, \tau_j] \quad \text{and} \quad f_j = 0 && \text{on } (\tau_j, \infty), \quad j = 1, \dots, k, \\ F_j &> F_{j+1} && \text{on } (0, \tau_j], \quad j = 1, \dots, k - 1, \\ G &&& \text{has Lebesgue density } g. \end{aligned}$$

*We allow that  $p_j \equiv P(T_j = \infty) > 0$  for  $j = j_0, \dots, k$  and  $j_0 \in \{1, \dots, k\}$ .*

*Then, the tangent space at  $P_{F,G}$  equals  $L_0^2(P_{F,G})$  and is thus saturated. This implies that an estimator of a real valued parameter of the distribution  $F$  which is regular and asymptotically linear at  $P_{F,G}$  is also asymptotically efficient if  $P_{F,G}$  satisfies (12). In particular, if  $\bar{G}(t) > 0$  and  $F, G$  satisfy (12), then the Kaplan–Meier estimator  $S_{k,KM}(t)$  of  $S_k(t) = P(T_k > t)$ , based on the i.i.d. data  $(\tilde{T}, \Delta)$ , is asymptotically efficient.*

**3.1. Regular and asymptotically linear estimators.** The important implication of Theorem 3.1 is that, if we can construct an estimator of  $\sqrt{n}$ -estimable parameters of  $F_j$  which is regular, then this estimator will be asymptotically efficient at any  $F$  satisfying (12),  $j = 1, \dots, k$ . In this subsection, we provide relatively simple regular and asymptotically linear estimators.

First, consider estimation of  $S_k(t) = P(T_k > t)$ . It is well known that  $S_{k,KM}(t)$  is a regular asymptotically linear estimator of  $S_k(t)$  whenever  $\bar{G}(t) > 0$ . Second, consider estimation of  $S_j(t) = P(T_j > t)$ ,  $j = 1, \dots, k - 1$ . Let  $\Delta_j \equiv I(T_j \leq C)$ . Under independent censoring (we can weaken this to noninformative censoring of  $T_k$ ), we have

$$(13) \quad E(1 - \Delta_j \mid C = c, T_k > c) = \frac{S_j(c)}{S_k(c)} \equiv R_j(c).$$

So

$$(14) \quad \begin{aligned} S_j(c) &= S_k(c)E(1 - \Delta_j | C = c, T_k > c) \\ &= E(S_k(c)(1 - \Delta_j) | C = c, T_k > c). \end{aligned}$$

In other words, estimating  $S_j$  can be viewed as estimating a monotonic regression of  $S_k(C)(1 - \Delta_j)$  on the observed  $C$ 's. This suggests replacing  $S_k$  by the efficient Kaplan–Meier estimator  $S_{k,KM}$  and minimizing

$$(15) \quad \frac{1}{n} \sum_{i=1}^n w_i \{S_{k,KM}(C_i)(1 - \Delta_{ji}) - S_j(C_i)\}^2 I(C_i \leq T_{ki})$$

over the vector  $(S_j(C_i) : i = 1, \dots, n)$ , under the constraint that  $S_j$  is monotone, where  $w_i, i = 1, \dots, n$ , is a given set of weights possibly assigning more mass to observations with smaller variance. The solution  $S_{j,n}$  of this problem can be obtained with the pool-adjacent-violator-algorithm (PAVA) [see, e.g., Barlow, Bartholomew, Bremner and Brunk (1972)].

A simple calculation shows that

$$(16) \quad \begin{aligned} \text{VAR}\{S_k(C)(1 - \Delta_j) | C = c, T_k > c\} \\ = S_k(c)^2 \text{VAR}\{1 - \Delta_j | C = c, T_k > c\} = S_k^2(c)R_j(c)\{1 - R_j(c)\}. \end{aligned}$$

Since  $R_j$  is not identified from the data at a better rate than  $S_j$ , a good set of weights is  $w_i = 1/S_{k,KM}^2(C_i), i = 1, \dots, n$  [see van der Laan, Jewell and Peterson (1997)].

It is beyond the scope of this paper to prove that smooth functionals of  $S_{j,n}$  are regular and asymptotically linear. Since it is straightforward to prove such a theorem for a standard histogram regression estimator of the regression of  $S_k(C)(1 - \Delta_j)$  on the observed  $C$ 's, one expects that the more sophisticated isotonic regression estimate  $S_{j,n}$  (which only differs because it selects its bins adaptively) is regular and asymptotically linear under the same conditions. We note that the choice of weights  $w_i, i = 1, \dots, n$ , has no effect on the limit distribution of smooth functionals of  $S_{j,n}$ .

3.2. *Proof of Theorem 3.1.* In the first part of the proof we establish that, if condition (12) holds, then the efficient score operator of  $F_k$  equals the efficient score operator of  $F_k$  in the reduced data model for  $(\tilde{T}_k, \Delta_k)$ , hereby establishing a proof of the efficiency of the Kaplan–Meier estimator  $S_{KM}(t)$ . Subsequently, exploiting this special form of the efficient score operator of  $F_k$ , we prove saturation of the tangent space and thus Theorem 3.1.

Consider the data structure  $(\tilde{T}_k = T_k \wedge C, N(\tilde{T}_k))$ , where  $N(t) = \sum_{j=1}^k I(T_j \leq t)$  and  $T_1 < T_2 < \dots < T_k$  are ordered random variables. Let  $R = N(\tilde{T}_k) + 1$ . The density of the data is given by

$$P(d\tilde{T}_k, R = j) = \prod_{m=1}^k (S_m - S_{m-1})(\tilde{T}_k)^{R=m} dF_k(\tilde{T}_k)^{R=k+1} dG(t)^{R \neq k+1} \bar{G}(t)^{R=k},$$

where  $S_0 \equiv 0$  and  $S_{k+1} \equiv 1$ . We refer to the beginning of the proof of Theorem 2.1 to show that the tangent space at a  $P_{F,G}$  satisfying condition (12) is the closure of the sum of the tangent spaces generated by  $F_j$ ,  $j = 1, \dots, k$  and the tangent space of  $G$ , treating  $F_j$  as locally variation-independent. We have that the score operators  $A_j : L_0^2(F_j) \rightarrow L_0^2(P_{F,G})$  for  $F_j$ ,  $j = 1, \dots, k - 1$ , are given by

$$A_j(h_j) = \frac{\int_c^\infty h_j dF_j}{(S_j - S_{j-1})(c)} I(R = j) - \frac{\int_c^\infty h_j dF_j}{(S_{j+1} - S_j)(c)} I(R = j + 1)$$

and

$$A_k(h_k) = h_k(\tilde{T}_k) I(R = k + 1) + \frac{\int_c^\infty h_k dF_k}{(S_k - S_{k-1})(c)} I(R = k).$$

*Derivation of efficient score operator of  $F_k$ .* We first determine the efficient score operator for  $F_k$ . For notational convenience, we consider the case  $k = 3$ . We have

$$A_3^*(h_3) = A_3(h_3) - \Pi\left(A_3(h_3) \mid \overline{R(A_2^1)}\right)$$

where

$$A_2^1 = A_2 - \Pi(A_2 \mid \overline{R(A_1)}).$$

Applying formula (11) gives

$$\Pi(A_2(h_2) \mid \overline{R(A_1)}) = -\frac{\int_c^\infty h_2 dF_2}{S_2(c)} I(R = 1) + \frac{\int_c^\infty h_2 dF_2}{(S_2 - S_1)^2 \phi_1(c)} I(R = 2),$$

where we need to assume that  $F_2 \ll F_1$  on  $[0, \tau_1]$ . Thus, an easy calculation shows that

$$A_2^1(h_2) = \frac{\int_c^\infty h_2 dF_2}{S_2(c)} I(R \in \{1, 2\}) - \frac{\int_c^\infty h_2 dF_2}{(S_3 - S_2)(c)} I(R = 3).$$

Another straightforward calculation shows that the adjoint  $A_2^{1\top} : L_0^2(P_{F,G}) \rightarrow L_0^2(F_2)$  of  $A_2^1 : L_0^2(F_2) \rightarrow L_0^2(P_{F,G})$  is given by

$$A_2^{1\top}(V) = \int_0^{T_2} \left\{ V(c, 1) \frac{S_1}{S_2}(c) + \frac{(S_2 - S_1)}{S_2}(c) V(c, 2) - V(c, 3) \right\} dG(c).$$

A straightforward calculation now shows that

$$A_2^{1\top} A_2^1(h_2) = \int_0^{T_2} \int_c^\infty h_2 dF_2 \frac{S_3}{S_2(S_3 - S_2)}(c) dG(c).$$

We also have

$$A_2^{1\top} A_3(h_3) = - \int_0^{T_2} \frac{\int_c^\infty h_3 dF_3}{(S_3 - S_2)(c)} dG(c).$$

This shows that  $h_{21,3} \equiv (A_2^{1\top} A_2^1)^{-1} A_2^{1\top} A_3(h_3)$  satisfies, on  $[0, \tau_2]$ ,

$$-\int_0^c h_{21,3} dF_2 = -\frac{S_2}{S_3}(c) \int_c^\infty h_3 dF_3,$$

and, if  $p_2 = P(T_2 = \infty) > 0$ , then  $h_{21,3}(\infty)$  is a simple function of  $h_{21,3}I_{[0, \tau_2]}$  as shown above (7). Here we need to assume that this equation can be solved in  $h_{21,3}$ . This is true if  $F_3 \ll F_2$  on  $[0, \tau_2]$ . Then

$$\begin{aligned} \Pi(A_3(h_3) | \overline{R(A_2^1)}) &= A_2^1(h_{21,3}) \\ &= -\frac{\int_c^\infty h_3 dF_3}{S_3(c)} I(R \in \{1, 2\}) \\ &\quad + \frac{S_2(c)}{S_3(S_3 - S_2)(c)} \int_c^\infty h_3 dF_3 I(R = 3). \end{aligned}$$

This proves that

$$\begin{aligned} A_3^*(h_3) &= h_3(\tilde{T}_3)I(R = 4) + \left\{ \frac{1}{S_3 - S_2} - \frac{S_2}{(S_3 - S_2)S_3} \right\}(c)I(R = 3) \\ &\quad + \frac{\int_c^\infty h_3 dF_3}{S_3(c)} I(R \in \{1, 2\}) \\ &= h_3(\tilde{T}_3)I(R = 4) + \frac{\int_c^\infty h_3 dF_3}{S_3(c)} I(R \in \{1, 2, 3\}). \end{aligned}$$

Thus, we have proved that, if  $F_k \equiv F_j$  on  $[0, \tau_j]$ ,  $j = 1, \dots, k - 1$ , then the efficient score  $A_k^*(h_k) = E(h_k(T_k) | \tilde{T}_k, \Delta_k)$ . The latter condition holds, in particular, if (12) holds. This proves the statement of Theorem 3.1 regarding efficiency of the Kaplan–Meier estimator  $S_{KM}$ .

*Saturated tangent space result.* Note that, for a random variable  $Y$ , we define  $L_0^2(Y) = \{h(Y) : Eh^2(Y) < \infty, Eh(Y) = 0\}$ . For simplicity, we prove saturation for  $k = 3$ . Let  $A : L_0^2(F_1) \times L_0^2(F_2) \rightarrow L_0^2(P_{F,G})$  be defined by  $A(h_1, h_2) = A_1(h_1) + A_2(h_2)$ . Then, the tangent space of  $F$  is given by  $\overline{R(A_1) + R(A_2) + R(A_3)} = \overline{R(A_1) + R(A_2)} \oplus \overline{R(A_3^*)}$ . Thus, the tangent space at  $P_{F,G}$  is given by  $\overline{R(A)} \oplus \overline{R(A_3^*)} \oplus \overline{R(B)}$ , where  $B : L_0^2(G) \rightarrow L_0^2(\tilde{T}_3, \Delta_3)$  is the score operator for the censoring mechanism  $G$ , given by  $B(h) = E(h(C) | \tilde{T}_3, \Delta_3)$ . By factorization of the likelihood into  $F$  and  $G$  parts, we have that  $R(B)$  is orthogonal to  $F$ -scores. It is well known that  $\overline{R(A_3^*)} \oplus \overline{R(B)} = L_0^2(\tilde{T}_3, \Delta_3)$ . The latter result simply states that the tangent space for the nonparametric right-censored data model for  $(\tilde{T}_3, \Delta_3)$ , only assuming that  $C$  is independent of  $T$ , is saturated [e.g., Bickel, Klaassen, Ritov and Wellner (1993)]. Thus, we need to prove that  $\overline{R(A)} \oplus L_0^2(\tilde{T}_3, \Delta_3) = L_0^2(P_{F,G})$  which is equivalent to proving

$N(A^\top) = L_0^2(\tilde{T}_3, \Delta_3)$ , where  $A^\top : L_0^2(P_{F,G}) \rightarrow L_0^2(F_1) \times L_0^2(F_2)$  is the adjoint of  $A$  and  $N(A^\top)$  denotes its null space.

First, we decompose  $A_1 + \dots + A_{k-1}$  into a sum of orthogonal operators (efficient score operators in the model with  $F_k$  known). Let  $A'_1 = A_1 - \Pi(A_1 | \overline{R(A_2)})$  and  $A'_2 = A_2 - \Pi(A_2 | \overline{R(A_1)})$ . By (4), it follows that

$$A'_1(h_1) = \frac{\int_c^\infty h_1 dF_1}{S_1(c)} I(R = 1) - \frac{\int_c^\infty h_1 dF_1}{(S_3 - S_1)} I(R \in \{2, 3\}),$$

$$A'_2(h_2) = \frac{\int_c^\infty h_2 dF_2}{S_2(c)} I(R \in \{1, 2\}) + \frac{\int_0^c h_2 dF_2}{(S_3 - S_2)(c)} I(R = 3),$$

where we need the equivalence assumptions  $F_j \equiv F_{j+1}$  on  $[0, \tau_j]$  for  $j = 1, \dots, k$ , again. A more compact manner of representing these operators  $A'_j : L_0^2(F_j) \rightarrow H(C, R) \equiv \{V(C, R)I(R < 4) \in L_0^2(P_{F,G}) : V\}$  is

$$(17) \quad A'_j(h_j) = E(h_j(T_j) | C, \Delta_j, T_3 > C)I(T_3 > C), \quad j = 1, 2.$$

Consider the operator  $A' : L_0^2(F_1) \times L_0^2(F_2) \rightarrow H(C, R)$  defined by  $A'(h_1, h_2) = A'_1(h_1) + A'_2(h_2)$ . Proving  $N(A^\top) = L_0^2(\tilde{T}_3, \Delta_3)$  is equivalent to proving  $N(A'^\top) = L_0^2(\tilde{T}_3, \Delta_3)$ , where  $A'^\top$  is the adjoint of  $A'$ .

From the representation (17), the adjoint  $A_j'^\top : H(C, R) \rightarrow L_0^2(F_j)$  is given by

$$A_j'^\top(V) = E(E(V(C, R)I(T_3 > C) | C, \Delta_j, T_3 > C) | T_j), \quad j = 1, 2,$$

and thus,  $N(A'^\top) = N(A_1'^\top) \cap N(A_2'^\top)$ .

Consider now a solution  $V I(T_3 > C) \in H(C, R)$  satisfying  $A_j'^\top(V I(T_3 > C)) = 0$ ,  $j = 1, 2$ . In order to prove  $V \in L_0^2(\tilde{T}_3, \Delta_3)$ , it suffices to show  $I(T_3 > C)V = I(T_3 > C)\phi(C)$  for some  $\phi$ . Using precisely the same proof as the proof of Lemma 2.2, it follows that, if  $F_j$  has a Lebesgue density  $f_j > 0$  on  $[0, \tau_j]$  and  $G$  has a Lebesgue density, then, for any function  $I(T_3 > C)\eta(C, \Delta_j)$ ,  $E(I(T_3 > C)\eta(C, \Delta_j) | T_j) = 0$  implies  $\eta(C, 1) = \eta(C, 0)$ . This proves that  $E(V(C, R)I(T_3 > C) | C, \Delta_j, T_3 > C) = E(V(C, R)I(T_3 > C) | C, T_3 > C) \equiv I(T_3 > C)\phi(C)$  does not depend on  $\Delta_j$ ,  $j = 1, 2$ .

Setting  $\Delta_1 = 0$  yields  $I(T_3 > C)\phi(C) = E(V(C, R)I(T_3 > C) | C, \Delta_j, T_3 > C) = V(C, 1)I(T_3 > C)$ . Now, we note that

$$P(R = m | \Delta_j = 1, C = c, T_3 > c)$$

$$= I(m \geq j + 1, m < 4) \frac{(S_m - S_{m-1})(c)}{(S_3 - S_j)(c)}, \quad j = 1, 2.$$

Thus,  $E(V(C, R)I(T_3 > C) | C, \Delta_j = 1, T_3 > C)$  is given by

$$I(T_3 > C) \sum_{m \geq j+1, m < 4} V(C, m) \frac{(S_m - S_{m-1})(C)}{(S_3 - S_j)(C)} = I(T_3 > C)\phi(C), \quad j = 1, 2.$$

For  $j = 2$ , this equality gives  $I(T_3 > C)V(C, 3) = I(T_3 > C)\phi(C)$ . For  $j = 1$ , this equality gives

$$I(T_3 > C) \left\{ V(C, 2) \frac{(S_2 - S_1)(c)}{(S_3 - S_1)(C)} + \phi(C) \frac{(S_3 - S_2)(C)}{(S_3 - S_1)(C)} \right\} = I(T_3 > C)\Phi(C),$$

so that  $I(T_3 > C)V(C, 2) = I(T_3 > C)\phi(C)$ . We have shown  $I(T_3 > C) \times V(C, 1) = I(T_3 > C)V(C, 2) = I(T_3 > C)V(C, 3)$  which proves that  $V = I(T_3 < C)V_1(T_3) + I(T_3 > C)\phi(C)$  for some functions  $V_1$  and  $\phi$ , and thus that  $V \in L_0^2(\tilde{T}_3, \Delta_3)$ . This completes the proof.  $\square$

**Acknowledgments.** The authors thank the referees and Associate Editor for their helpful comments.

## REFERENCES

- BARLOW, R. E., BARTHOLOMEW, D. J., BREMNER, J. M. and BRUNK, H. D. (1972). *Statistical Inference under Order Restrictions*. Wiley, New York.
- BICKEL, P. J., KLAASSEN, C. A. J., RITOV, Y. and WELLNER, J. A. (1993). *Efficient and Adaptive Estimation in Semi-Parametric Models*. Johns Hopkins Univ. Press.
- DIAMOND, I. D. and McDONALD, J. W. (1992). The analysis of current status data. In *Demographic Applications of Event History Analysis* (J. Trussell, R. Hankinson and J. Tilton, eds.) 231–252. Oxford Univ. Press.
- DIAMOND, I. D., McDONALD, J. W. and SHAH, I. H. (1986). Proportional hazards models for current status data: Application to the study of differentials in age at weaning in Pakistan. *Demography* **23** 607–620.
- DINSE, G. E. and LAGAKOS, S. W. (1982). Nonparametric estimation of lifetime and disease onset distributions from incomplete observations. *Biometrics* **38** 921–932.
- GILL, R. D., VAN DER LAAN, M. J. and ROBINS, J. M. (1997). Coarsening at random: Characterizations, conjectures and counterexamples. *Proc. First Seattle Symposium in Biostatistics. Lecture Notes in Statist.* **123** 255–294. Springer, New York.
- GROENEBOOM, P. J. (1998). Special topics course 593C: Nonparametric estimation for inverse problems: algorithms and asymptotics. Technical Report 344, Dept. Statistics, Univ. Washington. (For related software see [www.stat.washington.edu/jaw/RESEARCH/SOFTWARE/software.list.html](http://www.stat.washington.edu/jaw/RESEARCH/SOFTWARE/software.list.html).)
- GROENEBOOM, P. and WELLNER, J. A. (1992). *Information Bounds and Nonparametric Maximum Likelihood Estimation*. Birkhäuser, Basel.
- HUANG, J. and WELLNER, J. A. (1995). Asymptotic normality of the NPML of linear functionals for interval censored data, case I. *Statist. Neerlandica* **49** 153–163.
- JEWELL, N. P., MALANI, H. M. and VITTINGHOFF, E. (1994). Nonparametric estimation for a form of doubly censored data with application to two problems in AIDS. *J. Amer. Statist. Assoc.* **89** 7–18.
- JEWELL, N. P. and SHIBOSKI, S. C. (1990). Statistical analysis of HIV infectivity based on partner studies. *Biometrics* **46** 1133–1150.
- JEWELL, N. P. and VAN DER LAAN, M. J. (1995). Generalizations of current status data with applications. *Lifetime Data Analysis* **1** 101–109.
- JONGBLOED, G. (1995). Three statistical inverse problems. Ph.D. dissertation, Delft Univ. Technology.
- KEIDING, N. (1991). Age-specific incidence and prevalence: A statistical perspective (with discussion). *J. Roy. Statist. Soc. Ser. A* **154** 371–412.

- KODELL, R. L., SHAW, G. W. and JOHNSON, A. M. (1982). Nonparametric joint estimators for disease resistance and survival functions in survival/sacrifice experiments. *Biometrics* **38** 43–58.
- SUN, J. and KALBFLEISCH, J. D. (1993). The analysis of current status data on point processes. *J. Amer. Statist. Assoc.* **88** 1449–1454.
- TURNBULL, B. W. and MITCHELL, T. J. (1984). Nonparametric estimation of the distribution of time to onset for specific diseases in survival/sacrifice experiments. *Biometrics* **40** 41–50.
- VAN DER LAAN, M. J., JEWELL, N. P. and PETERSON, D. R. (1997). Efficient estimation of the lifetime and disease onset distribution. *Biometrika* **84** 539–554.

DIVISION OF BIostatISTICS  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720  
E-MAIL: laan@stat.berkeley.edu