

APPROXIMATE LIKELIHOOD AND PROBABILITY CALCULATIONS BASED ON TRANSFORMS¹

BY ROLLIN BRANT

University of Minnesota

Probability distributions, especially in applications, are generally specified via density functions; alternative representations, including the characteristic function, moment generating function, and sequence of moments, are most commonly encountered in theoretical settings. These alternative means of specification do, however, give rise to the construction of certain approximations that can facilitate the implementation of likelihood methods or the calculation of probabilities, even when the density is not available in closed form. The feasibility of a unified treatment of this topic stems from a number of properties shared by probability transforms in general.

1. Introduction. In applied settings, probability distributions are almost always specified by density or distribution functions. Occasionally, however, distributions arise which are only conveniently represented by transforms, such as the characteristic function or moment generating function. The absence of closed form expressions for the relevant densities complicates the implementation of standard methods of inference in even the simple case considered here, where we assume replicated observations on x , a real valued random variable with distribution P_θ , specified up to some unknown, real valued parameter θ . Likelihood methods, for example, require the evaluation of the density ($p(x; \theta)$) based forms, $\ell(x; \theta) = \log p(x; \theta)$, $S(x; \theta) = (\partial \log p(x; \theta))/\partial \theta$, and $\mathcal{I}(\theta) = \text{var}\{S(x; \theta)\}$. In this paper we shall explore certain transform-based approximations to these forms which facilitate likelihood estimation, and the associated asymptotic methods, when $p(x; \theta)$ is unavailable or intractable. Also, we shall consider the use of the exponential transformation of the approximation to $\ell(x; \theta)$ as an alternative to more familiar density approximations, such as the Edgeworth and Gram-Charlier expansions.

Approaches taken in earlier investigations have relied on either numerical inversion of the transform representations (Dumouchel, 1975; Feuerverger and McDunnough, 1981c) or on nonstandard estimation schemes based on the empirical version of the relevant transform (Brockwell and Brown, 1982; Bryant and Paulson, 1979; Chambers and Heathcote, 1981). The first approach leads to considerable computational complexity, while the second produces estimates with suboptimal properties. Feuerverger and McDunnough (1981a, 1981b, 1982) introduced methods which are statistically and computationally efficient. The methods

Received September 1982; revised March 1984.

¹This work originated while the author was attending the University of Toronto, under support from the Natural Sciences and Engineering Research Council of Canada.

AMS 1980 subject classifications. Primary 62A10, 60E10; secondary 65U05.

Key words and phrases. Least squares approximations, transform-based methods, likelihood, Fisher information, density expansions.

proposed here extend their work and establish links with more familiar modes of inference. A unified approach to the problem is possible based on certain general properties of transforms, outlined below.

2. Probability transforms. A probability transform, $\gamma(t)$, corresponding to a real valued random variable, x , is defined through an underlying class of real valued functions, $G = \{g(t, x), t \in T\}$, the kernel class, by $\gamma(t) = E\{g(t, x)\}$. The important transforms include the characteristic function (c.f.), the moment generating function (m.g.f.), and the sequence of moments, with kernel classes $G_H = \{\sin tx, \cos tx, t \in R^1\}$, $G_E = \{\exp tx, t \in R^1\}$, and $G_M = \{x^t, t = 0, 1, 2 \dots\}$ respectively, though others (e.g. the fractional moments) exist. Note that some transforms do not always completely characterize the underlying distribution, P , though we shall generally restrict ourselves to situations when this is the case. The common property of these kernel classes that facilitates the subsequent investigation is their closure under multiplication, which implies that γ yields the joint moment structure of G , in an explicit manner, so that the following approach to the construction of approximations is often available.

Suppose $h(x)$ is a function of interest which is not conveniently obtainable in closed form. The best linear approximation (predictor) of $h(x)$ in terms of the set of kernel functions indexed by $\mathbf{t} = (t_1, \dots, t_k)^T$, is defined to be the form $h_*(x) = \sum_{i=1}^k d_i g(t_i, x)$ minimizing the mean square error $\int_{R^1} [h_*(x) - h(x)]^2 dP$. For convenience, we shall define the vector-valued function, $\mathbf{g}(x)$, given by $\mathbf{g}(x) = g(\mathbf{t}, x) = (g(t_1, x), \dots, g(t_k, x))^T$, so that h_* may be written as $\mathbf{d}^T \mathbf{g}(x)$, where $\mathbf{d} = (d_1, \dots, d_k)^T$. Thus h_* is the projection (regression) of h on $\text{span}\{\mathbf{g}(x)\}$ in the Hilbert space, $L_2(dP)$, induced by the measure P . The coefficients of h_* are easily seen to be given by $\mathbf{d}_* = \Gamma^{-1} \boldsymbol{\eta}$, where $\Gamma = E\{\mathbf{g}(x) \cdot \mathbf{g}(x)^T\}$ and $\boldsymbol{\eta} = E\{h(x) \cdot \mathbf{g}(x)\}$. We shall adopt the notation, $h_*(x) = E^*\{h(x) | \mathbf{g}(x)\}$, motivated by the formal similarity to conditional expectation.

It is customary to include a constant term in the approximation, and this is most conveniently accomplished by centering the variables, leading to the form

$$h_*(x) = E^*\{h(x) | 1, \mathbf{g}(x)\} = E\{h(x)\} + \lambda^T \Sigma^{-1}[\mathbf{g}(x) - \boldsymbol{\gamma}],$$

where $\lambda = \text{cov}\{\mathbf{g}(x), h(x)\}$, $\Sigma = \text{var}\{\mathbf{g}(x)\}$, and $\boldsymbol{\gamma} = E\{\mathbf{g}(x)\} = \boldsymbol{\gamma}(\mathbf{t})$. By the foregoing remarks, $\boldsymbol{\gamma}$, Γ , and Σ are given explicitly by the transform $\gamma(t)$. We shall encounter a number of instances where the remaining coefficients in h_* are easily determined, facilitating the approximation of $h(x)$ and related functions (derivatives, antiderivatives). Note that the approximation, $h_*(x)$, can be made arbitrarily close to $h(x)$, in the sense considered, when $h(x)$ is contained in the L_2 closure of $\text{span}\{G\}$. In many cases G is complete, so that any $h(x) \in L_2(dP)$ may be so approximated.

3. The score function and Fisher's information. As previously mentioned, one obvious approach to the approximation of the score function, $S(x, \theta)$, or the information, $\mathcal{I}(\theta)$, is to proceed by numerically approximating $p(x; \theta)$, based on $\gamma(t; \theta)$, the corresponding transform. However, the following approach is more direct.

Observe that, under mild regularity conditions,

$$E_{\theta}\{S(x; \theta) \cdot g(x)\} = \partial E_{\theta}\{g(x)\}/\partial\theta$$

for any function, $g(x)$. Thus, $\lambda = \text{cov}_{\theta}\{S(x; \theta), \mathbf{g}(x)\} = \dot{\gamma}(\theta)$, where $\gamma(\theta) = E_{\theta}\{\mathbf{g}(x)\}$ and $\dot{\cdot}$ denotes differentiation (componentwise) with respect to θ . Since $E_{\theta}\{S(x; \theta)\} = 0$, we have that, for any selection of $\mathbf{g}(x)$ from G ,

$$S_{*}(x; \theta) = E^{*}\{S(x; \theta) \mid 1, \mathbf{g}(x)\} = \dot{\gamma}(\theta)^T \Sigma(\theta)^{-1}[\mathbf{g}(x) - \gamma(\theta)].$$

Since the expression is linear in $\mathbf{g}(x)$, the resulting approximation to the maximum likelihood equations based on a sample of size n , can be written as

$$(3.1) \quad \dot{\gamma}(\theta) \Sigma(\theta)^{-1}[\hat{\gamma} - \gamma(\theta)] = 0,$$

where $\hat{\gamma}$ is a vector of observations from $\hat{\gamma}(t) = n^{-1} \sum_{i=1}^n g(t, x_i)$, the empirical transform; cf. Feuerverger and McDunnough (1981a, b, and 1982). Later examples show that the dimension of \mathbf{g} need not be inordinately big for S_{*} to provide a useful approximation to S .

Observe, as well, that by the orthogonalities implied by the projection operations,

$$\text{var}_{\theta}\{S(x; \theta)\} = \text{var}_{\theta}\{S_{*}(x; \theta)\} + \text{var}_{\theta}\{S(x; \theta) - S_{*}(x; \theta)\}.$$

Since the latter term can be made arbitrarily small (given the conditions mentioned previously), $\mathcal{I}_{*}(\theta) = \dot{\gamma}(\theta) \Sigma(\theta)^{-1} \dot{\gamma}(\theta) = \text{var}_{\theta}\{S_{*}(x; \theta)\}$ can be regarded as a simply derived approximation to the Fisher information, $\mathcal{I}(\theta) = \text{var}_{\theta}\{S(x; \theta)\}$; see Jarrett (1973) and Brockwell and Brown (1982). In addition, standard arguments yield that the asymptotic variance of a consistent solution to (3.1) is $[n \mathcal{I}_{*}(\theta)]^{-1}$, which, owing to the relationship of $\mathcal{I}_{*}(\theta)$ and $\mathcal{I}(\theta)$, can be made arbitrarily close to the Cramer-Rao lower bound.

3.1. *Location and scale.* When $x \sim \theta + z$, z possessing a continuous distribution P , with density with respect to Lebesgue measure $p(z)$, the relevant score function has the form $S(x; \theta) = J(x - \theta)$, where $J(z) = -\partial \log p(z)/\partial z = -p'(z)/p(z)$. If the distribution of z is specified via $\gamma(t) = E\{g(t, z)\}$, then it is appropriate to consider $J_{*}(z) = E^{*}\{J(z) \mid 1, \mathbf{g}(z)\}$. In general, $E\{J(z)\} = 0$, and

$$(3.2) \quad \text{cov}\{J(z), g(z)\} = - \int g(z)p'(z) dz = -p(z)g(z) \Big|_{-\infty}^{\infty} + \int p(z)g'(z) dz,$$

assuming differentiability of the kernel function $g(z)$. The important kernel classes consist of differentiable functions, and are such that $\text{span}\{G\}$ is closed under this operation. Also, $\mathbf{g}(x)$ will typically be chosen so that $\text{span}\{\mathbf{g}(x)\}$ possesses the same closure. Since the first term in (3.2) will disappear in most cases, it follows that, in general, one may express $\lambda = \text{cov}\{\mathbf{g}(z), J(z)\}$ as $D\gamma$, where D is a matrix of constants and $\gamma = E\{\mathbf{g}(z)\}$. Thus it is possible to form $J_{*}(z)$, and take as an approximation to $S(x; \theta)$, $J_{*}(x - \theta) = [D\gamma]^T \Sigma^{-1}[\mathbf{g}(x - \theta) - \gamma]$, where $\Sigma = \text{var}\{\mathbf{g}(z)\}$. This form will in fact coincide with that derived from the more general approach in most cases, since it is generally the case that $\mathbf{g}(x - \theta) = M(\theta)\mathbf{g}(x)$, $M(\theta)$ a $k \times k$ matrix, i.e. $\text{span}\{\mathbf{g}(z)\}$ is invariant under translations.

This “pivotal” form is useful in that the translation invariance of procedures derived subsequently is made explicit (D , γ , and Σ being fixed), and since it also provides an approximation to $J(x; \theta) = \partial p(x; \theta)/\partial x$, a function that is useful in the more general context. Of course, $I_* = \gamma^T D^T \Sigma^{-1} D \gamma$ yields an approximation to the (constant) Fisher information.

When observations are of the form $x = \theta z$, z specified as before, the relevant score function has the form $S(x; \theta) = \theta^{-1} K(x/\theta)$, where $K(z) = z \cdot J(z) - 1$. Again $K_*(z) = E^*\{K(z) \mid 1, g(z)\}$ is generally available. Note that

$$K(z) = - \frac{zp'(z) + p(z)}{p(z)},$$

so that

$$\begin{aligned} \text{cov}\{g(z), K(z)\} &= - \int [zp'(z) + p(z)]g(z) dz \\ &= -zp(z)g(z) \Big|_{-\infty}^{\infty} + \int zg'(z)p(z) dz. \end{aligned}$$

Again the first term disappears in general, and the reader may check that the latter integral may be given explicitly in terms of γ in all the interesting cases. For example if $g(z) = \exp tz$,

$$\int zg'(z)p(z) dz = t \int z \exp(tz) p(z) dz = t \int \frac{\partial[\exp(tz) p(z)]}{\partial t} dz = t \cdot \gamma'(t),$$

assuming regularity. Thus $\lambda = \text{cov}\{g(z), K(z)\}$ is generally obtainable, so that $\theta^{-1}K_*(x/\theta) = \theta^{-1}\lambda^T \Sigma^{-1}[\mathbf{g}(x/\theta) - \gamma]$ provides a suitably invariant approximation to $S(x; \theta)$, while the quantity $I_*(\theta) = \theta^{-2}\lambda^T \Sigma^{-1} \lambda$ provides a corresponding approximation to the information. In general, the algebraic form of K_* will closely follow that of S_* ; K_* and S_* represent the projections of $S(x; \theta)$ on $\mathbf{g}(x/\theta)$ and $\mathbf{g}(x)$, respectively.

4. Choice of the kernel vector, \mathbf{g} . The convergence of the approximations considered thus far depend on the completeness properties of the kernel set G in the space $L_2(dP)$, and on the particular choice of \mathbf{g} from G . We therefore consider G_M , G_H and G_E individually.

In the case of $G_M = \{x^t, t = 1, 2, \dots\}$, uniqueness of the moments is sufficient to ensure completeness. The choice $\mathbf{g}(x) = \langle x, x^2, \dots, x^k \rangle^T$ is natural, though to achieve an adequate degree of convergence a fairly large k may be required. With regard to stability, it is the numerical difficulty associated with manipulating Σ , which involves theoretical moments up to order $2k$, rather than the stochastic instability of higher order sample moments, that is the most serious issue.

The completeness of $G_H = \{\cos tx, \sin tx, t \in R\}$ holds under completely arbitrary assumptions. A convenient choice for $\mathbf{g}(x)$ is to take $\mathbf{g}(x) = (\cos \tau x, \dots, \cos k\tau x, \sin \tau x, \dots, \sin k\tau x)^T$, in which case approximations reduce to trigonometric polynomials. Since $2\pi/\tau$ -periodic functions can be so approximated, if the domain of interest of the “target” function lies in an interval $M \pm$

L , the choice $\tau = 2\pi/L$ is indicated. The approximations considered will then depend on the characteristic function, $\phi(t) = E\{\cos tx\} + iE\{\sin tx\}$, at the points, $t = j\tau, j = 1, \dots, k$. This approach is equivalent to replacing P by a “wrapped” version, P_τ , the distribution induced by identifying points on R^1 on $2\pi/\tau$ spaced grids. The corresponding density $p_\tau(x)$ can be written as $p_\tau(x) = \sum_{j=-\infty}^{\infty} p(x + 2\pi j/\tau)$, and $\phi(j\tau)$ can be seen to yield the j th Fourier coefficient of $p_\tau(x)$,

$$a_j = \left[\frac{2\pi}{\tau} \right]^{-1} \int_{-\pi/\tau}^{\pi/\tau} \exp(ij\tau x) p_\tau(x) dx.$$

If τ is held fixed and $k \rightarrow \infty$, the forms h_* considered thus far will converge to wrapped versions h_τ . For instance $S_*(x; \theta)$ will tend to $S_\tau(x; \theta) = \partial \log p_\tau(x; \theta) / \partial \theta$, which can be viewed as the projection of $S(x; \theta)$ on $\{\exp(ij\tau x), j = -\infty, \dots, \infty\}$. Thus, in general, if τ can be chosen so that p_τ represents a reasonable approximation to P , so that h_τ approximates h , it is practicable to fix τ , and then choose k to obtain a reasonable degree of convergence. In the completely general parametric case, the range of interest of $S(x; \theta)$ may depend on θ , as in the case of a scale parameter. It is usually desirable to rescale the random variable x in accordance with θ , as is accomplished through the use of K_* in the scale case. Note that, due to the boundedness of the trigonometric functions, ill-conditioning problems associated with large k values are less prevalent than in the case of the moments.

It must be noted that the general strategy outline for choosing $g(x)$ can in no way be considered “optimal.” In individual instances, it is not, in fact, the completeness of G that is of the essence, but how the function of interest $h(x)$ may be best approximated by elements of G . For example, in the Gaussian location problem, arbitrarily good approximations to the score $S(x; \theta) = x - \theta$ are obtained by considering the pair $g(x) = (\cos \tau x, \sin \tau x)^T$, as $\tau \rightarrow 0$, since

$$x - \theta \approx \frac{\sin(\tau[x - \theta])}{\tau} = \tau^{-1}[\cos\tau\theta \sin\tau x - \sin\tau\theta \cos\tau x],$$

for τ small. In the case where the techniques considered are of interest, however, the form of $h(x)$ will be largely inaccessible, so that the general strategy entertained will be the only one feasible.

With regard to other kernel sets, such as G_E , yielding the m.g.f., less is known; numerical issues similar to those affecting G_M indicate that G_E is not a practical choice. Overall, the author’s experience has been that G_H leads to forms which are the most numerically reliable.

5. Examples

5.1. *Stable laws.* We consider aspects of inference regarding the family of symmetric stable laws, specified by the c.f.’s, $\phi(t; \alpha) = \exp\{-(|t|^\alpha/2)\}$, $1 \leq \alpha \leq 2$. Only two members of this family have closed form density representations, the Cauchy ($\alpha = 1$) and the Gaussian ($\alpha = 2$). These cases provide convenient

reference points, since the exact forms can be computed explicitly and compared with the approximations. This is done for J_* and K_* with k chosen to illustrate the rate of convergence. In accordance with the approach in Section 4, the domain of approximation in the Cauchy case (note nonstandard scale) is taken to be $-3\pi \leq x \leq 3\pi$, leading to the use of $\tau = 1/3$; in the Gaussian case we consider $-\pi < x < \pi$, with $\tau = 1$. (See Figures 1a - 1d.)

In addition, the forms, J_* , are illustrated for intermediate members of the stable family in Figure 2, based on the conservative choice of $k = 15$ and $\tau = .3$. The relevant efficiencies, $\mathcal{J}^*/\mathcal{J}$, were calculated, using the essentially exact results of Dumouchel (1975), and are given in Table 1.




FIG. 1(a). Gaussian model: J and J_* ($\tau = 1$; $k = 2, 4, 5$).




FIG. 1(b). Cauchy model: J and J_* ($\tau = 1/3$; $k = 5, 10, 15$).




FIG. 1(c). Gaussian model: K and K_* ($\tau = 1$; $k = 2, 4, 6$).




FIG. 1(d). Cauchy model: K and K_* ($\tau = 1/3$; $k = 5, 10, 15$).

5.2. *Binomial-Poisson convolution.* If $x = y + z$ where y is binomial, with parameters N and θ , and z is Poisson, with mean λ , independent of y , then the probability distribution function of x is $p(x; \theta, \lambda) = e^{-\lambda} \sum_y \binom{N}{y} \theta^y (1 - \theta)^{N-y} \cdot (\lambda^{x-y} / (x - y)!)$, whereas its c.f. has the simpler form

$$\phi(x; \theta, \lambda) = [(1 - \theta) + \theta \exp(it)]^n \exp\{\lambda[\exp(it) - 1]\}.$$

The problem of estimation for this family has been considered by Sclove and Van Ryzin (1969), who note that maximum likelihood is “computationally intractable.” Sprott (1983), however, points out certain recursive relationships




FIG. 2. Stable models ($\alpha = 1.1(.2)1.9$) J_* ($\tau = .3, k = 15$).

TABLE 1
Asymptotic efficiencies ($\mathcal{J}^*/\mathcal{J}$) for stable family location estimates

Stable exponent (α)	Order of trigonometric approximation (k)		
	5	10	15
1.1	.86225	.97722	.99158
1.3	.96261	.99465	.99514
1.5	.99426	.99717	.99718
1.7	.99889	.99929	.99929
1.9	.99863	.99888	.99889

in the derivatives of $p(x; \theta, \lambda)$ that facilitate essentially exact computations of $\mathcal{J}_\theta, \mathcal{J}_\lambda,$ and $\mathcal{J}_{\theta\lambda}$, the components of the joint information matrix for θ and λ . Alternatively, the straightforward multiparameter extensions of \mathcal{J}_* can be constructed to yield approximations. Some numerical results are given in Table 2 for various choices of $N, \theta,$ and $\lambda,$ with $\tau = 2\pi/(N + \lambda + 5\lambda^{1/2})$, based on a conservative bound on the domain of interest for x .

6. Fourier likelihood methods. We have seen that when distributions of a parametric family are specified by the characteristic function, $\phi(t; \theta) = E_\theta\{\cos tx\} + iE_\theta\{\sin tx\}$, the quantities $S(x; \theta)$ and $\mathcal{J}(\theta)$ may be approximated. In this section we see that $\ell(x; \theta) = \log p(x; \theta)$ may be similarly approximated and that, furthermore, the sense of approximation of the above forms may be strengthened, given regularity conditions. We consider first the approximation of $S(x; \theta)$.

6.1. *The score function.* If $S_*(x; \theta)$ is to provide a reliable basis for likelihood

TABLE 2
Information calculations for the binomial Poisson convolution model

Parameters			Exact (\mathcal{I}) and approximate (\mathcal{I}^*) calculations					
N	θ	λ	\mathcal{I}_θ^*	\mathcal{I}_θ	\mathcal{I}_λ^*	\mathcal{I}_λ	$\mathcal{I}_{\theta\lambda}^*$	$\mathcal{I}_{\theta\lambda}$
20	.50	10	26.64667	27.06326	.06581	.06691	1.30205	1.34342
20	.50	50	7.31844	7.32785	.01816	.01818	.36290	.36336
20	.90	10	37.28053	37.51228	.08471	.08504	1.65376	1.66239
20	.90	50	7.94860	7.95635	.01929	.01931	.38534	.38568
100	.50	10	286.04950	286.04951	.02878	.02878	2.84876	2.84876
100	.50	50	133.72869	133.72869	.01334	.01334	1.33136	1.33136
100	.90	10	538.53335	538.53335	.05362	.05362	5.15320	5.15320
100	.90	50	172.88912	172.88912	.01696	.01696	1.68880	1.68880

calculations, it is desirable to show some form of uniform convergence. This is demonstrated below; similar arguments apply to the other forms considered.

THEOREM 6.1. Let $P_\theta, \theta \in \Theta$, be a set of distributions on R^1 with associated densities $p(x; \theta)$ satisfying:

- $p(x; \theta)$ is bounded above and away from 0, on any compact subset of $R^1 \times \Theta$.
- $p(x; \theta)$, $\dot{p}(x; \theta)$, and $\dot{p}'(x; \theta) = \partial^2 p(x; \theta) / \partial x \partial \theta$ exist, are continuous in x and θ , and for compact subsets of Θ satisfy conditions of the form

$$h(x; \theta) \leq A |x|^{-\alpha} \text{ for } |x| > B$$

uniformly in θ , with $\alpha > 1$ and $A, B < \infty$.

Let $R_c \subseteq R^1 \times \Theta$ be compact. Then, for all $\epsilon > 0$ there exist k and τ such that for $\mathbf{t} = \langle \tau, 2\tau, \dots, k\tau \rangle$,

$$S_*(x; \theta) = E^*\{S(x; \theta) | 1, \sin \mathbf{t}x, \cos \mathbf{t}x\}$$

satisfies

$$|S_*(x; \theta) - S(x; \theta)| < \epsilon \text{ for all } (x, \theta) \in R_c.$$

PROOF. The approach to choosing τ and k described in Section 4 is reflected in the outline of the proof below. We consider first the "wrapped" version of $S(x; \theta)$, $S_\tau(x; \theta)$, choosing τ small enough that $|S_\tau(x; \theta) - S(x; \theta)| < \epsilon/2, (x, \theta) \in R_c$. To do this we require the following lemma, 6.2, and its corollaries, 6.3 and 6.4

LEMMA 6.2 Let $h(x)$ be a continuous, real valued function on R^1 , and suppose that there exist $A, B < \infty$ and $\alpha > 1$ such that

$$(6.1) \quad |h(x)| \leq A |x|^{-\alpha} \text{ for } |x| > B.$$

Define the $2\pi/\tau$ periodic function

$$h_\tau(x) = \sum_{j=-\infty}^{\infty} h\left(x + \frac{2\pi j}{\tau}\right).$$

Then

1. for $\tau < \pi/B$, $h_\tau(x)$ is a well defined, continuous function on $I_\tau = [-\pi/\tau, \pi/\tau]$;
2. $h_\tau(x)$ converges almost uniformly to $h(x)$ as $\tau \rightarrow 0$, in the sense that $\lim_{\tau \rightarrow 0} [\sup_{x \in I_\tau} |h_\tau(x) - h(x)|] \rightarrow 0$.

PROOF. For $\tau < \pi/B$ and $x \in \mathcal{I}_\tau$, consider the modulus of the partial sum,

$$\begin{aligned} |\sum_{|j| \geq k} h(x + 2\pi j/\tau)| &\leq A \sum_{|j| \geq k} |x + 2\pi j/\tau|^{-\alpha} \\ &\leq 2A \sum_{j=k}^\infty |(\pi/\tau)[2j - 1]|^{-\alpha} \leq A\tau^\alpha [2\pi^{-\alpha} \sum_{j=k}^\infty j^{-\alpha}] \end{aligned}$$

for $k > 0$. This confirms uniform convergence of the sum defining $h_\tau(x)$ and, hence, its existence and continuity. Setting $k = 1$ in the above yields, for $x \in I_\tau$,

$$|h_\tau(x) - h(x)| \leq A\tau^\alpha [2\pi^{-\alpha} \sum_{j=1}^\infty j^{-\alpha}],$$

from which conclusion 2 follows directly. (When h is defined and continuous on R^+ , similar results apply by considering $h_\tau(x)$ on $[\tau, \tau + 2\pi/\tau]$ as $\tau \rightarrow 0$.)

COROLLARY 6.3. *If h' exists and satisfies the conditions of the lemma as well, then $h'_\tau(x) = \partial h_\tau(x)/\partial x$ exists, and equals the termwise derivative, i.e.,*

$$h'_\tau(x) = \sum_{j=-\infty}^\infty h'(x + 2\pi j/\tau),$$

justifying the otherwise ambiguous notation.

COROLLARY 6.4. *If $h(x; \theta)$ depends continuously on $\theta \in \Theta$ and satisfies condition (6.1) uniformly in θ , $h_\tau(x; \theta)$ exists and is a continuous function on $I_\tau \times \Theta$. If $\dot{h}(x; \theta) = \partial h(x; \theta)/\partial \theta$ exists and satisfies a similar condition, $\dot{h}_\tau(x; \theta)$ exists and is well defined.*

PROOFS. The results follow by uniform convergence of the series defining each of the “wrapped” forms.

The following two theorems due to Jackson (1930) relate mean square and uniform convergence for trigonometric polynomials, and will allow us to construct $S_*(x; \theta)$, using the τ indicated above, so that

$$|S_*(x; \theta) - S_\tau(x; \theta)| < \epsilon/2.$$

Theorem 6.5 follows directly from Jackson’s Theorem 1, page 2 while Theorem 6.6 is a slight generalization of Theorem IIa, page 84.

THEOREM 6.5. *Let $h(x)$ be a differentiable function of period $2\pi/\tau$, such that $|h'(x)| < A$ for all x . Then there exists a trigonometric polynomial of degree k , $q(x) = a_0 + \sum_{j=1}^k (a_j \cos j\tau x + b_j \sin j\tau x)$, satisfying*

$$\sup_{x \in R^1} |q(x) - h(x)| \leq KA/k\tau,$$

where K is an absolute constant.

THEOREM 6.6. *Let $h(x)$ be continuous, and let $r(x)$ and $s(x)$ be k th degree trigonometric polynomials in τx such that*

$$\sup_{x \in I_\tau} |h(x) - r(x)| < \delta$$

and

$$\int_{I_\tau} |h(x) - s(x)|^2 dx \leq d.$$

Then

$$\sup_{x \in I_\tau} |h(x) - s(x)| \leq 4[k\tau d]^{1/2} + 5\delta.$$

We may now proceed directly to establish Theorem 6.1. We first consider $p_\tau(x; \theta)$, the wrapped version of $p(x; \theta)$, defined as in Lemma 6.2, as well as its derivative $\dot{p}_\tau(x; \theta)$, which exists by Corollary 6.4. Without loss of generality, we assume $R_c = [-C, C] \times \Theta_c$. Since $p_\tau(x; \theta)$ and $\dot{p}_\tau(x; \theta)$ tend uniformly on R_c to $p(x; \theta)$ and $\dot{p}(x; \theta)$, by appeal to Lemma 6.2 and its corollaries, and since $p(x; \theta)$ is bounded below, away from 0, on R_c , this implies that

$$S_\tau(x; \theta) = \partial \log p_\tau(x; \theta) / \partial \theta = \dot{p}_\tau(x; \theta) / p_\tau(x; \theta)$$

tends uniformly to $S(x; \theta)$ on the same set. Choose $\tau < \min[\pi/B, \pi/C]$ so that

$$|S_\tau(x; \theta) - S(x; \theta)| < \varepsilon/2,$$

for $(x, \theta) \in R_c$.

Now consider $S_\tau(x; \theta)$ and its derivative,

$$S'_\tau(x; \theta) = \frac{p'_\tau(x; \theta)\dot{p}_\tau(x; \theta) - p_\tau(x; \theta)\dot{p}'_\tau(x; \theta)}{p_\tau(x; \theta)^2}.$$

Again, by Lemma 6.2 and its corollaries, the functions in the numerator exist and are continuous on $I_\tau \times \Theta_c$, and hence bounded on that set. By assumption, $p_\tau(x; \theta) \geq p(x; \theta) \geq m_\tau > 0$ on $I_\tau \times \Theta_c$, so that $|S'_\tau(x; \theta)| < A_\tau$ on $I_\tau \times \Theta_c$ for some $A_\tau < \infty$. By appeal to Theorem 6.5 then, for each $k = 1, 2, 3, \dots$, there exist trigonometric polynomials, $q_k(x; \theta)$, satisfying

$$|q_k(x; \theta) - S_\tau(x; \theta)| < KA_\tau/k\tau.$$

Defining $S_*(x; \theta)$ as in the statement of the theorem, we let

$$\begin{aligned} d_k(\theta) &= \int_{I_\tau} [S_*(x; \theta) - S_\tau(x; \theta)]^2 dx \\ &\leq \frac{1}{m_\tau} \int_{I_\tau} [S_*(x; \theta) - S_\tau(x; \theta)]^2 p_\tau(x; \theta) dx. \end{aligned}$$

By construction, $S_*(x, \theta)$ is the best $L_2(dP_\theta)$ approximation to $S_\tau(x; \theta)$ as a k th degree trigonometric polynomial, so that the above is

$$\leq \frac{1}{m_\tau} \int_{I_\tau} [q_k(x; \theta) - S_\tau(x; \theta)]^2 p_\tau(x; \theta) dx \leq \frac{1}{m_\tau} \left[\frac{KA_\tau}{k\tau} \right]^2.$$

Application of Theorem 6.6 yields

$$\sup_{x \in I} |S_\tau(x; \theta) - S_*(x; \theta)| \leq (KA_\tau/\tau)[4(\tau/m_\tau)^{1/2}k^{-1/2} + 5k^{-1}],$$

uniformly in $\theta \in \Theta_c$. Since τ is fixed in the above, it is clearly possible to choose k so that $|S_\tau(x; \theta) - S_*(x; \theta)| \leq \varepsilon/2$, and the proof is complete.

It must be pointed out that the conditions on $p(x; \theta)$ and its derivatives, though not stringent, are nonetheless troublesome in that their verification depends on knowledge of their asymptotic behaviour. (The nonvanishing of $p(x; \theta)$ and its boundedness on compact sets will generally be easy to verify, and indeed will usually be implicit in the formulation of the model.) Ideally, these conditions should be restated in terms of $\phi(t; \theta)$, since this is what is known. Some progress in this direction can be made by noting that the Fourier transforms of $p(x; \theta)$, $\dot{p}(x; \theta)$, $p'(x; \theta)$, and $\dot{p}'(x; \theta)$ are, under regularity conditions $\phi(t; \theta)$, $\dot{\phi}(t; \theta)$, $t \cdot \phi(t; \theta)$, and $t \cdot \dot{\phi}(t; \theta)$. The following lemma is thus relevant.

LEMMA 6.7. *If $h(x)$ is an integrable function with Fourier transform, $\eta(t) = \int_R h(x)\exp(itx) dx$ and $\eta(t)$ is twice differentiable everywhere, then*

$$h(x) \leq \left[\int_{R^1} |\eta''(t)| dt \right] |x|^{-2}.$$

PROOF.

$$|x^2h(x)| = \left| \int_{R^1} x^2 \exp(-ixt)\eta(t) dt \right| \leq \int_{R^1} |\eta''(t)| dt,$$

by the Fourier inversion formula, and integration by parts. Thus, a strong version of the order condition, i.e., with $\alpha = 2$, can be verified easily, in terms of $\phi(t; \theta)$, $\dot{\phi}(t; \theta)$, et cetera. The condition that $\phi(t; \theta)$ be twice differentiable implies the existence of the second moment of x for all θ . Thus this result may not be applied in such interesting cases as x being stably distributed.

6.2. *The log likelihood function.* In the continuous case it is possible to approximate $\ell(x; \theta) = \log p(x; \theta)$ via a trigonometric polynomial, $\ell_*(x; \theta)$, facilitating examination of the log likelihood surface. The approximate form arises from considering $J(x; \theta) = (\partial \log p(x; \theta))/\partial x$, and $J_*(x; \theta) = E^*\{J(x; \theta) | 1, \exp itx\}$, which may be obtained explicitly since

$$E_\theta\{J(x; \theta)\exp itx\} = \int p'(x; \theta)\exp(itx) dx = it\phi(t; \theta),$$

integrating by parts. $J_*(x; \theta)$ can be written as

$$(6.2) \quad J_*(x; \theta) = [iD_t\phi(\theta)]^* \Sigma_k(\theta)^{-1} [\exp itx - \phi(\theta)],$$

where $\mathbf{t} = (-k\tau, \dots, -\tau, \tau, \dots, k\tau)$, $D_t = \text{diag}(\mathbf{t})$, $\phi(\theta) = \phi(\mathbf{t}; \theta)$ and $\Sigma_k(\theta) = \Gamma(\theta) - \phi(\theta)\phi(\theta)^*$. The switch to complex form is merely notational, since $\text{Im}(J_*) = 0$, whenever \mathbf{t} is as above. Note that in the complex domain it is customary to define $\text{cov}\{u, v\} = E\{u\bar{v}\} - E\{u\}E\{\bar{v}\}$, so that $\Gamma(\theta)$ above has entries $\Gamma_{ij} = \phi([i - j]\tau; \theta)$.

$\ell_*(x; \theta)$ is obtained by integrating the trigonometric terms of $J_*(x; \theta)$, yielding the intermediate form,

$$(6.3) \quad \tilde{\ell}_*(x; \theta) = \phi(\theta) * D_t \Sigma_k(\theta)^{-1} D_t^{-1} \exp(itx),$$

which approximates $\ell(x; \theta)$ up to a term in θ , which may be approximated by

$$(6.4) \quad b_*(\theta) = \frac{1}{2} \{ \log | \Sigma_k(\theta) | - \log | \Sigma_{k-1}(\theta) | - \log(2\pi/\tau) \}.$$

The result below pertains to

$$\ell_*(x; \theta) = \tilde{\ell}_*(x; \theta) + b_*(\theta).$$

THEOREM 6.8. *If $p(x; \theta) > 0$ on R^1 and is continuous and differentiable in x for all θ , then*

$$\lim_{\tau \rightarrow 0} \lim_{k \rightarrow \infty} | \ell_*(x; \theta) - \ell(x; \theta) | = 0$$

for all x and θ .

PROOF. We begin by noting that $J_*(x; \theta)$ is a slight generalization of $J_*(z)$ of Section 3.1, and converges in $L_2(dP_\theta)$ to $J(x; \theta)$ as long as $J(x; \theta)$ lies in this space. By the arguments of the previous section, $J_*(x; \theta)$ approximates $J_\tau(x; \theta)$, the form corresponding to $p_\tau(x; \theta)$. Let us consider the convergence of J_* more closely. $J_\tau(x; \theta)$ is periodic, with natural domain $I_\tau = [-\pi/\tau, \pi/\tau]$ and as such, possesses an ordinary Fourier expansion given by

$$J_\tau(x; \theta) = \sum_{j=-\infty}^{\infty} a_j(\theta) \exp(ij\tau x)$$

where

$$a_j(\theta) = \frac{\tau}{2\pi} \int_{I_\tau} J_\tau(x; \theta) \exp(-ij\tau x) dx.$$

Note that $a_0(\theta) = 0$, and that the partial sums $J_k(x; \theta) = \sum_{j=-k}^k a_j(\theta) \exp(ij\tau x)$ will tend in $L_2(dx_\tau)$ (the restriction of $L_2(dx)$ to I_τ) to $J_\tau(x; \theta)$, i.e.

$$\int_{I_\tau} [J_k(x; \theta) - J_\tau(x; \theta)]^2 dx \rightarrow 0,$$

as $k \rightarrow \infty$. Now $J_*(x; \theta)$ is itself of the form $J_*(x; \theta) = \sum_{j=-k}^k a_{jk}(\theta) \exp(ij\tau x)$, the coefficients being given implicitly by (6.2). Since $J_*(x; \theta)$ tends to $J(x; \theta)$ as well, let us consider the relationship of $a_{jk}(\theta)$ to $a_j(\theta)$ as $k \rightarrow \infty$. If $p(x; \theta) > 0$ on R^1 , and is continuous in x , then $p_\tau(x; \theta)$ will be bounded away from 0 in $I_\tau = [-\pi/\tau, \pi/\tau]$, say $p_\tau(x; \theta) \geq \lambda(\theta) > 0$. (If $p(x; \theta)$ has support on $(0, \infty)$ only, defining $p_\tau(x; \theta)$ on $[\tau, \tau + (2\pi/\tau)]$ permits a similar argument). This implies that

$$\int [J_*(x; \theta) - J_\tau(x; \theta)]^2 dP_\theta \geq \lambda(\theta) \int_{I_\tau} [J_*(x; \theta) - J_\tau(x; \theta)]^2 dx,$$

so that J_* tends to J_τ in $L_2(dx_\tau)$, as well. This fact can be expressed in terms of

coefficients as

$$\sum_{j=-k}^k |a_j(\theta) - a_{jk}(\theta)|^2 \rightarrow 0$$

as $k \rightarrow \infty$.

Now consider $\ell_\tau(x; \theta) = \log p_\tau(x; \theta)$, whose Fourier coefficients are given by $b_j(\theta) = a_j(\theta)/ij\tau$ for $j \neq 0$, and $b_0(\theta) = (\tau/2\pi) \int_{I_\tau} \ell_\tau(x; \theta) dx$. Whenever $J_\tau(x; \theta)$ exists and is bounded, the Fourier expansion of $\ell_\tau(x; \theta)$, $\ell_k(x; \theta) = \sum_{j=-k}^k b_j(\theta)\exp(ij\tau x)$, converges uniformly to $\ell_\tau(x; \theta)$ on I_τ , as well as in the $L_2(dx_\tau)$ sense. Consider then,

$$\tilde{\ell}_*(x; \theta) = -i \sum_{j=-k, j \neq 0}^k (a_{jk}(\theta)/j\tau)\exp(ij\tau x),$$

given previously by (6.3). Letting $\tilde{\ell}_k(x; \theta) = \ell_k(x; \theta) - b_0(\theta)$, we have

$$\begin{aligned} \frac{\tau}{2\pi} \int_{I_\tau} [\tilde{\ell}_*(x; \theta) - \tilde{\ell}_k(x; \theta)]^2 dx &= \sum_{j=-k, j \neq 0}^k \frac{|a_j(\theta) - a_{jk}(\theta)|^2}{(k\tau)^2} \\ &\leq (1/\tau^2) \sum_{j=-k}^k |a_j(\theta) - a_{jk}(\theta)|^2, \end{aligned}$$

which tends to 0 as $k \rightarrow \infty$, i.e., $\tilde{\ell}_*(x; \theta) - \tilde{\ell}_k(x; \theta)$ tends to 0 in $L_2(dx_\tau)$. Since $\tilde{\ell}_k(x; \theta)$ tends in $L_2(dx_\tau)$ to $\tilde{\ell}_\tau(x; \theta) = \ell_\tau(x; \theta) - b_0(\theta)$, by the above, the same holds for $\tilde{\ell}_*(x; \theta)$. Also, for all x in I_τ ,

$$\begin{aligned} |\tilde{\ell}_*(x; \theta) - \ell_k(x; \theta)| &= |\sum_{j=-k, j \neq 0}^k [a_j(\theta) - a_{jk}(\theta)](\exp(ij\tau x)/j\tau)| \\ &\leq (1/\tau)[\sum_{j=-k}^k |a_j(\theta) - a_{jk}(\theta)|^2 \cdot 2 \sum_{j=1}^k j^{-2}], \end{aligned}$$

by Cauchy-Schwarz. Since $\sum_{j=1}^\infty j^{-2}$ converges, and the first sum tends to 0, the above confirms the pointwise convergence of $\tilde{\ell}_*(x; \theta)$ to $\tilde{\ell}_\tau(x; \theta) - \ell(x; \theta) - b_0(\theta)$, as $k \rightarrow \infty$.

The term $b_0(\theta)$ may be approximated based on the result in Grenander and Szego (1958, page 45), which yields that

$$\lim_{k \rightarrow \infty} [|\Sigma_k(\theta)| / |\Sigma_{k-1}(\theta)|]^{1/2} = (2\pi/\tau)\exp(b_0(\theta)).$$

This establishes the convergence of $b_*(\theta)$ given in (6.4) to $b_0(\theta)$, completing the proof of the theorem.

Examples considered by the author show that, as in the case of $S_*(x; \theta)$, k need not be inordinately big for $\ell_*(x; \theta)$ to be virtually indistinguishable from $\ell(x; \theta)$. Thus $\ell_*(x; \theta)$ is of general utility. For example, if the empirical characteristic function $\hat{\phi}(t)$ is substituted for $\exp(itx)$ in $\ell_*(x; \theta)$, the result is a convenient transform based means of describing the log likelihood surface. Also, $\ell_*(x; \theta)$ may serve as a basis for approximating $p(x; \theta)$, a possibility which is further explored in Section 7, using the version of ℓ_* based on the moments.

6.3. *Related forms.* Other forms which are commonly encountered in connection with likelihood calculations are the Fisher Information, $\mathcal{I}(\theta)$, and the observed information, I_n , defined in terms of

$$I(x; \theta) = -(\partial^2 p(x; \theta)/\partial \theta^2).$$

Fundamental to all our results is the fact that $\mathcal{I}_*(\theta) = \dot{\phi}(\theta)^* \Sigma(\theta)^{-1} \dot{\phi}(\theta)$ provides

an approximation to $\mathcal{J}(\theta)$. Also, it may be noted that the behaviour of

$$I_{*n} = -(\partial S_{*n}/\partial\theta) |_{\theta=\hat{\theta}},$$

where $S_{*n} = \sum_{i=1}^n S_*(x_i; \theta)$, is analogous to that of I_n , and may be regarded as an approximation to it.

Alternately, it may be noted that

$$I(x; \theta) = -(\partial^2 \log p(x; \theta)/\partial\theta^2) = S(x; \theta)^2 - \bar{p}(x; \theta)/p(x; \theta),$$

which may be approximated by

$$S_*(x; \theta)^2 - \ddot{\phi}(\theta)^* \Sigma(\theta)^{-1} [\exp itx - \phi(\theta)].$$

Note that $S_*(x; \theta)^2$ is a trigonometric polynomial, and that we can write

$$S_*(x; \theta)^2 = \dot{\phi}(\theta)^* \Sigma(\theta)^{-1} \exp itx \cdot (\exp itx)^* \Sigma(\theta)^{-1} \dot{\phi}(\theta),$$

so that an alternative approximation to the observed information is

$$I_{*n} = n\{\dot{\phi}(\hat{\theta})^* \Sigma(\hat{\theta})^{-1} \hat{\Phi} \Sigma(\hat{\theta})^{-1} \dot{\phi}(\hat{\theta}) - \ddot{\phi}(\hat{\theta})^* \Sigma(\hat{\theta})^{-1} [\hat{\phi} - \phi(\hat{\theta})]\},$$

where $\hat{\Phi}$ has entries,

$$\hat{\Phi}_{ij} = \hat{\phi}([i - j] \cdot \tau).$$

7. Probability calculations based on transforms. As noted in Section 6, $p_*(x) = \exp \{\ell_*(x)\}$ yields a new approximation to $p(x)$ (in what follows we shall neglect the parameter θ , which is irrelevant). There ℓ_* was based on the c.f. $\phi(t)$, whereas here we will consider a version of ℓ_* based on the moments $\mu_k = E\{x^k\}$, $k = 1, 2, 3, \dots$, and will compare the resultant $p_*(x)$ with the Gram Charlier Type A series, and the closely related Edgeworth expansion, both of which provide moment based approximations to $p(x)$. Briefly, the latter expansions arise from best $L_2(\omega dx)$ polynomial approximations to $h(x) = p(x)/\omega(x)$, where $\omega(x)$ is the standard Gaussian density. Minimizing errors of approximation on the implied scale of measurement does not always yield satisfactory results. The approximations may fail to converge, or may be deficient in the tails, assuming negative values for example. The logarithmic scale, by comparison, is natural in any context where the relative, rather than absolute, error in approximation is relevant. The appearance of $J(x) = -p'(x)/p(x)$ in motivating ℓ_* and p_* is not unnatural either, relative to the central role the Gaussian distribution in density approximation. Hampel (1973) notes of the Gaussian density f that “ $f'(x)/f(x)$ is a linear function, and the simplest function which in turn determines the normal distribution”.

The possibility of approximating $J(x)$ by a polynomial arises since

$$\eta_k = \text{cov}\{J(x), x^k\} = k\mu_{k-1},$$

under mild regularity conditions.

Letting $\eta = (\eta_1, \eta_2, \dots, \eta_k)^T$, $\mathbf{t} = (1, 2, 3, \dots, k)$, $\mu = E\{x^t\}$ and $\Sigma = \text{var}(x^t)$, with entries $\Sigma_{ij} = \mu_{i+j} - \mu_i\mu_j$, we can write $J_*(x) = \eta^T \Sigma^{-1} (x^t - \mu)$. Integrating




FIG. 3. Lower and upper tails of Gamma(3) density and approximations.

termwise, we obtain the improper form

$$\tilde{\ell}_*(x) = \eta^T \Sigma^{-1} (D^{-1} x^{(t+1)} - \mu)$$

where $D = \text{diag}(t + \mathbf{1})$, where $\mathbf{1}$ is a k -vector of 1's. The convergence of $\ell_*(x)$ holds if $J(x)$ lies in $L_2(dP)$, since $J_* \rightarrow J$ in L_2 implies the L_1 result, and since

$$\begin{aligned} \left| \int_0^x [J(u) - J_*(u)] \cdot du \right| &= |\ell(x) - \ell(0) - \ell_*(x)| \\ &\leq \frac{1}{\lambda(x)} \int |J(x) - J_*(x)| p(x) dx \end{aligned}$$

where $\lambda(x) = \inf\{p(u); 0 \leq u \leq x\}$. This rather crude upper bound ($\lambda(x)$ may be rather small) is still sufficient, based on the L_1 convergence noted, to confirm the pointwise convergence of $\tilde{\ell}_*(x)$ to $\ell(x) - \ell(0)$, as long as $\lambda(x) \neq 0$. By taking $p_*(x) = c_* \exp(\tilde{\ell}_*(x))$, where c_* is the normalizing constant determined by numerical integration, we achieve an approximation to $p(x)$ that:

1. is easy to compute,
2. is always positive,
3. minimizes the error of approximation on an appropriate scale, and
4. exactly recovers $p(x)$ when J is a polynomial of order less than or equal to k , for example when P is exponential or Gaussian. (The Gram Charlier and Edgeworth expansions recover Gaussian distributions only after an appropriate standardization of the moments).

The approximation is applicable whenever the moments of P exist and

uniquely characterize P , and are known to sufficient precision to compute accurately the coefficients determining $\tilde{\ell}_*$.

The following examples give some indication of the quality of approximation afforded by p_* , in comparison with its competitors. In Figure 3, p_* , based on the least-squares 11th degree polynomial approximation to $J(x)$ (so that ℓ_* is 12th degree) for a Gamma(3) distribution, is compared to the 12th degree Edgeworth expansion. The corresponding Gram-Charlier expansion fared so poorly that it could not be plotted on the same scale, and hence was excluded. The plots indicate the superiority of p_* in the extreme regions of the distribution.

Acknowledgements. The major results of this paper form part of my Ph.D. thesis at the University of Toronto. I wish to thank my supervisors, Professor Andrey Feuerverger and Professor Philip McDunnough, for their insightful guidance. Also, I would like to thank the reviewers and the Associate Editor, whose comments led to substantial improvement in the organization of this paper.

REFERENCES

- BROCKWELL, P. J. and BROWN, B. M. (1982). High efficiency estimation for the positive stable laws. *J. Amer. Statist. Assoc.* **76** 626–631.
- BRYANT, J. L. and PAULSON, A. S. (1979). Some comments on characteristic function based estimators. *Sankhya A.* **41** 109–116.
- CHAMBERS, R. L. and HEATHCOTE, C. R. (1981). On the estimation of slope and the identification of outliers in linear regression. *Biometrika* **68** 21–33.
- DUMOUCHEL, W. H. (1975). Stable distributions in statistical inference 2: Information from stably distributed samples. *J. Amer. Statist. Assoc.* **70** 386–393.
- FEUERVERGER, A. and MCDUNNOUGH, P. (1981a). On the efficiency of empirical characteristic function procedures. *J. Roy. Statist. Soc. B* **43** 20–27.
- FEUERVERGER, A. and MCDUNNOUGH, P. (1981b). On some Fourier methods of inference. *J. Amer. Statist. Assoc.* **76** 379–387.
- FEUERVERGER, A. and MCDUNNOUGH, P. (1981c). On efficient inference in symmetric stable laws and processes. *Statistics and Related Topics: Proceedings of the International Symposium on Statistics 1980*. North-Holland, New York.
- FEUERVERGER, A. and MCDUNNOUGH, P. (1982). On statistical transform methods and their efficiency. Unpublished manuscript.
- GRENANDER, U. and SZEGO, G. (1958). *Toeplitz Forms and Their Applications*. Univ. of California Press, Berkeley.
- HAMPEL, F. (1973). Some small sample asymptotics. *Proceedings of the Prague Symposium on Asymptotic Statistics II* 109–126.
- JACKSON, D. (1930). *Theory of Approximation*. A.M.S. Colloq. Pub. 11.
- JARRETT, R. G. (1973). Efficiency and estimation in asymptotically normal distribution. Doctoral thesis, Univ. of London.
- SCLOVE, S. L. and VAN RYZIN, J. (1969). Estimating the parameters of a convolution. *J. Roy. Statist. Soc. B* 181–191.
- SPROTT, D. A. (1983). Estimating the parameters of a convolution by maximum likelihood. *J. Amer. Statist. Assoc.* **78** 457–463.

DEPARTMENT OF APPLIED STATISTICS
UNIVERSITY OF MINNESOTA
1994 BUFORD AVENUE
ST. PAUL, MINNESOTA 55108.