

GLUING SILTING OBJECTS

QUNHUA LIU, JORGE VITÓRIA, AND DONG YANG

Abstract. Recent results by Keller and Nicolás and by Koenig and Yang have shown bijective correspondences between suitable classes of t-structures and co-t-structures with certain objects of the derived category: silting objects. On the other hand, the techniques of gluing (co-)t-structures along a recollement play an important role in the understanding of derived module categories. Using the above correspondence with silting objects, we present explicit constructions of gluing of silting objects, and, furthermore, we answer the question of when the glued silting is tilting.

§1. Introduction

In a given triangulated category, the study of its torsion pairs helps to understand its structure. Two kinds of torsion pairs have been considered with particular emphasis in the literature. These are the notions of t-structure introduced by Beilinson, Bernstein, and Deligne in [5] and co-t-structure introduced independently by Bondarko in [8] and by Pauksztello in [22]. These are torsion pairs with an additional property concerning the suspension functor of the underlying triangulated category, and they give rise to additive (or even abelian in the case of t-structures) subcategories which are of interest. In this paper, we work with correspondences that classify these torsion pairs in terms of objects of the triangulated category.

Keller and Vossieck established in [17] a bijection between bounded t-structures and equivalence classes of silting objects in the bounded derived category of modules over the path algebra of a Dynkin quiver over a field. Recently, this bijection has been extended by Keller and Nicolás in [16] for the bounded derived categories of homologically smooth nonpositive differential graded algebras, and by Koenig and Yang in [18] for bounded derived categories of finite-dimensional algebras over a field. Indeed, they show that in such a category, there is a bijection between silting objects

Received July 10, 2012. Revised November 10, 2013. Accepted February 5, 2014.
First published online January 9, 2015.

2010 Mathematics Subject Classification. Primary 16E35; Secondary 18E30.

and bounded t-structures whose hearts are length categories. It turns out that Keller and Vossieck’s result is a corollary, since for algebras of finite representation type all hearts of bounded t-structures are length categories. A new correspondence between silting objects and bounded co-t-structures was proved in [21] and [16]. This bijection will be central in our approach.

Silting objects play, thus, a more general role than tilting objects. They describe all hearts which are length categories and these turn out to be precisely those which are module categories over some finite-dimensional algebra over a field. This algebra, although not in general derived equivalent to the one we started with, is obtained as the endomorphism algebra of the silting object, just like in the tilting setting. Indeed, it is easy to observe that a silting object is tilting if and only if it lies in the heart of the corresponding t-structure.

Gluing techniques with respect to a recollement, due to Beilinson, Bernstein, and Deligne, have been intensively studied in [5] for t-structures and, recently, in [8] for co-t-structures. This leads to the natural question of how to glue silting objects and which silting objects are glued from *smaller* ones. Indeed, recent work by the two first authors has shown that, in the piecewise hereditary case, all bounded t-structures whose heart is a length category are glued with respect to a nontrivial recollement (see [20]). In this setting it is then clear that every silting object can be decomposed by this process into as many pieces as derived simple factors of the algebra (check [3] and [4] for terminology). It turns out, however, that an answer to the problem of gluing silting objects can be given more easily when the focus is on co-t-structures rather than on t-structures. Our main result is as follows.

THEOREM A (Theorem 3.1). *Let \mathcal{R} be a recollement of a triangulated category \mathcal{D} of the form*

$$\mathcal{Y} \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D} \begin{array}{c} \xleftarrow{j_!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{X}.$$

*Let X and Y be, respectively, silting objects of \mathcal{X} and \mathcal{Y} , and let $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ be, respectively, the associated co-t-structures in \mathcal{X} and \mathcal{Y} . Then the induced co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ in \mathcal{D} is associated with the silting object $Z = i_*Y \oplus K_X$ in \mathcal{D} , with K_X defined by the triangle*

$$i_*\beta_{\geq 1}i^!j_!X \rightarrow j_!X \rightarrow K_X \rightarrow (i_*\beta_{\geq 1}i^!j_!X)[1],$$

where $\beta_{\geq 1}$ is a (nonfunctorial) choice of truncation for the co-t-structure $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in \mathcal{Y} .

Furthermore, the question of when we can glue derived equivalences—that is, tilting objects—comes as a particular setting of the general context of gluing silting. Similar constructions of tilting objects have been discussed in [19] and [2]. In particular, we will show that the construction in [2] is a particular case of the preceding construction. The following is our main theorem concerning tilting (see Theorem 4.5).

THEOREM B (Theorem 4.5). *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ by $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$, where R , C , and B are finite-dimensional algebras over a field of finite global dimension. Let X and Y be tilting objects of \mathcal{X} and \mathcal{Y} , respectively. Then $Z = i_*Y \oplus K_X$ is tilting in \mathcal{D} if and only if the following conditions hold:*

- (a) $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k]) = 0$ for all $k < -1$,
- (b) $\text{Hom}_{\mathcal{Y}}(i^*j_*X, Y[k]) = 0$ for all $k < 0$,
- (c) $\text{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0$ for all $k < -1$.

This paper is structured as follows. In the next section we discuss some preliminary results on recollements, t-structures, co-t-structures, and silting needed for the later sections. In Section 3 we show how to glue silting objects and we use this in Section 4 to give necessary and sufficient conditions for the glued silting to be tilting. These conditions are particularly nice in the hereditary case and they are made explicit in Section 5. Finally, in Section 6, we make a few observations on the gluing of Happel–Reiten–Smalø (HRS)-tilts with a view towards a result on the compatibility of silting mutation with gluing (see Corollary 6.9).

§2. Preliminaries

Throughout, \mathbb{K} denotes a fixed field, and R , B , and C denote finite-dimensional \mathbb{K} -algebras. The bounded derived category of finitely generated right R -modules and the bounded homotopy category of finitely generated projective right R -modules will be denoted, respectively, by $\mathcal{D}^b(R)$ and $\mathcal{K}^b(\text{proj-}R)$. The symbols \mathcal{X} , \mathcal{Y} , and \mathcal{D} denote triangulated categories.

2.1. Recollements

A *recollement* of \mathcal{D} by \mathcal{X} and \mathcal{Y} is a diagram of six triangle functors

$$(2.1) \quad \begin{array}{ccc} \xleftarrow{i^*} & & \xleftarrow{j!} \\ \mathcal{Y} & \xrightarrow{i_*} \mathcal{D} & \xrightarrow{j^*} \mathcal{X} \\ \xleftarrow{i^!} & & \xleftarrow{j_*} \end{array}$$

satisfying the following properties:

- (1) (i^*, i_*) , $(i_*, i^!)$, $(j_!, j^*)$, (j^*, j_*) are adjoint pairs,
- (2) i_* , j_* , $j_!$ are full embeddings,
- (3) $i^! \circ j_* = 0$ (and thus also $j^* \circ i_* = 0$ and $i^* \circ j_! = 0$),
- (4) for each $Z \in \mathcal{D}$, the units and counits of the adjunctions yield triangles

$$\begin{aligned} i_* i^! Z &\rightarrow Z \rightarrow j_* j^* Z \rightarrow i_* i^! Z[1], \\ j_! j^* Z &\rightarrow Z \rightarrow i_* i^* Z \rightarrow j_! j^* Z[1]. \end{aligned}$$

The following result allows us to change the sides of a recollement.

THEOREM 2.1 ([7, Propositions 3.6–3.7], [15, Proposition 5, Theorem 7]). *Let \mathcal{R} be a recollement of \mathcal{D} of the form (2.1). If \mathcal{D} has a Serre functor S , then both \mathcal{X} and \mathcal{Y} have Serre functors (resp., $S_{\mathcal{X}}$ and $S_{\mathcal{Y}}$) and there are reflected recollements \mathcal{R}_U and \mathcal{R}_L (resp., upper and lower reflection)*

$$\mathcal{R}_U: \quad \mathcal{X} \begin{array}{c} \xleftarrow{j^\#} \\ \xrightarrow{j_!} \\ \xleftarrow{j^*} \end{array} \mathcal{D} \begin{array}{c} \xleftarrow{i_\#} \\ \xrightarrow{i^*} \\ \xleftarrow{i_*} \end{array} \mathcal{Y},$$

where $i_\# = S^{-1}i_*S_{\mathcal{Y}}$ and $j^\# = S_{\mathcal{X}}^{-1}j^*S$ are left adjoints of i^* and $j_!$, respectively, and

$$\mathcal{R}_L: \quad \mathcal{X} \begin{array}{c} \xleftarrow{j^*} \\ \xrightarrow{j_*} \\ \xleftarrow{j^+} \end{array} \mathcal{D} \begin{array}{c} \xleftarrow{i_*} \\ \xrightarrow{i^!} \\ \xleftarrow{i_+} \end{array} \mathcal{Y},$$

where $i_+ = Si_*S_{\mathcal{Y}}^{-1}$ and $j^+ = S_{\mathcal{X}}j^*S^{-1}$ are right adjoints of $i^!$ and j_* , respectively.

2.2. t-structures and co-t-structures

A *torsion pair* in \mathcal{D} is a pair of strict full subcategories $(\mathcal{D}', \mathcal{D}'')$, closed under taking direct summands and direct sums, such that

- (1) $\mathrm{Hom}_{\mathcal{D}}(X, Y) = 0$ for all $X \in \mathcal{D}'$ and $Y \in \mathcal{D}''$,
- (2) for each Z in \mathcal{D} , there are $X \in \mathcal{D}'$, $Y \in \mathcal{D}''$ and a triangle

$$(2.2) \quad X \rightarrow Z \rightarrow Y \rightarrow X[1].$$

By definition, it is easy to see that

$$\begin{aligned} \mathcal{D}' &= \{X \in \mathcal{D} : \mathrm{Hom}_{\mathcal{D}}(X, Y) = 0, \forall Y \in \mathcal{D}''\} =: {}^\perp\mathcal{D}'', \\ \mathcal{D}'' &= \{Y \in \mathcal{D} : \mathrm{Hom}_{\mathcal{D}}(X, Y) = 0, \forall X \in \mathcal{D}'\} =: \mathcal{D}'^\perp. \end{aligned}$$

A *t-structure* in \mathcal{D} is a pair $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ such that $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 1})$ is a torsion pair and $\mathcal{D}^{\leq 0} \subseteq \mathcal{D}^{\leq 1}$, where $\mathcal{D}^{\leq n} := \mathcal{D}^{\leq 0}[-n]$ and $\mathcal{D}^{\geq n} := \mathcal{D}^{\geq 0}[-n]$ for $n \in \mathbb{Z}$. The subcategory $\mathcal{D}^{\leq 0}$ is called the *t-aisle*, and $\mathcal{D}^{\geq 0}$ is called the *t-co-aisle*. The *heart* $\mathcal{D}^{\leq 0} \cap \mathcal{D}^{\geq 0}$ is always an abelian category. We say that the heart is a *length category* if it has only finitely many simple objects (up to isomorphism) and each object in the heart has finite length. A t-structure is said to be *bounded* if it satisfies

$$\bigcup_{n \in \mathbb{Z}} \mathcal{D}^{\geq n} = \mathcal{D} = \bigcup_{n \in \mathbb{Z}} \mathcal{D}^{\leq n}.$$

For all $n \in \mathbb{Z}$, there is a right adjoint to the inclusion of the subcategory $\mathcal{D}^{\leq n}$ in \mathcal{D} , called the *truncation at n* and denoted by $\tau^{\leq n}$. Similarly, there is a left adjoint to the inclusion of $\mathcal{D}^{\geq n}$, denoted by $\tau^{\geq n}$. In fact, the triangle (2.2) can be written functorially, for any $Z \in \mathcal{D}$, as

$$\tau^{\leq 0} Z \rightarrow Z \rightarrow \tau^{\geq 1} Z \rightarrow (\tau^{\leq 0} Z)[1].$$

Moreover, it is possible to define associated cohomological functors

$$H^i : \mathcal{D} \rightarrow \mathcal{D}^{\leq 0} \cap \mathcal{D}^{\geq 0}, \quad H^i(X) = (\tau^{\leq i} \tau^{\geq i} X)[i], \quad \forall X \in \mathcal{D}.$$

If a t-structure is bounded, then for any object $X \in \mathcal{D}$, $H^i(X) = 0$ for all but finitely many $i \in \mathbb{Z}$. A well-known example is the standard t-structure in $\mathcal{D}^b(R)$. The associated cohomological functors are given by the usual cohomology of complexes in $\mathcal{D}^b(R)$.

Similarly, a *co-t-structure* in \mathcal{D} is a pair $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ such that $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq -1})$ is a torsion pair and $\mathcal{D}_{\geq 0} \subseteq \mathcal{D}_{\geq -1}$, where $\mathcal{D}_{\geq n} := \mathcal{D}_{\geq 0}[-n]$ and $\mathcal{D}_{\leq n} := \mathcal{D}_{\leq 0}[-n]$, for all $n \in \mathbb{Z}$. The subcategory $\mathcal{D}_{\geq 0}$ is called the *co-t-aisle*, and $\mathcal{D}_{\leq 0}$ is called the *co-t-co-aisle*. The *co-heart* $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ is not, in general, an abelian category, as in the case of t-structures, but it still has the structure of an additive category. A co-t-structure is said to be *bounded* if it satisfies

$$\bigcup_{n \in \mathbb{Z}} \mathcal{D}_{\geq n} = \mathcal{D} = \bigcup_{n \in \mathbb{Z}} \mathcal{D}_{\leq n}.$$

Contrarily to t-structures, the triangle obtained from the torsion pair $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq -1})$ is not functorial. However, a choice of $X \in \mathcal{D}_{\geq 0}$ (resp., of $Y \in \mathcal{D}_{\leq -1}$) as in triangle (2.2) will be denoted by $\beta_{\geq 0} Z$ (resp., by $\beta_{\leq -1} Z$). A well-known example (see [8], [22]) is the standard co-t-structure in $\mathcal{K}^b(\text{proj-}R)$. Here, the nonfunctorial choices of $\beta_{\geq 0} Z$ or $\beta_{\leq 0} Z$ for some object

$Z \in \mathcal{K}^b(\text{proj-}R)$ are usually given by the *stupid truncations*, consisting in setting suitable entries of the complex Z to be zero.

It is known that t-structures and co-t-structures can be glued (or induced) with respect to a recollement (see [5] for t-structures and [8] for co-t-structures). The same arguments can be used to prove the following result.

THEOREM 2.2. *Let \mathcal{R} be a recollement of \mathcal{D} of the form (2.1). Let $(\mathcal{X}', \mathcal{X}'')$ and $(\mathcal{Y}', \mathcal{Y}'')$ be torsion pairs in \mathcal{X} and \mathcal{Y} , respectively.*

(1) *There is a glued torsion pair $(\mathcal{D}', \mathcal{D}'')$ in \mathcal{D} defined by*

$$\mathcal{D}' = \{Z \in \mathcal{D} : j^*Z \in \mathcal{X}', i^*Z \in \mathcal{Y}'\}, \quad \mathcal{D}'' = \{Z \in \mathcal{D} : j^*Z \in \mathcal{X}'', i^!Z \in \mathcal{Y}''\}.$$

- (2) ([5, Théorème 1.4.10]) *If $(\mathcal{X}', \mathcal{X}''[1])$ and $(\mathcal{Y}', \mathcal{Y}''[1])$ are t-structures, then $(\mathcal{D}', \mathcal{D}''[-1])$ is a t-structure in \mathcal{D} .*
- (3) ([8, Theorem 8.2.3]) *If $(\mathcal{X}', \mathcal{X}''[-1])$ and $(\mathcal{Y}', \mathcal{Y}''[-1])$ are co-t-structures, then $(\mathcal{D}', \mathcal{D}''[1])$ is a co-t-structure in \mathcal{D} .*
- (4) ([8], [20], [28]) *The gluing of bounded t-structures whose heart is a length category is a bounded t-structure whose heart is a length category. Also, the gluing of bounded co-t-structures is still bounded.*

If \mathcal{D} admits a Serre functor S , if \mathcal{R} is a recollement of \mathcal{D} of the form (2.1), and if $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ are t-structures in \mathcal{X} and \mathcal{Y} , respectively, then there are three naturally associated t-structures in \mathcal{D} , obtained by gluing these with respect to \mathcal{R} , \mathcal{R}_U , and \mathcal{R}_L —we will denote them by $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{D}_{\bar{U}}^{\leq 0}, \mathcal{D}_{\bar{U}}^{\geq 0})$, and $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$, respectively. Similarly, given co-t-structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in \mathcal{X} and \mathcal{Y} , we get three glued co-t-structures $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$, $(\mathcal{D}_{\bar{U}}^U, \mathcal{D}_{\bar{U}}^L)$, and $(\mathcal{D}_{\geq 0}^L, \mathcal{D}_{\leq 0}^L)$.

2.3. Correspondences with silting objects

An object $M \in \mathcal{D}$ is *silting* if $\text{Hom}_{\mathcal{D}}(M, M[i]) = 0$, for all $i > 0$, and if \mathcal{D} is the smallest triangulated subcategory containing M which is closed under direct summands, extensions, and shifts. It is *tilting* if, in addition, $\text{Hom}_{\mathcal{D}}(M, M[i]) = 0$, for all $i < 0$. We say that two such objects M and N are *equivalent* if $\text{add}(M) = \text{add}(N)$. The following result is a consequence of [21, Corollary 5.9] (see also [16]).

THEOREM 2.3. *There is a bijection between bounded co-t-structures in \mathcal{D} whose co-heart is additively generated by one object and equivalence classes of silting objects of \mathcal{D} .*

This bijection can be described as follows.

- Given a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$, the associated equivalence class is given by a silting object M which is an additive generator of the co-heart, that is, $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0} = \text{add}(M)$.
- Given a silting object $M \in \mathcal{D}$, the associated co-t-structure is defined as follows: $\mathcal{D}_{\geq 0}$ is the smallest full subcategory closed under direct summands, direct sums, and extensions containing $\{M[i] : i \leq 0\}$; similarly, $\mathcal{D}_{\leq 0}$ is the smallest full subcategory closed under direct summands, direct sums, and extensions containing $\{M[i] : i \geq 0\}$.

THEOREM 2.4 ([18, Theorem 7.1]). *In $\mathcal{D}^b(R)$, there are bijections between*

- *the set of bounded t-structures in $\mathcal{D}^b(R)$ whose heart is a length category;*
- *the set of equivalence classes of silting objects in $K^b(\text{proj-}R)$;*
- *the set of bounded co-t-structures in $K^b(\text{proj-}R)$.*

Recall that R is of finite global dimension if and only if $\mathcal{D}^b(R)$ admits a Serre functor (see [7], [23]). Also, in this case, we have $K^b(\text{proj-}R) \cong \mathcal{D}^b(R)$ and, thus, the correspondence between t-structures and co-t-structures occurs in the same category. Under this assumption, the bijections of the theorem can be made explicit as follows (see [18, Section 6] for details).

- Given a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, we can associate a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ by taking $\mathcal{D}_{\leq 0} = \mathcal{D}^{\leq 0}$ and $\mathcal{D}_{\geq 0} = {}^{\perp} \mathcal{D}^{\leq -1}$; in the language of [8], this is the *left adjacent co-t-structure* to the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$.
- Given a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ in $\mathcal{D}^b(R)$, we can associate a t-structure by taking $\mathcal{D}^{\leq 0} = \mathcal{D}_{\leq 0}$ and $\mathcal{D}^{\geq 0} = \mathcal{D}_{\leq -1}^{\perp}$; in the language of [8], this is the *right adjacent t-structure* to the co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$.
- Given a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, the associated silting object is the direct sum of the indecomposable Ext-projective objects in the aisle, that is, the objects X of $\mathcal{D}^{\leq 0}$ such that $\text{Hom}_{\mathcal{D}^b(R)}(X, Y[1]) = 0$ for all $Y \in \mathcal{D}^{\leq 0}$.
- Given a silting object M in $\mathcal{D}^b(R)$, the associated t-structure is defined as follows:

$$\begin{aligned} \mathcal{D}^{\leq 0} &= \{Z \in \mathcal{D}^b(R) : \text{Hom}_{\mathcal{D}^b(R)}(M, Z[i]) = 0, \forall i > 0\}, \\ \mathcal{D}^{\geq 0} &= \{Z \in \mathcal{D}^b(R) : \text{Hom}_{\mathcal{D}^b(R)}(M, Z[i]) = 0, \forall i < 0\}. \end{aligned}$$

REMARK 2.5. If R is of finite global dimension, as before, then the co-t-co-aisle $\mathcal{D}_{\leq 0}$ associated with a silting object M in $\mathcal{D}^b(R)$ coincides with

the aisle $\mathcal{D}^{\leq 0}$ associated with the same silting object. In other words, the co-t-structure associated to M is left adjacent to the t-structure associated to M .

REMARK 2.6. The structures glued from adjacent t-structures and co-t-structures are not directly related by adjacency, as observed by Bondarko [8, Remark 8.2.4.4]. Suppose, however, that \mathcal{D} has a Serre functor and that $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ (resp., $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$) is a left adjacent co-t-structure to the t-structure $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ (resp., $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$). Using the descriptions provided in Theorem 2.2 and the functors of Theorem 2.1 for a fixed recollement \mathcal{R} , we have

$$\mathcal{D}_{\leq 0}^U = \mathcal{D}^{\leq 0} = \{Z \in \mathcal{D} : j^*Z \in \mathcal{X}^{\leq 0} = \mathcal{X}_{\leq 0}, i^*Z \in \mathcal{Y}^{\leq 0} = \mathcal{Y}_{\leq 0}\}.$$

This means that the co-structure $(\mathcal{D}_{\geq 0}^U, \mathcal{D}_{\leq 0}^U)$ glued by the upper reflected recollement \mathcal{R}_U is left adjacent to the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ glued by the recollement \mathcal{R} . Similarly, the t-structure $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$ glued by \mathcal{R}_L is right adjacent to the co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ glued by \mathcal{R} .

REMARK 2.7. A tilting object T in $\mathcal{K}^b(\text{proj-}R)$ yields equivalences between $\mathcal{D}^b(R)$ and $\mathcal{D}^b(\text{End}(T))$ and between $\mathcal{K}^b(\text{proj-}R)$ and $\mathcal{K}^b(\text{proj-End}(T))$ (see [24, Theorem 6.4]). Under these equivalences, the t-structure and the co-t-structure associated to T in $\mathcal{D}^b(R)$ correspond to the standard ones in $\mathcal{D}^b(\text{End}(T))$ and $\mathcal{K}^b(\text{proj-End}(T))$, respectively.

2.4. HRS-tilts and silting mutation

There are mutation operations on both t-structures and silting objects. Recall that the definition of a torsion pair in an abelian category is analogous to that of a torsion pair in a triangulated category, replacing the triangle axiom by a short exact sequence.

THEOREM 2.8 ([14, Proposition 2.1], [9, Proposition 2.5]). *Let $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ be a bounded t-structure in a triangulated category \mathcal{D} with heart \mathcal{A} and associated cohomology functors H^i , $i \in \mathbb{Z}$. Suppose that $(\mathcal{T}, \mathcal{F})$ is a torsion pair in \mathcal{A} . Then $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ is a t-structure in \mathcal{D} , where*

$$\begin{aligned} \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0} &= \{E \in \mathcal{D} : H^i(E) = 0, \forall i > 0, H^0(E) \in \mathcal{T}\}, \\ \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0} &= \{E \in \mathcal{D} : H^i(E) = 0, \forall i < -1, H^{-1}(E) \in \mathcal{F}\}. \end{aligned}$$

The t-structure $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ is called the *HRS-tilt* of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$.

REMARK 2.9. HRS-tilts span an important class of t-structures. In fact, it is known from [6, Theorem 3.1] and [29, Proposition 2.1] that, for a bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, the bounded t-structures $(\mathcal{C}^{\leq 0}, \mathcal{C}^{\geq 0})$ such that $\mathcal{D}^{\leq -1} \subseteq \mathcal{C}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$ are precisely those obtained from $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ by an HRS-tilt with respect to a torsion pair.

Assume that the heart \mathcal{A} of the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is a length category, and let S be a simple object of \mathcal{A} without self-extensions. The left mutation of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to S is defined by

$$\mu_S^-(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0}) = (\mathcal{D}_{-,S}^{\leq 0}, \mathcal{D}_{-,S}^{\geq 0}) := (\mathcal{D}_{(\perp S, \text{add}(S))}^{\leq 0}, \mathcal{D}_{(\perp S, \text{add}(S))}^{\geq 0}),$$

and its right mutation by

$$\mu_S^+(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0}) = (\mathcal{D}_{+,S}^{\leq 0}, \mathcal{D}_{+,S}^{\geq 0}) := (\mathcal{D}_{(\text{add}(S), S^\perp)}^{\leq 0}[-1], \mathcal{D}_{(\text{add}(S), S^\perp)}^{\geq 0}[-1]).$$

The torsion pairs $(\text{add}(S), S^\perp)$ and $(\perp S, \text{add}(S))$ will be called *mutation torsion pairs*.

On the other hand, silting mutation was introduced and studied by Buan, Reiten, and Thomas in [10] and, independently, by Aihara and Iyama in [1]. We recall its definition.

DEFINITION 2.10. Let $M = X \oplus Y$ be a silting object in $\mathcal{D}^b(R)$. The *left mutation of M at X* , denoted by $\mu_X^-(M)$, is defined as the direct sum $\tilde{X} \oplus Y$, where \tilde{X} is the cone of a left $\text{add}(Y)$ -approximation of X (i.e., of a morphism $\phi: X \rightarrow L$, with L in $\text{add}(Y)$ such that for any Z in $\text{add}(Y)$, $\text{Hom}_{\mathcal{D}^b(R)}(\phi, Z)$ is surjective).

Similarly, the *right mutation of M at X* , denoted by $\mu_X^+(M)$, is defined as the sum $\bar{X} \oplus Y$, where \bar{X} is the cone of a right $\text{add}(Y)$ -approximation of X (i.e., of a morphism $\psi: K \rightarrow X$, with K in $\text{add}(Y)$ such that for any Z in $\text{add}(Y)$, $\text{Hom}_{\mathcal{D}^b(R)}(Z, \psi)$ is surjective).

We say that a mutation is *irreducible* if X is indecomposable.

THEOREM 2.11 ([18, Theorem 7.12]). *Irreducible silting mutations are compatible with the bijections established in Theorem 2.4. More precisely, if S_1, \dots, S_n are the simple objects in the heart of a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ and X_1, \dots, X_n are indecomposable Ext-projective objects in $\mathcal{D}^{\leq 0}$ such that $\text{Hom}(X_i, S_i) \neq 0$, for all $1 \leq i \leq n$, then the aisles corresponding to the silting objects $\mu_{X_i}^+(M)$ and $\mu_{X_i}^-(M)$ are $\mathcal{D}_{+,S_i}^{\leq 0}$ and $\mathcal{D}_{-,S_i}^{\leq 0}$, respectively.*

The question of whether these mutations are compatible with gluing is discussed in Section 6.

§3. Gluing of silting objects

The results in this section make explicit in a more general setting the bijection established in [1, Theorem 2.37], for Krull–Schmidt triangulated categories \mathcal{D} , between the equivalence classes of silting objects in \mathcal{D} containing i_*Y as a direct summand and the equivalence classes of silting objects in the triangle quotient $\mathcal{D}/\text{thick}(i_*Y)$. Here $\text{thick}(i_*Y)$ is the smallest thick triangulated subcategory of \mathcal{D} containing i_*Y . More precisely, the theorem below shows how to glue two silting objects of \mathcal{X} and \mathcal{Y} into a silting object of \mathcal{D} with respect to a recollement of the form (2.1) in a way that is compatible with the bijection between silting objects and co-t-structures of Theorem 2.3.

THEOREM 3.1. *Let \mathcal{R} be a recollement of \mathcal{D} of the form (2.1). Let X and Y be silting objects corresponding to co-t-structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in \mathcal{X} and \mathcal{Y} , respectively. Then the induced co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ in \mathcal{D} is associated with the silting object $Z = i_*Y \oplus K_X$, with K_X defined by the following triangle:*

$$i_*\beta_{\geq 1}i^!j_!X \rightarrow j_!X \rightarrow K_X \rightarrow (i_*\beta_{\geq 1}i^!j_!X)[1],$$

where $\beta_{\geq 1}$ is a (nonfunctorial) choice of truncation for the co-t-structure $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in \mathcal{Y} .

Proof. First observe that, since i_* is a fully faithful functor, i_*Y is partial silting and, moreover, it is easily checked to lie in the co-heart of the glued co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ in \mathcal{D} .

Let us consider the following (nonfunctorial) triangle associated with the co-t-structure in \mathcal{Y}

$$i_*\beta_{\geq 1}i^!j_!X \rightarrow i_*i^!j_!X \rightarrow i_*\beta_{\leq 0}i^!j_!X \rightarrow (i_*\beta_{\geq 1}i^!j_!X)[1]$$

and the following universal triangle of the recollement (applied to $j_!X$)

$$i_*i^!j_!X \rightarrow j_!X \rightarrow j_*X \rightarrow (i_*i^!j_!X)[1].$$

Hence, we get the following commutative diagram where the rows are triangles

$$\begin{array}{ccccccc}
i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & i_*i^!j_!X & \longrightarrow & i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\
\downarrow & & \downarrow & & & & \downarrow \\
i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & j_!X & \longrightarrow & K_X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\
\downarrow & & \downarrow & & & & \downarrow \\
i_*i^!j_!X & \longrightarrow & j_!X & \longrightarrow & j_*X & \longrightarrow & (i_*i^!j_!X)[1]
\end{array}$$

By the octahedral axiom, this can be completed to a commutative diagram

$$\begin{array}{ccccccc}
i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & i_*i^!j_!X & \longrightarrow & i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & j_!X & \longrightarrow & K_X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
i_*i^!j_!X & \longrightarrow & j_!X & \longrightarrow & j_*X & \longrightarrow & (i_*i^!j_!X)[1] \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & K_X & \longrightarrow & j_*X & \longrightarrow & (i_*\beta_{\leq 0}i^!j_!X)[1]
\end{array}$$

the rows of which are again triangles. We will now show that K_X lies in the co-heart of $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$. Indeed, it is clear that $i_*\beta_{\geq 1}i^!j_!X \in \mathcal{D}_{\geq 1}$ and, therefore, that $(i_*\beta_{\geq 1}i^!j_!X)[1]$ lies in $\mathcal{D}_{\geq 0}$. Clearly we also have that $j_!X$ lies in $\mathcal{D}_{\geq 0}$ and, thus, the second row of the diagram shows that K_X lies in $\mathcal{D}_{\geq 0}$. Similarly, since $i_*\beta_{\leq 0}i^!j_!X$ lies in $\mathcal{D}_{\leq 0}$ and j_*X lies in $\mathcal{D}_{\leq 0}$, it follows that K_X lies in $\mathcal{D}_{\leq 0}$, proving that it lies in the co-heart.

Finally, it is enough to observe that Z generates \mathcal{D} . It is clear that i_*Y and $j_!X$ generate \mathcal{D} since X and Y generate \mathcal{X} and \mathcal{Y} , respectively, and i_* and $j_!$ are fully faithful functors. But, by the second triangle of the diagram, $j_!X$ can be generated by K_X and an object in the image of i_* , which is generated by i_*Y . Therefore Z generates \mathcal{D} and thus it is silting. \square

REMARK 3.2. Note that the leftmost morphism $i_*\beta_{\geq 1}i^!j_!X \rightarrow j_!X$ of a triangle in Theorem 3.1 corresponds to the chosen morphism $\beta_{\geq 1}i^!j_!X \rightarrow i^!j_!X$ via the adjunction morphism

$$\mathrm{Hom}(i_*\beta_{\geq 1}i^!j_!X, j_!X) \xrightarrow{\simeq} \mathrm{Hom}(\beta_{\geq 1}i^!j_!X, i^!j_!X).$$

REMARK 3.3. In Krull–Schmidt categories (such as $\mathcal{D}^b(R)$ for an algebra, finite-dimensional over a field), the object Z constructed in the proof of the theorem is not necessarily a basic object (i.e., the indecomposable summands of Z can appear with multiplicities) even if both X and Y are basic. We can, however, obtain a basic silting object in the corresponding co-heart by ignoring the multiplicities of the direct summands of Z .

Similarly, we can describe a gluing of silting objects that is compatible with the bijection between silting objects and t-structures for $\mathcal{D} = \mathcal{D}^b(R)$, where R has finite global dimension.

COROLLARY 3.4. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be silting objects corresponding to t-structures $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ in \mathcal{X} and \mathcal{Y} , respectively. Suppose that R has finite global dimension. Then the glued t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in \mathcal{D} is associated with the silting object $Z = j_!X \oplus K_Y$, with K_Y defined by the triangle*

$$j_!\alpha_{\geq 1}j^*i_{\#}Y \rightarrow i_{\#}Y \rightarrow K_Y \rightarrow (j_!\alpha_{\geq 1}j^*i_{\#}Y)[1],$$

where $\alpha_{\geq 1}$ is a (nonfunctorial) choice of truncation for the left adjacent co-t-structure of the t-structure $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ in \mathcal{X} .

Proof. This follows from Theorem 3.1 by observing, as in Remark 2.6 that $\mathcal{D}_{\leq 0}^U = \mathcal{D}^{\leq 0}$ and that the left adjacent co-t-structure of a t-structure corresponds to the same silting object. \square

The following definition settles what we will mean by *gluing silting objects*. As shown by Theorem 3.1, it is more natural, in this setting, to consider co-t-structures rather than t-structures.

DEFINITION 3.5. Let $\mathcal{D}, \mathcal{X}, \mathcal{Y}$ be triangulated categories, and let \mathcal{R} be a recollement of the form (2.1). We say that a silting object $Z \in \mathcal{D}$ is *glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to \mathcal{R}* if Z is obtained by the construction of Theorem 3.1, that is, Z corresponds to the co-t-structure glued from the co-t-structures associated to X and Y with respect to \mathcal{R} .

The object K_X of Theorem 3.1 can be described in a nonconstructive way as follows.

PROPOSITION 3.6. *Let \mathcal{R} be a recollement of a triangulated category \mathcal{D} of the form (2.1). Let X and Y be silting objects corresponding to co- t -structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in \mathcal{X} and \mathcal{Y} , respectively. Let $Z = i_*Y \oplus K_X$ be the silting object in \mathcal{D} glued from X and Y with respect to \mathcal{R} , compatible with the glued co- t -structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$. Then the following hold.*

- (1) *Object K_X is a right $(\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0})$ -approximation of j_*X .*
- (2) *Object K_X is a left $(\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0})$ -approximation of $j_!X$.*
- (3) *Up to summands in $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$, K_X is uniquely determined by the following conditions:*
 - (i) $j^*K_X = X$,
 - (ii) $i^*K_X \in \mathcal{Y}_{\geq 0}$,
 - (iii) $i^!K_X \in \mathcal{Y}_{\leq 0}$.

Proof. To prove (1), let C be an object of $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$, and let f be a morphism in $\text{Hom}_{\mathcal{D}}(C, j_*X)$. Since $(i_*\beta_{\leq 0}i^!j_!X)[1]$ lies in $\mathcal{D}_{\leq -1}$, it is clear that $\text{Hom}_{\mathcal{D}}(C, (i_*\beta_{\leq 0}i^!j_!X)[1]) = 0$ and thus, by using the defining triangle

$$i_*\beta_{\leq 0}i^!j_!X \rightarrow K_X \rightarrow j_*X \rightarrow (i_*\beta_{\leq 0}i^!j_!X)[1],$$

f factors through K_X , proving (1). Analogously, (2) can be shown using the defining triangle

$$i_*\beta_{\geq 1}i^!j_!X \rightarrow j_!X \rightarrow K_X \rightarrow (i_*\beta_{\geq 1}i^!j_!X)[1].$$

In order to prove (3) observe that, for any object L satisfying properties (i), (ii), and (iii), we have a canonical triangle coming from the recollement \mathcal{R}

$$i_*i^!L \rightarrow L \rightarrow j_*X \rightarrow i_*i^!L[1]$$

and, since $i_*(\mathcal{Y}_{\leq 0}) \subseteq \mathcal{D}_{\leq 0}$, we have that $i_*i^!L[1]$ lies in $\mathcal{D}_{\leq -1}$. Therefore, for any object C in $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$, any map from C to j_*X factors through L , proving that L is a right $(\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0})$ -approximation of j_*X . Hence, by (1), L differs from K_X by a summand in $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$. \square

§4. Gluing of tilting objects

In this section we investigate necessary and sufficient conditions for the glued silting to be tilting. These conditions will be expressed exclusively in terms of the functors of the recollement and of vanishing conditions on \mathcal{X} and \mathcal{Y} rather than in \mathcal{D} . In this section, R has finite global dimension and, thus, $\mathcal{D}^b(R)$ has a Serre functor. Under this assumption we can use Theorem 2.1 and the functors therein.

PROPOSITION 4.1. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be silting objects of \mathcal{X} and \mathcal{Y} , respectively. Then $Z = i_*Y \oplus K_X$ is tilting if and only if the following conditions are satisfied:*

- (1) Y is tilting,
- (2) $\mathrm{Hom}_{\mathcal{Y}}(Y, i^!j_!X[k]) = 0$ for all $k < 0$,
- (3) $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, Y[k]) = 0$ for all $k < 0$,
- (4) $\mathrm{Hom}_{\mathcal{X}}(X, j^+K_X[k]) = 0$ for all $k < 0$.

Proof. The induced silting object Z corresponds to the co-t-structure glued along \mathcal{R} and to the t-structure glued along \mathcal{R}_L from the respective structures on \mathcal{X} and \mathcal{Y} associated with X and Y , respectively. The statement that Z is tilting is equivalent to the statement that Z lies in the heart of $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$ which, by the description of the glued structures, translates into the conditions

- (i) $i^!Z \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$ or, equivalently, $Y, i^!K_X \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$,
- (ii) $j^*Z \in \mathcal{X}^{\leq 0}$,
- (iii) $j^+Z \in \mathcal{X}^{\geq 0}$.

To examine these conditions we will use the following triangle defining K_X

$$(4.1) \quad i_*\beta_{\leq 0}i^!j_!X \rightarrow K_X \rightarrow j_*X \rightarrow (i_*\beta_{\leq 0}i^!j_!X)[1].$$

Condition (i) is equivalent to Y being tilting (corresponding to condition (1) of the proposition) and $i^!K_X$ lying in $\mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$. By applying $i^!$ to the triangle (4.1), we get that $i^!K_X \cong \beta_{\leq 0}i^!j_!X$ lies in $\mathcal{Y}_{\leq 0} = \mathcal{Y}^{\leq 0}$. By Remark 2.7, $\beta_{\leq 0}$ corresponds to the stupid truncation in $\mathcal{D}^b(\mathrm{End}(Y))$ and, therefore, the fact that $i^!K_X$ lies in $\mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$ is equivalent to the condition that $i^!j_!X$ lies in $\mathcal{Y}^{\geq 0}$. This can then be translated to condition (2) of the proposition, by definition of the t-structure associated with a silting object.

Clearly, since j^* maps the co-heart in \mathcal{D} to the co-heart in \mathcal{X} , (ii) is automatically satisfied for any silting Z and, thus, this condition is irrelevant.

Condition (iii) is equivalent to $\text{Hom}_{\mathcal{X}}(X, j^+Z[k]) = \text{Hom}_{\mathcal{D}}(j_*X, Z[k]) = 0$ for all $k \leq 0$. By splitting Z into its summands i_*Y and K_X and applying adjunction, we get precisely conditions (3) and (4) of the proposition. \square


REMARK 4.2. To simplify our notation, we will use the fact (see [5, Section 1.4]) that $i^*j_* = i^!j_![1]$.

In many important cases, the left side of the recollement is just given by $\mathcal{D}^b(\mathbb{K})$.

COROLLARY 4.3. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$. Let X be a sifting object in \mathcal{X} , and let $Y = \mathbb{K}$. If $Z = i_*Y \oplus K_X$ is tilting, then there are finite-dimensional \mathbb{K} -vector spaces X'_{-1}, X'_0 such that $i^*j_*X \cong X'_{-1}[1] \oplus X'_0$.*

Proof. In view of Remark 4.2, condition (2) of Proposition 4.1 is equivalent to $H^k(i^*j_*X) = 0$ for all $k < -1$, since $Y = B = \mathbb{K}$. Since the Serre functor of $\mathcal{D}^b(\mathbb{K})$ is isomorphic to the identity functor, condition (3) of Proposition 4.1 is equivalent to $\text{Hom}_{\mathcal{Y}}(Y[k], i^*j_*X) = 0$ for all $k < 0$, which is equivalent to $H^k(i^*j_*X) = 0$ for all $k > 0$. To summarize, if Z is a tilting object, then it follows from Proposition 4.1 that $H^k(i^*j_*X) = 0$ for $k \neq -1, 0$, thus proving the corollary. \square

EXAMPLE 4.4. Let R be the \mathbb{K} -algebra given by the quiver with relations



The simple module S_1 supported at 1 is a partial tilting module of projective dimension 2. It has a minimal projective resolution over R given by the exact sequence

$$0 \rightarrow P_2 \xrightarrow{\beta} P_3 \xrightarrow{\delta} P_1 \rightarrow S_1 \rightarrow 0.$$

Rickard and Schofield showed in [25] that S_1 cannot be completed to a tilting module over R . We will strengthen this result by showing that S_1 cannot be completed to a tilting object in $\mathcal{D}^b(R)$.

If e is the idempotent $e_2 + e_3$ then, as a right A -module, R/ReR is isomorphic to S_1 , which does not have self-extensions. It therefore follows from [11, Theorem 3.1] and [12, Section 2] that there is a recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j!} & \\ \mathcal{D}^b(R/ReR) & \xrightarrow{i_*} & \mathcal{D}^b(R) & \xrightarrow{j^*} & \mathcal{D}^b(eRe), \\ & \xleftarrow{i!} & & \xleftarrow{j_*} & \end{array}$$

where

$$(4.2) \quad \begin{aligned} i^* &= -\bigotimes_R^{\mathbb{L}} R/ReR, & j! &= -\bigotimes_{eRe}^{\mathbb{L}} eR, \\ i_* &= -\bigotimes_{R/ReR}^{\mathbb{L}} R/ReR, & j^* &= -\bigotimes_R^{\mathbb{L}} Re, \\ i! &= \mathbb{R}\mathrm{Hom}_R(R/ReR, -), & j_* &= \mathbb{R}\mathrm{Hom}_{eRe}(Re, -). \end{aligned}$$

Fix $Y = R/ReR$, and let $X \in \mathcal{D}^b(eRe)$ be a silting object (e.g., $X = eRe$). Then Theorem 3.1 yields a completion $Z = i_*Y \oplus K_X$ of the partial silting object $i_*Y = S_1$ into a silting object. However, as we will show, Z is never tilting and, thus, S_1 cannot be completed to a tilting object.


Let T be a basic tilting object of $\mathcal{D}^b(R)$ which contains S_1 as a direct summand. The functor j^* factors through the canonical projection functor $\pi : \mathcal{D}^b(R) \rightarrow \mathcal{D}^b(R)/\mathrm{thick}(S_1)$, as follows:

$$\begin{array}{ccc} \mathcal{D}^b(R) & \xrightarrow{j^*} & \mathcal{D}^b(eRe) \\ & \searrow \pi & \nearrow \simeq \\ & \mathcal{D}^b(R)/\mathrm{thick}(S_1) & \end{array}$$

As $\mathrm{thick}(S_1)$ is a silting subcategory of $\mathcal{D}^b(R)$, by [1, Theorem 2.37], the object $X = j^*(T)$ is silting in $\mathcal{D}^b(eRe)$ and T is equivalent to $S_1 \oplus K_X$. Observe that eRe is the path algebra of the Kronecker quiver

$$2 \begin{array}{c} \xrightarrow{\gamma} \\ \xrightarrow{\beta} \end{array} 3.$$

We draw the component of the AR quiver of $\mathcal{D}^b(eRe)$ containing the pre-projective modules



where P'_2 and P'_3 are, respectively, the indecomposable projective eRe -module at the vertex 2 and 3. Let $\tau = \tau_{eRe}$ denote the Auslander–Reiten translation of $\mathcal{D}^b(eRe)$. Then according to [26, Lemma 3.1], there exist p and q such that $\tau^p P'_2[q]$ is a direct summand of K_X . We claim that

$$H^n(i^* j_*(\tau^p P'_2[q])) = \begin{cases} \mathbb{K} & \text{if } n = -q - 1, -q + 1, \\ 0 & \text{otherwise.} \end{cases}$$

As a consequence of Corollary 4.3, $S_1 \oplus K_X$ cannot be a tilting object, which is a contradiction.

The claim follows from a direct computation, which we show only for $P'_{2,1} = \tau^{-1} P'_2$ (it is analogous for others). For $n \in \mathbb{Z}$, we have

$$\begin{aligned}
 H^n(i^* j_*(P'_{2,1})) &= H^{n+1}(i^! j_!(P'_{2,1})) \\
 &= H^{n+1}(\mathbb{R}\mathrm{Hom}_R(R/ReR, j_!(P'_{2,1}))) \\
 &= H^{n+1}(\mathbb{R}\mathrm{Hom}_R(S_1, j_!(P'_{2,1}))) \\
 &= \mathrm{Hom}_{\mathcal{D}^b(R)}(S_1, j_!(P'_{2,1})[n+1]).
 \end{aligned}$$

We take a minimal projective resolution of $P'_{2,1}$

$$P'_2 \xrightarrow{\begin{pmatrix} \gamma \\ \beta \end{pmatrix}} \underline{P'_3 \oplus P'_3},$$

where the underlined term is in degree 0. Applying $j_!$ we obtain the following object in $\mathcal{D}^b(R)$:

$$P_2 \xrightarrow{\begin{pmatrix} \gamma \\ \beta \end{pmatrix}} \underline{P_3 \oplus P_3}.$$

The above differential is injective and its cokernel has the following minimal injective resolution:

$$\underline{I_2 \oplus I_2 \oplus I_1} \rightarrow I_3 \oplus I_3 \oplus I_3 \rightarrow I_1 \rightarrow I_2 \rightarrow I_3.$$

Therefore $j_!(P'_{2,1})$ is isomorphic to this complex in $\mathcal{D}^b(R)$ and it follows that

$$\mathrm{Hom}_R(S_1, j_!(P'_{2,1})[n+1]) = \begin{cases} \mathbb{K} & \text{if } n = -1, 1, \\ 0 & \text{otherwise.} \end{cases}$$

In using Proposition 4.1 we do not seem to be dealing with computations in \mathcal{D} . Still, these occur in the construction of K_X . This can be avoided by introducing the assumption that X is tilting.

THEOREM 4.5. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of \mathcal{X} and \mathcal{Y} , respectively. Then $Z = i_*Y \oplus K_X$ is tilting in \mathcal{D} if and only if the following conditions are satisfied:*

- (a) $\mathrm{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k]) = 0$ for all $k < -1$,
- (b) $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, Y[k]) = 0$ for all $k < 0$,
- (c) $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0$ for all $k < -1$.

Proof. Condition (a) is obtained from Proposition 4.1(2) and from Remark 4.2, while condition (b) corresponds exactly to Proposition 4.1(3). We will show that if X is tilting and (a) and (b) are satisfied, then Proposition 4.1(4) is equivalent to (c). Indeed, if X is tilting, $\mathrm{Hom}_{\mathcal{D}}(j_*X, j_*X[k]) = 0$ for all $k \neq 0$ and thus, applying the functor $\mathrm{Hom}_{\mathcal{D}}(j_*X, -)$ to the k th shift of triangle (4.1), we get $\mathrm{Hom}_{\mathcal{D}}(j_*X, K_X[k]) \cong \mathrm{Hom}_{\mathcal{D}}(j_*X, (i_*\beta_{\leq 0}i^!j_!X)[k])$ for all $k \neq 0, 1$. Since (j_*, j^+) is an adjoint pair, this shows that Proposition 4.1(4) is equivalent to

$$\mathrm{Hom}_{\mathcal{D}}(j_*X, (i_*\beta_{\leq 0}i^!j_!X)[k]) = \mathrm{Hom}_{\mathcal{D}}(i^*j_*X, (\beta_{\leq 0}i^!j_!X)[k]) = 0, \quad \forall k < 0.$$

Applying the functor $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, -)$ to the triangle

$$(\beta_{\geq 1}i^!j_!X)[k] \rightarrow i^!j_!X[k] \rightarrow (\beta_{\leq 0}i^!j_!X)[k] \rightarrow (\beta_{\geq 1}i^!j_!X)[k+1],$$

we get that $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, \beta_{\leq 0}i^!j_!X[k]) \cong \mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, i^!j_!X[k])$, for all $k < 0$. Indeed, this follows from (b) after recalling that $\beta_{\geq 1}i^!j_!X$ lies in $\mathcal{Y}_{\geq 1}$, which by construction (see Section 2.3) is suitably generated by $(Y[n])_{n < 0}$, showing that $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, (\beta_{\geq 1}i^!j_!X)[k+1]) = 0$ for all $k < 0$. Using again Remark 4.2, we obtain condition (c), thus finishing the proof. \square

Before showing applications of this theorem, we discuss the behavior of the Serre functor with respect to a t-structure corresponding to a tilting object. Recall from [5, Proposition 3.1.10] that for any bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$ with heart \mathcal{A} , there is a triangle functor $\mathrm{real} :$

$\mathcal{D}^b(\mathcal{A}) \rightarrow \mathcal{D}^b(R)$, t-exact for the standard t-structure $(\mathcal{D}_{\text{st}}^{\leq 0}, \mathcal{D}_{\text{st}}^{\geq 0})$ in $\mathcal{D}^b(\mathcal{A})$ and $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, that is, it is both right t-exact ($\text{real}(\mathcal{D}_{\text{st}}^{\leq 0}) \subseteq \mathcal{D}^{\leq 0}$) and left t-exact ($\text{real}(\mathcal{D}_{\text{st}}^{\geq 0}) \subseteq \mathcal{D}^{\geq 0}$).

LEMMA 4.6. *Let $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ be a bounded t-structure in $\mathcal{D}^b(R)$ with associated basic sifting object T . Let S_R be the Serre functor of $\mathcal{D}^b(R)$. Then the following are equivalent:*

- (1) T is tilting,
- (2) S_R is right t-exact,
- (3) $S_R(T)$ lies in the heart.

Proof. (1) \Rightarrow (2): Suppose that T is tilting. Let $\Gamma = \text{End}_{\mathcal{D}^b(R)}(T)$, and let $(\mathcal{D}_{\text{st}}^{\leq 0}, \mathcal{D}_{\text{st}}^{\geq 0})$ be the standard t-structure in $\mathcal{D}^b(\Gamma)$. Then, by [27, Theorem 6.6], the realization functor is a t-exact equivalence. Therefore, $\text{real} \circ S_\Gamma = S_R \circ \text{real}$ (since Serre functors are unique) and $\text{real}(\mathcal{D}_{\text{st}}^{\leq 0}) = \mathcal{D}^{\leq 0}$. Note also that S_Γ preserves the standard aisle in $\mathcal{D}^b(\Gamma)$ since it can be realized as a derived tensor product with a bimodule. This is enough to observe that S_R is right t-exact, since

$$S_R(\mathcal{D}^{\leq 0}) = S_R(\text{real}(\mathcal{D}_{\text{st}}^{\leq 0})) = \text{real}(S_\Gamma(\mathcal{D}_{\text{st}}^{\leq 0})) \subseteq \text{real}(\mathcal{D}_{\text{st}}^{\leq 0}) = \mathcal{D}^{\leq 0}.$$

(2) \Rightarrow (3): For any $K \in \mathcal{D}^{\leq -1}$, $\text{Hom}_{\mathcal{D}^b(R)}(K, S_RT) = \text{Hom}_{\mathcal{D}^b(R)}(T, K) = 0$, by definition of the t-structure associated to T (see Section 2.3). This shows that $S_RT \in \mathcal{D}^{\geq 0}$. If S_R is right t-exact, then $S_R(T) \in \mathcal{D}^{\leq 0}$ and, thus, $S_R(T)$ lies in the heart.

(3) \Rightarrow (1): Suppose that $S_R(T)$ lies in the heart. By the definition of the t-structure associated to T , it is clear that $\text{Hom}(T, T[k]) = \text{Hom}(T, S_RT[-k]) = 0$ for all $k \neq 0$. Therefore T is tilting. \square

PROPOSITION 4.7. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of \mathcal{X} and \mathcal{Y} , respectively. If i^*j_*X is an element of $\mathcal{D}^b(B)$ such that $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k])$ is zero except for two consecutive values of $k \in \mathbb{Z}$, then the family of sifting objects $(Z_n = i_*Y \oplus K_{X[n]})_{n \in \mathbb{Z}}$ contains at least one tilting object.*

Proof. Let $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ be the t-structure associated with Y . By definition of the t-structure associated to Y , $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k])$ is nonzero exactly for two consecutive values of $k \in \mathbb{Z}$ if and only if there is a in \mathbb{Z} such that $i^*j_*X \in \mathcal{Y}^{\geq a} \cap \mathcal{Y}^{\leq a+1}$. Since Y is tilting, by Lemma 4.6 we have that $S_{\mathcal{Y}}i^*j_*X$ lies in $\mathcal{Y}^{\leq a+1}$. In Theorem 4.5, condition (a) can be reformulated as $i^*j_*X \in \mathcal{Y}^{\geq -1}$

and condition (b) as $S_{\mathcal{Y}}i^*j_*X \in \mathcal{Y}^{\leq 0}$. Note that once condition (c) is satisfied for some $X[n]$, it is satisfied for all its shifts. By Theorem 4.5, Z_n is tilting if and only if

- (a) $i^*j_*X[n] \in \mathcal{Y}^{\geq -1}$, which means that $a - n \geq -1$ or, equivalently, $n \leq a + 1$;
- (b) $S_{\mathcal{Y}}i^*j_*X[n] \in \mathcal{Y}^{\leq 0}$, which is guaranteed if $a + 1 - n \leq 0$ or, equivalently, $n \geq a + 1$;
- (c) $\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0$ for all $k < -1$.

Therefore, take $n = a + 1$, and we only need to check condition (c). This condition is, however, automatic from our assumption that i^*j_*X has cohomologies with respect to $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ concentrated in exactly two consecutive degrees, thus finishing the proof. \square

REMARK 4.8. In the family $(Z_n)_{n \in \mathbb{Z}}$ there may be more than one tilting object. Let a be the maximal integer such that $i^*j_*X \in \mathcal{Y}^{\geq a}$, and let b be the minimal integer such that $S_{\mathcal{Y}}i^*j_*X \in \mathcal{Y}^{\leq b}$. Then Z_n is tilting if and only if $b \leq n \leq a + 1$.

From the proof above, we see that, for X and Y tilting, whenever i^*j_*X lies in $\mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq -1}$ (i.e., the cohomologies of i^*j_*X with respect to Y lie in degrees -1 and 0), the induced silting $Z = i_*Y \oplus K_X$ is tilting. Since $i^*j_* = i^!j_![1]$, the condition is equivalent to that $i^!j_!X \in \mathcal{Y}^{\leq 1} \cap \mathcal{Y}^{\geq 0}$ which, in its turn, is equivalent to $\mathrm{Hom}_{\mathcal{Y}}(Y, i^!j_!X[k]) \cong \mathrm{Hom}_{\mathcal{D}}(i_*Y, j_!X[k]) = 0$ whenever $k \neq 0, 1$. Under this assumption, we can apply the construction from [2, Theorem 2.5] to the pair $T_1 = j_!X$ and $T_2 = i_*Y$. A similar construction was studied in [19] in the setting of triangular matrix rings.

We briefly recall the construction in [2]. Since $(j_!, j^*)$ is an adjoint pair in the recollement (of the form (2.1)), $\mathrm{Hom}_{\mathcal{D}}(j_!X, i_*Y[k]) \cong \mathrm{Hom}_{\mathcal{X}}(X, j^*i_*Y[k]) = 0$ for all $k \in \mathbb{Z}$. Let m be the dimension of $\mathrm{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$, and take a basis $\alpha_1, \dots, \alpha_m : i_*Y \rightarrow j_!X[1]$. Consider the universal maps

$$\begin{aligned} \alpha &= (\alpha_1, \dots, \alpha_m)^{\mathrm{tr}} : i_*Y^{\oplus m} \rightarrow j_!X[1], \\ \beta &= (\alpha_1, \dots, \alpha_m) : i_*Y \rightarrow j_!X[1]^{\oplus m}. \end{aligned}$$

The map α is left-universal, that is, the induced map

$$\mathrm{Hom}(i_*Y, \alpha) : \mathrm{Hom}_{\mathcal{D}}(i_*Y, i_*Y^{\oplus m}) \rightarrow \mathrm{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$$

is surjective. Similarly, the map β is right-universal, that is, the induced map

$$\mathrm{Hom}(\beta, j_!X[1]) : \mathrm{Hom}_{\mathcal{D}}(j_!X[1]^{\oplus m}, j_!X[1]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$$

is surjective. Consider the triangles determined by α and β :

$$\begin{aligned} T_\alpha : j_!X &\rightarrow C_1 \rightarrow i_*Y^{\oplus m} \xrightarrow{\alpha} j_!X[1], \\ T_\beta : j_!X^{\oplus m} &\rightarrow C_2 \rightarrow i_*Y \xrightarrow{\beta} j_!X[1]^{\oplus m}. \end{aligned}$$

Theorem 2.5 in [2] asserts that $i_*Y \oplus C_1$ and $j_!X \oplus C_2$ are tilting objects in \mathcal{D} .

The next proposition proves that the tilting object $i_*Y \oplus C_1$ is precisely the silting (indeed tilting) glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to the recollement \mathcal{R} , and $j_!X \oplus C_2$ is the silting (indeed tilting) glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to the upper reflection \mathcal{R}_U . As a consequence, under the conditions of Theorem 4.5, we are not only able to glue tilting objects in the original recollement \mathcal{R} , but also in the upper reflection \mathcal{R}_U .

PROPOSITION 4.9. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of \mathcal{X} and \mathcal{Y} , respectively, such that $\mathrm{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k]) = 0$ for all $k \neq 0, -1$. Then, up to multiplicity, the tilting object $i_*Y \oplus C_1$ coincides with $Z = i_*Y \oplus K_X$ of Theorem 3.1, and the tilting object $j_!X \oplus C_2$ coincides with $Z_U = j_!X \oplus K_Y$ of Corollary 3.4.*

Proof. By hypothesis, i^*j_*X lies in $\mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq -1}$, for the t-structure $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ associated with Y . We will prove that C_1 satisfies the conditions (i)–(iii) for K_X in Proposition 3.6(3). It follows then that $i_*Y \oplus C_1$ lies in the glued co-heart of \mathcal{D} and that it generates $i_*Y \oplus j_!X$, thus generating the whole of \mathcal{D} . Therefore, up to multiplicities, it coincides with the glued silting $Z = i_*Y \oplus K_X$.

For condition (i), we apply j^* to the defining triangle T_α of C_1 and obtain that $j^*C_1 = j^*j_!X = X$, since $j^*i_* = 0$. For condition (ii), we apply i^* to T_α and obtain that $i^*C_1 = i^*i_*Y^{\oplus m} = Y^{\oplus m} \in \mathcal{Y}_{\geq 0}$, since $i^*j_! = 0$. For condition (iii), we apply $i^!$ to T_α and obtain a triangle

$$i^!j_!X \rightarrow i^!C_1 \rightarrow i^!i_*Y^{\oplus m} \xrightarrow{i^!\alpha} i^!j_!X[1].$$

By applying $\mathrm{Hom}_{\mathcal{Y}}(Y, -)$ to this triangle, we obtain a long exact sequence

$$\begin{aligned} \cdots &\rightarrow \mathrm{Hom}_{\mathcal{Y}}(Y, i^!i_*Y^{\oplus m}) \xrightarrow{(i^!\alpha)_*} \mathrm{Hom}_{\mathcal{Y}}(Y, i^!j_!X[1]) \rightarrow \mathrm{Hom}_{\mathcal{Y}}(Y, i^!C_1[1]) \\ &\rightarrow \mathrm{Hom}_{\mathcal{Y}}(Y, i^!i_*Y[1]^{\oplus m}) \rightarrow \cdots \end{aligned}$$

For $k > 0$, we have $\mathrm{Hom}_{\mathcal{Y}}(Y, i^! i_* Y[k]^{\oplus m}) \simeq \mathrm{Hom}_{\mathcal{Y}}(Y, Y[k]^{\oplus m}) = 0$ (since, by assumption, Y is tilting) and for $k > 1$, $\mathrm{Hom}_{\mathcal{Y}}(Y, i^! j_! X[k]) = 0$ (by our assumption on $i^* j_* X = i^! j_! X[1]$). Hence, $\mathrm{Hom}_{\mathcal{Y}}(Y, i^! C_1[k]) = 0$, for all $k > 1$. By adjunction, there is a commutative diagram

$$\begin{array}{ccc} (i^! \alpha)_* : & \mathrm{Hom}_{\mathcal{Y}}(Y, i^! i_* Y^{\oplus m}) & \longrightarrow & \mathrm{Hom}_{\mathcal{Y}}(Y, i^! j_! X[1]) \\ & \simeq \uparrow & & \simeq \uparrow \\ \alpha_* : & \mathrm{Hom}_{\mathcal{D}}(i_* Y, i_* Y^{\oplus m}) & \longrightarrow & \mathrm{Hom}_{\mathcal{D}}(i_* Y, j_! X[1]) \end{array}$$

Since, by construction, α is left-universal, that is, α_* is surjective, it follows that $(i^! \alpha)_*$ is surjective. Hence, we get $\mathrm{Hom}_{\mathcal{Y}}(Y, i^! C_1[1]) = 0$ and, thus, $i^! C_1$ lies in $\mathcal{Y}^{\leq 0} = \mathcal{Y}_{\leq 0}$. The proof for the statement about $j_! X \oplus C_2$ is analogous. \square

§5. The hereditary case

In this section we assume that R has finite global dimension, and we apply Theorem 4.5 when B is a hereditary algebra. The conditions for the glued silting to be tilting are then easier to handle.

PROPOSITION 5.1. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$, $\mathcal{Y} = \mathcal{D}^b(B)$, and B hereditary. Let X be a tilting object in \mathcal{X} , and let $Y = B$. Then $Z = i_* Y \oplus K_X$ is tilting in \mathcal{D} if and only if there are finitely generated B -modules X'_{-1} , X'_0 , and X'_1 such that*

- $i^* j_* X$ is isomorphic to $X'_{-1}[1] \oplus X'_0 \oplus X'_1[-1]$,
- X'_1 is either zero or not projective,
- $\mathrm{Hom}_B(X'_1, X'_{-1}) = 0$.

Proof. We analyze the conditions of Theorem 4.5 when B is hereditary and $Y = B$. Since B is hereditary, we may assume that $i^* j_* X = \bigoplus_{n \in \mathbb{Z}} X'_n[-n]$, with X'_n a finitely generated right B -module for all n in \mathbb{Z} . In this setting, $\mathrm{Hom}_{\mathcal{Y}}(Y, i^* j_* X[k]) = 0$, for all $k < -1$, is equivalent to

$$\mathrm{Hom}_{\mathcal{D}^b(B)}\left(B, \bigoplus_{n \in \mathbb{Z}} X'_n[-n+k]\right) = 0 \quad \text{for all } k < -1,$$

which happens if and only if $X'_k = 0$, for all $k < -1$. On the other hand, the condition

$$\mathrm{Hom}_{\mathcal{Y}}(i^* j_* X, Y[k]) = \mathrm{Hom}_{\mathcal{D}^b(B)}(i^* j_* X, B[k]) = 0, \quad \text{for all } k < 0$$

can be reformulated, using the Serre functor S_B in $\mathcal{D}^b(B)$, by

$$\mathrm{Hom}_{\mathcal{D}^b(B)}(B, S_B(i^*j_*X)[k]) = 0, \quad \text{for all } k > 0.$$

It is well known that $S_B = [1] \circ \tau_B$, where τ_B is the Auslander–Reiten translation in $\mathcal{D}^b(B)$ and, thus, the statement above is equivalent to

$$\mathrm{Hom}_{\mathcal{D}^b(B)}\left(B, \bigoplus_{n \in \mathbb{Z}} \tau_B(X'_n)[-n + k + 1]\right) = 0, \quad \forall k > 0,$$

that is, $X'_k = 0$, for all $k > 1$, and X'_1 is not a projective B -module. So, (a) and (b) of Theorem 4.5 hold if and only if $i^*j_*X = X'_{-1}[1] \oplus X'_0 \oplus X'_1[-1]$, with X'_1 not projective as a B -module. Assuming this, we can easily unfold the last condition of Theorem 4.5 as

$$\mathrm{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0, \quad \forall k < -1 \quad \Leftrightarrow \quad \mathrm{Hom}_{\mathcal{D}^b(B)}(X'_1, X'_{-1}) = 0,$$

thus finishing the proof. □

As a nice corollary, we obtain a necessary and sufficient condition for the case $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$.

COROLLARY 5.2. *Let \mathcal{R} be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$. Let X be a tilting object in \mathcal{X} , and let $Y = \mathbb{K}$. Then $Z = i_*Y \oplus K_X$ is tilting if and only if there are finite-dimensional \mathbb{K} -vector spaces X'_{-1}, X'_0 such that $i^*j_*X \cong X'_{-1}[1] \oplus X'_0$.*

EXAMPLE 5.3. Let R be the path algebra over \mathbb{K} of the quiver $1 \rightarrow 2$, of type A_2 . Let $e = e_1$ be the trivial path at the vertex 1. Consider the following standard recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j!} & \\ \mathcal{D}^b(R/ReR) & \xrightarrow{i_*} & \mathcal{D}^b(R) & \xrightarrow{j^*} & \mathcal{D}^b(eRe), \\ & \xleftarrow{i!} & & \xleftarrow{j_*} & \end{array}$$

where the six functors are defined as in (4.2). Let D denote the functor $\mathrm{Hom}_{\mathbb{K}}(-, \mathbb{K})$. It is easy to observe that

- (1) as algebras, both R/ReR and eRe are isomorphic to \mathbb{K} ;
- (2) as an R -module, eR is the simple module S_1 , supported at the vertex 1, and has an injective resolution ($D(Re) \rightarrow D(R(1 - e))$);
- (3) as an R -module, R/ReR is the simple module S_2 supported at the vertex 2.

Fix $Y = R/ReR$ and $X = eRe$. Then, in $\mathcal{D}^b(R)$, we have

$$\begin{aligned} i^!j_!X &= i^!(eR) = \mathbb{R}\mathrm{Hom}_R(R/ReR, eR) \\ &\cong (\mathrm{Hom}_R(R/ReR, D(Re)) \rightarrow \mathrm{Hom}_R(R/ReR, D(R(1-e)))) \\ &\cong R/ReR[-1]. \end{aligned}$$

By Corollary 5.2, there are exactly two tilting objects in $\{Z_n = i_*Y \oplus K_{X[n]} : n \in \mathbb{Z}\}$. We construct this family using Theorem 3.1. For $n \in \mathbb{Z}$, we can choose the map $\beta_{\geq 1}(i^!j_!X[n]) \rightarrow i^!j_!X[n]$ to be

$$\begin{cases} R/ReR[n-1] \xrightarrow{\mathrm{id}} R/ReR[n-1] & \text{if } n \leq 0, \\ 0 \rightarrow R/ReR[n-1] & \text{if } n > 0. \end{cases}$$

By adjunction, we have a bijection

$$\mathrm{Hom}(i_*\beta_{\geq 1}(i^!j_!X[n]), j_!X[n]) \xrightarrow{\cong} \mathrm{Hom}(\beta_{\geq 1}(i^!j_!X[n]), i^!j_!X[n]).$$

If $n \geq 0$, this adjunction morphism is

$$\mathrm{Hom}_{\mathcal{D}^b(R)}(S_2[n-1], S_1[n]) \xrightarrow{\cong} \mathrm{Hom}_{\mathcal{D}^b(R)}(R/ReR[n-1], R/ReR[n-1]).$$

Both these vector spaces are 1-dimensional over \mathbb{K} . Let f be the preimage of $\mathrm{id}_{R/ReR[n-1]}$. Then the cone of $f[-n]$ is $(1-e)R$, the projective cover P_2 of S_2 . If $n < 0$, the above adjunction morphism is

$$\mathrm{Hom}_{\mathcal{D}^b(R)}(0, S_1[n]) \xrightarrow{\cong} \mathrm{Hom}_{\mathcal{D}^b(R)}(0, R/ReR[n-1]).$$

Thus a morphism $i_*\beta_{\geq 1}(i^!j_!X[n]) \rightarrow j_!X[n]$ as in Theorem 3.1 is (see Remark 3.2)

$$\begin{cases} S_2[n-1] \xrightarrow{f} S_1[n] & \text{if } n \leq 0, \\ 0 \rightarrow S_1[n] & \text{if } n > 0. \end{cases}$$

Hence, we have

$$K_{X[n]} = \begin{cases} P_2[n] & \text{if } n \leq 0, \\ S_1[n] & \text{if } n > 0, \end{cases} \quad \text{and} \quad Z_n = \begin{cases} S_2 \oplus P_2[n] & \text{if } n \leq 0, \\ S_2 \oplus S_1[n] & \text{if } n > 0. \end{cases}$$

Among these silting objects, $Z_0 = S_2 \oplus P_2$ and $Z_1 = S_2 \oplus S_1[1]$ are tilting objects.

§6. HRS-tilts and recollements

In this section we show that HRS-tilts of t-structures with respect to torsion theories (see [14]) are *compatible* with the gluing of t-structures via recollements. The main results of this section are Theorem 6.4 and Proposition 6.5. Our notation is fixed as follows:

- \mathcal{D} is a triangulated category admitting a recollement \mathcal{R} of the form (2.1),
- $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ are bounded t-structures in \mathcal{Y} and \mathcal{X} , respectively,
- $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued from $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ (see Theorem 2.2).

In fact (see [5], [20]), $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued with respect to \mathcal{R} if and only if $j_!j^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{D}^{\leq 0}$ (or, equivalently, if and only if $j_*j^*(\mathcal{D}^{\geq 0}) \subseteq \mathcal{D}^{\geq 0}$) in which case the *restrictions* to \mathcal{Y} and \mathcal{X} satisfy

$$\begin{aligned} \mathcal{Y}^{\leq 0} &= i^*(\mathcal{D}^{\leq 0}), & \mathcal{Y}^{\geq 0} &= i^!(\mathcal{D}^{\geq 0}), \\ \mathcal{X}^{\leq 0} &= j^*(\mathcal{D}^{\leq 0}), & \mathcal{X}^{\geq 0} &= j^*(\mathcal{D}^{\geq 0}). \end{aligned}$$

Let $\mathcal{A}_{\mathcal{D}}$, $\mathcal{A}_{\mathcal{Y}}$, and $\mathcal{A}_{\mathcal{X}}$ be the hearts of these t-structures. In [5, Section 1.4], it is shown that there is a *recollement of abelian categories* at the level of hearts

$$(6.1) \quad \begin{array}{ccc} & \xleftarrow{p_{i^*}} & \xleftarrow{p_{j_!}} \\ \mathcal{A}_{\mathcal{Y}} & \xrightarrow{p_{i_*}} \mathcal{A}_{\mathcal{D}} & \xrightarrow{p_{j^*}} \mathcal{A}_{\mathcal{X}} \\ & \xleftarrow{p_{i^!}} & \xleftarrow{p_{j_*}} \end{array}$$

We describe these functors explicitly. Let $\epsilon_{\mathcal{D}} : \mathcal{A}_{\mathcal{D}} \hookrightarrow \mathcal{D}$ denote the full embedding (similarly $\epsilon_{\mathcal{Y}} : \mathcal{A}_{\mathcal{Y}} \hookrightarrow \mathcal{Y}$ and $\epsilon_{\mathcal{X}} : \mathcal{A}_{\mathcal{X}} \hookrightarrow \mathcal{X}$), and let $H_{\mathcal{D}}^i : \mathcal{D} \rightarrow \mathcal{A}_{\mathcal{D}}$, $i \in \mathbb{Z}$, denote the cohomological functors with respect to the fixed t-structure in \mathcal{D} (similarly $H_{\mathcal{Y}}^i$ and $H_{\mathcal{X}}^i$ for the t-structures in \mathcal{Y} and \mathcal{X}). Then the functors in the recollement (6.1) are given by

$$\begin{aligned} p_{i^*} &= H_{\mathcal{Y}}^0 \circ i^* \circ \epsilon_{\mathcal{D}}, & p_{i^!} &= H_{\mathcal{Y}}^0 \circ i^! \circ \epsilon_{\mathcal{D}}, & p_{i_*} &= H_{\mathcal{D}}^0 \circ i_* \circ \epsilon_{\mathcal{Y}}, \\ p_{j_!} &= H_{\mathcal{D}}^0 \circ j_! \circ \epsilon_{\mathcal{X}}, & p_{j_*} &= H_{\mathcal{D}}^0 \circ j_* \circ \epsilon_{\mathcal{X}}, & p_{j^*} &= H_{\mathcal{X}}^0 \circ j^* \circ \epsilon_{\mathcal{D}}. \end{aligned}$$

See [13] for more on recollements of abelian categories.

REMARK 6.1. Since for our fixed t-structures, i_* and j^* are t-exact (see, e.g., [20] for details), we have that $p_{i_*} = i_* \circ \epsilon_{\mathcal{Y}}$ and $p_{j^*} = j^* \circ \epsilon_{\mathcal{D}}$. As a consequence, these two functors are exact.

Moreover, for any object A in $\mathcal{A}_{\mathcal{D}}$ there are exact sequences

$$(6.2) \quad 0 \rightarrow {}^p i_* H_{\mathcal{Y}}^{-1} {}^p i^* A \rightarrow {}^p j_! {}^p j^* A \rightarrow A \rightarrow {}^p i_* {}^p i^* A \rightarrow 0,$$

$$(6.3) \quad 0 \rightarrow {}^p i_* {}^p i^! A \rightarrow A \rightarrow {}^p j_* {}^p j^* A \rightarrow {}^p i_* H_{\mathcal{Y}}^1 {}^p i^* A \rightarrow 0.$$

Torsion pairs in $\mathcal{A}_{\mathcal{D}}$ satisfying certain conditions can be *restricted* to torsion pairs in $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$.

LEMMA 6.2. *Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$. Then*

- (1) $({}^p j^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{Y}}$;
- (2) the following are equivalent:
 - (i) $({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{X}}$,
 - (ii) ${}^p j_! {}^p j^*(\mathcal{T}) \subseteq \mathcal{T}$,
 - (iii) ${}^p j_* {}^p j^*(\mathcal{F}) \subseteq \mathcal{F}$.

Proof. Item (1): We first check the orthogonality condition for $({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$. We have that

$$\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F})) = \mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i_* {}^p i^*(\mathcal{T}), {}^p i_* {}^p i^!(\mathcal{F})),$$

since ${}^p i_*$ is fully faithful. Consider the exact sequence (6.2) with A in \mathcal{T} and apply to it the functor $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}(-, B)$ for some B in \mathcal{F} . Since $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}(A, B) = 0$, we conclude that $\mathrm{Hom}({}^p i_* {}^p i^*(A), B) = 0$, that is, ${}^p i_* {}^p i^*(\mathcal{T}) \subseteq \mathcal{T}$. Similarly, we have ${}^p i_* {}^p i^!(\mathcal{F}) \subseteq \mathcal{F}$ and, thus, $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i_* {}^p i^*(\mathcal{T}), {}^p i_* {}^p i^!(\mathcal{F})) = 0$, as wanted.

Second, we produce a suitable short exact sequence for any object in $\mathcal{A}_{\mathcal{D}}$. By [5, Remarque 1.4.17.1], ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ is a Serre subcategory of $\mathcal{A}_{\mathcal{D}}$, that is, for any short exact sequence in $\mathcal{A}_{\mathcal{D}}$

$$0 \rightarrow K \rightarrow M \rightarrow L \rightarrow 0,$$

M lies in the ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ if and only if K and L lie there also. Let M lie in $\mathcal{A}_{\mathcal{Y}}$, and consider a short exact sequence

$$0 \rightarrow \widetilde{M}_1 \rightarrow {}^p i_*(M) \rightarrow \widetilde{M}_2 \rightarrow 0$$

in $\mathcal{A}_{\mathcal{D}}$ with $\widetilde{M}_1 \in \mathcal{T}$ and $\widetilde{M}_2 \in \mathcal{F}$ (which exists since $(\mathcal{T}, \mathcal{F})$ is a torsion pair in \mathcal{D}). Since ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ is a Serre subcategory of $\mathcal{A}_{\mathcal{D}}$, there exist M_1 and M_2 in $\mathcal{A}_{\mathcal{Y}}$ such that $\widetilde{M}_i = {}^p i_*(M_i)$ for $i = 1, 2$. Now, ${}^p i_*$ is an exact full embedding (see Remark 6.1) and, thus, we get a short exact sequence in $\mathcal{A}_{\mathcal{Y}}$

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0,$$

with $M_1 \cong {}^{p_i^*}p_i^*(M_1) = {}^{p_i^*}(\widetilde{M}_1)$ lying in ${}^{p_i^*}(\mathcal{T})$ and $M_2 \cong {}^{p_i^!}p_i^*(M_2) = {}^{p_i^!}(\widetilde{M}_2)$ lying in ${}^{p_i^!}(\mathcal{F})$.

Item (2): By Remark 6.1, the functor ${}^{p_j^*} : \mathcal{A}_{\mathcal{D}} \rightarrow \mathcal{A}_{\mathcal{X}}$ is exact. Hence, $({}^{p_j^*}(\mathcal{T}), {}^{p_j^*}(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{X}}$ if and only these classes are orthogonal, that is, $\text{Hom}_{\mathcal{A}_{\mathcal{X}}}({}^{p_j^*}(\mathcal{T}), {}^{p_j^*}(\mathcal{F})) = 0$. Using the adjunctions $({}^{p_j!}, {}^{p_j^*})$ and $({}^{p_j^*}, {}^{p_j*})$, this holds if and only if $\text{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^{p_j!}{}^{p_j^*}(\mathcal{T}), \mathcal{F}) = 0$ (equivalent to (ii)) and if and only if $\text{Hom}_{\mathcal{A}_{\mathcal{D}}}(\mathcal{T}, {}^{p_j*}{}^{p_j^*}(\mathcal{F})) = 0$ (equivalent to (iii)). \square

When the conditions of item (2) of the lemma are fulfilled, we say that $(\mathcal{T}, \mathcal{F})$ is *compatible* with the recollement (6.1). We will see that this compatibility condition is precisely the requirement on a torsion pair so that the corresponding HRS-tilt of a glued t-structure is also obtained by gluing.

PROPOSITION 6.3. *A torsion pair $(\mathcal{T}, \mathcal{F})$ in $\mathcal{A}_{\mathcal{D}}$ is compatible with respect to (6.1) if and only if the corresponding HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is a t-structure glued with respect to the recollement (2.1).*

Proof. Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$, and let $(\widetilde{\mathcal{D}}^{\leq 0}, \widetilde{\mathcal{D}}^{\geq 0})$ be the corresponding HRS-tilt, that is,

$$\begin{aligned} \widetilde{\mathcal{D}}^{\leq 0} &= \{X \in \mathcal{D} \mid H^0(X) \in \mathcal{T}, H^i(X) = 0, \forall i > 0\}, \\ \widetilde{\mathcal{D}}^{\geq 0} &= \{X \in \mathcal{D} \mid H^{-1}(X) \in \mathcal{F}, H^i(X) = 0, \forall i < -1\}, \end{aligned}$$

where $H^i, i \in \mathbb{Z}$, denote the cohomological functors of the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$. By [5], the HRS-tilt is glued if and only if $j_!j^*(\widetilde{\mathcal{D}}^{\leq 0}) \subseteq \widetilde{\mathcal{D}}^{\leq 0}$. Denote by $\tau^{\leq -1}$ the truncation at -1 associated to $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$. For any $X \in \widetilde{\mathcal{D}}^{\leq 0}$, there is a triangle in \mathcal{D} of the form

$$\tau^{\leq -1}X \rightarrow X \rightarrow H^0X \rightarrow (\tau^{-1}X)[1],$$

since, by definition, $\widetilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$. Applying to it the functor $j_!j^*$, we get another triangle

$$j_!j^*(\tau^{\leq -1}X) \rightarrow j_!j^*(X) \rightarrow j_!j^*(H^0X) \rightarrow j_!j^*(\tau^{\leq -1}X)[1].$$

Since $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued, $j_!j^*(\mathcal{D}^{\leq -1}) \subseteq \mathcal{D}^{\leq -1} \subseteq \widetilde{\mathcal{D}}^{\leq 0}$. Hence, $j_!j^*(\tau^{\leq -1}X)$ lies in $\widetilde{\mathcal{D}}^{\leq 0}$ and, thus, $j_!j^*(X)$ lies in $\widetilde{\mathcal{D}}^{\leq 0}$ if and only if $j_!j^*(H^0X)$ lies in $\widetilde{\mathcal{D}}^{\leq 0}$. Since X is arbitrary, this means precisely that $(\widetilde{\mathcal{D}}^{\leq 0}, \widetilde{\mathcal{D}}^{\geq 0})$ can be restricted if and only if the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with (6.1). \square

The question that naturally follows is whether, under the compatibility condition on the torsion pair in \mathcal{D} , the restrictions of the HRS-tilt of a glued t-structure in \mathcal{D} are precisely the HRS-tilts of the restricted t-structures on \mathcal{Y} and \mathcal{X} with respect to the restricted torsion pairs.

THEOREM 6.4. *Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$, compatible with (6.1) and with restrictions $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$ and $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ in $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$, respectively. Let $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ be the corresponding HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$, respectively. Then the HRS-tilt $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ is obtained by gluing $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ and $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ with respect to the recollement (2.1).*

Proof. Let $H_{\mathcal{D}}^i$, $H_{\mathcal{X}}^i$, and $H_{\mathcal{Y}}^i$, $i \in \mathbb{Z}$, denote the cohomological functors associated with $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$, and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$, respectively. We have to show that the following hold:

- (i) $j^*(\tilde{\mathcal{D}}^{\leq 0}) = \tilde{\mathcal{X}}^{\leq 0}$, $j^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{X}}^{\geq 0}$,
- (ii) $i^*(\tilde{\mathcal{D}}^{\leq 0}) = \tilde{\mathcal{Y}}^{\leq 0}$, $i^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{Y}}^{\geq 0}$.

By definition, we have

$$j^*(\tilde{\mathcal{D}}^{\leq 0}) = \{j^*X \mid X \in \mathcal{D}, H_{\mathcal{D}}^0(X) \in \mathcal{T}, H_{\mathcal{D}}^i(X) = 0, \forall i > 0\},$$

and, since j^* is t-exact (see [5]), $H_{\mathcal{X}}^i(j^*X) = j^*(H_{\mathcal{D}}^i(X))$, thus showing that

$$j^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \{j^*X \mid X \in \mathcal{D}, H_{\mathcal{X}}^0(j^*X) \in j^*\mathcal{T}, H_{\mathcal{X}}^i(j^*X) = 0, \forall i > 0\} = \tilde{\mathcal{X}}^{\leq 0}.$$

Similarly, we get that

$$j^*(\tilde{\mathcal{D}}^{\geq 0}) \subseteq \{j^*X \mid X \in \mathcal{D}, H_{\mathcal{X}}^{-1}(j^*X) \in j^*\mathcal{T}, H_{\mathcal{X}}^i(j^*X) = 0, \forall i < -1\} = \tilde{\mathcal{X}}^{\geq 0}.$$

Since $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement, by Proposition 6.3, the HRS-tilt $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ can be restricted and, thus, $(j^*(\tilde{\mathcal{D}}^{\leq 0}), j^*(\tilde{\mathcal{D}}^{\geq 0}))$ is a t-structure in \mathcal{X} . Since $j^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \tilde{\mathcal{X}}^{\leq 0}$, we also have that $j^*(\tilde{\mathcal{D}}^{\geq 0}) \supseteq \tilde{\mathcal{X}}^{\geq 0}$ and, thus, $j^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{X}}^{\geq 0}$. This proves (i).

To prove (ii), observe that i^* is right t-exact (i.e., $i^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{Y}^{\leq 0}$) and that

$$i^*(\tilde{\mathcal{D}}^{\leq 0}) = \{i^*X \mid X \in \mathcal{D}, H_{\mathcal{D}}^0(X) \in \mathcal{T}, H_{\mathcal{D}}^i(X) = 0, \forall i > 0\}.$$

By [5, Proposition 1.3.17(ii)], for X in $\tilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$, we have $H_{\mathcal{Y}}^0(i^*X) = {}^p i^* H_{\mathcal{D}}^0(X)$ which belongs to ${}^p i^*(\mathcal{T})$. Since $i^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq i^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{Y}^{\leq 0}$, we get that $H_{\mathcal{Y}}^i(i^*\tilde{\mathcal{D}}^{\leq 0}) = 0$, for all $i > 0$, and thus

$$i^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \{i^*X \mid X \in \mathcal{D}, H_{\mathcal{Y}}^0(i^*X) \in {}^p i^*\mathcal{T}, H_{\mathcal{Y}}^i(i^*X) = 0, \forall i > 0\} = \tilde{\mathcal{Y}}^{\leq 0}.$$

Analogously, we may conclude that

$$i^*(\tilde{\mathcal{D}}^{\geq 0}) \subseteq \{i^*X \mid X \in \mathcal{D}, H_{\tilde{\mathcal{Y}}}^{-1}(i^*X) \in {}^p i^* \mathcal{F}, H_{\tilde{\mathcal{Y}}}^i(i^*X) = 0, \forall i < -1\} = \tilde{\mathcal{Y}}^{\geq 0}.$$

The same argument as before shows that an equality holds, as wanted. \square

The previous result showed that an HRS-tilt with respect to a compatible torsion pair is glued. We may now ask whether HRS-tilts on the sides of the recollement (2.1) glue to an HRS-tilt in the middle. The following proposition answers this question positively.

PROPOSITION 6.5. *Let $(\mathcal{T}_Y, \mathcal{F}_Y)$ and $(\mathcal{T}_X, \mathcal{F}_X)$ be torsion pairs in \mathcal{A}_Y and \mathcal{A}_X , respectively. Then the pair $(\mathcal{T}, \mathcal{F})$ defined by*

$$\begin{aligned} \mathcal{T} &= \{A \in \mathcal{A}_{\mathcal{D}} \mid {}^p i^* A \in \mathcal{T}_Y, {}^p j^* A \in \mathcal{T}_X\}, \\ \mathcal{F} &= \{A \in \mathcal{A}_{\mathcal{D}} \mid {}^p i^! A \in \mathcal{F}_Y, {}^p j^* A \in \mathcal{F}_X\} \end{aligned}$$

is a torsion pair in $\mathcal{A}_{\mathcal{D}}$, compatible with (6.1). Moreover, its restrictions are given by $(\mathcal{T}_X, \mathcal{F}_X)$ and $(\mathcal{T}_Y, \mathcal{F}_Y)$, and the HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$ is the gluing of the HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ with respect to $(\mathcal{T}_Y, \mathcal{F}_Y)$ and $(\mathcal{T}_X, \mathcal{F}_X)$, respectively.

Proof. We will use Remark 2.9 to show that $(\mathcal{T}, \mathcal{F})$ is a torsion pair, that is, we will show that the pair $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ formally defined as the HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$ is a t-structure (satisfying $\mathcal{D}^{\leq -1} \subseteq \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$). Denote by $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ the HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ at $(\mathcal{T}_Y, \mathcal{F}_Y)$ and $(\mathcal{T}_X, \mathcal{F}_X)$, respectively, and let $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ be the gluing of $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ with respect to the fixed recollement, that is,

$$\begin{aligned} \tilde{\mathcal{D}}^{\leq 0} &= \{X \in \mathcal{D} \mid i^*(X) \in \tilde{\mathcal{Y}}^{\leq 0}, j^*(X) \in \tilde{\mathcal{X}}^{\leq 0}\}, \\ \tilde{\mathcal{D}}^{\geq 0} &= \{X \in \mathcal{D} \mid i^!(X) \in \tilde{\mathcal{Y}}^{\geq 0}, j^*(X) \in \tilde{\mathcal{X}}^{\geq 0}\}. \end{aligned}$$

We will show that $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0}) = (\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$. As usual, $H_{\mathcal{D}}^i$, $H_{\mathcal{X}}^i$, and $H_{\mathcal{Y}}^i$, $i \in \mathbb{Z}$, denote the cohomological functors associated with $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$, and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$, respectively.

First we show that $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$ is a subcategory of $\tilde{\mathcal{D}}^{\leq 0}$. Take X in $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$ ($\subseteq \mathcal{D}^{\leq 0}$). Since i^* is right t-exact, $i^*(X)$ lies in $\mathcal{Y}^{\leq 0}$, that is, $H_{\mathcal{Y}}^i(i^*X) = 0$, for all $i > 0$. By [5, Proposition 1.3.17(ii)], $H_{\mathcal{Y}}^0(i^*X) = {}^p i^* H_{\mathcal{D}}^0(X)$ lies

in ${}^p i^*(\mathcal{T}) \subseteq \mathcal{T}_y$, showing that $i^*(X) \in \tilde{\mathcal{Y}}^{\leq 0}$. Since j^* is t-exact, we have $H_{\mathcal{X}}^i(j^*X) = j^*H_{\mathcal{D}}^i(X) = 0$, for all $i > 0$, and $H_{\mathcal{X}}^0(j^*X) = j^*H_{\mathcal{D}}^0(X)$ lies in $j^*(\mathcal{T}) \subseteq \mathcal{T}_{\mathcal{X}}$, that is, j^*X lies in $\tilde{\mathcal{X}}^{\leq 0}$ and, hence, X lies in $\tilde{\mathcal{D}}^{\leq 0}$.

Now we prove that $\tilde{\mathcal{D}}^{\leq 0}$ is a subcategory of $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$. Take X in $\tilde{\mathcal{D}}^{\leq 0}$. By definition of HRS-tilt, we have $\tilde{\mathcal{Y}}^{\leq 0} \subset \mathcal{Y}^{\leq 0}$ and $\tilde{\mathcal{X}}^{\leq 0} \subseteq \mathcal{X}^{\leq 0}$. This implies that $\tilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$ and, hence, $H_{\mathcal{D}}^i(X) = 0$, for all $i > 0$. By [5, Proposition 1.3.17(ii)], we have ${}^p i^*H_{\mathcal{D}}^0(X) = H_{\mathcal{Y}}^0(i^*X)$ which belongs to \mathcal{T}_y . By the t-exactness of j^* , we have ${}^p j^*H_{\mathcal{D}}^0(X) = H_{\mathcal{X}}^0(j^*X)$ which, thus, belongs to $\mathcal{T}_{\mathcal{X}}$. This means that, by definition of \mathcal{T} , $H_{\mathcal{D}}^0(X)$ belongs to \mathcal{T} . Thus X belongs to $\tilde{\mathcal{D}}^{\leq 0}$ and $\tilde{\mathcal{D}}^{\leq 0} = \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$, as wanted.


It remains to show that $(\mathcal{T}, \mathcal{F})$ is compatible and that the restrictions are as expected. By Lemma 6.2, we need to show that ${}^p j_! {}^p j^* \mathcal{T} \subseteq \mathcal{T}$. This is true since, for any A in \mathcal{T} , ${}^p i^* {}^p j_! {}^p j^* A = 0$ and ${}^p j^* {}^p j_! {}^p j^* A = {}^p j^* A \subseteq \mathcal{T}_{\mathcal{X}}$. It follows that $({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$ and $({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F}))$ are torsion pairs in \mathcal{A}_y and $\mathcal{A}_{\mathcal{X}}$, respectively. On the other hand, by definition of $(\mathcal{T}, \mathcal{F})$, we have

$$\begin{aligned} {}^p i^*(\mathcal{T}) &\subseteq \mathcal{T}_y, & {}^p i^!(\mathcal{F}) &\subseteq \mathcal{F}_y, \\ {}^p j^*(\mathcal{T}) &\subseteq \mathcal{T}_{\mathcal{X}}, & {}^p j^*(\mathcal{F}) &\subseteq \mathcal{F}_{\mathcal{X}}, \end{aligned}$$

and, thus, the inclusions are actually equalities, as desired. □

We illustrate Proposition 6.5 with a simple example.


EXAMPLE 6.6. We adopt the notation in Example 5.3 and consider the standard recollement there. Let $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ be the standard t-structure on $\mathcal{D}^b(R)$, which is depicted in the Auslander–Reiten quiver of $\mathcal{D}^b(R)$ as



where the objects in the boxes belong to the aisle $\mathcal{D}^{\leq 0}$. This t-structure restricts to standard t-structures $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ on $\mathcal{D}^b(eRe)$ and $\mathcal{D}^b(R/ReR)$, respectively. Recall that both eRe and R/ReR are isomorphic to \mathbb{K} . Consequently, all bounded t-structures on $\mathcal{D}^b(eRe)$ and $\mathcal{D}^b(R/ReR)$ are shifts of the standard ones, and all mutation torsion pairs of $\text{mod-}eRe$ and $\text{mod-}R/ReR$ are trivial torsion pairs.

The HRS-tilt of $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ with respect to the torsion pair $(0, \text{mod-}eRe)$ is $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$, and the HRS-tilt of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ with respect to the torsion

pair $(\text{mod-}R/ReR, 0)$ is $(\mathcal{Y}^{\leq 0}[1], \mathcal{Y}^{\geq 0}[1])$. These two new t-structures are glued, via the standard recollement, to a t-structure $(\mathcal{D}'^{\leq 0}, \mathcal{D}'^{\geq 0})$ on $\mathcal{D}^b(R)$, which is depicted as



where the objects in the boxes belong to the aisle $\mathcal{D}'^{\leq 0}$. The heart of $(\mathcal{D}'^{\leq 0}, \mathcal{D}'^{\geq 0})$ is $\text{add}(S_1 \oplus S_2[1])$. It is easy to check that $(\mathcal{D}'^{\leq 0}, \mathcal{D}'^{\geq 0})$ is the HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ at the torsion pair $(\text{add}(S_1), \text{add}(S_2))$, which is glued from the torsion pairs $(0, \text{mod-}eRe)$ and $(\text{mod-}R/ReR, 0)$.

By Section 2.4, we know that irreducible silting mutations correspond to HRS-tilts (see Theorem 2.11) with respect to mutation torsion pairs in the heart (i.e., torsion pairs in which either the torsion class or the torsion-free class are given by the additive closure of a simple object). We analyze when such torsion pairs are compatible with (6.1).

REMARK 6.7. Recall that there is a functor $j_{1*} : \mathcal{A}_{\mathcal{X}} \rightarrow \mathcal{A}_{\mathcal{D}}$, called the *intermediate image functor*, such that every simple object in $\mathcal{A}_{\mathcal{D}}$ is either of the form i_*S for some simple object in $\mathcal{A}_{\mathcal{Y}}$ or of the form $j_{1*}S$ for some simple object in $\mathcal{A}_{\mathcal{X}}$. (For details, check [5, Section 1.4].)

PROPOSITION 6.8. *Suppose that $\mathcal{A}_{\mathcal{D}}$ is a length category. Let $(\mathcal{T}, \mathcal{F})$ be a mutation torsion pair in $\mathcal{A}_{\mathcal{D}}$. Then $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement (6.1) if and only if one of the following holds:*

- (1) $\mathcal{T} = \text{add}(i_*S_Y)$ for some simple object S_Y in $\mathcal{A}_{\mathcal{Y}}$,
- (2) $\mathcal{F} = \text{add}(i_*S_Y)$ for some simple object S_Y in $\mathcal{A}_{\mathcal{Y}}$,
- (3) there is a simple object S_X in $\mathcal{A}_{\mathcal{X}}$ such that ${}^p j_{!}S_X$ is simple and $\mathcal{T} = \text{add}({}^p j_{!}S_X)$,
- (4) there is a simple object S_X in $\mathcal{A}_{\mathcal{X}}$ such that ${}^p j_{*}S_X$ is simple and $\mathcal{F} = \text{add}({}^p j_{*}S_X)$.

Moreover, in each case, the restrictions are a trivial torsion pair on one side and a mutation torsion pair on the other.

Proof. This can be shown using Lemma 6.2 on a case-by-case analysis of the possible mutation torsion pairs in $\mathcal{A}_{\mathcal{D}}$. Using Remark 6.7 we get the following four cases, where S_Y is a simple object in $\mathcal{A}_{\mathcal{Y}}$ and S_X is a simple object in $\mathcal{A}_{\mathcal{X}}$.

- (1) Suppose that $\mathcal{T} = \text{add}(i_*S_Y)$. Since $j^*i_* = 0$, we have that ${}^p j_! j^* \mathcal{T} = 0$ and, thus, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement. The restriction to \mathcal{X} is $(0, \mathcal{X})$ and the restriction to \mathcal{Y} is the mutation torsion pair with torsion class $\mathcal{T}_{\mathcal{Y}} = \text{add}(S_Y)$.
- (2) Suppose that $\mathcal{F} = \text{add}(i_*S_Y)$. Since $j^*i_* = 0$, we have that ${}^p j_* j^* \mathcal{T} = 0$ and, thus, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement. The restriction to \mathcal{X} is $(\mathcal{X}, 0)$ and the restriction to \mathcal{Y} is the mutation torsion pair with torsion-free class $\mathcal{F}_{\mathcal{Y}} = \text{add}(S_Y)$.
- (3) Suppose that $\mathcal{T} = \text{add}(j_!S_X)$. By Lemma 6.2, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement if and only if ${}^p j_! j^*(j_!S_X)$ lies in $\text{add}(j_!S_X)$. Since $j^*j_!$ is naturally equivalent to the identity functor, this amounts to ${}^p j_!S_X$ lying in $\text{add}(j_!S_X)$, which is equivalent to ${}^p j_!S_X \cong j_!S_X$ because $j^*{}^p j_!$ is also naturally equivalent to the identity functor. The last condition holds if and only if ${}^p j_!S_X$ is simple. This follows from the fact that there is always a natural epimorphism ${}^p j_!S_X \rightarrow j_!S_X$. The restriction to \mathcal{Y} is the pair $(0, \mathcal{Y})$ and the restriction to \mathcal{X} is the mutation torsion pair given by the torsion class $\mathcal{T}_{\mathcal{X}} = \text{add}(S_X)$.
- (4) Suppose that $\mathcal{F} = \text{add}(j_!S_X)$. By Lemma 6.2, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement if and only if ${}^p j_* j^*(j_!S_X)$ lies in $\text{add}(j_!S_X)$. Since $j^*j_!$ is naturally equivalent to the identity functor, this amounts to ${}^p j_*S_X$ lying in $\text{add}(j_!S_X)$, which is equivalent to ${}^p j_*S_X \cong j_!S_X$ because $j^*{}^p j_*$ is also naturally equivalent to the identity functor. The last condition holds if and only if ${}^p j_*S_X$ is simple. This follows from the fact that there is always a natural monomorphism $j_!S_X \rightarrow {}^p j_*S_X$. The restriction to \mathcal{Y} is the pair $(\mathcal{Y}, 0)$ and the restriction to \mathcal{X} is the mutation torsion pair given by the torsion-free class $\mathcal{F}_{\mathcal{X}} = \text{add}(S_X)$.

The proof is complete. \square

Suppose now that \mathcal{D} has a Serre functor. We finish this section with an observation on the compatibility of irreducible silting mutation with gluing (via the recollement \mathcal{R}_U , since our focus is on t-structures rather than co-t-structures). Note that the items (3) and (4) of Proposition 6.8 rarely occur, because, in general, neither ${}^p j_*S_X$ nor ${}^p j_!S_Y$ are simple in $\mathcal{A}_{\mathcal{D}}$. Translating items (1) and (2) of Proposition 6.8 in terms of irreducible silting mutation, we get the following corollary, which also appears in [1, Lemma 2.40].

COROLLARY 6.9 ([1, Lemma 2.40]). *Let Y and X be the silting objects associated to the fixed bounded t-structures in \mathcal{Y} and \mathcal{X} , respectively. Suppose*

that Y' is an indecomposable direct summand of Y . Let Z be the glued silting object with respect to the recollement \mathcal{R}_U (i.e., compatible with the gluing of t -structures via \mathcal{R}) of Y and X . Then there is an indecomposable direct summand Z' of K_Y such that the glued silting (with respect to \mathcal{R}_U) of μ_Y^+, Y (resp., μ_Y^-, Y) and X is precisely $\mu_{Z'}^+, Z$ (resp., $\mu_{Z'}^-, Z$).

Acknowledgments. The authors would like to thank David Pauksztello and Lutz Hille for helpful remarks and for a discussion which led to Example 4.4. Vitória's work was supported by DFG-Schwerpunktprogramm Darstellungstheorie 1388 grant KO1281/7-1. Yang's work was supported by DFG-Schwerpunktprogramm Darstellungstheorie 1388 grant KO1281/9-1.

REFERENCES

- [1] T. Aihara and O. Iyama, *Silting mutation in triangulated categories*, J. Lond. Math. Soc. (2) **85** (2012), 633–668. MR 2927802. DOI 10.1112/jlms/jdr055.
- [2] L. Angeleri Hügel, S. Koenig, and Q. Liu, *Recollements and tilting objects*, J. Pure Appl. Algebra **215** (2011), 420–438. MR 2738361. DOI 10.1016/j.jpaa.2010.04.027.
- [3] ———, *Jordan–Hölder theorems for derived module categories of piecewise hereditary algebras*, J. Algebra **352** (2012), 361–381. MR 2862193. DOI 10.1016/j.jalgebra.2011.09.041.
- [4] ———, *On the uniqueness of stratifications of derived module categories*, J. Algebra **359** (2012), 120–137. MR 2914629. DOI 10.1016/j.jalgebra.2012.02.022.
- [5] A. A. Beilinson, J. Bernstein, and P. Deligne, “Faisceaux pervers” in *Analysis and Topology on Singular Spaces, I (Luminy, 1981)*, Astérisque **100**, Soc. Math. France, Paris, 1982, 5–171. MR 0751966.
- [6] A. Beligiannis and I. Reiten, *Homological and Homotopical Aspects of Torsion Theories*, Mem. Amer. Math. Soc. **188**, 2007. MR 2327478. DOI 10.1090/memo/0883.
- [7] A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions* (in Russian), Izv. Akad. Nauk SSSR Ser. Mat. **53**, no. 6 (1989), 1183–1205, 1337; English translation in Math. USSR-Izv. **35**, no. 3 (1990), 519–541. MR 1039961.
- [8] M. V. Bondarko, *Weight structures vs. t -structures; weight filtrations, spectral sequences, and complexes (for motives and in general)*, J. K-Theory **6** (2010), 387–504. MR 2746283. DOI 10.1017/is010012005jkt083.
- [9] T. Bridgeland, *t -structures on some local Calabi–Yau varieties*, J. Algebra **289** (2005), 453–483. MR 2142382. DOI 10.1016/j.jalgebra.2005.03.016.
- [10] A. B. Buan, I. Reiten, and H. Thomas, *Three kinds of mutation*, J. Algebra **339** (2011), 97–113. MR 2811314. DOI 10.1016/j.jalgebra.2011.04.030.
- [11] E. Cline, B. Parshall, and L. L. Scott, *Algebraic stratification in representation categories*, J. Algebra **117** (1988), 504–521. MR 0957457. DOI 10.1016/0021-8693(88)90123-8.
- [12] ———, *Stratifying Endomorphism Algebras*, Mem. Amer. Math. Soc. **124** (1996), no. 591. MR 1350891. DOI 10.1090/memo/0591.
- [13] V. Franjou and T. Pirashvili, *Comparison of abelian categories recollements*, Doc. Math. **9** (2004), 41–56. MR 2054979.

- [14] D. Happel, I. Reiten, and S. O. Smalø, *Tilting in Abelian Categories and Quasi-tilted Algebras*, Mem. Amer. Math. Soc. **120** (1996), no. 575. MR 1327209. DOI 10.1090/memo/0575.
- [15] P. Jørgensen, *Reflecting recollements*, Osaka J. Math. **47** (2010), 209–213. MR 2666132.
- [16] B. Keller and P. Nicolás, *Cluster hearts and cluster tilting objects*, in preparation.
- [17] B. Keller and D. Vossieck, “Aisles in derived categories” in *Deuxième Contact Franco-Belge en Algèbre (Faulx-les-Tombes, 1987)*, Bull. Soc. Math. Belg. Sér. A **40** (1988), 239–253. MR 0976638.
- [18] S. Koenig and D. Yang, *Silting objects, simple-minded collections, t-structures and co-t-structures for finite-dimensional algebras*, Doc. Math. **19** (2014), 403–438. MR 3178243.
- [19] S. Ladkani, *Derived equivalences of triangular matrix rings arising from extensions of tilting modules*, Algebr. Represent. Theory **14** (2011), 57–74. MR 2763291. DOI 10.1007/s10468-009-9175-0.
- [20] Q. Liu and J. Vitória, *t-structures via recollements for piecewise hereditary algebras*, J. Pure Appl. Algebra **216** (2012), 837–849. MR 2864858. DOI 10.1016/j.jpaa.2011.08.010.
- [21] O. Mendoza Hernández, E. C. Sáenz Valadez, V. Santiago Vargas, and M. J. Souto Salorio, *Auslander-Buchweitz context and co-t-structures*, Appl. Categ. Structures **21** (2013), 417–440. MR 3097052. DOI 10.1007/s10485-011-9271-2.
- [22] D. Pauksztello, *Compact corigid objects in triangulated categories and co-t-structures*, Cent. Eur. J. Math. **6** (2008), 25–42. MR 2379950. DOI 10.2478/s11533-008-0003-2.
- [23] I. Reiten and M. Van den Bergh, *Noetherian hereditary abelian categories satisfying Serre duality*, J. Amer. Math. Soc. **15** (2002), 295–366. MR 1887637. DOI 10.1090/S0894-0347-02-00387-9.
- [24] J. Rickard, *Morita theory for derived categories*, J. Lond. Math. Soc. (2) **39** (1989), 436–456. MR 1002456. DOI 10.1112/jlms/s2-39.3.436.
- [25] J. Rickard and A. Schofield, *Cocovers and tilting modules*, Math. Proc. Cambridge Philos. Soc. **106** (1989), 1–5. MR 0994075. DOI 10.1017/S0305004100067931.
- [26] M. J. Souto Salorio and S. Trepode, *T-structures on the bounded derived category of the Kronecker algebra*, Appl. Categ. Structures **20** (2012), 513–529. MR 2957314. DOI 10.1007/s10485-011-9248-1.
- [27] L. Alonso Tarrío, A. Jeremías López, and M. J. Souto Salorio, *Construction of t-structures and equivalences of derived categories*, Trans. Amer. Math. Soc. **355**, no. 6 (2003), 2523–2543. MR 1974001. DOI 10.1090/S0002-9947-03-03261-6.
- [28] Z. Wang, *Nondegeneration and boundedness of t-structure induced by recollement*, Xiamen Daxue Xuebao Ziran Kexue Ban **45** (2006), 10–13. MR 2193734.
- [29] J. Woolf, *Stability conditions, torsion theories and tilting*, J. Lond. Math. Soc. (2) **82** (2010), 663–682. MR 2739061. DOI 10.1112/jlms/jdq035.

Qunhua Liu
Institute of Mathematics
School of Mathematical Sciences
Nanjing Normal University
Nanjing 210023
People's Republic of China
05402@njnu.edu.cn

Jorge Vitória
Department of Computer Science—Sector of Mathematics
University of Verona
I-37134 Verona
Italy
jorge.vitoria@univr.it

Dong Yang
Department of Mathematics
Nanjing University
Nanjing 210093
People's Republic of China
dongyang2002@gmail.com