

Cohomology of wheels on toric varieties

Alastair CRAW and Alexander QUINTERO VÉLEZ

(Received November 13, 2012; Revised February 2, 2014)

Abstract. We describe explicitly the cohomology of the total complex of certain diagrams of invertible sheaves on normal toric varieties. These diagrams, called wheels, arise in the study of toric singularities associated to dimer models. Our main tool describes the generators in a family of syzygy modules associated to the wheel in terms of walks in a family of graphs.

Key words: Cohomology of complexes, toric varieties, syzygies.

1. Introduction

A standard tool in homological algebra is to study a finitely generated module over a ring in terms of a free resolution, or more generally, a coherent sheaf on a variety in terms of a resolution by locally free sheaves. Conversely, given a complex T^\bullet of locally free sheaves on a variety X , it is natural to ask whether the cohomology of the complex is nonzero in one degree only, say $k \in \mathbb{Z}$, in which case T^\bullet is quasi-isomorphic to the pure sheaf $H^k(T^\bullet)[-k]$. In particular, it is important to have an explicit understanding of the cohomology sheaves of a complex of locally free sheaves. Our main result achieves this for a class of four-term complexes of locally free sheaves on normal toric varieties.

Our motivation comes from the study of derived categories of toric varieties associated to consistent dimer model algebras (see Bocklandt–Craw–Quintero-Vélez [2, Section 2.4] for a brief introduction). The best-known example of a consistent dimer model algebra is the skew group algebra $\mathbb{C}[x, y, z] * G$ for a finite abelian subgroup $G \subset \mathrm{SL}(3, \mathbb{C})$, in which case the relevant toric variety is the G -Hilbert scheme $X = G\text{-Hilb}(\mathbb{C}^3)$ introduced by Nakamura [10]. In their study of the equivalence of derived categories induced by the universal family on the G -Hilbert scheme, Cautis–Logvinenko [3] describes explicitly the cohomology sheaves of certain four-term com-

plexes T^\bullet on X and hence shows that with only one exception, every such complex is quasi-isomorphic to a pure sheaf $H^k(T^\bullet)[-k]$ for $k = 0, 1$ (see also Cautis–Craw–Logvinenko [4]). Our main result (see Theorem 1.1 below) can be applied to a broader class of four-term complexes, including those arising in the study of the derived equivalences induced by the universal family of fine moduli spaces X associated to any consistent dimer model algebra. As an application, joint work with Raf Bocklandt [2] establishes the dimer model analogue of the Cautis–Logvinenko result, namely, that for a special choice of moduli space generalising the G -Hilbert scheme, all but one of the four-term complexes T^\bullet on X obtained from the derived equivalence is quasi-isomorphic to a pure sheaf $H^k(T^\bullet)[-k]$ for $k = 0, 1$.

The complexes T^\bullet that we consider in this paper are four-term complexes of the form

$$L \xrightarrow{d^3} \bigoplus_{j=1}^m L_{j,j+1} \xrightarrow{d^2} \bigoplus_{j=1}^m L_j \xrightarrow{d^1} L \quad (1.1)$$

for some $m \geq 2$, where L , $L_{j,j+1}$ and L_j ($1 \leq j \leq m$) are invertible sheaves on any normal toric variety X , where each differential is equivariant with respect to the torus-action on X , and where the right-hand copy of L lies in degree zero. Assume in addition that for $1 \leq j \leq m$, the restriction of the differential d^2 to the summand $L_{j,j+1}$ has image in $L_j \oplus L_{j+1}$ (with indices modulo m). This means that if we separate vertically the summands in the terms of T^\bullet and hence break the matrices defining the differentials into their constituent maps between summands, the complex can be presented as a diagram of the form

$$\begin{array}{ccccc}
 & & L_{1,2} & \xrightarrow{D_1^2} & L_1 & & \\
 & & & \searrow^{D_2^2} & & & \\
 & & L_{2,3} & \xrightarrow{D_2^2} & L_2 & \xrightarrow{D^1} & \\
 & & & \searrow^{D_3^2} & & & \\
 & & L_{3,4} & \xrightarrow{D_3^2} & L_3 & \xrightarrow{D^2} & L \\
 & & \vdots & & \vdots & & \\
 & & L_{m,1} & \xrightarrow{D_m^2} & L_m & \xrightarrow{D^3} & \\
 & & & \searrow^{D_1^2} & & &
 \end{array} \quad (1.2)$$

The maps between invertible sheaves in this diagram are multiplication by a torus-invariant section of an invertible sheaf on X . We illustrate this and fix notation by writing on each arrow in diagram (1.2) the Cartier divisor of zeros of the corresponding section so, for example, the effective divisor $D_2^1 \in H^0(L_2 \otimes L_{1,2}^{-1}) \cong \text{Hom}(L_{1,2}, L_2)$ denotes the Cartier divisor of zeros of the section that defines the map from $L_{1,2}$ to L_2 . This diagram can be represented equally well in a planar picture that is reminiscent of a bicycle wheel (see Figure 4 in Section 3), and we refer to any such four-term complex T^\bullet as a ‘wheel’ on X .

To state our main result we choose once and for all a rather special order on the set of transpositions of m letters (see Section 2), giving $\tau_1 = (\mu_1, \nu_1), \dots, \tau_n = (\mu_n, \nu_n)$ where $n = \binom{m}{2}$ and $\mu_k < \nu_k$ for $1 \leq k \leq n$. In addition, for every index $1 \leq k \leq n$ we define a subscheme $Z_k \subset X$ to be the scheme-theoretic intersection of certain torus-invariant divisors in X . To be more precise, let $\mathcal{D} := \{D_\lambda\}_{\lambda \in \Lambda}$ be a set of torus-invariant divisors in X . Define the greatest common divisor and the least common multiple of the set \mathcal{D} to be the torus-invariant divisors

$$\begin{aligned} \gcd(\mathcal{D}) &= \max\{D \mid D_\lambda - D \geq 0 \forall \lambda \in \Lambda\} \quad \text{and} \\ \text{lcm}(\mathcal{D}) &= \min\{D \mid D - D_\lambda \geq 0 \forall \lambda \in \Lambda\} \end{aligned}$$

respectively; here max/min means choose the maximal/minimal values for the coefficients of each prime divisor in the expression for D . Define subschemes $Z_k \subset X$ for $1 \leq k \leq n$ in terms of the Cartier divisors labelling the arrows in diagram (1.2) as follows:

- (i) for $1 \leq k \leq m$, define Z_k to be the scheme-theoretic intersection of $\gcd(D_{k+1}^k, D_k^{k+1})$ and the divisor $\text{lcm}(D^1, \dots, D^m, \gcd(D_{k+2}^{k+1}, D_{k+1}^{k+2}), \dots, \gcd(D_1^m, D_m^1)) - \text{lcm}(D^k, D^{k+1})$;
- (ii) for $m+1 \leq k \leq 2m-3$, define Z_k to be the scheme-theoretic intersection of the divisors $\text{lcm}(D^1, D^{\nu_k}, D^{\nu_k+1}, \dots, D^m) - \text{lcm}(D^1, D^{\nu_k})$ and $\text{lcm}(D^1, D^{\nu_k-1}, D^{\nu_k}) - \text{lcm}(D^1, D^{\nu_k})$;
- (iii) for $2m-2 \leq k \leq n$, define Z_k to be the scheme-theoretic intersection of the divisors $\text{lcm}(D^\mu, D^{\mu_k}, D^{\nu_k}) - \text{lcm}(D^{\mu_k}, D^{\nu_k})$ for $\mu \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\}$.

The subschemes $Z_k \subset X$ are torus-invariant, though some (possibly all) may be empty, see Example 3.5 for an explicit calculation.

Theorem 1.1 *Let X be a normal toric variety and let T^\bullet be the complex from (1.1), with differentials determined by the Cartier divisors shown in (1.2). Then:*

- (1) $H^0(T^\bullet) \cong \mathcal{O}_Z \otimes L$ where Z is the scheme-theoretic intersection of D^1, \dots, D^m ;
- (2) $H^{-1}(T^\bullet)$ has an n -step filtration

$$\mathrm{im}(d^2) = F^0 \subseteq F^1 \subseteq \dots \subseteq F^{n-1} \subseteq F^n = \ker(d^1)$$

where, for $1 \leq k \leq n$ and for the permutation $\tau_k = (\mu_k, \nu_k)$, we have

$$F^k / F^{k-1} \cong \mathcal{O}_{Z_k} \otimes L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\mathrm{gcd}(D^{\mu_k}, D^{\nu_k})); \quad (1.3)$$

- (3) $H^{-2}(T^\bullet) \cong \mathcal{O}_D \otimes L(D)$ where $D = \mathrm{gcd}(D_{1,2}, D_{2,3}, \dots, D_{m,1})$;
- (4) $H^{-3}(T^\bullet) \cong 0$.

To prove Theorem 1.1 we lift the complex T^\bullet to a complex of $\mathrm{Cl}(X)$ -graded S -modules using the functor of Cox [5], where $\mathrm{Cl}(X)$ and S denote the class group and Cox ring of X respectively. Explicitly, if $S(L)$ denotes the free S -module with generator in degree $L \in \mathrm{Cl}(X)$, then T^\bullet can be lifted to the complex

$$S(L) \xrightarrow{\varphi^3} \bigoplus_{j=1}^m S(L_{j,j+1}) \xrightarrow{\varphi^2} \bigoplus_{j=1}^m S(L_j) \xrightarrow{\varphi^1} S(L). \quad (1.4)$$

This translates the problem to one from commutative algebra. The lion's share of the effort in proving Theorem 1.1 goes into proving part (2). For this, the image of φ^2 is generated by elements $\alpha_1, \dots, \alpha_m$, and our chosen order on the set of transpositions on m letters determines an order on the generators β_1, \dots, β_n of $\ker(\varphi^1)$ which in turn defines a filtration

$$\mathrm{im}(\varphi^2) = F^0 \subseteq F^1 \subseteq F^2 \subseteq \dots \subseteq F^{n-1} \subseteq F^n = \ker(\varphi^1).$$

We give a presentation for each successive quotient F^k / F^{k-1} as a cyclic $\mathrm{Cl}(X)$ -graded S -module of the form $(S/I_k)(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\mathrm{gcd}(D^{\mu_k}, D^{\nu_k})))$ for some monomial ideal I_k whose generators are defined via the Cartier divisors D^1, \dots, D^m labelling the right-hand arrows in the diagram (1.2)

illustrating the wheel (see Proposition 3.1). This calculation can be performed in any given example using Macaulay2 [7], but we present a unified description for all $1 \leq k \leq n$. (Warning: M2 may choose an order on the generators β_1, \dots, β_n that differs from ours, see Remark 3.6.)

Our main tool, which may be of independent interest, is a description of the syzygy module of $\ker(\varphi^1)$ in terms of walks in the complete graph Γ on m vertices. In fact, for each $1 \leq k \leq n$ we introduce a subgraph Γ_k of Γ that enables us to describe uniformly the module of syzygies $\text{syz}(F^k)$ in terms of certain walks in Γ_k . To state the result, recall that a circuit in Γ_k is a closed walk that does not pass through a given vertex twice. It is straightforward to associate a syzygy to every such circuit (see Lemma 2.3). A circuit is said to be minimal if it admits no chords (see (2.4)). We prove the following result (see Theorem 2.5).

Theorem 1.2 *For $m \leq k \leq n$, the module $\text{syz}(F^k)$ is generated by the set of syzygies associated to the minimal circuits of Γ_k .*

The precise description of the syzygies from Theorem 1.2 allows us to read off directly a set of monomial generators for each ideal I_k , and this feeds into the proof of Theorem 1.1 above. Generating sets for toric ideals arising from graphs were studied by Hibi–Ohsugi [11], and some of the graph-theoretic tools that we use here were also employed there. Properties of \mathbb{k} -algebras arising from graphs have also been studied widely by Villarreal, see for example [12].

Our main result was motivated by the statement of Cautis–Logvinenko [3, Lemma 3.1] which asserts that in the special case $m = 3$, a version of Theorem 1.1 holds for the complex T^\bullet from (1.1) arising from a diagram (1.2) on an arbitrary smooth separated scheme. However, this is not true in general: the assertion [3, Proof of Lemma 3.1(2)] that certain elements $\beta_1, \beta_2, \beta_3$ generate $\ker(d^1)$ may fail if the maps from diagram (1.2) are not monomial maps.

Example 1.3 For a counterexample in the notation of *loc.cit.* (we write the signs explicitly), suppose the maps $L_1 \rightarrow L$, $L_2 \rightarrow L$ and $L_3 \rightarrow L$ from (1.2) are defined locally near a point $p \in X$ as multiplication by $f_1 := x$, $f_2 := x + y$, $f_3 := y \in \mathcal{O}_{X,p}$. Then $(1, -1, 1)$ lies in $\ker(d^1)$, but it does not lie in the submodule generated by $\beta_1 = (f_2, -f_1, 0)$, $\beta_2 = (-f_3, 0, f_1)$, $\beta_3 = (0, f_3, -f_2)$.

The assumption in Theorem 1.1 that X is toric and the maps from (1.2) are torus-equivariant ensures that each map arises from multiplication by a monomial in the Cox ring of X , in which case standard Gröbner theory shows that analogous elements $\beta_1, \beta_2, \beta_3$ generate the appropriate kernel (see Lemma 2.1). Under these additional assumptions, Remark 3.4 explains how the statement of Cautis–Logvinenko [3, Lemma 3.1] can be recovered as a special case of Theorem 1.1 when X is smooth. The main results of both Cautis–Logvinenko [3] and Cautis–Craw–Logvinenko [4] require the statement of [3, Lemma 3.1] only when X is a smooth toric variety and the maps from (1.2) are torus-equivariant, so Theorem 1.1 holds at the level of generality required for both of those papers.

In fact, Theorem 1.1 provides a unified description of the sheaves (1.3) in the filtration on $H^{-1}(T^\bullet)$ even for $m = 3$, improving slightly on the statement from [3, Lemma 3.1]. More generally, for $m > 3$, the schemes Z_k ($1 \leq k \leq n$) divide naturally into three families determined by the intervals (i) $1 \leq k \leq m$; (ii) $m + 1 \leq k \leq 2m - 3$; and (iii) $2m - 2 \leq k \leq n$, leading to a more involved filtration in this case. That the statement is considerably more complicated for $m > 3$ stems from the simple fact that any pair of vertices of a triangle are adjacent, while the same statement is not true for a polygon with $m > 3$ vertices.

2. Syzygies from walks in a complete graph

Let $S = \mathbb{k}[x_1, \dots, x_d]$ be a polynomial ring over a field \mathbb{k} and let $f^1, \dots, f^m \in S$ be monomials for some $m \geq 2$. Consider the free S -module with basis $\mathbf{e}_1, \dots, \mathbf{e}_m$ and define an S -module homomorphism $\varphi: \bigoplus_{\mu=1}^m S\mathbf{e}_\mu \rightarrow S$ by setting $\varphi(\mathbf{e}_\mu) = f^\mu$ for $1 \leq \mu \leq m$. For every pair of indices $1 \leq \mu < \nu \leq m$ we define monomials $f^{\mu, \nu} = \text{lcm}(f^\mu, f^\nu)$ and set

$$\beta_{(\mu, \nu)} = \frac{f^{\mu, \nu}}{f^\nu} \mathbf{e}_\nu - \frac{f^{\mu, \nu}}{f^\mu} \mathbf{e}_\mu. \quad (2.1)$$

The module of syzygies of $M := \langle f^1, \dots, f^m \rangle$ is defined to be the S -module $\text{syz}(M) := \ker(\varphi)$. The following result is well known; see for example Eisenbud [6, Lemma 15.1].

Lemma 2.1 *The kernel of φ is generated by the elements $\beta_{(\mu, \nu)}$ for $1 \leq \mu < \nu \leq m$.*

It is convenient to order the set $\{(\mu, \nu) \mid 1 \leq \mu < \nu \leq m\}$ of transpositions of m letters. First list the transpositions of adjacent letters $\tau_j = (j, j+1)$ for $1 \leq j \leq m-1$. Set $\tau_m = (1, m)$, then list all remaining transpositions that involve 1 as $\tau_j = (1, j-m+2)$ for $m+1 \leq j \leq 2m-3$, and finally list all remaining transpositions lexicographically, so $\tau_i = (\mu_i, \nu_i)$ precedes $\tau_j = (\mu_j, \nu_j)$ if and only if $\mu_i < \mu_j$ or $\mu_i = \mu_j$ and $\nu_i < \nu_j$. We may therefore list the generators of $\ker(\varphi)$ from Lemma 2.1 by setting $\beta_j := \beta_{(\mu_j, \nu_j)}$ for all $1 \leq j \leq n$, where $n = \binom{m}{2}$. This choice of order enables us to define for each $1 \leq k \leq n$ an S -module

$$F^k = \langle \beta_1, \dots, \beta_k \rangle.$$

Our primary goal is to provide for each $1 \leq k \leq n$ an explicit set of generators for the module of syzygies $\text{syz}(F^k)$ that encodes the relations between β_1, \dots, β_k . Recall that this module is defined to be the kernel of the surjective S -module homomorphism

$$\psi: \bigoplus_{j=1}^k S\varepsilon_j \longrightarrow F^k$$

satisfying $\psi(\varepsilon_j) = \beta_j$ for $1 \leq j \leq k$. We compute this module directly for $1 \leq k \leq m$.

Lemma 2.2 *The S -module $\text{syz}(F^k)$ is the zero module for $1 \leq k \leq m-1$, and it is a free module of rank one for $k = m$.*

Proof. Our choice of order on transpositions ensures that for $1 \leq k \leq m-1$, there can be no relations between β_1, \dots, β_k . For $k = m$, let $\sigma = \sum_{j=1}^m s_j \varepsilon_j$ be a syzygy on β_1, \dots, β_m where $s_1, \dots, s_m \in S$. By comparing coefficients of each \mathbf{e}_i in the expression

$$0 = \psi(\sigma) = s_m \left(\frac{f^{1,m}}{f^m} \mathbf{e}_m - \frac{f^{1,m}}{f^1} \mathbf{e}_1 \right) + \sum_{j=1}^{m-1} s_j \left(\frac{f^{j,j+1}}{f^{j+1}} \mathbf{e}_{j+1} - \frac{f^{j,j+1}}{f^j} \mathbf{e}_j \right)$$

we obtain the following equations

$$s_1 f^{1,2} = s_2 f^{2,3} = \dots = s_{m-1} f^{m-1,m} = -s_m f^{1,m}. \quad (2.2)$$

It's easy to see (or see Lemma 2.3 below for a proof) that the element

$$\sigma_0 := -\frac{\text{lcm}(f^1, \dots, f^m)}{f^{1,m}} \varepsilon_m + \sum_{j=1}^{m-1} \frac{\text{lcm}(f^1, \dots, f^m)}{f^{j,j+1}} \varepsilon_j \quad (2.3)$$

is a syzygy. Moreover, equations (2.2) imply that

$$\sigma = \frac{s_1 f^{1,2}}{\text{lcm}(f^1, \dots, f^m)} \sigma_0,$$

so $\text{syz}(F^m)$ is the free S -module with basis σ_0 . \square

We study the module $\text{syz}(F^k)$ for $m+1 \leq k \leq n$ by studying walks in a graph. Let Γ be the complete graph on m vertices, with vertex set $\{1, 2, \dots, m\}$. Assign an orientation to each edge $e = (\mu, \nu)$ by directing it from μ to ν if $\mu < \nu$. Regard every such edge as being labelled by the corresponding generator $\beta_{(\mu, \nu)}$ of $\ker(\varphi)$. The order on the generators β_1, \dots, β_n introduced above determines an order on the set of edges e_1, \dots, e_n of Γ . A *walk* γ of length ℓ in Γ is a walk in the undirected graph that traverses precisely ℓ edges. Every such walk is characterised by the sequence of vertices $\gamma = (\mu_1, \mu_2, \dots, \mu_{\ell+1})$ in Γ that it touches. A walk γ is *closed* if $\mu_1 = \mu_{\ell+1}$, and a *circuit* is a closed walk for which μ_1, \dots, μ_ℓ are distinct. Each circuit γ defines uniquely a subgraph of Γ , and we let $\text{supp}(\gamma)$ denote its set of edges. Given a circuit γ and an edge $e \in \text{supp}(\gamma)$, set $\text{sign}_\gamma(e) = +1$ if γ traverses e according to the orientation in Γ , and set $\text{sign}_\gamma(e) = -1$ if γ traverses e against orientation.




Figure 1. Directed graph Γ illustrating generators β_1, \dots, β_n for $m = 5$.

Given that the elements β_j for $1 \leq j \leq n$ correspond to edges in Γ , we may index the basis elements ε_j for $1 \leq j \leq n$ by edges e_1, \dots, e_n in Γ . Thus, for the edge $e = e_j$ for $1 \leq j \leq n$, we write $\varepsilon_e := \varepsilon_j$. For any vertices $\mu_1, \dots, \mu_{\ell+1}$ in Γ , set

$$f^{\mu_1, \dots, \mu_{\ell+1}} = \text{lcm}(f^{\mu_1}, \dots, f^{\mu_{\ell+1}}).$$

For a walk $\gamma = (\mu_1, \mu_2, \dots, \mu_{\ell+1})$ in Γ we define the monomial $f^\gamma := f^{\mu_1, \dots, \mu_{\ell+1}}$. In particular, for an edge e in Γ joining vertex μ to ν , we obtain $f^e = f^{\mu, \nu}$.

Lemma 2.3 *For any circuit γ of length at least three in Γ , the vector*

$$\sigma_\gamma = \sum_{e \in \text{supp}(\gamma)} \text{sign}_\gamma(e) \frac{f^\gamma}{f^e} \varepsilon_e$$

is a syzygy on β_1, \dots, β_n .

Proof. If γ has length two then $\sigma_\gamma = \varepsilon_e - \varepsilon_e = 0$ which is not in fact a syzygy by definition. For any circuit γ of length at least three we must show that

$$\psi(\sigma_\gamma) = \sum_{e \in \text{supp}(\gamma)} \text{sign}_\gamma(e) \frac{f^\gamma}{f^e} \beta_e = 0.$$

For an edge e that γ traverses in the direction from vertex μ to vertex μ' , we have that

$$\text{sign}_\gamma(e) \frac{f^\gamma}{f^e} \beta_e = \frac{f^\gamma}{f^e} \left(\frac{f^e}{f^{\mu'}} \mathbf{e}_{\mu'} - \frac{f^e}{f^\mu} \mathbf{e}_\mu \right) = \frac{f^\gamma}{f^{\mu'}} \mathbf{e}_{\mu'} - \frac{f^\gamma}{f^\mu} \mathbf{e}_\mu.$$

The sum of all such terms over $e \in \text{supp}(\gamma)$ collapses as a telescoping series since γ is closed. \square

For $1 \leq k \leq n$, let Γ_k denote the spanning subgraph of Γ that has vertex set $\{1, \dots, m\}$, and which includes only the first k edges of Γ (see Figure 2(a) below for the case $k = m + 3$). Clearly $\Gamma = \Gamma_n$. Let $\gamma = (\mu_1, \dots, \mu_\ell, \mu_1)$ be a circuit in Γ_k for some k . A *chord* of γ in Γ_k is any edge of the form $c = (\mu_r, \mu_s)$ for some $1 \leq r < s \leq \ell$ that does not lie in $\text{supp}(\gamma)$. Every such

chord c splits γ into two circuits:

$$\gamma_1 = (\mu_r, \dots, \mu_s, \mu_r) \quad \text{and} \quad \gamma_2 = (\mu_1, \dots, \mu_r, \mu_s, \dots, \mu_\ell, \mu_1). \quad (2.4)$$

A circuit must have length at least four if it is to admit a chord. We define a *minimal circuit* of Γ_k to be a circuit of length at least three that has no chords.

Lemma 2.4 *Let γ be a circuit in Γ_k admitting a chord in Γ_k that splits γ into circuits γ_1 and γ_2 as in (2.4). Then the syzygy σ_γ is contained in the module generated by σ_{γ_1} and σ_{γ_2} .*

Proof. Let c be the chord. For $i = 1, 2$, let $\gamma_i \setminus c$ denote the walk obtained from γ_i by removing the edge c . Since $\text{sign}_{\gamma_1}(c) = -\text{sign}_{\gamma_2}(c)$ we may rewrite

$$\begin{aligned} \sigma_\gamma &= \text{sign}_{\gamma_1}(c) \frac{f^\gamma}{f^c} \varepsilon_c + \text{sign}_{\gamma_2}(c) \frac{f^\gamma}{f^c} \varepsilon_c + \sum_{e \in \text{supp}(\gamma)} \text{sign}_\gamma(e) \frac{f^\gamma}{f^e} \varepsilon_e \\ &= \text{sign}_{\gamma_1}(c) \frac{f^\gamma}{f^c} \varepsilon_c + \sum_{e \in \text{supp}(\gamma_1 \setminus c)} \text{sign}_{\gamma_1}(e) \frac{f^\gamma}{f^e} \varepsilon_e + \text{sign}_{\gamma_2}(c) \frac{f^\gamma}{f^c} \varepsilon_c \\ &\quad + \sum_{e \in \text{supp}(\gamma_2 \setminus c)} \text{sign}_{\gamma_2}(e) \frac{f^\gamma}{f^e} \varepsilon_e \\ &= \frac{f^\gamma}{f^{\gamma_1}} \sigma_{\gamma_1} + \frac{f^\gamma}{f^{\gamma_2}} \sigma_{\gamma_2}. \end{aligned}$$

It remains to note that $f^{\gamma_1} = f^{\mu_r, \dots, \mu_s}$ divides $f^\gamma = f^{\mu_1, \dots, \mu_\ell}$, and similarly, f^{γ_2} divides f^γ . \square

We are now in a position to establish the main result of this section.

Theorem 2.5 *For $1 \leq k \leq n$, the S -module $\text{syz}(F^k)$ is generated by the syzygies σ_γ , where γ is a minimal circuit of Γ_k .*

Proof. We distinguish three cases. The first case, in which $1 \leq k \leq m - 1$, is straightforward: the graph Γ_k admits no circuits and $\text{syz}(F^k) = 0$ by Lemma 2.2, so the result holds.

We prove the second case, in which $m \leq k \leq 2m - 3$, by induction. For $k = m$, Lemma 2.2 shows that the S -module $\text{syz}(F^m)$ is free with basis




Figure 2. The graph Γ_k for $m \leq k \leq 2m - 3$ illustrated for $k = m + 3$.

σ_0 from (2.3). The syzygy σ_{γ_0} associated to the unique minimal circuit $\gamma_0 = (1, 2, \dots, m, 1)$ in Γ_m coincides with σ_0 , so the statement holds for $k = m$. Assume the statement for Γ_{k-1} for any $m + 1 \leq k \leq 2m - 3$, and let

$$\sigma = \sum_{j=1}^k s_j \mathcal{E}_j$$

be a syzygy on β_1, \dots, β_k where $s_1, \dots, s_k \in S$.

As a first step we reduce to the case in which the coefficients satisfy $s_j = 0$ for $k - m + 2 \leq j \leq m$ (these indices determine the edges to the left of β_k in Figure 2(a)). Indeed, suppose otherwise, so $s_i \neq 0$ for some $k - m + 2 \leq i \leq m$. By comparing the coefficient of \mathbf{e}_μ for each index $k - m + 3 \leq \mu \leq m$ in the equation

$$0 = \psi(\sigma) = \sum_{j=1}^k s_j \left(\frac{f^{\mu_j, \nu_j}}{f^{\nu_j}} \mathbf{e}_{\nu_j} - \frac{f^{\mu_j, \nu_j}}{f^{\mu_j}} \mathbf{e}_{\mu_j} \right),$$

we obtain a collection of equations

$$\begin{aligned} & s_{k-m+2} f^{k-m+2, k-m+3} \\ &= s_{k-m+3} f^{k-m+3, k-m+4} = \dots = s_{m-1} f^{m-1, m} = -s_m f^{1, m} \end{aligned} \quad (2.5)$$

which imply that $s_j \neq 0$ for all $k - m + 2 \leq j \leq m$. As illustrated in Figure 2(b) for $k = m + 3$, the circuit $\gamma_1 := (1, k - m + 2, k - m + 3, \dots, m, 1)$ is mini-

mal in Γ_k , and it determines both the monomial $f^{\gamma_1} = f^{1,k-m+2,k-m+3,\dots,m}$ and the syzygy

$$\sigma_{\gamma_1} = -\frac{f^{\gamma_1}}{f^{1,m}}\varepsilon_m + \frac{f^{\gamma_1}}{f^{1,k-m+2}}\varepsilon_k + \sum_{j=k-m+2}^{m-1} \frac{f^{\gamma_1}}{f^{j,j+1}}\varepsilon_j. \quad (2.6)$$

Equations (2.5) and the fact that $s_m \neq 0$ imply that f^{γ_1} divides $s_m f^{1,m}$, and a straightforward computation shows that

$$\sigma_1 := \sigma + \frac{s_m f^{1,m}}{f^{\gamma_1}} \sigma_{\gamma_1} = \left(s_k + \frac{s_m f^{1,m}}{f^{1,k-m+2}} \right) \varepsilon_k + \sum_{j=1}^{k-m+1} s_j \varepsilon_j + \sum_{j=m+1}^{k-1} s_j \varepsilon_j.$$

In particular, if we expand $\sigma_1 = \sum_{j=1}^k t_j \varepsilon_j$ for $t_1, \dots, t_k \in S$, then $t_j = 0$ for $k-m+2 \leq j \leq m$, and it suffices to prove the result for σ_1 as claimed.

The second step is to repeat the above, comparing the coefficient of ε_{k-m+2} in the equation $\psi(\sigma_1) = 0$, and since $t_{k-m+2} = 0$ we obtain

$$t_{k-m+1} f^{k-m+1,k-m+2} + t_k f^{1,k-m+2} = 0. \quad (2.7)$$

If $t_k \neq 0$ then the minimal circuit $\gamma_2 := (1, k-m+2, k-m+1, 1)$ in Γ_k from Figure 2(b) determines both the monomial $f^{\gamma_2} = f^{1,k-m+1,k-m+2}$ and the syzygy

$$\sigma_{\gamma_2} = \frac{f^{\gamma_2}}{f^{1,k-m+2}}\varepsilon_k - \frac{f^{\gamma_2}}{f^{k-m+1,k-m+2}}\varepsilon_{k-m+1} - \frac{f^{\gamma_2}}{f^{1,k-m+1}}\varepsilon_{k-1}. \quad (2.8)$$

Equation (2.7) implies that f^{γ_2} divides $t_k f^{1,k-m+2}$ and again, a straightforward computation, this time using equation (2.7), shows that the coefficients of both ε_k and ε_{k-m+1} in the syzygy

$$\sigma_2 := \sigma_1 - \frac{t_k f^{1,k-m+2}}{f^{\gamma_2}} \sigma_{\gamma_2}$$

are zero. This means that $\sigma_2 \in \text{syz}(F^{k-1})$, and we deduce from the inductive hypothesis that σ_2 is generated by the elements σ_γ associated to minimal circuits γ in Γ_{k-1} . Among all minimal circuits in Γ_{k-1} , only $\gamma = (1, k-m+1, k-m+2, \dots, m, 1)$ is not minimal in Γ_k ; indeed, the edge labelled β_k is

a chord. However, this edge splits γ into the circuits γ_1, γ_2 defined earlier in the current proof that *are* minimal in Γ_k , and Lemma 2.4 writes σ_γ as an S -linear combination of σ_{γ_1} and σ_{γ_2} . Thus, the syzygy σ_2 , and hence both σ_1 and σ , are generated by the elements σ_γ associated to minimal circuits γ in Γ_k . This completes the proof for $m \leq k \leq 2m - 3$.

Finally, consider $2m - 2 \leq k \leq n$. Given any monomial order on S , let $>$ denote the term over position order on the free S -module $\bigoplus_{\mu=1}^m S\mathbf{e}_\mu$, that is, $>$ is the monomial order defined for $g, g' \in S$ and $1 \leq \mu, \nu \leq m$ by taking $g'\mathbf{e}_\nu > g\mathbf{e}_\mu$ if and only if $g'f^\nu > gf^\mu$ with respect to the monomial order on S , or $g'f^\nu = gf^\mu$ and $\nu > \mu$. It follows that for $1 \leq j \leq k$, the leading term of β_j with respect to this order is $f^{\mu_j, \nu_j} / f^{\nu_j} \mathbf{e}_{\nu_j}$. This implies that the S -vectors of critical pairs are the elements

$$S(\beta_i, \beta_j) = \frac{f^{\mu_i, \mu_j, \nu_j}}{f^{\mu_j, \nu_j}} \beta_j - \frac{f^{\mu_i, \mu_j, \nu_j}}{f^{\mu_i, \nu_j}} \beta_i$$

arising from all elements in $\mathbb{B}_k := \{(i, j) \mid 1 \leq i < j \leq k, \nu_i = \nu_j\}$ (see Kreuzer–Robbiano [8, Definition 2.5.1]). Substituting (2.1) into every S -vector ensures that the leading terms cancel by definition. Since any critical pair (i, j) corresponds to a pair of directed edges (μ_i, ν_j) and (μ_j, ν_j) in Γ_k , the S -vector can then be written as a multiple of the generator $\beta_{(\mu_i, \mu_j)}$ corresponding to the third directed edge from Figure 3. Indeed, if we choose the index $1 \leq h \leq k$ so that $\beta_h = \beta_{(\mu_i, \mu_j)}$, then we compute explicitly that the ‘standard expressions’ are

$$S(\beta_i, \beta_j) = -\frac{f^{\mu_i, \mu_j, \nu_i}}{f^{\mu_i, \mu_j}} \beta_h.$$




Figure 3. Minimal circuit in Γ_k for $2m - 2 \leq k \leq n$ where $i < j$.

Moreover, we deduce from Buchberger's Criterion [6, Theorem 15.8] that β_1, \dots, β_k are a Gröbner basis of F^k . Every standard expression determines a syzygy, namely

$$\sigma_{(i,j)} = \frac{f^{\mu_i, \mu_j, \nu_j}}{f^{\mu_j, \nu_j}} \varepsilon_j - \frac{f^{\mu_i, \mu_j, \nu_j}}{f^{\mu_i, \nu_j}} \varepsilon_i + \frac{f^{\mu_i, \mu_j, \nu_i}}{f^{\mu_i, \mu_j}} \varepsilon_h. \quad (2.9)$$

Schreyer's theorem [6, Theorem 15.10] implies that the set of syzygies $\{\sigma_{(i,j)} \mid (i,j) \in \mathbb{B}_k\}$ is a system of generators for $\text{syz}(F^k)$. Let $\gamma(i,j) := (\mu_i, \mu_j, \nu_j, \mu_i)$ denote circuit in Γ_k obtained by traversing the edges labelled β_h, β_j according to orientation followed by the edge labelled β_i against orientation (see Figure 3). Then $\sigma_{(i,j)}$ coincides with the syzygy $\sigma_{\gamma(i,j)}$ from Lemma 2.3, and the result is a consequence of the following Lemma. \square

Lemma 2.6 *For $2m-3 \leq k \leq n$, the minimal circuits in the graph Γ_k are precisely those of the form $\gamma(i,j) = (\mu_i, \mu_j, \nu_j, \mu_i)$ arising from pairs (i,j) in $\mathbb{B}_k = \{(i,j) \mid 1 \leq i < j \leq k, \nu_i = \nu_j\}$.*

Proof. We proceed by induction. Let γ be a minimal circuit in Γ_{2m-3} that is not of the form $\gamma(i,j)$ for any $(i,j) \in \mathbb{B}_{2m-3}$. Since γ is a circuit, it must traverse an edge e of the subgraph Γ_m , and since $\gamma \neq \gamma(i,j)$, then either the edge that follows e in γ , or that preceding e in γ , must lie in Γ_m . In either case, γ traverses two edges from Γ_m that share a common vertex μ . The special nature of Γ_{2m-3} then forces the edge $(1, \mu)$ to be a chord of γ , a contradiction. Assume now that the result holds for Γ_{k-1} and let γ be a minimal circuit in Γ_k that is not of the form $\gamma(i,j)$ for any $(i,j) \in \mathbb{B}_k$. If the edge $e_k = (\mu_k, \nu_k)$ does not lie in $\text{supp}(\gamma)$ then the result holds by induction, so we suppose otherwise. Let e be the unique edge in $\text{supp}(\gamma) \setminus \{e_k\}$ that has ν_k as a vertex. There are three cases:

- (i) $e = (\nu_k - 1, \nu_k)$, in which case $(\mu_k, \nu_k - 1)$ is a chord because $\gamma \neq \gamma(\nu_k - 1, k)$;
- (ii) $e = (\nu_k, \nu_k + 1)$, in which case γ must pass through a vertex of the form $1 \leq \mu \leq \mu_k$ since it is a circuit, but then (μ, ν_k) is a chord;
- (iii) $e = (\mu, \nu_k)$ for some $1 \leq \mu < \mu_k$. Since $\gamma \neq \gamma(j, k)$ for any $j < k$, the circuit γ must pass through another vertex of the form $1 \leq \mu' < \mu_k$, but then (μ', ν_k) is a chord.

Thus, the minimal circuit γ cannot exist. \square

- Remark 2.7** (1) If for $2m - 2 \leq k \leq n$ we draw the vertices of Γ_k spaced evenly around a circle centred at the origin in \mathbb{R}^2 , then each minimal circuit γ has length three and hence determines a triangle as in Figure 3. In the spirit of the Taylor resolution of a monomial ideal (see, for example, Bayer–Peeva–Sturmfels [1]), the triangle can be viewed as a 2-cell that defines f^{μ_i, μ_j, ν_j} , and the edges are 1-cells defining f^{μ_i, μ_j} , f^{μ_i, ν_j} and f^{μ_j, ν_j} . The coefficients of the syzygy $\sigma_{(i,j)}$ are then simply the quotients of the monomial for the 2-cell divided by the monomial for the corresponding 1-cell. An analogous statement holds for $m \leq k \leq 2m - 3$, where the syzygies σ_0 and σ_1 from the proofs of Lemma 2.2 and Theorem 2.5 respectively define polygons with more than three sides.
- (2) We emphasise that our choice of order on the set of transpositions of m letters is imposed on us by the geometry: the filtration in Proposition 3.1 below requires that the S -module F^k contains F^0 for $1 \leq k \leq n$. Without this constraint one could choose an alternative order in which each minimal circuit of Γ_k for $m \leq k \leq n$ determines a triangle, leading to a more unified proof of Theorem 2.5. Indeed, since f^1, \dots, f^m are monomials, the modules $\text{syz}(F^k)$ can be read off directly from the Taylor resolution for $1 \leq k \leq m$.

As an application of Theorem 2.5, we introduce a filtration of the module S -module $\ker(\varphi) = \text{syz}(M)$ that feeds into the proof of our main result. For $1 \leq k \leq n$, the S -modules F^k define a filtration

$$0 \subseteq F^1 \subseteq F^2 \subseteq \dots \subseteq F^{n-1} \subseteq F^n = \text{syz}(M)$$

in which the successive quotients are cyclic S -modules

$$\frac{F^k}{F^{k-1}} \cong \frac{\langle \beta_k \rangle}{\langle \beta_1, \dots, \beta_{k-1} \rangle \cap \langle \beta_k \rangle}. \quad (2.10)$$

The next result gives an explicit description of these quotient modules.

Proposition 2.8 *For each $1 \leq k \leq n$, the quotient F^k/F^{k-1} is isomorphic to the cyclic S -module S/I_k , where the monomial ideal I_k depends on k as follows:*

- (i) for $1 \leq k \leq m - 1$, the ideal I_k is the zero ideal;
- (ii) for $k = m$, the ideal I_k is principal with generator $f^{1, \dots, m}/f^{1, m}$;

(iii) for $m + 1 \leq k \leq 2m - 3$, the ideal is

$$I_k = \left\langle \frac{f^{1,k-m+2,k-m+3,\dots,m}}{f^{1,k-m+2}}, \frac{f^{1,k-m+1,k-m+2}}{f^{1,k-m+2}} \right\rangle;$$

(iv) for $2m - 2 \leq k \leq n$, the corresponding transposition is $\tau_k = (\mu_k, \nu_k)$, and the ideal is

$$I_k = \left\langle \frac{f^{\mu_k, \nu_k}}{f^{\mu_k, \nu_k}} \mid \mu \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\} \right\rangle.$$

Proof. For $1 \leq k \leq n$, let $\{\sigma_1, \dots, \sigma_r\}$ be a set of generators for the S -module $\text{syz}(F^k)$. If we write $\sigma_\nu = \sum_{j=1}^k s_{\nu j} \varepsilon_j$ with $s_{\nu 1}, \dots, s_{\nu k} \in S$ for $1 \leq \nu \leq r$, then [8, Proposition 3.2.3] implies that the coefficients s_{1k}, \dots, s_{rk} of ε_k give the generators $s_{1k}\beta_k, \dots, s_{rk}\beta_k$ of the S -module $\langle \beta_1, \dots, \beta_{k-1} \rangle \cap \langle \beta_k \rangle$, so we obtain

$$\frac{F^k}{F^{k-1}} \cong \frac{S}{\langle s_{1k}, \dots, s_{rk} \rangle}.$$

It remains to compute $I_k := \langle s_{1k}, \dots, s_{rk} \rangle$. Parts (i) and (ii) now follow from Lemma 2.2 and equation (2.3). For part (iii), the proof of Theorem 2.5 shows that the only minimal circuits γ in Γ_k with $m + 1 \leq k \leq 2m - 3$ for which the associated syzygy σ_γ has a nonzero coefficient for ε_k are $\gamma_1 := (1, k - m + 2, k - m + 3, \dots, m, 1)$ and $\gamma_2 := (1, k - m + 2, k - m + 1, 1)$. These nonzero coefficients are presented in equations (2.6) and (2.8), namely

$$\frac{f^{\gamma_1}}{f^{1,k-m+2}} = \frac{f^{1,k-m+2,k-m+3,\dots,m}}{f^{1,k-m+2}} \quad \text{and} \quad \frac{f^{\gamma_2}}{f^{1,k-m+2}} = \frac{f^{1,k-m+1,k-m+2}}{f^{1,k-m+2}}.$$

For part (iv), we deduce from Theorem 2.5 and Lemma 2.6 that $\text{syz}(F^k)$ is generated by the syzygies $\sigma_{(i,j)} = \sigma_{\gamma(i,j)}$ associated to pairs $(i, j) \in \mathbb{B}_k$. By equation (2.9), such syzygies have a nonzero coefficient of ε_k if and only if $(i, j) = (i, k)$ for those $1 \leq i < k$ satisfying $\nu_i = \nu_k$. The i th edge (μ_i, ν_i) in Γ_k has $\nu_i = \nu_k$ if and only if $\mu_i \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\}$, that is, we must consider all pairs of the form (μ, ν_k) for $\mu \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\}$. Equation (2.9) shows that the coefficient of ε_k in this case is $f^{\mu, \mu_k, \nu_k} / f^{\mu_k, \nu_k}$ as required. \square

Remark 2.9 The generators of I_k listed in Proposition 2.8 need not be minimal for $m + 1 \leq k \leq n$. For example (though not the simplest), a straightforward calculation for the module M over $S = \mathbb{k}[x_1, \dots, x_7]$ with generators

$$f^1 = x_1x_6, \quad f^2 = x_1x_2x_7, \quad f^3 = x_2x_3, \quad f^4 = x_3x_4, \quad f^5 = x_4x_5x_7, \quad f^6 = x_5x_6$$

gives $I_k = S$ for $k = 9, 10, 12, 13$. Thus, I_k is principal even though this ideal is listed as having more than one generator in Proposition 2.8.

3. Cohomology of wheels on toric varieties

Let X be a normal variety over \mathbb{k} . The divisor class group $\text{Cl}(X)$ is defined to be the group of linear equivalence classes of Weil divisors on X . Since X is normal, two divisors D and D' are linearly equivalent if and only if the associated rank-one reflexive sheaves $\mathcal{O}_X(D)$ and $\mathcal{O}_X(D')$ are isomorphic. We may therefore identify elements of the class group of X with (isomorphism classes of) sheaves of the form $\mathcal{O}_X(D)$. In particular, for a Cartier divisor D on X defining an invertible sheaf $L := \mathcal{O}_X(D)$, we sometimes write $L \in \text{Cl}(X)$.

Let X be a normal toric variety over \mathbb{k} defined by a fan Σ in the real vector space $N \otimes_{\mathbb{Z}} \mathbb{R}$ with underlying lattice N of rank n . Write $\Sigma(1)$ for the set of one-dimensional cones in Σ , set $d := |\Sigma(1)|$, and let $v_\rho \in N$ denote the primitive lattice point on the cone ρ . Each $\rho \in \Sigma(1)$ determines a torus-invariant Weil divisor D_ρ in X , and we let \mathbb{Z}^d denote the free abelian group of torus-invariant Weil divisors. Assume that X has no torus factors. The map $\text{deg}: \mathbb{Z}^d \rightarrow \text{Cl}(X)$ sending D to the sheaf $\mathcal{O}_X(D)$ fits into a short exact sequence of abelian groups

$$0 \rightarrow M \xrightarrow{\text{div}} \mathbb{Z}^d \xrightarrow{\text{deg}} \text{Cl}(X) \rightarrow 0,$$

where M is the lattice dual to N and where $m \in M$ maps to $\text{div}(m) = \sum_{\rho \in \Sigma(1)} \langle m, v_\rho \rangle D_\rho$. The restriction of the map $\text{deg}: \mathbb{Z}^d \rightarrow \text{Cl}(X)$ to the subsemigroup \mathbb{N}^d defines a $\text{Cl}(X)$ -grading of the Cox ring of X which is the semigroup ring $S := \mathbb{k}[x_1, \dots, x_d]$ of \mathbb{N}^d . Explicitly, the degree of a monomial $\prod_{\rho \in \Sigma(1)} x_\rho^{a_\rho} \in S$ is $\mathcal{O}_X(\sum_{\rho \in \Sigma(1)} a_\rho D_\rho) \in \text{Cl}(X)$. Armed with this $\text{Cl}(X)$ -grading of the ring S , Cox [5, Proposition 3.1] introduced an

exact covariant functor

$$\{\text{Cl}(X)\text{-graded } S\text{-modules}\} \longrightarrow \{\text{quasicoherent } \mathcal{O}_X\text{-modules}\} : \\ F \longmapsto \tilde{F} \quad (3.1)$$

from the category of $\text{Cl}(X)$ -graded S -modules to the category of quasi-coherent sheaves on X , and Mustață [9, Theorem 1.1] subsequently showed that the functor is essentially surjective, i.e., that every quasi-coherent sheaf (up to isomorphism) on X lies in the image of this functor. If X is smooth, two such graded modules determine isomorphic sheaves if and only if they agree upto saturation by Cox’s irrelevant ideal $B = (\prod_{\rho \notin \sigma} x_\rho \mid \sigma \in \Sigma)$, but we do not use this fact (until Remark 3.6). The important point for us is that the functor enables us to lift a complex of quasi-coherent sheaves on X to obtain a complex of $\text{Cl}(X)$ -graded S -modules which we can study, and then push down again to the original complex of sheaves.

As described in the introduction, our primary motivation is to study four-term complexes T^\bullet on X of the form (1.1) for some integer $m \geq 2$. In fact, we take as the primary object of study the corresponding diagram of torus-equivariant maps between invertible sheaves on X :

$$(3.2)$$

Every torus-equivariant map is multiplication by a torus-invariant section of an invertible sheaf on X , and we illustrate on each arrow the torus-invariant Cartier divisor of zeros of the corresponding section. Thus, for example, the effective divisor $D_2^1 \in H^0(L_2 \otimes L_{1,2}^{-1}) \cong \text{Hom}(L_{1,2}, L_2)$ denotes the Cartier divisor of zeros of the section that defines the map from $L_{1,2}$ to L_2 . One

can think of any such diagram as a representation of a quiver (arising as the skeleton of a three-dimensional rhombic polyhedron) in the category of invertible sheaves on X .

Throughout, we impose relations on this quiver, whereby each of the two-dimensional rhombic faces of this quiver forms a commutative square, i.e.

$$D_{j+1}^j + D^{j+1} = D_j^{j+1} + D^j, \quad (3.3)$$

$$D_j^{j-1} + D_{j-1,j} = D_j^{j+1} + D_{j,j+1}, \quad (3.4)$$

for $1 \leq j \leq m$ (working modulo m , with indices in the range $1, \dots, m$). We now describe how a diagram of the form (3.2) gives rise to a complex of $\text{Cl}(X)$ -graded S -modules precisely when (3.3) and (3.4) hold. Indeed, let $S(L)$ denote the free S -module with generator \mathbf{e}_L in degree L , and for $1 \leq j \leq m$ let $S(L_j)$ and $S(L_{j,j+1})$ denote the free S -modules with generators \mathbf{e}_j in degree L_j and $\mathbf{e}_{j,j+1}$ in degree $L_{j,j+1}$ respectively. In addition, let f^j , f_{j+1}^j , f_j^{j+1} , $f_{j,j+1}$ denote the monomials in the Cox ring S whose divisors of zeroes are the torus-invariant Cartier divisors D^j , D_{j+1}^j , D_j^{j+1} , $D_{j,j+1}$ from (3.2). Consider the sequence of $\text{Cl}(X)$ -graded S -modules

$$S(L) \xrightarrow{\varphi^3} \bigoplus_{j=1}^m S(L_{j,j+1}) \xrightarrow{\varphi^2} \bigoplus_{j=1}^m S(L_j) \xrightarrow{\varphi^1} S(L), \quad (3.5)$$

with maps

$$\varphi^3(\mathbf{e}_L) = \sum_{j=1}^m f_{j,j+1} \mathbf{e}_{j,j+1}, \quad \varphi^2(\mathbf{e}_{j,j+1}) = f_{j+1}^j \mathbf{e}_{j+1} - f_j^{j+1} \mathbf{e}_j, \quad \varphi^1(\mathbf{e}_j) = f^j \mathbf{e}_L.$$

We claim that the sequence (3.5) is a complex if and only if the relations (3.3) and (3.4) hold. Indeed, (3.5) is a complex if and only if we have

$$(\varphi^2 \circ \varphi^3)(\mathbf{e}_L) = 0 \quad \text{and} \quad (\varphi^1 \circ \varphi^2)(\mathbf{e}_{j,j+1}) = 0 \quad \text{for } 1 \leq j \leq m,$$

which is the case if and only if $f_{j+1}^j f^{j+1} - f_j^{j+1} f^j = 0$ and $f_j^{j-1} f_{j-1,j} - f_j^{j+1} f_{j,j+1} = 0$ for all $1 \leq j \leq m$, and these equations hold if and only if (3.3) and (3.4) hold for all $1 \leq j \leq m$. In summary, the diagram (3.2)

of invertible sheaves in which the relations (3.3) and (3.4) hold determines a complex of $\text{Cl}(X)$ -graded S -modules of the form (3.5). Conversely, to any complex of the form (3.5), one can reverse this procedure to obtain a diagram (3.2) of invertible sheaves on X in which the relations (3.3) and (3.4) hold.

Applying the exact functor (3.1) to the complex (3.5) of $\text{Cl}(X)$ -graded S -modules determines a complex T^\bullet of locally free sheaves on X of the form

$$L \xrightarrow{d^3} \bigoplus_{j=1}^m L_{j,j+1} \xrightarrow{d^2} \bigoplus_{j=1}^m L_j \xrightarrow{d^1} L,$$

where each differential is torus-equivariant, and where the right-hand copy of L lies in degree zero. Moreover, for each $1 \leq j \leq m$ the restriction of the differential d^2 to the summand $L_{j,j+1}$ has image in $L_j \oplus L_{j+1}$ (with indices modulo m). This is the *total chain complex* T^\bullet of the diagram (3.2). The complexes studied by Cautis–Logvinenko [3], Cautis–Craw–Logvinenko [4] and Bocklandt–Craw–Quintero–Vélez [2] that motivated our main result all take this form. The invertible sheaves at the left and right of diagram (3.2) coincide, so the sheaves and the maps between them in diagram (3.2) can be represented equally well in a planar picture as in Figure 4; we call this the *wheel* of invertible sheaves on X .

We now use the results of the previous section to compute the coho-




Figure 4. Wheel of invertible sheaves on X .

mology of the complex T^\bullet . For this purpose, we first note that the map φ^1 is of the form considered in the preceding section, so we may list the generators of its kernel in a sequence β_1, \dots, β_n with $n = \binom{m}{2}$. We also list the generators of the image of φ^2 as

$$\alpha_j := f_{j+1}^j \mathbf{e}_{j+1} - f_j^{j+1} \mathbf{e}_j$$

for $1 \leq j \leq m$. The next proposition is central to the main result of this paper.

Proposition 3.1 *The S -modules*

$$F^k = \begin{cases} \langle \beta_1, \dots, \beta_k, \alpha_{k+1}, \dots, \alpha_m \rangle & \text{for } 1 \leq k \leq m, \\ \langle \beta_1, \dots, \beta_m, \beta_{m+1}, \dots, \beta_j \rangle & \text{for } m+1 \leq j \leq n, \end{cases}$$

define a filtration

$$\text{im}(\varphi^2) = F^0 \subseteq F^1 \subseteq \dots \subseteq F^{n-1} \subseteq F^n = \ker(\varphi^1).$$

Moreover, for $1 \leq k \leq n$ and for the transposition is $\tau_k = (\mu_k, \nu_k)$, the quotient F^k/F^{k-1} is isomorphic to the cyclic $\text{Cl}(X)$ -graded S -module $(S/I_k)(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\text{gcd}(D^{\mu_k}, D^{\nu_k})))$, where the monomial ideal I_k depends on k as follows:

(1) for $1 \leq k \leq m$, the ideal is

$$I_k = \left\langle \text{gcd}(f_{k+1}^k, f_k^{k+1}), \frac{\text{lcm}(f^1, \dots, f^m, \text{gcd}(f_{k+2}^{k+1}, f_{k+1}^{k+2}), \dots, \text{gcd}(f_1^m, f_m^1))}{f^{k, k+1}} \right\rangle;$$

(2) for $m+1 \leq k \leq 2m-3$, the ideal is

$$I_k = \left\langle \frac{f^{1, k-m+2, k-m+3, \dots, m}}{f^{1, k-m+2}}, \frac{f^{1, k-m+1, k-m+2}}{f^{1, k-m+2}} \right\rangle;$$

(3) for $2m-2 \leq k \leq n$, the ideal is

$$I_k = \left\langle \frac{f^{\mu, \mu_k, \nu_k}}{f^{\mu_k, \nu_k}} \mid \mu \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\} \right\rangle.$$

Proof. To prove that the S -modules F^k define a filtration, we need only show that $\alpha_k \in F^k$ for all $1 \leq k \leq m$. For this, relation (3.3) gives

$$D^k - \gcd(D^k, D^{k+1}) = D_{k+1}^k - \gcd(D_{k+1}^k, D_k^{k+1}), \quad (3.6)$$

and hence

$$\frac{f^{k,k+1}}{f^{k+1}} = \frac{\text{lcm}(f^k, f^{k+1})}{f^{k+1}} = \frac{f^k}{\gcd(f^k, f^{k+1})} = \frac{f_{k+1}^k}{\gcd(f_{k+1}^k, f_k^{k+1})}.$$

Similarly, we have $f^{k,k+1}/f^k = f_k^{k+1}/\gcd(f_{k+1}^k, f_k^{k+1})$. Therefore

$$\begin{aligned} \alpha_k &= \gcd(f_{k+1}^k, f_k^{k+1}) \left(\frac{f^{k,k+1}}{f^{k+1}} \mathbf{e}_{k+1} - \frac{f^{k,k+1}}{f^k} \mathbf{e}_k \right) \\ &= \gcd(f_{k+1}^k, f_k^{k+1}) \beta_k \end{aligned} \quad (3.7)$$

for $1 \leq k \leq m$ as required. To prove part (1), we first note that

$$\frac{F^k}{F^{k-1}} \cong \frac{\langle \beta_k \rangle / (\langle \beta_1, \dots, \beta_{k-1}, \alpha_{k+1}, \dots, \alpha_m \rangle \cap \langle \beta_k \rangle)}{\langle \alpha_k \rangle / (\langle \beta_1, \dots, \beta_{k-1}, \alpha_{k+1}, \dots, \alpha_m \rangle \cap \langle \alpha_k \rangle)}.$$

In order to compute this quotient, it suffices, in view of (3.7) and the remarks at the beginning of the proof of Proposition 2.8, to determine a set of generators for the module of syzygies on $\beta_1, \dots, \beta_k, \alpha_{k+1}, \dots, \alpha_m$ for $1 \leq k \leq m$. Proceeding exactly as in the proof of Lemma 2.2, we find that this module is cyclic with generator

$$\begin{aligned} \sigma_0 &:= - \frac{\text{lcm}(f^{1,\dots,m}, g^{k+1,k+2}, \dots, g^{m,1})}{f^{1,m}} \varepsilon_m \\ &\quad + \sum_{j=1}^{m-1} \frac{\text{lcm}(f^{1,\dots,m}, g^{k+1,k+2}, \dots, g^{m,1})}{f^{j,j+1}} \varepsilon_j, \end{aligned} \quad (3.8)$$

where we have set $g^{i,i+1} := \gcd(f_{i+1}^i, f_i^{i+1})$ for $k+1 \leq i \leq m$. Ignoring for now the $\text{Cl}(X)$ -grading, we deduce from this that

$$\begin{aligned} & \frac{\langle \beta_k \rangle}{\langle \beta_1, \dots, \beta_{k-1}, \alpha_{k+1}, \dots, \alpha_m \rangle \cap \langle \beta_k \rangle} \\ & \cong \frac{S}{\langle \text{lcm}(f^1, \dots, m, g^{k+1, k+2}, \dots, g^{m, 1}) / f^{k, k+1} \rangle}. \end{aligned}$$

and therefore, by virtue of (3.7),

$$\frac{F^k}{F^{k-1}} \cong \frac{S}{\langle \text{gcd}(f_{k+1}^k, f_k^{k+1}), \text{lcm}(f^1, \dots, m, g^{k+1, k+2}, \dots, g^{m, 1}) / f^{k, k+1} \rangle}$$

which gives the ideal I_k in part (1). For parts (2) and (3), Proposition 2.8(iii) and (iv) respectively determine the ideals I_k for which F^k/F^{k-1} is isomorphic to S/I_k as ungraded rings.

It remains to establish the isomorphism as $\text{Cl}(X)$ -graded rings. In light of the above and isomorphism (2.10), it suffices to show that the degree of β_k is $L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\text{gcd}(D^{\mu_k}, D^{\nu_k}))$ for $1 \leq k \leq n$. For each $1 \leq k \leq n$, multiplication by the monomials f^{μ_k} and f^{ν_k} define $\text{Cl}(X)$ -graded maps $S \rightarrow S(L \otimes L_{\mu_k}^{-1})$ and $S \rightarrow S(L \otimes L_{\nu_k}^{-1})$ respectively. Tensoring each map with $S(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1})$ yields $\text{Cl}(X)$ -graded maps $S(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}) \rightarrow S(L_{\nu_k})$ and $S(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}) \rightarrow S(L_{\mu_k})$ which, in turn, can be combined to form a $\text{Cl}(X)$ -graded map

$$S(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}) \longrightarrow \bigoplus_{j=1}^m S(L_j),$$

whose image in $\bigoplus_{j=1}^m S(L_j)$ is generated by the element $f^{\mu_k} \mathbf{e}_{\nu_k} - f^{\nu_k} \mathbf{e}_{\mu_k}$. Twisting further by $S(\mathcal{O}_X(\text{gcd}(D^{\mu_k}, D^{\nu_k})))$ determines a $\text{Cl}(X)$ -graded map

$$S(L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\text{gcd}(D^{\mu_k}, D^{\nu_k}))) \longrightarrow \bigoplus_{j=1}^m S(L_j)$$

whose image is generated by the element

$$\frac{f^{\mu_k}}{\text{gcd}(f^{\mu_k}, f^{\nu_k})} \mathbf{e}_{\nu_k} - \frac{f^{\nu_k}}{\text{gcd}(f^{\mu_k}, f^{\nu_k})} \mathbf{e}_{\mu_k}. \quad (3.9)$$

To prove the claim it remains to show that (3.9) coincides with β_k ,

but this is immediate since $f^{\mu_k} / \gcd(f^{\mu_k}, f^{\nu_k}) = \text{lcm}(f^{\mu_k}, f^{\nu_k}) / f^{\nu_k}$ and $f^{\nu_k} / \gcd(f^{\mu_k}, f^{\nu_k}) = \text{lcm}(f^{\mu_k}, f^{\nu_k}) / f^{\mu_k}$. \square

For $1 \leq k \leq n$, each of the generators of I_k listed in Proposition 3.1 is a monomial in the Cox ring S of X , so its divisor of zeros is an effective torus-invariant Weil divisor in X . Notice that while $f^j, f_{j+1}^j, f_j^{j+1}, f_{j,j+1}$ define torus-invariant Cartier divisors $D^j, D_{j+1}^j, D_j^{j+1}, D_{j,j+1}$ in X , the generators of the ideals I_k are Weil divisors in general.

Definition 3.2 For each $1 \leq k \leq n$, define a subscheme $Z_k \subset X$ to be the scheme-theoretic intersection of a set of effective Weil divisors depending on k as follows:

- (i) for $1 \leq k \leq m$, define Z_k to be the scheme-theoretic intersection of $\gcd(D_{k+1}^k, D_k^{k+1})$ and the divisor $\text{lcm}(D^1, \dots, D^m, \gcd(D_{k+2}^{k+1}, D_{k+1}^{k+2}), \dots, \gcd(D_1^m, D_m^1)) - \text{lcm}(D^k, D^{k+1})$;
- (ii) for $m+1 \leq k \leq 2m-3$, define Z_k to be the scheme-theoretic intersection of the divisors $\text{lcm}(D^1, D^{\nu_k}, D^{\nu_k+1}, \dots, D^m) - \text{lcm}(D^1, D^{\nu_k})$ and $\text{lcm}(D^1, D^{\nu_k-1}, D^{\nu_k}) - \text{lcm}(D^1, D^{\nu_k})$;
- (iii) for $2m-2 \leq k \leq n$, define Z_k to be the scheme-theoretic intersection of the divisors $\text{lcm}(D^\mu, D^{\mu_k}, D^{\nu_k}) - \text{lcm}(D^{\mu_k}, D^{\nu_k})$ for $\mu \in \{1, \dots, \mu_k - 1\} \cup \{\nu_k - 1\}$.

The subschemes $Z_k \subset X$ are torus-invariant, though some (possibly all) may be empty, see Example 3.5 for an explicit calculation. These subschemes enable us to formulate and prove the main result of this paper (this is Theorem 1.1 from the introduction).

Theorem 3.3 *Let X be a normal toric variety and let T^\bullet be the complex from (1.1), with differentials determined by the Cartier divisors shown in (1.2). Then:*

- (1) $H^0(T^\bullet) \cong \mathcal{O}_Z \otimes L$ where Z is the scheme-theoretic intersection of D^1, \dots, D^m ;
- (2) $H^{-1}(T^\bullet)$ has an n -step filtration

$$\text{im}(d^2) = F^0 \subseteq F^1 \subseteq \dots \subseteq F^{n-1} \subseteq F^n = \ker(d^1)$$

where, for $1 \leq k \leq n$ and for the permutation $\tau_k = (\mu_k, \nu_k)$, we have

$$F^k/F^{k-1} \cong \mathcal{O}_{Z_k} \otimes L_{\mu_k} \otimes L_{\nu_k} \otimes L^{-1}(\gcd(D^{\mu_k}, D^{\nu_k})); \quad (3.10)$$

- (3) $H^{-2}(T^\bullet) \cong \mathcal{O}_D \otimes L(D)$ where $D = \gcd(D_{1,2}, D_{2,3}, \dots, D_{m,1})$;
(4) $H^{-3}(T^\bullet) \cong 0$.

Proof. As described at the beginning of this section, the complex T^\bullet arises from a diagram (3.2) of invertible sheaves on X in which the relations (3.3) and (3.4) hold, and every such diagram determines a complex of $\text{Cl}(X)$ -graded S -modules of the form (3.5), where one can reproduce the original complex T^\bullet by applying the exact functor (3.1). In particular, one can calculate the cohomology sheaves of T^\bullet by computing the cohomology modules of (3.5) and applying the Cox functor. The statement of part (2) then follows from Proposition 3.1 and Definition 3.2.

For part (1), note that $H^0(T^\bullet)$ is the cokernel of $\bigoplus_i \mathcal{O}_X(-D^i) \otimes L \hookrightarrow \mathcal{O}_X \otimes L$, namely the sheaf $\mathcal{O}_Z \otimes L$ where Z is the scheme-theoretic intersection of D^1, \dots, D^m . For part (4), every nonzero map between invertible sheaves is injective, so $H^{-3}(T^\bullet) \cong 0$. It remains to prove part (3). The proof of the analogous statement from [3, Lemma 3.1] does not immediately extend to our setting, as was the case with parts (1) and (4) above, but we can nevertheless adapt the argument as follows. We claim first that if the greatest common divisor D is zero then $H^{-2}(T^\bullet) \cong 0$. We need only show that complex (3.5) has no cohomology in degree -2 . Indeed, suppose $\boldsymbol{\eta} = \sum_{j=1}^m u_j \mathbf{e}_{j,j+1}$ lies in the kernel of φ^2 , so

$$0 = \varphi^2(\boldsymbol{\eta}) = \sum_{j=1}^m u_j (f_{j+1}^j \mathbf{e}_{j+1} - f_j^{j+1} \mathbf{e}_j).$$

This translates into the following set of equations:

$$u_{j-1} f_j^{j-1} = u_j f_j^{j+1} \quad 1 \leq j \leq m.$$

By relation (3.4) we have $f_j^{j-1} f_{j-1,j} = f_j^{j+1} f_{j,j+1}$ for $1 \leq j \leq m$. Consequently, we find that

$$u_{j-1} f_{j,j+1} = u_j f_{j-1,j}, \quad 1 \leq j \leq m. \quad (3.11)$$

We claim that $f_{j,j+1}$ divides u_j for all $1 \leq j \leq m$. It suffices to prove that

$f_{1,2}$ divides u_1 by virtue of (3.11). Let x_i be a prime factor of $f_{1,2}$ with multiplicity p . Since by assumption $\gcd(f_{1,2}, f_{2,3}, \dots, f_{m,1}) = 1$, it follows that x_i^p does not divide $f_{\nu, \nu+1}$ for some $\nu \neq 1$. Appealing to (3.11) once again, we find that $u_1 f_{\nu, \nu+1} = u_\nu f_{1,2}$, and thus x_i^p divides $u_1 f_{\nu, \nu+1}$. Since S is a unique factorisation domain, this means that x_i^p divides u_1 , which in turn implies that $f_{1,2}$ divides u_1 . If we now set $u := u_1/f_{1,2}$, then equations (3.11) give

$$u = \frac{u_1}{f_{1,2}} = \frac{u_2}{f_{2,3}} = \dots = \frac{u_m}{f_{m,1}},$$

from which it follows that $\boldsymbol{\eta} = u \sum_{j=1}^m f_{j,j+1} \mathbf{e}_{j,j+1}$. Thus, $\boldsymbol{\eta}$ lies in the image of φ^3 , so the complex (3.5) has no cohomology in degree -2 as required.

To complete the proof of part (3), suppose $D \neq 0$. We can factor $d^3: T^{-3} \rightarrow T^{-2}$ as a map $L \rightarrow L(D)$ followed by a map with no common divisors. By the above argument, the image of $L(D)$ under this map equals the kernel of $d^2: T^{-2} \rightarrow T^{-1}$. Therefore $H^{-2}(T^\bullet)$ can be identified with the cokernel of $L \rightarrow L(D)$, which is $\mathcal{O}_D \otimes L(D)$. \square

Remark 3.4 For $m = 3$, Theorem 1.1 agrees with the statement of the main technical result from Cautis–Logvinenko [3, Lemma 3.1] (recall from the discussion surrounding Example 1.3 above that the assumptions from *loc. cit.*, namely that X is an arbitrary smooth separated scheme, should be replaced by the assumptions of Theorem 1.1). Parts (1), (3), (4) of Theorem 1.1 clearly generalise the analogues from [3, Lemma 3.1]. As for $H^{-1}(T^\bullet)$, we have $m = 3$ and hence $n = 3$, so Theorem 1.1(2) gives a 3-step filtration

$$\mathrm{im}(d^2) = F^0 \subseteq F^1 \subseteq F^2 \subseteq F^3 = \ker(d^1),$$

and we claim that the successive quotients agree with those of *loc. cit.*. To justify this we first compute F^2/F^1 . Since $\tau_2 = (2, 3)$, Theorem 1.1(2) shows that

$$F^2/F^1 \cong \mathcal{O}_{Z_2} \otimes L_2 \otimes L_3 \otimes L^{-1}(\gcd(D^2, D^3)),$$

where Z_2 is the intersection of $\gcd(D_3^2, D_2^3)$ and $\mathrm{lcm}(D^1, D^2, D^3, \gcd(D_1^3, D_3^1)) - \mathrm{lcm}(D^2, D^3)$. A direct computation shows that the relation defined by the generator σ_0 from (3.8) is

$$\frac{f_1^3}{\gcd(f_1^3, \tilde{f}_1^2)} \beta_1 + \frac{\tilde{f}_2^1 f_1^3}{\gcd(f_1^3, \tilde{f}_1^2) \tilde{f}_2^3} \beta_2 - \frac{\tilde{f}_1^2}{\gcd(f_1^3, \tilde{f}_1^2)} \alpha_3 = 0,$$

where $\tilde{f}_j^i = f_j^i / \gcd(f_j^i, f_i^j)$. Since $k = 2$, the coefficient of β_2 coincides with the generator $\text{lcm}(f^{1,2,3}, \gcd(f_1^3, f_3^1)) / f^{2,3}$ of the ideal I_2 . In particular, the scheme Z_2 is the intersection of $\gcd(D_3^2, D_2^3)$ and $\tilde{D}_2^1 + D_1^3 - \tilde{D}_2^3 - \gcd(D_1^3, \tilde{D}_1^2)$, where \tilde{D}_j^i is the divisor of zeros of the function \tilde{f}_j^i . Permutations are listed as $\tau_1 = (1, 2)$, $\tau_2 = (3, 1)$, $\tau_3 = (2, 3)$ in [3], so after applying permutation $(1, 2, 3)$ to our indices, we need only invoke the identity

$$\tilde{D}_3^2 + D_2^1 - \tilde{D}_3^1 - \gcd(D_2^1, \tilde{D}_2^3) = D^2 + \text{lcm}(D_2^1, \tilde{D}_2^3) - D^3 - \tilde{D}_3^1$$

from [3, p206] to see that Z_2 is the scheme in the second bullet point of [3, Lemma 3.1(2)]. In order to compare the sheaves, equation (3.6) gives $\gcd(D^2, D^3) = D^2 + \gcd(D_3^2, D_2^3) - D_3^2$, and $\mathcal{O}_X(D^2) = L_2^{-1} \otimes L$ and $\mathcal{O}_X(-D_3^2) \cong L_3^{-1} \otimes L_{2,3}$ hence

$$\begin{aligned} & L_2 \otimes L_3 \otimes L^{-1}(\gcd(D^2, D^3)) \\ & \cong L_2 \otimes L_3 \otimes L^{-1}(\gcd(D_3^2, D_2^3)) \otimes L_2^{-1} \otimes L \otimes L_3^{-1} \otimes L_{2,3} \\ & \cong L_{2,3}(\gcd(D_2^3, D_3^2)). \end{aligned}$$

Again, applying the permutation $(1, 2, 3)$ to the indices recovers the sheaf from the second bullet point of [3, Lemma 3.1(2)], so our description of F^2/F^1 agrees with that from *loc.cit.*. A very similar calculation shows that our unified description of the quotients F^k/F^{k-1} for $k = 1, 3$ agrees with those of F^3/F^2 and F^1/F^0 from [3, Lemma 3.1(2)].

Example 3.5 Let X be the smooth toric threefold determined by the fan Σ in \mathbb{R}^3 whose one-dimensional cones are generated by the vectors

$$\begin{aligned} v_1 &= (1, 0, 1), & v_2 &= (0, 1, 1), & v_3 &= (-1, 1, 1), \\ v_4 &= (-1, 0, 1), & v_6 &= (1, -1, 1), & v_7 &= (0, 0, 1), \end{aligned}$$

where the cones in higher dimension are best illustrated by the height one slice of Σ as shown in Figure 5. In particular, the Cox ring of X is $S = \mathbb{k}[x_1, \dots, x_7]$ and the Cox irrelevant ideal is the monomial ideal $B =$

$$\begin{aligned}
\beta_3 &= -x_4\mathbf{e}_3 + x_2\mathbf{e}_4, & \beta_{11} &= -x_4x_5\mathbf{e}_2 + x_1x_2\mathbf{e}_5, \\
\beta_4 &= -x_5x_7\mathbf{e}_4 + x_3\mathbf{e}_5, & \beta_{12} &= -x_5x_6\mathbf{e}_2 + x_1x_2x_7\mathbf{e}_6, \\
\beta_5 &= -x_6\mathbf{e}_5 + x_4x_7\mathbf{e}_6, & \beta_{13} &= -x_4x_5x_7\mathbf{e}_3 + x_2x_3\mathbf{e}_5, \\
\beta_6 &= x_5\mathbf{e}_1 - x_1\mathbf{e}_6, & \beta_{14} &= -x_5x_6\mathbf{e}_3 + x_2x_3\mathbf{e}_6, \\
\beta_7 &= -x_2x_3\mathbf{e}_1 + x_1x_6\mathbf{e}_3, & \beta_{15} &= -x_5x_6\mathbf{e}_4 + x_3x_4\mathbf{e}_6. \\
\beta_8 &= -x_3x_4\mathbf{e}_1 + x_1x_6\mathbf{e}_4,
\end{aligned}$$

It is easy to see that the relations

$$\begin{aligned}
\beta_9 &= -x_4x_7\beta_6 - x_1\beta_5, & \beta_{10} &= x_4\beta_2 + x_1x_7\beta_3, \\
\beta_{12} &= -x_5\beta_1 - x_2x_7\beta_6, & \beta_{13} &= x_5x_7\beta_3 + x_2\beta_4
\end{aligned}$$

hold, so the successive quotients F^k/F^{k-1} vanish for $k = 9, 10, 12, 13$. In addition, the generators $\alpha_1, \dots, \alpha_6$ of $\text{im}(d^2)$ satisfy $\alpha_1 = \beta_1$, $\alpha_2 = \beta_2$, $\alpha_3 = x_7\beta_3$, $\alpha_4 = \beta_4$, $\alpha_5 = \beta_5$ and $\alpha_6 = x_7\beta_6$, so F^k/F^{k-1} also vanishes for $k = 1, 2, 4, 5$.

We now analyse three nonvanishing quotients F^k/F^{k-1} to illustrate part (2) of Theorem 1.1. First consider $k = 3$. The transposition $\tau_3 = (3, 4)$ determines $\text{gcd}(D^3, D^4) = E_3$, so

$$F^3/F^2 \cong \mathcal{O}_{Z_3} \otimes L_3 \otimes L_4 \otimes L^{-1}(E_3)$$

where, according to Definition 3.2(i), Z_3 is the scheme-theoretic intersection of the effective torus-invariant divisors $\text{gcd}(D_4^3, D_3^4) = E_7$ and

$$\begin{aligned}
&\text{lcm}(D^1, D^2, D^3, D^4, D^5, D^6, \\
&\text{gcd}(D_5^4, D_4^5), \text{gcd}(D_6^5, D_5^6), \text{gcd}(D_1^6, D_6^1)) - \text{lcm}(D^3, D^4) = E_{1567}.
\end{aligned}$$

In particular, $\text{supp}(\mathcal{O}_{Z_3}) = E_7$. Now consider the case $k = 7$. The corresponding transposition $\tau_7 = (1, 3)$ determines $\text{gcd}(D^1, D^3) = 0$, so

$$F^7/F^6 \cong \mathcal{O}_{Z_7} \otimes L_1 \otimes L_3 \otimes L^{-1}$$

where, according to Definition 3.2(ii), Z_7 is the scheme-theoretic intersection of the divisors $\text{lcm}(D^1, D^2, D^3) - \text{lcm}(D^1, D^3) = E_7$ and $\text{lcm}(D^1, D^3, D^4)$,

$D^5, D^6) - \text{lcm}(D^1, D^3) = E_{457}$, giving $Z_7 = E_7 \cap E_{457}$ and $\text{supp}(\mathcal{O}_{Z_7}) = E_7$. Finally, consider the case $k = 15$ for which the corresponding transposition $\tau_{15} = (4, 6)$ determines $\text{gcd}(D^4, D^6) = 0$, so

$$F^{15}/F^{14} \cong \mathcal{O}_{Z_{15}} \otimes L_4 \otimes L_6 \otimes L^{-1}$$

where, according to Definition 3.2(iii), Z_{15} is the scheme-theoretic intersection of the divisors $\text{lcm}(D^\mu, D^4, D^6) - \text{lcm}(D^4, D^6)$ for $\mu = 1, 2, 3, 5$, giving $Z_{15} = E_1 \cap E_{127} \cap E_2 \cap E_7$. In particular, the support of $\mathcal{O}_{Z_{15}}$ is the torus-invariant point $E_1 \cap E_2 \cap E_7$ in X .

As for $H^k(T^\bullet)$ for $k \neq -1$, notice that the scheme theoretic intersection of D^1, \dots, D^6 is contained in $D^1 \cap D^4 = (E_1 + E_6) \cap (E_3 + E_4) = \emptyset$, so $H^0(T^\bullet) \cong 0$ by Theorem 1.1(1). Similarly, $\text{gcd}(D_{1,2}, D_{2,3}, D_{3,4}, D_{4,5}, D_{5,6}, D_{6,1}) = 0$ so $H^{-2}(T^\bullet) \cong 0$ by Theorem 1.1(3). It follows that the complex T^\bullet has cohomology concentrated in degree -1 .

Remark 3.6 One can carry out much of the above calculation using Macaulay2 [7] in any given example, though the final description of F^k/F^{k-1} is less user-friendly and geometric than ours. To give the flavour, we reproduce some of the calculations from Example 3.5, omitting for brevity the information on the degree in the $\text{Cl}(X)$ -grading of each S -module generator¹.

```
S = QQ[x_1,x_2,x_3,x_4,x_5,x_6,x_7];
d1 = matrix{{x_1*x_6,x_1*x_2*x_7,x_2*x_3,x_3*x_4,x_4*x_5*x_7,
             x_5*x_6}}
d2 = matrix{{-x_2*x_7,0,0,0,0,-x_5*x_7},{x_6,x_3,0,0,0,0},
             {0,-x_1*x_7,x_4*x_7,0,0,0},
             {0,0,-x_2*x_7,-x_5*x_7,0,0},{0,0,0,x_3,x_6,0},
             {0,0,0,0,-x_4*x_7,x_1*x_7}}
d3 = matrix{ {-x_3*x_4*x_5},{x_4*x_5*x_6},{x_1*x_5*x_6},
             {-x_1*x_2*x_6},{x_1*x_2*x_3},{x_2*x_3*x_4}}
T = chainComplex(d1,d2,d3)
```

The minimal generators $\{\beta_j \mid j \in \{1, \dots, 15\} \setminus \{9, 10, 12, 13\}\}$ can be

¹Macaulay2 require the $\text{Cl}(X)$ -degree information in order to create the chain complex T , so for convenience we include the complete M2 commands at the end of the latex source file.

obtained using

```
ker d1
```

though Macaulay2 chooses an order on these generators that differs from ours. To obtain the cohomology sheaf $H^{-k}(T^\bullet)$ we compute the k th cohomology of T and saturate by the irrelevant ideal. For example, the commands

```
B = ideal(x_3*x_4*x_5*x_6,x_2*x_3*x_4*x_7,x_2*x_3*x_4*x_6,
          x_1*x_5*x_6*x_7,x_1*x_3*x_5*x_6,x_1*x_2*x_3*x_6 )
H0 = prune HH_0(T)
prune (H0/ saturate(O_S*H0,B))
```

show that $H^0(T^\bullet) \cong 0$. Similarly $H^{-2}(T^\bullet) = 0$. As for the filtration on $H^{-1}(T^\bullet)$, we input the submodules F^k by hand and compute the quotients, for example,

```
F2=image matrix{{-x_2*x_7,0,0,0,0,-x_5*x_7}, {x_6,x_3,0,0,0,0},
                {0,-x_1*x_7,x_4*x_7,0,0,0},{0,0,-x_2*x_7,-x_5*x_7,0,0},
                {0,0,0,x_3,x_6,0},{0,0,0,0,-x_4*x_7,x_1*x_7}}
F3=image matrix{{-x_2*x_7,0,0,0,0,-x_5*x_7}, {x_6,x_3,0,0,0,0},
                {0,-x_1*x_7,x_4,0,0,0}, {0,0,-x_2,-x_5*x_7,0,0},
                {0,0,0,x_3,x_6,0},{0,0,0,0,-x_4*x_7,x_1*x_7}}
Q3 = F3/F2
prune Q3
```

In this case, the output is

```
cokernel | x_7 |
```

so we reproduce our result that F^3/F^2 is supported on the divisor E_7 . Similar, input

```
F15=image matrix{
  {-x_2*x_7,0,0,0,0,x_5,-x_2*x_3,-x_3*x_4,-x_4*x_5*x_7,
   0,0,0,0,0,0},
  {x_6,-x_3,0,0,0,0,0,0,-x_3*x_4,-x_4*x_5,-x_5*x_6,
   0,0,0},
  {0,x_1*x_7,-x_4,0,0,0,x_1*x_6,0,0,0,0,0,-x_4*x_5*x_7,
   -x_5*x_6,0},
```

$$\{0, 0, x_2, -x_5x_7, 0, 0, 0, x_1x_6, 0, x_1x_2x_7, 0, 0, 0, 0, -x_5x_6\},$$

$$\{0, 0, 0, x_3, -x_6, 0, 0, 0, x_1x_6, 0, x_1x_2, 0, x_2x_3, 0, 0\},$$

$$\{0, 0, 0, 0, x_4x_7, -x_1, 0, 0, 0, 0, 0, x_1x_2x_7, 0, x_2x_3, x_3x_4\}$$

and F14 (simply delete the final column in the above), then compute

```
Q15 = F15/F14
prune Q15
```

In this case, the output is

```
cokernel | x_7 x_2 x_1 |
```

This confirms our calculation from Example 3.5 that F^{15}/F^{14} is supported on the torus-invariant point $E_1 \cap E_2 \cap E_7$.

Acknowledgements. Thanks to Raf Bocklandt for generating Example 3.5 and to Sonja Petrovic for comments on an earlier version of this paper. Thanks also to the anonymous referees for many helpful remarks. Our results owe much to experiments made with Macaulay2 [7]. Both authors were supported by EPSRC grant EP/G004048/1.

References

- [1] Bayer D., Peeva I. and Sturmfels B., *Monomial resolutions*. Math. Res. Lett., **5(1–2)** (1998), 31–46.
- [2] Bocklandt R., Craw A. and Quintero Vélez A., 2013. *Geometric Reid’s recipe for dimer models*. Math. Ann., doi:10.1007/s00208-014-1085-8.
- [3] Cautis S. and Logvinenko T., *A derived approach to geometric McKay correspondence in dimension three*. J. Reine Angew. Math., **636** (2009), 193–236.
- [4] Cautis S., Craw A. and Logvinenko T., 2012. *Derived Reid’s recipe for abelian subgroups of $SL(3, \mathbb{C})$* . J. Reine Angew. Math., doi:10.1515/crelle-2014-0086.
- [5] Cox D., *The homogeneous coordinate ring of a toric variety*. J. Algebraic Geom., **4(1)** (1995), 17–50.
- [6] Eisenbud D., *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.

- [7] Grayson D. and Stillman M., *Macaulay 2, a software system for research in algebraic geometry*. Available at www.math.uiuc.edu/Macaulay2/.
- [8] Kreuzer M. and Robbiano L., *Computational commutative algebra. 1*. Springer-Verlag, Berlin, 2000.
- [9] Mustața M., *Vanishing theorems on toric varieties*. Tohoku Math. J. (2), **54(3)** (2002), 451–470.
- [10] Nakamura I., *Hilbert schemes of abelian group orbits*. J. Algebraic Geom., **10(4)** (2001), 757–779.
- [11] Ohsugi H. and Hibi T., *Toric ideals generated by quadratic binomials*. J. Algebra, **218(2)** (1999), 509–527.
- [12] Villarreal R., *Monomial algebras*, volume 238 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 2001.

Alastair Craw
Department of Mathematical Sciences
University of Bath
Bath BA2 7AY
UK
E-mail: A.Craw@bath.ac.uk

Alexander Quintero Vélez
Departamento de Matemáticas
Universidad del Valle
Calle 13 No. 100-00, Edificio 320-3190
Ciudadela Universitaria Meléndez
Cali, Colombia
E-mail: alexander.quintero@correounivalle.edu.co