



## THE SEQUENCE SPACE $E_n^q(M, p, s)$ AND $N_k$ - LACUNARY STATISTICAL CONVERGENCE

NAIM L. BRAHA<sup>1\*</sup> AND MIKÂIL ET<sup>2</sup>

Communicated by F. Kittaneh

ABSTRACT. In this paper we define the sequence space  $E_n^q(M, p, s)$  by using an Orlicz function and we study various properties and obtain some inclusion relations involving this space. We give some relations between  $N_k$ -lacunary statistical convergence and strongly  $N_k$ -lacunary convergence.

### 1. INTRODUCTION AND PRELIMINARIES

Let  $\sum_{k=0}^{\infty} a_k$  be an infinite series with sequence of partial sums  $(s_k)$  and  $q > 0$  any real number. The Euler means  $(E, q)$  of the sequences  $(s_n)$  are defined to be

$$E_n^q = \frac{1}{(1+q)^n} \sum_{v=0}^n \binom{n}{v} q^{n-v} s_v.$$

The series  $\sum_{n=0}^{\infty} a_n$  is said to be summable  $(E, q)$  to the number  $s$  if

$$E_n^q = \frac{1}{(1+q)^n} \sum_{v=0}^n \binom{n}{v} q^{n-v} s_v \rightarrow s \quad \text{as } n \rightarrow \infty,$$

and is said to be absolutely summable  $(E, q)$  or summable  $|E, q|$ , if

$$\sum_k |E_k^q - E_{k-1}^q| < \infty.$$

*Date:* Received: 8 May 2012; Accepted: 19 July 2012.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 40A05; Secondary 40C05, 46A45.

*Key words and phrases.* Orlicz function, sequence space, Euler mean, statistical convergence.

Let  $a = (a_n)$  be a sequence of scalars, for  $k \geq 1$  we will denote by  $N_n(a) = E_n^q - E_{n-1}^q$ , where  $E_n^q$  is defined as above. After applications of Abel's transform we have

$$N_n(a) = -\frac{1}{(1+q)^{n-1}} \sum_{k=0}^{n-2} a_{k+1} A_k + \frac{s_{n-1} A_{n-1}}{(1+q)^{n-1}} + \frac{s_n}{(1+q)^n} - \frac{q^{n-1}}{(1+q)^n} s_0,$$

where  $A_k = \sum_{i=0}^k \left[ \frac{q}{1+q} \binom{n}{i} - \binom{n-1}{i} \right] q^{n-i-1}$ .

Note that for any sequences  $a = (a_n)$ ,  $b = (b_n)$  and scalar  $\lambda$ , we have:

$$N_n(a+b) = N_n(a) + N_n(b) \quad \text{and} \quad N_n(\lambda a) = \lambda N_n(a).$$

An Orlicz function is a function  $M : [0, \infty) \rightarrow [0, \infty)$  which is continuous, nondecreasing and convex with  $M(0) = 0$ ,  $M(x) > 0$  for  $x > 0$  and  $M(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . It is well known that if  $M$  is a convex function and  $M(0) = 0$ ; then  $M(\lambda x) \leq \lambda M(x)$  for all  $\lambda$  with  $0 < \lambda \leq 1$ . Two Orlicz functions  $M_1$  and  $M_2$  are said to be equivalent if there are positive constants  $\alpha$  and  $\beta$ , and  $x_0$  such that  $M_1(\alpha x) \leq M_2(x) \leq M_1(\beta x)$  for all  $x$  with  $0 \leq x \leq x_0$ . (see [15, 16]).

Lindentrauss and Tzafirir [17] used the idea of Orlicz function and defined the following sequence space

$$l_M = \left\{ x = (x_i) \in w : \sum_{i=1}^{\infty} M\left(\frac{|x_i|}{\varrho}\right) < \infty, \varrho > 0 \right\}$$

which is called an Orlicz sequence space. The space  $l_M$  is a Banach space with the norm

$$\|x\| = \inf \left\{ \varrho > 0 : \sum_{i=1}^{\infty} M\left(\frac{|x_i|}{\varrho}\right) \leq 1 \right\}.$$

The space  $l_M$  is closely related to the space  $l_p$  which is an Orlicz sequence space with,  $M(x) = x^p$ ,  $1 \leq p < \infty$ .

In the later stage different Orlicz sequence spaces were introduced and studied by Altın et al. [1], Altun and Bilgin [2], Bhardwaj and Singh [3], Braha [7], Et et al. [9], Mursaleen et al. [18] and many others.

## 2. MAIN RESULTS

Let  $M$  be an Orlicz function and  $p = (p_k)$  be a sequence of positive real numbers. We define the following sequence space:

$$E_n^q(M, p, s) = \left\{ a = (a_k) : \sum_{k=1}^{\infty} k^{-s} \left[ M\left(\frac{|N_k(a)|}{\varrho}\right) \right]^{p_k} < \infty, s \geq 0, \text{ for some } \varrho > 0 \right\}.$$

If we take  $s = 0$ , then we have

$$E_n^q(M, p) = \left\{ a = (a_k) : \sum_{k=1}^{\infty} \left[ M\left(\frac{|N_k(a)|}{\varrho}\right) \right]^{p_k} < \infty, \text{ for some } \varrho > 0 \right\},$$

if  $p_k = 1$  for all  $k \in \mathbb{N}$ , then we have

$$E_n^q(M, s) = \left\{ a = (a_k) : \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{\varrho} \right) \right] < \infty, s \geq 0, \text{ for some } \varrho > 0 \right\}.$$

In what follows we will prove the structure properties of the spaces defined above.

**Theorem 2.1** Let the sequence  $(p_k)$  be bounded. Then the spaces  $E_n^q(M, s)$ ,  $E_n^q(M, p)$  and  $E_n^q(M, p, s)$  are linear spaces over the field  $\mathbb{C}$  of complex numbers.

**Proof.** Clear.

**Theorem 2.2** Let  $p = (p_i)$  be a bounded sequence of strictly positive real numbers. Then the sequence space  $E_n^q(M, p, s)$  is a paranormed (need not total paranorm) space with

$$g(a) = \inf \left\{ \rho^{p_n/H} : \left( \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{\varrho} \right) \right]^{p_k} \right)^{\frac{1}{H}} \leq 1, n = 1, 2, \dots \right\}$$

where  $H = \max(1, \sup p_k)$ .

**Proof.** Clearly  $g(a) = g(-a)$  and  $g(a+b) \leq g(a) + g(b)$ . Since  $M(0) = 0$ , we get  $\inf \{\rho^{r_n/H}\} = 0$  for  $a = 0$ .

Finally, we prove that scalar multiplication is continuous. Let  $\lambda$  be any number. Since

$$g(\lambda a) = \inf \left\{ \rho^{p_n/H} : \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|\lambda N_k(a)|}{\varrho} \right) \right]^{p_k} \leq 1, n = 1, 2, \dots \right\},$$

we may write

$$g(\lambda a) = \inf \left\{ (\lambda s)^{p_n/H} : \left( \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{s} \right) \right]^{p_k} \right)^{\frac{1}{H}} \leq 1, n = 1, 2, \dots \right\},$$

where  $s = \varrho/|\lambda|$ . Since  $|\lambda|^{p_k} \leq \max(1, |\lambda|^H)$ , then  $|\lambda|^{p_k/H} \leq \left( \max(1, |\lambda|^H) \right)^{\frac{1}{H}}$ . Hence

$$g(\lambda a) \leq \left( \max(1, |\lambda|^H) \right)^{\frac{1}{H}} \cdot \inf \left\{ (s)^{p_n/H} : \left( \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{s} \right) \right]^{p_k} \right)^{\frac{1}{H}} \leq 1, n = 1, 2, \dots \right\}$$

and therefore  $g(a)$  converges to zero when  $g(a)$  converges to zero in  $E_n^q(M, p, s)$ . Now suppose that  $\lambda_n \rightarrow 0$  as  $n \rightarrow \infty$  and  $a$  in  $E_n^q(M, p, s)$ . For arbitrary  $\varepsilon > 0$ , let  $n_0$  be a positive integer such that

$$\sum_{k=n_0+1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{\varrho} \right) \right]^{p_k} < \frac{\varepsilon}{2}$$

for some  $\varrho > 0$ . This implies that

$$\left( \sum_{k=n_0+1}^{\infty} k^{-s} \left[ M \left( \frac{|N_k(a)|}{\varrho} \right) \right]^{p_k} \right)^{\frac{1}{H}} \leq \frac{\varepsilon}{2}.$$

Let  $0 < |\lambda| < 1$ , then using convexity of  $M$  we get

$$\sum_{k=n_0+1}^{\infty} k^{-s} \left[ M \left( \frac{|\lambda N_k(a)|}{\varrho} \right) \right]^{p_k} < \sum_{k=n_0+1}^{\infty} k^{-s} \left[ |\lambda| \frac{|N_k(a)|}{\varrho} \right]^{p_k} < \left( \frac{\varepsilon}{2} \right)^H.$$

Since  $M$  is continuous everywhere in  $[0, \infty)$ , then

$$f(t) = \sum_{k=1}^{n_0} k^{-s} \left[ M \left( \frac{|t N_k(a)|}{\rho} \right) \right]^{p_k}$$

is continuous at 0. So there is  $0 < \delta < 1$  such that  $|f(t)| < \frac{\varepsilon}{2}$  for  $0 < t < \delta$ . Let  $K$  be such that  $|\lambda_n| < \delta$  for  $n > K$ , then for  $n > K$  we have

$$\left( \sum_{k=1}^{n_0} k^{-s} \left[ M \left( \frac{|\lambda_n N_k(a)|}{\rho} \right) \right]^{p_k} \right)^{\frac{1}{H}} < \frac{\varepsilon}{2}.$$

Thus

$$\left( \sum_{k=1}^{\infty} k^{-s} \left[ M \left( \frac{|\lambda_n N_k(a)|}{\rho} \right) \right]^{p_k} \right)^{\frac{1}{H}} < \varepsilon, \text{ for } n > K.$$

Hence  $g(\lambda a) \rightarrow 0$ , as  $\lambda \rightarrow 0$ . This completes the proof of the Theorem.

**Theorem 2.3** Let  $M_1$  and  $M_2$  be two Orlicz functions and  $s, s_1, s_2$  be non-negative real numbers, then we have

- i)  $E_n^q(M_1, p, s) \cap E_n^q(M_2, p, s) \subseteq E_n^q(M_1 + M_2, p, s)$ ,
- ii) If  $s_1 \leq s_2$ , then  $E_n^q(M_1, p, s_1) \subseteq E_n^q(M_1, p, s_2)$ .
- iii) If  $M_1$  and  $M_2$  are equivalent, then  $E_n^q(M_1, p, s) = E_n^q(M_2, p, s)$ .

**Proof.** Clear.

**Corollary 2.4** Let  $s > 1$  and  $M$  be Orlicz function.

- i)  $E_n^q(M, p) \subset E_n^q(M, p, s)$
- ii)  $E_n^q(M) \subset E_n^q(M, s)$

**Theorem 2.5** Suppose that  $0 < u_k \leq p_k < \infty$  for each  $k$ . Then  $E_n^q(M, u) \subseteq E_n^q(M, p)$ .

**Proof.** Let  $a \in E_n^q(M, u, s)$ . Then there exists some  $\varrho > 0$  such that

$$\sum_{k=1}^{\infty} \left[ M \left( \frac{|N_k(a)|}{\varrho} \right) \right]^{u_k} < \infty.$$

This implies that  $M\left(\frac{|N_i(a)|}{\varrho}\right) \leq 1$  for sufficiently large values of  $i$ , say  $i \geq k_0$  for some fixed  $k_0 \in \mathbb{N}$ . Since  $M$  is non-decreasing, we get

$$\sum_{k \geq k_0}^{\infty} \left[ M\left(\frac{|N_k(a)|}{\varrho}\right) \right]^{p_k} \leq \sum_{k \geq k_0}^{\infty} \left[ M\left(\frac{|N_k(a)|}{\varrho}\right) \right]^{u_k} < \infty.$$

Hence  $a \in E_n^q(M, p)$ .

The following results are consequence of the above result.

**Corollary 2.6** (i) If  $0 < p_k \leq 1$  for each  $k$ , then  $E_n^q(M, p) \subseteq E_n^q(M)$ .

(ii) If  $p_k \geq 1$  for all  $k$ , then  $E_n^q(M) \subseteq E_n^q(M, p)$ .

### 3. RESULTS ON STATISTICAL CONVERGENCE

The idea of statistical convergence was given by Zygmund [26] in the first edition of his monograph published in Warsaw in 1935. The concept of statistical convergence was introduced by Steinhaus [25] and Fast [11] and later reintroduced by Schoenberg [24] independently. Over the years and under different names statistical convergence has been discussed in the theory of Fourier analysis, ergodic theory, number theory, measure theory, trigonometric series, turnpike theory and Banach spaces. Later on it was further investigated from the sequence space point of view and linked with summability theory by Connor [4], Çolak [5], Fridy [13], Et et al. [10], Mursaleen [19], Rath and Tripathy [20], Salat [21], Savaş [22] and many others. In recent years, generalizations of statistical convergence have appeared in the study of strong integral summability and the structure of ideals of bounded continuous functions on locally compact spaces. Statistical convergence and its generalizations are also connected with subsets of the Stone-Čech compactification of the natural numbers. Moreover, statistical convergence is closely related to the concept of convergence in probability.

Let  $\theta = (k_r)$  be the sequence of positive integers such that  $k_0 = 0$ ,  $0 < k_r < k_{r+1}$  and  $h_r = k_r - k_{r-1} \rightarrow \infty$  as  $r \rightarrow \infty$ . Then  $\theta$  is called a lacunary sequence. The intervals determined by  $\theta$  will be denoted by  $I_r = (k_{r-1}, k_r]$  and the ratio  $k_r/k_{r-1}$  will be denoted by  $q_r$ . Lacunary sequences have been studied by ([3],[6],[8],[12],[14],[23]).

In this section we give some results about  $N_k$ -lacunary statistical convergence and give some relations between the set of  $N_k$ -lacunary statistical convergence sequences and other spaces which defined with respect to an Orlicz function.

**Definition 3.1** Let  $\theta = (k_r)$  be a lacunary sequence, then the sequence  $a = (a_k)$  is said to be  $N_k$ -lacunary statistically convergent to the number  $\ell$  provided that for every  $\varepsilon > 0$ ,

$$\lim_r \frac{1}{h_r} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}| = 0.$$

In this case we write  $[N_k, S]_{\theta} - \lim a = \ell$  or  $a_k \rightarrow \ell ([N_k, S]_{\theta})$ .

**Definition 3.2** Let  $\theta = (k_r)$  be a lacunary sequence,  $M$  be an Orlicz function and  $p = (p_k)$  be any sequence of strictly positive real numbers. A sequence  $a = (a_k)$  is said to be strongly  $N_k$ -lacunary convergent to the number  $\ell$  with respect to the Orlicz function  $M$ , provided that

$$\lim_r \frac{1}{h_r} \sum_{k \in I_r} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} = 0.$$

The set of all strongly  $N_k$ -lacunary convergent sequences to the number  $\ell$  with respect to the Orlicz function  $M$  is denoted by  $[N_k, w, M, p]_\theta$ . In this case we write  $a_k \rightarrow \ell([N_k, w, M, p]_\theta)$ . In the special case  $M(x) = x$ ,  $p_k = p_0$  for all  $k \in \mathbb{N}$  we shall write  $[N_k, w]_\theta$  instead of  $[N_k, w, M, p]_\theta$ .

**Theorem 3.3** Let  $\theta = (k_r)$  be a lacunary sequence.

i) If a sequence  $a = (a_k)$  is strongly  $N_k$ -lacunary convergent to  $\ell$ , then it is  $N_k$ -lacunary statistically convergent to  $\ell$ .

ii) If a bounded sequence  $a = (a_k)$  is  $N_k$ -lacunary statistically convergent to  $\ell$ , then it is strongly  $N_k$ -lacunary convergent to  $\ell$ .

**Proof.** (i) Let  $\varepsilon > 0$  and  $a_k \rightarrow \ell([N_k, w]_\theta)$ . Then we can write

$$\sum_{k \in I_r} |N_k(a) - \ell|^{p_0} \geq \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| \geq \varepsilon}} |N_k(a) - \ell|^{p_0} \geq \varepsilon^{p_0} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}|.$$

Hence  $a_k \rightarrow \ell([N_k, S]_\theta)$ .

ii) Suppose that  $a_k \rightarrow \ell([N_k, S]_\theta)$  and let  $a \in \ell_\infty$ . Let  $\varepsilon > 0$  be given and select  $N_\varepsilon$  such that

$$\frac{1}{h_r} \left| \left\{ k \in I_r : |N_k(a) - \ell| \geq \left( \frac{\varepsilon}{2} \right)^{\frac{1}{p_0}} \right\} \right| \leq \frac{\varepsilon}{2K^{p_0}}$$

for all  $r > N_\varepsilon$  and set  $L_r = \left\{ k \in I_r : |N_k(a) - \ell| \geq \left( \frac{\varepsilon}{2} \right)^{\frac{1}{p_0}} \right\}$ , where  $K = \sup_k |a_k| < \infty$ . Now for all  $r > N_\varepsilon$  we have

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} |N_k(a) - \ell|^{p_0} &= \frac{1}{h_r} \sum_{\substack{k \in I_r \\ k \in L_r}} |N_k(a) - \ell|^{p_0} + \frac{1}{h_r} \sum_{\substack{k \in I_r \\ k \notin L_r}} |N_k(a) - \ell|^{p_0} \\ &\leq \frac{1}{h_r} \left( \frac{h_r \varepsilon}{2K^{p_0}} \right) K^{p_0} + \frac{\varepsilon}{2h_r} h_r = \varepsilon. \end{aligned}$$

Thus  $(a_k) \in [N_k, w]_\theta$ . This completes the proof.

**Theorem 3.4** For any lacunary sequence  $\theta$ , if  $\liminf_{r \rightarrow \infty} q_r > 1$ , then  $[N_k, S] \subset [N_k, S]_\theta$ .

**Proof.** If  $\liminf_{r \rightarrow \infty} q_r > 1$ , then there exists a  $\delta > 0$  such that  $1 + \delta \leq q_r$  for sufficiently large  $r$ . Since  $h_r = k_r - k_{r-1}$ , we have  $\frac{k_r}{h_r} \leq \frac{1+\delta}{\delta}$ . Let  $a_k \rightarrow \ell([N_k, S])$ .

Then for every  $\varepsilon > 0$  we have

$$\begin{aligned} \frac{1}{k_r} |\{k \leq k_r : |N_k(a) - \ell| \geq \varepsilon\}| &\geq \frac{1}{k_r} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}| \\ &\geq \frac{\delta}{1 + \delta} \frac{1}{h_r} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}|. \end{aligned}$$

Hence  $[N_k, S] \subset [N_k, S]_\theta$ .

**Theorem 3.5** For any lacunary sequence  $\theta$ , if  $\limsup_r q_r < \infty$ , then  $[N_k, S]_\theta \subset [N_k, S]$ .

**Proof.** Suppose that  $\limsup_r q_r < \infty$ . Then there exists a  $\beta > 0$  such that  $q_r < \beta$  for all  $r$ . Let  $a_k \rightarrow \ell$  ( $[N_k, S]_\theta$ ), and set  $E_r = |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}|$ . Then there exists an  $r_0 \in \mathbb{N}$  such that  $\frac{E_r}{h_r} < \varepsilon$  for all  $r > r_0$ . Let  $K = \max\{E_r : 1 \leq r \leq r_0\}$  and choose  $n$  such that  $k_{r-1} < n \leq k_r$ , then we have

$$\begin{aligned} \frac{1}{n} |\{k \leq n : |N_k(a) - \ell| \geq \varepsilon\}| &\leq \frac{1}{k_{r-1}} |\{k \leq k_r : |N_k(a) - \ell| \geq \varepsilon\}| \\ &\leq \frac{1}{k_{r-1}} \{E_1 + E_2 + \dots + E_{r_0} + E_{(r_0+1)} + \dots + E_r\} \\ &\leq \frac{K}{k_{r-1}} r_0 + \frac{1}{k_{r-1}} \left\{ \frac{E_{(r_0+1)}}{h_{r_0+1}} h_{r_0+1} + \dots + \frac{E_r}{h_r} h_r \right\} \\ &\leq \frac{K}{k_{r-1}} r_0 + \frac{1}{k_{r-1}} \left( \sup_{r>r_0} \frac{E_r}{h_r} \right) \{h_{r_0+1} + \dots + h_r\} \\ &\leq \frac{K}{k_{r-1}} r_0 + \varepsilon \frac{k_r - k_{r_0}}{k_{r-1}} \\ &\leq \frac{K}{k_{r-1}} r_0 + \varepsilon q_r \\ &\leq \frac{K}{k_{r-1}} r_0 + \varepsilon \beta. \end{aligned}$$

This completes the proof.

**Theorem 3.6** Let  $\theta = (k_r)$  be a lacunary sequence with  $1 < \liminf_r q_r \leq \limsup_r q_r < \infty$ , then we have  $[N_k, S]_\theta = [N_k, S]$ .

**Proof.** The proof follows from Theorem 3.4 and Theorem 3.5.

**Theorem 3.7** Let  $\theta = (k_r)$  be a lacunary sequence,  $M$  be an Orlicz function and  $0 < h = \inf_k p_k \leq p_k \leq \sup_k p_k = H$ . Then  $[N_k, w, M, p]_\theta \subset [N_k, S]_\theta$ .

**Proof.** Let  $a \in [N_k, w, M, p]_\theta$ . Then there exists a number  $\rho > 0$  such that

$$\frac{1}{h_r} \sum_{k \in I_r} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} \rightarrow 0, \text{ as } r \rightarrow \infty.$$

Then given  $\varepsilon > 0$  we have

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} &\geq \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| \geq \varepsilon}} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} \\ &\geq \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| \geq \varepsilon}} [M(\varepsilon_1)]^{p_k}, \text{ where } \varepsilon/\rho = \varepsilon_1 \\ &\geq \frac{1}{h_r} \sum \min \left\{ [M(\varepsilon_1)]^h, [M(\varepsilon_1)]^H \right\} \\ &\geq \frac{1}{h_r} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}| \cdot \min \left\{ [M(\varepsilon_1)]^h, [M(\varepsilon_1)]^H \right\}. \end{aligned}$$

Hence  $x \in [N_k, S]_\theta$ .

**Theorem 3.8** Let  $\theta = (k_r)$  be a lacunary sequence,  $M$  be an Orlicz function and  $a = (a_k)$  be bounded sequence, then  $[N_k, S]_\theta \subset [N_k, w, M, p]_\theta$ .

**Proof.** Suppose that  $a \in \ell_\infty$  and  $a_k \rightarrow \ell$  ( $[N_k, S]_\theta$ ). Since  $a \in \ell_\infty$ , there is a constant  $T > 0$  such that  $|N_k(a) - \ell| \leq T$ . Given  $\varepsilon > 0$  we have

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} &= \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| \geq \varepsilon}} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} \\ &+ \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| < \varepsilon}} \left[ M \left( \frac{|N_k(a) - \ell|}{\rho} \right) \right]^{p_k} \\ &\leq \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| \geq \varepsilon}} \max \left\{ \left[ M \left( \frac{T}{\rho} \right) \right]^h, \left[ M \left( \frac{T}{\rho} \right) \right]^H \right\} + \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |N_k(a) - \ell| < \varepsilon}} \left[ M \left( \frac{\varepsilon}{\rho} \right) \right]^{p_k} \\ &\leq \max \left\{ [M(K)]^h, [M(K)]^H \right\} \frac{1}{h_r} |\{k \in I_r : |N_k(a) - \ell| \geq \varepsilon\}| \\ &+ \max \left\{ [M(\varepsilon_1)]^h, [M(\varepsilon_1)]^H \right\}, \quad \frac{T}{\rho} = K, \quad \frac{\varepsilon}{\rho} = \varepsilon_1. \end{aligned}$$

Hence  $a \in [N_k, w, M, p]_\theta$ .

**Acknowledgment.** The authors thank the referees for their comments.

## REFERENCES

1. Y. Altın, M. Et and B.C. Tripathy, *The sequence space  $|\overline{N}_p|(M, r, q, s)$  on seminormed spaces*, Appl. Math. Comput. **154** (2004), no. 2, 423–430.
2. Y. Altın and T. Bilgin, *On a new class of sequences related to the  $l_p$  space defined by Orlicz function*, Taiwanese J. Math. **13** (2009), no. 4, 1189–1196.
3. V.K. Bhardwaj and N. Singh, *Some sequence spaces defined by Orlicz functions*, Demonstratio Math. **33** (2000), no. 3, 571–582.

4. J.S. Connor, *The Statistical and strong  $p$ -Cesaro convergence of sequences*, Analysis **8** (1988), 47–63.
5. R. Çolak, *Statistical convergence of order  $\alpha$* , *Modern Methods in Analysis and its Applications*, Anamaya Publishers, New Delhi, India, 2010.
6. R. Çolak, B.C. Tripathy and M. Et, *Lacunary strongly summable sequences and  $q$ -lacunary almost statistical convergence*, Vietnam J. Math. **34** (2006), no. 2, 129–138.
7. N.L. Braha, *A new class of sequences related to the  $l_p$  spaces defined by sequences of Orlicz functions*, J. Inequal. Appl. 2011, Article ID 539745, 10 p. (2011).
8. G. Das and S.K. Mishra, *Banach limits and lacunary strong almost convergence*, J. Orissa Math. Soc. **2** (1983), 61–70.
9. M. Et, Y. Altin, Choudhary, B. and B.C. Tripathy, *On some classes of sequences defined by sequences of Orlicz functions*, Math. Inequal. Appl. **9** (2006), no. 2, 335–342.
10. M. Et, A. Gökhan and H. Altinok, *On statistical convergence of vector-valued sequences associated with multiplier sequences*, Ukrainian Math. J. **58** (2006), no. 1, 139–146
11. H. Fast, *Sur la convergence statistique*, Colloquium Math. **2** (1951), 241–244.
12. A.R. Freedman, J.J. Sember and M. Raphael, *Some Cesaro-type summability spaces*, Proc. London Math. Soc. **37** (1978), 508–520.
13. J.A. Fridy, *On the statistical convergence*, Analysis **5** (1985), 301–313.
14. J.A. Fridy and C. Orhan, *Lacunary statistical convergence*, Pacific J. Math. **160** (1993), 43–51.
15. P.K. Kamptan and M. Gupta, *Sequence Spaces and Series*, Marcel Dekker Inc., New York, 1981.
16. M.A. Krasnoselskii and Y.B. Rutitsky, *Convex function and Orlicz spaces*, Groningen, Netherland, 1961.
17. J. Lindenstrauss and L. Tzafriri, *On Orlicz sequence spaces*, Israel J. Math. **1** (1971), 379–390.
18. M. Mursaleen, Q.A. Khan and T.A. Chishti, *Some new convergent sequences spaces defined by Orlicz functions and statistical convergence*, Ital. J. Pure Appl. Math. **9** (2001), 25–32.
19. M. Mursaleen,  *$\lambda$ -statistical convergence*, Math. Slovaca **50** (2000), no. 1, 111–115.
20. D. Rath and B.C. Tripathy, *On statistically convergent and statistically Cauchy sequences*, Indian J. Pure. Appl. Math., **25** (1994), no. 4, 381–386.
21. T. Šalát, *On statistically convergent sequences of real numbers*, Math. Slovaca **30** (1980), 139–150.
22. E. Savaş, *Strong almost convergence and almost  $\lambda$ -statistically convergence*, Hokkaido Math. J. **29** (2000), 531–536.
23. E. Savas and B.E. Rhoades, *On some new sequence spaces of invariant means defined by Orlicz functions*, Math. Inequal. Appl. **5** (2002), no. 2, 271–281.
24. I.J. Schoenberg, *The integrability of certain functions and related summability methods*, Amer. Math. Monthly **66** (1959), 361–375.
25. H. Steinhaus, *Sur la convergence ordinaire et la convergence asymptotique*, Colloquium Math. **2** (1951), 73–74.
26. A. Zygmund, *Trigonometric Series*, Cambridge University Press, Cambridge, UK, 1979.

<sup>1</sup> DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, AVENUE "MOTHER THERESA" 5, PRISHTINË, 10000, KOSOVA.

*E-mail address:* [nbraha@yahoo.com](mailto:nbraha@yahoo.com)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, FIRAT UNIVERSITY, 23119, ELAZIĞ-TURKEY.

*E-mail address:* [mikaillet@yahoo.com](mailto:mikaillet@yahoo.com)