

## ON THE SOLVABILITY OF CONVOLUTION EQUATIONS IN $\mathcal{K}'_M$

By

D. H. PAHK and B. K. SOHN

**Abstract.** Let  $\mathcal{K}'_M$  be the space of distributions on  $R^n$  which grow no faster than  $e^{M(kx)}$  for some  $k > 0$  where  $M$  is an increasing continuous function on  $R^n$ , and let  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  be the space of convolution operators in  $\mathcal{K}'_M$ . We show that, for  $S \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$ ,  $S * \mathcal{K}'_M = \mathcal{K}'_M$  is equivalent to the following: Every distribution  $u \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  with  $S * u \in \mathcal{K}_M$  is in  $\mathcal{K}_M$ .

### 1. Introduction.

Let  $\mathcal{K}'_M$  be the space of distributions on  $R^n$  which grow no faster than  $e^{M(kx)}$  for some  $k > 0$ , where  $M$  is an increasing continuous functions on  $R^n$ ;  $\mathcal{K}'_M$  is the dual space of  $\mathcal{K}_M$ , which we describe later. We denote by  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  the space of convolution operators in  $\mathcal{K}'_M$ .

In [1], S. Abdullah proved that, if  $S$  is a distributions in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $\hat{S}$  is its Fourier transform, the following conditions are equivalent:

(a) There exist positive constants  $A, C$  and a positive integer  $N$  such that

$$\sup_{\substack{z \in \mathcal{O}^n \\ |z| \leq A\Omega^{-1}(\log(2+|\xi|))}} |\hat{S}(z+\xi)| \geq \frac{C}{(1+|\xi|)^N}, \quad \xi \in R^n$$

where  $\Omega^{-1}$  is the inverse of  $\Omega$ , which is the dual to  $M$  in the sense of Young.

(b)  $S * \mathcal{K}'_M = \mathcal{K}'_M$ .

In this paper we prove that, for  $S \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$ , the statements (a) and (b) are equivalent to the following: Every distribution  $u \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  satisfying  $S * u \in \mathcal{K}_M$  is in  $\mathcal{K}_M$ .

The motivation for this problem comes from the paper [5]. Here S. Sznajder and Z. Zielezny proved that, if  $S$  is a distribution in  $\mathcal{O}'_C(\mathcal{K}'_1; \mathcal{K}'_1)$  and  $\hat{S}$  is its Fourier transform, the following statements are equivalent:

(i) There exist positive constants  $N, r, C$  such that

$$\sup_{z \in \mathbb{C}^n, |z| \leq r} |\hat{S}(\xi+z)| \geq \frac{C}{(1+|\xi|)^N}, \quad \xi \in \mathbb{R}^n,$$

(ii)  $S*\mathcal{K}'_1 = \mathcal{K}'_1$

(iii) If  $u \in \mathcal{O}'_C(\mathcal{K}'_1; \mathcal{K}'_1)$  and  $S*u \in \mathcal{K}_1$ , then  $u \in \mathcal{K}_1$ .

In view of this result it is natural to think the property (iii) in the space  $\mathcal{K}'_M$  of distributions on  $\mathbb{R}^n$  which grow no faster than  $\exp(M(kx))$  for some  $k > 0$ . Before presenting our theorems we recall briefly the basic facts about the spaces  $\mathcal{K}'_M, \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $\mathcal{K}'_M$ , for further details, we refer to [3].

**The space  $\mathcal{K}'_M$ .** Let  $\mu(\xi)(0 \leq \xi \leq \infty)$  denote a continuous increasing function such that  $\mu(0)=0, \mu(\infty)=\infty$ . For  $x \geq 0$ , we define

$$M(x) = \int_0^x \mu(\xi) d\xi.$$

The functions  $M(x)$  is an increasing, convex and continuous function with  $M(0)=0, M(\infty)=\infty$ . For  $x < 0$ , we define  $M(x)$  to be  $M(-x)$  and for  $x = (x_1, \dots, x_n) \in \mathbb{R}^n, n \geq 2$ , we define  $M(x)$  to be  $M(x_1) + \dots + M(x_n)$ .

Now we list some properties of  $M(x)$  which will be used later;

- (i)  $M(x) + M(y) \leq M(x+y)$  for all  $x, y \geq 0$
- (ii)  $M(x+y) \leq M(2x) + M(2y)$  for all  $x, y \geq 0$ .

Let  $\mathcal{K}_M$  be the space of all  $C^\infty$ -functions  $\phi$  in  $\mathbb{R}^n$  such that

$$\nu_k(\phi) = \sup_{\substack{x \in \mathbb{R}^n \\ |\alpha| \leq k}} e^{M(kx)} |D^\alpha \phi(x)| < \infty, \quad k=0, 1, 2, \dots,$$

where  $D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}$  and  $D_j = i^{-1}(\partial/\partial x_j)$ . Provided with the topology defined by the seminorms  $\nu_k, \mathcal{K}_M$  is a Frechet space. The dual  $\mathcal{K}'_M$  of  $\mathcal{K}_M$  is the space of all continuous linear functionals on  $\mathcal{K}_M$ . Then a distribution  $u$  is in  $\mathcal{K}'_M$  if and only if there exist  $m \in \mathbb{N}^n, k \in \mathbb{N}$  and a bounded continuous function  $f(x)$  on  $\mathbb{R}^n$  such that

$$u = D^m(e^{M(kx)} f(x)).$$

$\mathcal{K}'_M$  is endowed with the topology of uniform convergence on all bounded sets in  $\mathcal{K}_M$ .

**The space  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$ .** If  $u \in \mathcal{K}'_M$  and  $\phi \in \mathcal{K}_M$ , then the convolution  $u * \phi$  is a  $C^\infty$ -function defined by

$$u * \phi(x) = \langle u, \phi(x-y) \rangle,$$

where  $\langle u, \phi \rangle = u(\phi)$ .

The space  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  of convolution operators in  $\mathcal{K}'_M$  consists of distributions  $S \in \mathcal{K}'_M$  such that  $S * u \in \mathcal{K}'_M$  for every  $u \in \mathcal{K}'_M$ , where  $\langle S * u, \phi \rangle = \langle u, \check{S} * \phi \rangle$  for every  $\phi \in \mathcal{K}_M$ . Then the space is the set of distributions  $S$  which satisfy the following equivalent conditions [3]:

(i) The distributions  $S_k = \gamma_k S$ ,  $k=1, 2, \dots$  are in tempered distribution space, where  $\gamma_k = e^{M(kx)}$ .

(ii) For every integer  $k \geq 0$ , there exists an integer  $m \geq 0$  such that

$$S = \sum_{|\alpha| \leq m} D^\alpha f_\alpha$$

where  $f_\alpha$  are continuous functions in  $R^n$  whose products with  $e^{M(kx)}$  are bounded.

(iii) For every  $\phi \in \mathcal{K}_M$ , the convolution  $S * \phi$  is in  $\mathcal{K}_M$ .

**The space  $K'_M$ .** For  $\phi \in \mathcal{K}_M$ , the Fourier transform

$$\hat{\phi}(\xi) = \int_{R^n} e^{-i\langle x, \xi \rangle} \phi(x) dx$$

can be continued in  $C^n$  as an entire function of  $\zeta = \xi + i\eta$  such that

$$(1) \quad \omega_k(\hat{\phi}) = \sup_{\zeta \in C^n} (1 + |\xi|)^k e^{-\Omega(\eta/k)} |\hat{\phi}(\zeta)| < \infty, \quad k=1, 2, \dots$$

where  $\Omega(y)$  is the dual of  $M(x)$  in the sense of Young. If  $K_M$  is the space of all entire functions with the property (1) and the topology in  $K_M$  is defined by the seminorms  $\omega_k$ , then the Fourier transform is an isomorphism of  $\mathcal{K}_M$  onto  $K_M$ . The dual  $K'_M$  of  $K_M$  is the space of the Fourier transforms of distributions in  $\mathcal{K}'_M$ . The Fourier transform  $\hat{u}$  of a distribution  $u \in \mathcal{K}'_M$  is defined by the Parseval formula

$$\langle \hat{u}, \hat{\phi} \rangle = (2\pi)^n \langle u_x, \phi(-x) \rangle.$$

Also if  $S \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $u \in \mathcal{K}'_M$ , we have the formula

$$\widehat{S * u} = \hat{S} \cdot \hat{u},$$

where the product on the right-hand side is defined by

$$\langle \hat{S} \hat{u}, \phi \rangle = \langle \hat{u}, \hat{S} \phi \rangle, \quad \phi \in K_M.$$

The following lemma will be used in the next section. It's proof can be found in [3].

**LEMMA (Paley-Wiener type theorem).** *Let  $\zeta = \xi + i\eta \in C^n$ . An entire function  $F(\zeta)$  is the Fourier transform of a distribution  $S$  in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  if and only if for every  $\varepsilon > 0$ , there exist constants  $N$  and  $C$  such that*

$$|F(\xi+i\eta)| \leq C(1+|\zeta|)^N e^{\Omega(\varepsilon\eta)}.$$

**2. Main Theorem**

THEOREM. *If  $S$  is a distribution in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $\hat{S}$  be its Fourier transform, then the following conditions are equivalent:*

(a) *There exist positive constants  $A, C$  and a positive integer  $N$  such that*

$$\sup_{\substack{z \in \mathbb{C}^n \\ |z| \leq A\Omega^{-1}(\log(2+|\xi|))}} |\hat{S}(z+\xi)| \geq \frac{C}{(1+|\xi|)^N}, \quad \xi \in \mathbb{R}^n$$

(b)  $S*\mathcal{K}'_M = \mathcal{K}'_M$

(c) *If  $u \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $S*u \in \mathcal{K}_M$ , then  $u \in \mathcal{K}_M$ .*

PROOF. It suffices to show that (b) $\Rightarrow$ (c) $\Rightarrow$ (a).

(b) $\Rightarrow$ (c). The proof goes along exactly the same lines as proof of Theorem 1 in [5]. For the completeness we give the proof. If  $S$  is a distribution in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$ , then so is  $T=\check{S}$  and, by (1), the mapping  $S^*: u \rightarrow S*u$  of  $\mathcal{K}'_M$  into  $\mathcal{K}'_M$  is the transpose of the mapping  $T^*: \phi \rightarrow T*\phi$  of  $\mathcal{K}_M$  into  $\mathcal{K}_M$ . Condition (b) is satisfied if and only if  $T^*$  an isomorphism of  $\mathcal{K}_M$  onto  $T*\mathcal{K}_M$  (see e. g., [2, Corollary on p. 92]). In particular the inverse  $T*\phi \rightarrow \phi$  must be continuous.

Suppose now that  $S*u = \phi$  where  $u \in \mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and  $\phi \in \mathcal{K}_M$ . Since  $\langle S*u, \varphi \rangle = \langle T*\check{u}, \check{\varphi} \rangle$  for  $\varphi \in \mathcal{K}_M$ , then

$$(2) \quad T*\check{u} = (-1)^n \check{\phi}$$

and for the proof it suffices to show that  $\check{u} \in \mathcal{K}_M$ . If  $\phi$  is a  $C^\infty$ -function with  $\text{supp } \phi \subset B(0, 1) = \{x \in \mathbb{R}^n : |x| \leq 1\}$  and  $\hat{\phi}(0) = 1$ , we define  $\phi_k(x) = k^n \phi(kx)$ ,  $k = 1, 2, \dots$ . From (2) it follows that

$$T*(\check{u}*\phi_k) = (-1)^n \check{\phi}*\phi_k,$$

and the convolutions  $\check{u}*\phi_k$  and  $(-1)^n \check{\phi}*\phi_k$  are in  $\mathcal{K}_M$ . Moreover, the sequence  $\{\phi_k\}$  converges in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  to  $\delta$ , the Dirac measure at the origin. Hence  $(-1)^n \check{\phi}*\phi \rightarrow (-1)^n \check{\phi}$  in  $\mathcal{K}_M$  and  $\check{u}*\phi_k \rightarrow \check{u}$  in  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$ . On the other hand, the sequence  $\{\check{u}*\phi_k\}$  converges in  $\mathcal{K}_M$ , by the assumption that the inverse of  $T^*$  is continuous. The limit must be again  $\check{u}$ , and so  $\check{u}$  is a function in  $\mathcal{K}_M$ .

(c) $\Rightarrow$ (a). Let  $\mathcal{F}$  be the space all functions  $u \in C(\mathbb{R}^n)$  such that

$$\sup_{x \in \mathbb{R}^n} e^{M(kx)} |u(x)| < \infty, \quad \text{for all } k$$

and  $S*u \in \mathcal{K}_M$ . We provide  $\mathcal{F}$  with the topology defined by the seminorms

$$\|u\|_k = \sup_{x \in \mathbb{R}^n} e^{M(kx)} |u(x)| + \nu_k(S*u), \quad k=0, 1, 2, \dots$$

Then  $\mathcal{F}$  becomes a Frechet space. Further, let  $\mathcal{G}$  be the space of all functions  $u \in C^1(\mathbb{R}^n)$  such that

$$\|u\| = \sup_{x \in \mathbb{R}^n, |\alpha| \leq 1} |D^\alpha u(x)| < \infty$$

with the norm  $\|\cdot\|$ ,  $\mathcal{G}$  is a Banach space.

By the fact  $\mathcal{F} \subset \mathcal{O}'_c(\mathcal{K}'_M; \mathcal{K}'_M)$  and the assumption (c), each function  $u \in \mathcal{F}$  is in  $\mathcal{G}$ . Also, the natural mapping  $\mathcal{F} \rightarrow \mathcal{G}$  is closed and therefore continuous. Consequently there exist an integer  $\mu > 0$  and a constant  $C$  such that

$$\|u\| \leq C \|u\|_\mu = C \left\{ \sup_{x \in \mathbb{R}^n} e^{M(\mu x)} |u(x)| + \nu_\mu(S*u) \right\}$$

for all  $u \in \mathcal{F}$ . Since the Fourier transformation is an isomorphism from  $\mathcal{K}_M$  onto  $K_M$ , there exist another integer  $\nu > 0$  and a constant  $C_0$  such that

$$(3) \quad \|u\| - C \sup_{x \in \mathbb{R}^n} e^{M(\mu x)} |u(x)| \leq C_0 \omega_\nu(\hat{S} \cdot \hat{u}), \quad \text{for all } u \in \mathcal{K}_M.$$

Suppose now that the condition (a) is not satisfied. Then there exists a sequence  $\{\xi_j\}$  such that  $|\xi_j| \rightarrow +\infty$  as  $j \rightarrow \infty$  and

$$(4) \quad \sup_{\substack{z \in \mathbb{C}^n \\ |z - \xi_j| \leq j \Omega^{-1}(\log(2 + |\xi_j|))}} |\hat{S}(z)| < \frac{1}{(1 + |\xi_j|)^j},$$

For each  $j$ , we define  $k_j$  to be the greatest integer equal or less than  $\alpha_j = \Omega^{-1}(\log(2 + |\xi_j|))$ . Let  $\phi \geq 0$  in  $C_c^\infty$ ,  $\text{supp } \phi \subset B(0, 1)$  and  $\hat{\phi}(0) = 1$ . We also define

$$\phi_j^1(x) = e^{i \langle \xi_j, x \rangle} (\phi_j * \dots * \phi_j)(x),$$

and

$$\phi_j^2(x) = (\phi * (\phi_j * \dots * \phi_j))(x)$$

where  $\phi_j(x) = \alpha_j^n \phi(\alpha_j x)$  and the convolution product in the parenthesis is being taken  $k_j$ -times. Now we define

$$\psi_j(x) = (\phi * \phi_j^1)(x)$$

Since  $\text{supp } \psi_j \subset B(0, 2)$ , clearly  $\psi_j \in \mathcal{F}$ .

Substituting  $\psi_j$ 's into the inequality (3), we will show that the left side of (3) goes to  $\infty$  and the right to 0, as  $j \rightarrow \infty$ , which gives the desired contradiction.

To show this, we first estimate

$$\begin{aligned}
 (5) \quad \|\phi_j\| &= \sup_{x \in \mathbb{R}^n, |\alpha| \leq 1} |D^\alpha \phi_j(x)| \\
 &\geq \sup_{x \in \mathbb{R}^n, |\alpha| = 1} |\phi_* \{e^{i\langle x, \xi_j \rangle} D^\alpha(\phi_j * \dots * \phi_j) \\
 &\quad + D^\alpha(i\langle x, \xi_j \rangle) e^{i\langle x, \xi_j \rangle} (\phi_j * \dots * \phi_j)\}(x)| \\
 &\geq \sup_{x \in \mathbb{R}^n, |\alpha| = 1} |D^\alpha(\langle x, \xi_j \rangle)(\phi_* (\phi_j * \dots * \phi_j))(x)| \\
 &\geq \frac{|\xi_j|}{n} \sup_{x \in \mathbb{R}^n} |\phi_j^2(x)|
 \end{aligned}$$

and since  $\text{supp } \phi_j, \text{supp } \phi_j^2 \subset B(0, 2)$ ,

$$(6) \quad \sup_{x \in \mathbb{R}^n} e^{M(\mu x)} |\phi_j(x)| = \sup_{|x| \leq 2} e^{M(\mu x)} |\phi_j^2(x)| \leq C' \sup_{x \in \mathbb{R}^n} |\phi_j^2(x)|,$$

where  $C' = e^{nM(2\mu)}$ .

Viewing

$$1 = \int_{\mathbb{R}^n} \phi_j^2(x) dx \leq C'' \sup_{x \in \mathbb{R}^n} |\phi_j^2(x)|,$$

where  $C''$  is the volume of  $B(0, 2)$ , we have

$$(7) \quad \sup_{x \in \mathbb{R}^n} |\phi_j^2(x)| \geq \frac{1}{C''}.$$

Substituting (5), (6) and (7) into (3), the left hand side of (3) behaves, as  $j \rightarrow \infty$ ,

$$\begin{aligned}
 \lim_{j \rightarrow \infty} \{ \|\phi_j\| - C \sup_{x \in \mathbb{R}^n} e^{M(\mu x)} |\phi_j(x)| \} \\
 \geq \lim_{j \rightarrow \infty} \left\{ \frac{|\xi_j|}{n} - CC' \right\} \frac{1}{C''} = \infty
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 (8) \quad \omega_\nu(\hat{S} \cdot \hat{\phi}_j) &= \sup_{\zeta \in \mathcal{C}^n} (1 + |\zeta|)^\nu e^{-\Omega(\eta/\nu)} |\hat{S}(\zeta)| |\hat{\phi}_j(\zeta)| \\
 &\leq \sup_{|\zeta - \xi_j| \leq j\alpha_j} (1 + |\zeta|)^\nu e^{-\Omega(\eta/\nu)} |\hat{S}(\zeta)| |\hat{\phi}_j(\zeta)| \\
 &\quad + \sup_{|\zeta - \xi_j| > j\alpha_j} (1 + |\zeta|)^\nu e^{-\Omega(\eta/\nu)} |\hat{S}(\zeta)| |\hat{\phi}_j(\zeta)|,
 \end{aligned}$$

where  $\zeta = \xi + i\eta$ .

It now suffices to prove that both terms in the right side of (8) go to 0, as  $j \rightarrow \infty$ . We first observe that, by the Paley-Wiener theorem for  $\phi$  as element of  $C_c^\infty$  with  $\text{supp } \phi \subset B(0, 1)$ , there exist a  $C_m \geq 0, m=0, 1, 2, \dots$ , such that

$$(9) \quad |\hat{\phi}(\zeta)| \leq C_m (1 + |\zeta|)^{-m} e^{|\eta|}.$$

Also, we observe that

$$\hat{\phi}_j^1(\zeta) = [\hat{\phi}_j(\zeta - \xi_j)]^{k_j} = \left[ \hat{\phi} \left( \frac{\zeta - \xi_j}{\alpha_j} \right) \right]^{k_j}$$

and, by (9),

$$(10) \quad |\hat{\phi}_j^1(\zeta)| \leq \left[ C_1 \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-1} e^{|\eta|/\alpha_j} \right]^{k_j}.$$

Also we observe that  $\Omega$  grows faster than any linear function of  $|\eta|$  as  $|\eta|$  goes large and  $\hat{\phi}_j(\zeta) = \hat{\phi}(\zeta) \cdot \hat{\phi}_j^1(\zeta)$ .

From these observations, the first term of the last estimate in (8) is bounded by

$$\begin{aligned} & \sup_{|\zeta - \xi_j| \leq j\alpha_j} (1 + |\zeta|)^\nu e^{\Omega(\eta/\nu)} |\hat{S}(\zeta)| (C_1(1 + |\zeta|)^{-1} e^{|\eta|}) \\ & \quad \times \left( C_1 \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-1} e^{|\eta|/\alpha_j} \right)^{k_j} \\ & \leq C_2 \sup_{|\zeta - \xi_j| \leq j\alpha_j} (1 + |\zeta - \xi_j|)^{\nu-1} (1 + |\xi_j|)^{\nu-1} C_1^{k_j} \\ & \quad \times \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-k_j} |\hat{S}(\zeta)| \\ & \leq C'_2 \sup_{|\zeta - \xi_j| \leq j\alpha_j} (1 + |\xi_j|)^{2\nu-2+d} \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{\nu-1-k_j} |\hat{S}(\zeta)| \\ & \leq C'_2 (1 + |\xi_j|)^{2\nu-2+d-j}, \end{aligned}$$

where we used that  $e^{-\Omega(\eta/\nu) + |\eta| + (k_j/\alpha_j)|\eta|}$  is bounded in  $R^n$  and  $d = \log C_1$ .

Therefore the first term of the last part in (8) approaches to 0 as  $j \rightarrow \infty$ .

From the lemma for  $S$  as element of  $\mathcal{O}'_C(\mathcal{K}'_M; \mathcal{K}'_M)$  and (9), (10), the second term of the last estimate in (8) is bounded by

$$\begin{aligned} & \sup_{|\zeta - \xi_j| > j\alpha_j} C_{S, 2\nu+N} (1 + |\zeta|)^\nu e^{-\Omega(\eta/\nu)} (1 + |\zeta|)^N e^{\Omega(\eta/2\nu)} \\ & \quad \times (1 + |\zeta|)^{-(2\nu+N)} e^{|\eta|} \left( C_1 \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-1} e^{|\eta|/\alpha_j} \right)^{k_j} \\ & \leq \sup_{|\zeta - \xi_j| > j\alpha_j} C'_{S, 2\nu+N} (1 + |\zeta|)^{\nu+N-(2\nu+N)} C_1^{k_j} \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-k_j} \\ & \leq \sup_{|\zeta - \xi_j| > j\alpha_j} C'_{S, 2\nu+N} C_1^{k_j} \left( 1 + \left| \frac{\zeta - \xi_j}{\alpha_j} \right| \right)^{-k_j} \\ & \leq C'_{S, 2\nu+N} \left( \frac{1+j}{C_1} \right)^{-k_j}, \end{aligned}$$

where we used that  $e^{-\Omega(\eta/\nu) + \Omega(\eta/2\nu) + |\eta| + (k_j/\alpha_j)|\eta|}$  is bounded in  $R^n$ . Here  $C_{S, 2\nu+N}$  and  $C'_{S, 2\nu+N}$  are constants which depend on  $S$ ,  $\phi$ ,  $\nu$  and  $N$  only.

Hence the second term of the last part in (8) approaches to 0 as  $j \rightarrow \infty$ .

Combining both estimates we have

$$\lim_{j \rightarrow \infty} \omega_\nu(\hat{S} \cdot \hat{\varphi}_j) = 0,$$

which gives the desired contradiction.

### References

- [ 1 ] Abdullah, S., Solvability of convolution equations in  $\mathcal{K}'_{\mathcal{M}}$ , Hokkaido Math. J. 17 (1988), 197-209.
- [ 2 ] Dieudonné, J. et Schwartz, L., La dualité dans les espaces  $(\mathcal{D})$  et  $(\mathcal{L}\mathcal{D})$ , Ann. Inst. Fourier Gremoble 1 (1949), 61-101.
- [ 3 ] Pahk, D.H., Structure theorem and Fourier transform for distributions with restricted growth, Kyungpook Math. J. 23(2) (1983), 129-146.
- [ 4 ] Pahk, D.H. and Sohn, B.K., Relation between solvability and a regularity of convolution operators in  $\mathcal{K}'_p$ ,  $p > 1$ , J. Math. Anal. Appl. 185 (1994), 207-214.
- [ 5 ] Sznajder, S. and Zielezny, Z., Solvability of convolution equations in  $\mathcal{K}'_1$ , Proc. Amer. Math. Soc. 57 (1976), 103-106.
- [ 6 ] ———, On some properties of convolution operators in  $\mathcal{K}'_1$  and  $\mathcal{S}'$ , J. Math. Anal. Appl. 65 (1978), 543-554.

Dae Hyeon Pahk  
 Department of Mathematics  
 Yonsei University  
 Seoul 120-749, Korea

Byung Keun Sohn  
 Department of Mathematics  
 Inje University  
 Kyungnam Kimhae 621-749, Korea