

*Communications in  
Applied  
Mathematics and  
Computational  
Science*

AN URN MODEL ASSOCIATED WITH JACOBI  
POLYNOMIALS

F. ALBERTO GRÜNBAUM

vol. 5    no. 1    2010

# AN URN MODEL ASSOCIATED WITH JACOBI POLYNOMIALS

F. ALBERTO GRÜNBAUM

We consider an urn model leading to a random walk that can be solved explicitly in terms of the well-known Jacobi polynomials.

## 1. Urn models and orthogonal polynomials

There are two simple and classical models in statistical mechanics which have recently been associated with very important classes of orthogonal polynomials. The oldest one of these models is due to D. Bernoulli (1770) and S. Laplace (1810), while the more recent model is due to Paul and Tatiana Ehrenfest (1907) [6]. While both of these models are featured in very classical texts in probability theory, such as [7], the connection with orthogonal polynomials is of much more recent vintage. In fact, the polynomials in question due to Krawtchouk and Hahn had not been recognized as basic objects with rich properties until around 1950. For a few pertinent and useful references, see [1; 2; 4; 5; 11; 14; 16; 18].

From these comments one could get the impression that the relations between orthogonal polynomials — especially some well-known classes of them — are only a matter of historical interest. Nothing could be further from the truth: there are several areas of probability and mathematical physics where recent important progress hinges on the connections with orthogonal polynomials.

The entire area of random matrix theory starts with the work of E. Wigner and F. Dyson and reaches a new stage in the hands of M. Mehta who brought the power of orthogonal polynomials into the picture.

In the area of random growth processes, the seminal work of K. Johansson depends heavily on orthogonal polynomials, specifically Laguerre and Meixner ones; see [15].

The connection between birth-and-death processes and orthogonal polynomials has many parents, but the people that made the most of it are S. Karlin and J. McGregor [17]. We will have a chance to go back to their work in connection with our model here. The ideas of using the spectral analysis of the corresponding

---

*MSC2000:* primary 33C45, 60J10; secondary 60G99.

*Keywords:* random walks, urn models, Jacobi polynomials, orthogonal polynomials.

The author was supported in part by NSF Grant DMS-0603901 and AFOSR Contract FA9550-08-1-0169.

one-step transition matrix have been pushed recently in the case of quantum random walks, an area where physics, computer science and mathematics could make important contributions. See [3; 12].

The study of the so-called ASEP (asymmetric simple exclusion processes), going back to F. Spitzer [21] and very much connected with the work of K. Johansson mentioned earlier, has recently profited from connections with the Askey–Wilson polynomials. All of this has important and deep connections with combinatorics and a host of other areas of mathematics; for example, the study of nonintersecting or noncolliding random processes, which goes back to F. Dyson.

There are many interrelations among these areas. For one example: the Hahn polynomials that were mentioned in connection with the Bernoulli–Laplace model were studied by Karlin and McGregor [18] in connection with a model in genetics due to Moran. They have also been found to be useful in discussing random processes with nonintersecting paths [8].

All of these areas are places where orthogonal polynomials have been put to very good use. For a review of several of these items, see [19]. Orthogonal polynomials of several variables, as well as matrix-valued orthogonal polynomials, have recently been connected to certain random walks. For three examples, see [9; 10; 13].

## 2. The Jacobi polynomials

The classical Jacobi polynomials are usually considered either over the interval  $[-1, 1]$  or, as we will do, over  $[0, 1]$ .

These polynomials are orthogonal with respect to the weight function

$$W(x) = x^\alpha (1 - x)^\beta.$$

Here we assume that  $\alpha, \beta > -1$ ; in fact it will be assumed throughout that  $\alpha, \beta$  are nonnegative integers.

These polynomials are eigenfunctions of the differential operator

$$x(1-x) \frac{d^2}{dx^2} + (\alpha + 1 + x(\alpha + \beta + 2)) \frac{d}{dx},$$

a fact that will not play any role in our discussion but which is crucial in most physical and geometrical applications of Jacobi polynomials. These applications cover a vast spectrum including potential theory, electromagnetism and quantum mechanics.

Neither the orthogonality, nor the fact that our polynomials are eigenfunctions of this differential operator are enough to determine them uniquely. One can multiply each polynomial by a constant and preserve these properties. We chose to normalize

our polynomials by the condition

$$Q_n(1) = 1.$$

For us it will be important that these polynomials satisfy (and in fact be defined by) the three-term recursion relation

$$x Q_n(x) = A_n Q_{n+1}(x) + B_n Q_n(x) + C_n Q_{n-1}(x),$$

with  $Q_0 = 1$  and  $C_0 Q_{-1} = 0$ .

The coefficients  $A_n, B_n, C_n$  are given by

$$A_n = \frac{(n + \beta + 1)(n + \alpha + \beta + 1)}{(2n + \alpha + \beta + 1)(2n + \alpha + \beta + 2)}, \quad n \geq 0,$$

$$B_n = 1 + \frac{n(n + \beta)}{2n + \alpha + \beta} - \frac{(n + 1)(n + \beta + 1)}{2n + \alpha + \beta + 2}, \quad n \geq 0,$$

$$C_n = \frac{n(n + \alpha)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 1)}, \quad n \geq 1.$$

The coefficient  $B_n$  can be rewritten as

$$B_n = \frac{2n(n + \alpha + \beta + 1) + (\alpha + 1)\beta + \alpha(\alpha + 1)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 2)},$$

which makes it clear that, along with the other coefficients, it is nonnegative.

Since we insist on the condition  $Q_n(1) = 1$ , we can see, for instance by induction and using the recursion relation, that

$$A_n + B_n + C_n = 1, \quad n \geq 1.$$

We also have

$$A_0 + B_0 = 1.$$

There are, of course, several explicit expressions for the different variants of the Jacobi polynomials, and they can be used, for instance, in computing the integrals that appear in the last section of this paper.

In terms of hypergeometric functions, the usual Jacobi polynomials are given by

$$((\alpha + 1)_n/n!) {}_2F_1(-n, n + \alpha + \beta + 1; (1 - x)/2),$$

while our polynomials  $Q_n(x)$  are obtained by multiplying these standard Jacobi polynomials by  $(-1)^n n! / (\beta + 1)_n$  and replacing  $x$  by  $1 - 2x$ .

The normalization chosen above is natural when one thinks of these polynomials (at least for some values of  $\alpha, \beta$ ) as the spherical functions for some appropriate

symmetric space, and insists that these functions take the value 1 at the north pole of the corresponding sphere. The simplest of all cases is the one where  $\alpha = \beta = 0$ ; one gets the Legendre polynomials and the usual two-dimensional sphere sitting in  $\mathbb{R}^3$  (see [23]).

The fact that the coefficients are nonnegative and add up to one cries out for a probabilistic interpretation of these quantities. This is the purpose of this paper. We have not seen in the literature concrete models of random walks where the Jacobi polynomials play this role. The urn model we give is admittedly a bit contrived, but it is quite concrete. Hopefully it will motivate other people to find a more natural and simpler model that goes along with this recursion relation.

### 3. The model

Here we consider a discrete-time random walk on the nonnegative integers whose one-step transition probability matrix coincides with the one that gives the three-term recursion relation satisfied by the Jacobi polynomials.

At times  $t = 0, 1, 2, \dots$ , an urn contains  $n$  blue balls and this determines the state of our random walk on  $\mathbb{Z} \geq 0$ .

The urn sits in a “bath” consisting of an infinite number of red balls. The transition mechanism is made up of a few steps which are described now, leaving some of the details for later.

In the first step a certain number of red balls from the surrounding bath are mixed with the  $n$  blue balls in the urn.

In the second step a ball is selected (with uniform distribution) from among the balls in the urn. This “chosen ball” can be blue or red. In either case an experiment is performed in a parallel world, using an appropriate “auxiliary urn”, to determine if this chosen ball will retain its color or have it changed (from red to blue or vice versa).

Once this is settled, and the possible change of color has taken place, the main urn contains the initial  $n$  balls plus a certain number of balls taken from the bath in the first step, and we are ready for the third and last step. This final step consists of having all red balls in the main urn removed and dropped back into the bath.

The state of the system at time  $t + 1$  is given by the number of blue balls in the urn after these three steps are completed. Clearly, the new state can take any of the values  $n - 1, n, n + 1$ .

A more detailed description of the three steps above is given in the next section.

### 4. The details of the model

If at time  $t$  the urn contains  $n$  blue balls, with  $n = 0, 1, 2, \dots$ , we pick

$$n + \alpha + \beta + 1$$

red balls from the bath to get a total of  $2n + \alpha + \beta + 1$  balls in the urn at the end of step one.

We now perform step 2: this gives us a blue ball with probability

$$\frac{n}{2n + \alpha + \beta + 1}$$

and a red ball with probability

$$\frac{n + \alpha + \beta + 1}{2n + \alpha + \beta + 1}.$$

If the chosen ball is blue, then we throw  $\alpha$  blue balls and  $n + \beta$  red balls into an “auxiliary urn” with  $n$  blue balls, mix all these balls, and pick one with uniform distribution. We imagine the auxiliary urn surrounded by a bath of an infinite number of blue and red balls which are used to augment the  $n$  blue balls in this auxiliary urn.

The probability of getting a blue ball in the auxiliary urn is

$$\frac{n + \alpha}{2n + \alpha + \beta},$$

and if this is the outcome, the chosen ball in the main urn has its color changed from blue to red. If we get a red ball in this auxiliary urn, then the chosen ball retains its blue color.

On the other hand, if in step 2 we had chosen a red ball, then we throw  $\alpha + 1$  blue balls and  $n + \beta + 1$  red balls into a different auxiliary urn with  $n$  blue balls. This auxiliary urn is also surrounded by a bath of an infinite number of blue and red balls.

These balls are mixed and one is chosen with the uniform distribution. The probability that this ball in the auxiliary urn is red is given by

$$\frac{n + \beta + 1}{2n + \alpha + \beta + 2},$$

and if this is the case, the chosen ball in the main urn has its color changed from red to blue. Otherwise the chosen ball retains its red color.

Notice that the chosen ball in the main urn has a change of color only when we get a match of colors for the balls drawn in the main and an auxiliary urn: blue followed by blue or red followed by red.

In either case once the possible change of color of the chosen ball in the main urn has been decided upon, we remove all the red balls from the main urn.

We see that the state of the system goes from  $n$  to  $n - 1$  when the chosen ball is blue and then its color gets changed into red. This event has probability

$$\frac{n}{2n + \alpha + \beta + 1} \times \frac{n + \alpha}{2n + \alpha + \beta}.$$

Observe that this coincides with the value of  $C_n$  in the recursion relation satisfied by our version of the Jacobi polynomials.

The state increases from  $n$  to  $n + 1$  if the chosen ball is red and its color gets changed into blue. This event has probability

$$\frac{n + \alpha + \beta + 1}{2n + \alpha + \beta + 1} \times \frac{n + \beta + 1}{2n + \alpha + \beta + 2}.$$

This coincides with the values of  $A_n$  given earlier.

As we noticed earlier, when the chosen ball is blue and the ball in the corresponding auxiliary urn is red then the chosen ball retains its color. Likewise if the chosen ball is red and the ball in the corresponding auxiliary urn is blue then the chosen one retains its color. In either case, the total number of blue balls in the main urn remains unchanged and the state goes from  $n$  to  $n$ .

Recall the basic property of the coefficients  $A_n, B_n, C_n$ , namely

$$A_n + B_n + C_n = 1.$$

This shows that the probability of going from state  $n$  to state  $n$  is given by  $B_n$ .

In summary, we have built a random walk whose one-step transition probability is the one given by the three-term relation satisfied by our version of the Jacobi polynomials.

## 5. Birth-and-death processes and orthogonal polynomials

A Markov chain with state space given by the nonnegative integers and a tridiagonal one-step transition probability matrix  $\mathbb{P}$  is called a birth-and-death process. Our model given above clearly fits in this framework.

One of the most important connections between orthogonal polynomials and birth-and-death processes, such as the one considered here, is given by the Karlin–McGregor formula [17].

If the polynomials satisfy

$$\pi_j \int_0^1 Q_i(x) Q_j(x) W(x) dx = \delta_{ij},$$

one gets the following representation formula for the entries of the powers of the one-step transition probability matrix

$$(\mathbb{P}^n)_{ij} = \pi_j \int_0^1 x^n Q_i(x) Q_j(x) W(x) dx.$$

This compact expression gives the solution to the dynamics of our random walk and allows for the study of many of its properties.

In the case of our version of the Jacobi polynomials, the squares of the norms of the polynomials  $Q_i$  are given by

$$\frac{\Gamma(i+1)\Gamma(i+\alpha+1)\Gamma(\beta+1)^2}{\Gamma(i+\beta+1)\Gamma(i+\alpha+\beta+1)(2i+\alpha+\beta+1)}.$$

In our case, when  $\alpha, \beta$  are assumed to be nonnegative integers, this expression can of course be written without any reference to the Gamma function.

We recall how one can compute in the case of our transition matrix  $P$  its invariant (stationary) distribution, that is, the (unique up to scalars) row vector

$$\boldsymbol{\pi} = (\pi_0, \pi_1, \pi_2, \dots)$$

such that

$$\boldsymbol{\pi} P = \boldsymbol{\pi}.$$

It is a simple matter of using the recursion relation for the polynomials  $Q_i$  to show that the components  $\pi_i$  are given, up to a common multiplicative constant, by the inverses of the integrals

$$\int_0^1 Q_i^2(x) W(x) dx$$

mentioned above. This justifies the notation  $\pi_i$  for these two apparently unrelated quantities, and in our case furnishes an explicit expression for an invariant distribution.

We close this paper with a note of historical interest. One of the referees suggested that I contact Dick Askey, who reportedly had pointed out that the Legendre polynomials had surfaced for the first time in connection with a problem in probability theory.

Askey recalls that Arthur Erdélyi told him once that this occurred in a work by J. L. Lagrange. Indeed in [20], Lagrange considers such a problem. In the process of solving it he needs to find the power series expansion of the expression

$$\frac{1}{\sqrt{1 - 2az + (a^2 - 4b^2)z^2}}$$

in powers of  $z$ .



The three-term recursion for the coefficients in this expansion is explicitly written down and considered well-known by Lagrange. The account of Lagrange's work given in the very complete and scholarly book [22] has a derivation of this recursion. The case  $a^2 - 4b^2 = 1$  gives the Legendre polynomials in the variable  $a$ . The work of Lagrange took place in the period 1770–1773, and predates the work of Legendre and Laplace. I am thankful to the anonymous referee and to Askey for pointing me in the correct direction.

## References

- [1] G. E. Andrews, R. Askey, and R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, no. 71, Cambridge University Press, 1999. [MR 2000g:33001](#) [Zbl 0920.33001](#)
- [2] R. Askey, *Evaluation of Sylvester type determinants using orthogonal polynomials*, Advances in analysis (H. G. W. Begehr, R. P. Gilbert, M. E. Muldoon, and M. W. Wong, eds.), World Sci. Publ., Hackensack, NJ, 2005, pp. 1–16. [MR 2006i:15013](#) [Zbl 1090.15007](#)
- [3] M. J. Cantero, F. A. Grünbaum, L. Moral, and L. Velazquez, *Matrix valued Szegő polynomials and quantum random walks*, Comm. Pure Appl. Math. (2009). [arXiv 0901.2244](#)
- [4] T. S. Chihara, *An introduction to orthogonal polynomials*, Mathematics and its Applications, no. 13, Gordon and Breach, New York, 1978. [MR 58 #1979](#) [Zbl 0389.33008](#)
- [5] P. Diaconis and M. Shahshahani, *Time to reach stationarity in the Bernoulli–Laplace diffusion model*, SIAM J. Math. Anal. **18** (1987), no. 1, 208–218. [MR 88c:60014](#) [Zbl 0617.60009](#)
- [6] P. Ehrenfest and T. Ehrenfest, *über zwei bekannte Einwände gegen das Boltzmannsche H-Theorem*, Phys. Z. **8** (1907), 311–314. [JFM 38.0931.01](#)
- [7] W. Feller, *An introduction to probability theory and its applications*, 3rd ed., vol. I, Wiley, New York, 1968. [MR 37 #3604](#) [Zbl 0155.23101](#)
- [8] V. E. Gorin, *Nonintersecting paths and the Hahn orthogonal polynomial ensemble*, Funktsional. Anal. i Prilozhen. **42** (2008), no. 3, 23–44, 96. [MR 2010a:60027](#) [Zbl 05519876](#)
- [9] F. A. Grünbaum, *The Karlin–McGregor formula for a variant of a discrete version of walsh's spider*, J. Phys. **45**, art. 454010. [Zbl 05638242](#)
- [10] F. A. Grünbaum, *The Rahman polynomials are bispectral*, Symmetry, Integrability Geom. Methods Appl. **3** (2007), Paper 065, 11. [MR 2009m:33019](#) [Zbl 05241578](#)
- [11] ———, *Random walks and orthogonal polynomials: some challenges*, Probability, geometry and integrable systems (M. Pinsky and B. Birnir, eds.), Math. Sci. Res. Inst. Publ., no. 55, Cambridge Univ. Press, 2008, pp. 241–260. [MR 2009i:33011](#)
- [12] M. Hamada, N. Konno, and W. Mlotkowski, *Orthogonal polynomials induced by discrete-time quantum walks in one dimension*, preprint, 2009. [arXiv 0903.4047](#)
- [13] M. R. Hoare and M. Rahman, *A probabilistic origin for a new class of bivariate polynomials*, Symmetry Integrability Geom. Methods Appl. **4** (2008), Paper 089, 18 pp. [MR 2010a:33028](#)
- [14] M. E. H. Ismail, D. R. Masson, J. Letessier, and G. Valent, *Birth and death processes and orthogonal polynomials*, Orthogonal polynomials (P. Nevai, ed.), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., no. 294, Kluwer, Dordrecht, 1990, pp. 229–255. [MR 93c:42021](#) [Zbl 0704.60084](#)
- [15] K. Johansson, *Shape fluctuations and random matrices*, Comm. Math. Phys. **209** (2000), no. 2, 437–476. [MR 2001h:60177](#) [Zbl 0969.15008](#)

- [16] M. Kac, *Random walk and the theory of Brownian motion*, Amer. Math. Monthly **54** (1947), 369–391. [MR 9,46c](#) [Zbl 0031.22604](#)
- [17] S. Karlin and J. McGregor, *Random walks*, Illinois J. Math. **3** (1959), 66–81. [MR 20 #7352](#) [Zbl 0104.11804](#)
- [18] S. Karlin and J. L. McGregor, *The Hahn polynomials, formulas and an application*, Scripta Math. **26** (1961), 33–46. [MR 25 #2249](#) [Zbl 0104.29103](#)
- [19] W. König, *Orthogonal polynomial ensembles in probability theory*, Probab. Surv. **2** (2005), 385–447. [MR 2007e:60007](#)
- [20] J. L. Lagrange, *Mémoire sur l'utilité de la méthode de prendre le milieu entre les résultats de plusieurs observations*, Miscellanea Taurinensia **5**, Reprinted in his Œuvres, volume 2, pages 173–185, Min. de l'Instr. Pub. Paris.
- [21] F. Spitzer, *Interaction of Markov processes*, Advances in Math. **5** (1970), 246–290 (1970). [MR 42 #3856](#) [Zbl 0312.60060](#)
- [22] I. Todhunter, *A history of the mathematical theory of probability*, MacMillan, 1865.
- [23] N. J. Vilenkin and A. U. Klimyk, *Representation of Lie groups and special functions, Vol 3: Classical and quantum groups and special functions*, Mathematics and its Applications (Soviet Series), no. 75, Kluwer, Dordrecht, 1992. [MR 96f:33043](#) [Zbl 0778.22001](#)

Received August 26, 2009. Revised December 13, 2009.

F. ALBERTO GRÜNBAUM: [grunbaum@math.berkeley.edu](mailto:grunbaum@math.berkeley.edu)

*Department of Mathematics, University of California, Berkeley, CA 94720, United States*

# Communications in Applied Mathematics and Computational Science

[pjm.math.berkeley.edu/camcos](http://pjm.math.berkeley.edu/camcos)

## EDITORS

MANAGING EDITOR

John B. Bell

Lawrence Berkeley National Laboratory, USA

[jbbell@lbl.gov](mailto:jbbell@lbl.gov)

BOARD OF EDITORS

Marsha Berger	New York University <a href="mailto:berger@cs.nyu.edu">berger@cs.nyu.edu</a>	Ahmed Ghoniem	Massachusetts Inst. of Technology, USA <a href="mailto:ghoniem@mit.edu">ghoniem@mit.edu</a>
Alexandre Chorin	University of California, Berkeley, USA <a href="mailto:chorin@math.berkeley.edu">chorin@math.berkeley.edu</a>	Raz Kupferman	The Hebrew University, Israel <a href="mailto:raz@math.huji.ac.il">raz@math.huji.ac.il</a>
Phil Colella	Lawrence Berkeley Nat. Lab., USA <a href="mailto:pcollella@lbl.gov">pcollella@lbl.gov</a>	Randall J. LeVeque	University of Washington, USA <a href="mailto:rjl@amath.washington.edu">rjl@amath.washington.edu</a>
Peter Constantin	University of Chicago, USA <a href="mailto:const@cs.uchicago.edu">const@cs.uchicago.edu</a>	Mitchell Luskin	University of Minnesota, USA <a href="mailto:luskin@umn.edu">luskin@umn.edu</a>
Maksymilian Dryja	Warsaw University, Poland <a href="mailto:maksymilian.dryja@acn.waw.pl">maksymilian.dryja@acn.waw.pl</a>	Yvon Maday	Université Pierre et Marie Curie, France <a href="mailto:maday@ann.jussieu.fr">maday@ann.jussieu.fr</a>
M. Gregory Forest	University of North Carolina, USA <a href="mailto:forest@amath.unc.edu">forest@amath.unc.edu</a>	James Sethian	University of California, Berkeley, USA <a href="mailto:sethian@math.berkeley.edu">sethian@math.berkeley.edu</a>
Leslie Greengard	New York University, USA <a href="mailto:greengard@cims.nyu.edu">greengard@cims.nyu.edu</a>	Juan Luis Vázquez	Universidad Autónoma de Madrid, Spain <a href="mailto:juanluis.vazquez@uam.es">juanluis.vazquez@uam.es</a>
Rupert Klein	Freie Universität Berlin, Germany <a href="mailto:rupert.klein@pik-potsdam.de">rupert.klein@pik-potsdam.de</a>	Alfio Quarteroni	Ecole Polytech. Féd. Lausanne, Switzerland <a href="mailto:alfio.quarteroni@epfl.ch">alfio.quarteroni@epfl.ch</a>
Nigel Goldenfeld	University of Illinois, USA <a href="mailto:nigel@uiuc.edu">nigel@uiuc.edu</a>	Eitan Tadmor	University of Maryland, USA <a href="mailto:etadmor@cscamm.umd.edu">etadmor@cscamm.umd.edu</a>
	Denis Talay	INRIA, France <a href="mailto:denis.talay@inria.fr">denis.talay@inria.fr</a>	

## PRODUCTION

[apde@mathscipub.org](mailto:apde@mathscipub.org)

Paulo Ney de Souza, Production Manager      Sheila Newbery, Production Editor      Silvio Levy, Senior Production Editor

---

See inside back cover or [pjm.math.berkeley.edu/camcos](http://pjm.math.berkeley.edu/camcos) for submission instructions.

The subscription price for 2010 is US \$70/year for the electronic version, and \$100/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

Communications in Applied Mathematics and Computational Science, at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

CAMCoS peer-review and production is managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY  
 **mathematical sciences publishers**  
<http://www.mathscipub.org>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2010 by Mathematical Sciences Publishers

# *Communications in Applied Mathematics and Computational Science*

vol. 5

no. 1

2010

---

FETI and BDD preconditioners for Stokes–Mortar–Darcy Systems JUAN GALVIS and MARCUS SARKIS	1
A cut-cell method for simulating spatial models of biochemical reaction networks in arbitrary geometries WANDA STRYCHALSKI, DAVID ADALSTEINSSON and TIMOTHY ELSTON	31
An urn model associated with Jacobi polynomials F. ALBERTO GRÜNBAUM	55
Ensemble samplers with affine invariance JONATHAN GOODMAN and JONATHAN WEARE	65
A second-order accurate method for solving the signed distance function equation PETER SCHWARTZ and PHILLIP COLELLA	81
On the second-order accuracy of volume-of-fluid interface reconstruction algorithms: convergence in the max norm ELBRIDGE GERRY PUCKETT	99