

The degree of the Alexander polynomial is an upper bound for the topological slice genus

PETER FELLER

We use the famous knot-theoretic consequence of Freedman’s disc theorem — knots with trivial Alexander polynomial bound a locally flat disc in the 4–ball — to prove the following generalization: the degree of the Alexander polynomial of a knot is an upper bound for twice its topological slice genus. We provide examples of knots where this determines the topological slice genus.

57M25, 57M27

1 Introduction

For a knot K — a smooth and oriented embedding of the unit circle S^1 into the unit 3–sphere S^3 — the *topological slice genus* $g_4^{\text{top}}(K)$ is the minimal genus of locally flat oriented surfaces S in the closed unit 4–ball B^4 with oriented boundary $K \subset \partial B^4 = S^3$. A celebrated theorem of Freedman [7, Theorem 1.13] asserts that every knot K with trivial Alexander polynomial is *topologically slice*, ie $g_4^{\text{top}}(K)$ equals 0; see also Freedman and Quinn [8, Section 11.7B] and Garoufalidis and Teichner [9, Appendix]. Here, the *Alexander polynomial* Δ_K , first introduced by Alexander [1], is the Laurent polynomial with integer coefficients in the indeterminate t defined by

$$\det\left(M\sqrt{t} - M^T \frac{1}{\sqrt{t}}\right),$$

where M is any Seifert matrix for K and M^T is its transpose. The *degree* $\deg \Delta_K$ of the Alexander polynomial Δ_K is the difference of the largest and the smallest exponent among the exponents of the monomials of Δ_K ; ie $\deg \Delta_K$ is the breadth of Δ_K .

Theorem 1 *For every knot, the degree of its Alexander polynomial is greater than or equal to twice its topological slice genus.*

An appealing way of summarizing **Theorem 1** and the classical genus bound of the Alexander polynomial is the following. For all knots K , we have

$$2g_4^{\text{top}}(K) \leq \deg \Delta_K \leq 2g(K),$$

where $g(K)$ denotes the genus of K , ie the minimal genus of Seifert surfaces for K . [Theorem 1](#) determines g_4^{top} for many knots including some of small crossing number; examples are provided in [Section 2](#).

The following purely 3-dimensional proposition reduces [Theorem 1](#) to the genus-zero case. The proof of this proposition, detailed in [Section 3](#), is completely elementary whereas the genus-zero case uses the entire Freedman machine of infinite constructions in topological 4-manifold theory.¹

Proposition 2 *Let K be a knot. Every Seifert surface S of K contains a separating simple closed curve L with the following properties:*

- *The Alexander polynomial of L (as a knot in S^3) is trivial.*
- *The connected component C of $S \setminus L$ that does not contain K is a Seifert surface for L with*

$$2 \text{ genus}(C) = 2 \text{ genus}(S) - \text{deg } \Delta_K.$$

The reduction of [Theorem 1](#) to [Proposition 2](#) and Freedman’s result is rather direct. The same idea was used by Rudolph to provide examples of torus knots whose topological slice genus is smaller than their genus [[12](#), [Theorem 2](#)].

Proof of Theorem 1 For a given knot K , let $L \subset S$ be a simple closed curve in some Seifert surface S of K with the properties described in [Proposition 2](#). By removing the connected component C of $S \setminus L$ that does not contain K , one obtains a surface $\overline{S \setminus C} \subset S^3$ with boundary $K \cup L$. By Freedman’s work [[7](#), [Theorem 1.13](#)], L bounds a topological locally flat disc D in B^4 . Gluing $\overline{S \setminus C}$ and D along L yields a locally flat surface S^{top} of genus $\frac{1}{2} \text{deg } \Delta_K$ in B^4 with boundary $K \subset S^3 = \partial B^4$. To be explicit, S^{top} can be given as follows: shrink D by a factor of 2, yielding a disc in the 4-ball $B_{1/2}^4$ of radius $\frac{1}{2}$ with boundary L viewed as a knot in the 3-sphere of radius $\frac{1}{2}$. Then embed $S \setminus C$ in $B^4 \setminus B_{1/2}^4 = S^3 \times (\frac{1}{2}, 1]$ via the map

$$S \setminus C \rightarrow S^3 \times (\frac{1}{2}, 1], \quad x \mapsto \left(x, \frac{1}{2} + \frac{\text{dist}(L, x)}{2 \text{dist}(K, x) + 2 \text{dist}(L, x)} \right),$$

and set $S^{\text{top}} \subset B^4$ to be the union of the shrunken D and the embedded $S \setminus C$. □

We conclude the introduction by describing previous work relating $\text{deg } \Delta_K$ and g_4^{top} . Borodzik and Friedl proved that $\text{deg } \Delta_K + 1$ is an upper bound for the algebraic unknotting number u_a , which follows from combining their results [[3](#), [Lemma 2.3](#)] and [[2](#), [Theorem 1.1](#)]. Since $g_4^{\text{top}} \leq u_a$, this yields that $g_4^{\text{top}} \leq \text{deg } \Delta_K + 1$.

¹The editorial board of *Geometry & Topology* encourages the reader to refer to [Theorem 1](#) as the “Freedman–Feller Theorem”.

Acknowledgements The present article grew out of questions that arose during a joint effort with Sebastian Baader, Lukas Lewark and Livio Liechti to determine the topological slice genus for families of torus knots and positive knots. I thank them for their helpful remarks; in particular, I thank Lukas for improving the statement of [Proposition 2](#). Thanks also to Josh Greene and the referee for valuable inputs and Stefan Friedl for pointing me to work on the algebraic unknotting number.

The author gratefully acknowledges support by the Swiss National Science Foundation Grant 155477.

2 Applications

Combining [Theorem 1](#) with classical bounds for the topological slice genus, eg Kauffman and Taylor's signature bound [[10](#), Theorem 3.13], yields simple criteria to determine the topological slice genus. Indeed, let $\sigma(K)$ denote the signature of a knot K as introduced by Trotter [[15](#)], ie the signature $\sigma(M + M^T)$ of the symmetrization $M + M^T$ of any Seifert matrix M for the knot K .

Corollary 3 *For every knot K , we have*

$$|\sigma(K)| \leq 2g_4^{\text{top}}(K) \leq \deg \Delta_K.$$

In particular, if $|\sigma(K)| = \deg \Delta_K$, then

$$g_4^{\text{top}}(K) = \left| \frac{1}{2} \sigma(K) \right| = \frac{1}{2} \deg \Delta_K. \quad \square$$

Example 4 We describe an infinite family of knots for which g_4^{top} is arbitrarily large, while being arbitrarily smaller than the smooth slice genus g_4 .

For any positive integer g , any integer $2g \times 2g$ matrix M for which $M - M^T$ has determinant 1 describes the Seifert form on a Seifert surface S bounded by some knot K ; in fact, S can be chosen to be a quasipositive Seifert surface, as proven by Rudolph [[11](#); [13](#), Theorem 1.2]. If one chooses M to satisfy

$$|\sigma(M + M^T)| = \deg \left(\det \left(M \sqrt{t} - M^T \frac{1}{\sqrt{t}} \right) \right) < 2g,$$

then one has examples of knots K for which

$$\deg \Delta_K = 2g_4^{\text{top}}(K) < 2g = 2g_4(K),$$

by [Corollary 3](#) and the fact that quasipositive surfaces realize the smooth slice genus; see Rudolph's slice-Bennequin inequality [[14](#)]. Of course, the above examples include

knots for which g_4^{top} is determined by Freedman's result; eg if K has trivial Alexander polynomial, or if K is a connected sum of knots with trivial Alexander polynomial and knots for which $g = \lfloor \frac{1}{2}\sigma \rfloor$. However, for most knots K as above, we do not know of a method that determines the topological slice genus and that does not use [Theorem 1](#).

Next, we apply [Corollary 3](#) to knots with small crossing number:

Example 5 We determine the topological slice genus of the following knots, which can be represented by diagrams with 12 crossings. With designations as in *KnotInfo* [4], we have:

- $\sigma = -\deg \Delta_K = -4$ for the two knots $12n830$ and $12n750$.
- $\sigma = \deg \Delta_K = 2$ for the two knots $12n519$ and $12n411$.
- $\sigma = -\deg \Delta_K = -2$ for the two knots $12n321$ and $12n293$.

Previously, the topological slice genus appears to have been unknown for all these knots; compare [4]. We remark that, for $12n830$ and $12n750$, the smooth slice genus is known to be 3, and for $12n321$ and $12n293$ it is known to be 2; in particular, it is strictly larger than the topological slice genus; while for $12n519$ and $12n411$ the smooth slice genus appears to be unknown (it is either 1 or 2). In [Section 4](#), we discuss the knot $12n750$ explicitly.

3 Proof of [Proposition 2](#)

We provide a sketch of our proof of [Proposition 2](#). For a given knot K , we fix (in all of [Section 3](#)) a Seifert surface S and denote its genus by g . We find a basis for $H_1(S, \mathbb{Z})$ for which the corresponding Seifert matrix M is of the following form: $M - M^T$ is the standard symplectic form on \mathbb{Z}^{2g} and the bottom right corner of M is a square matrix N of size $2g - \deg \Delta_K$ which represents the Seifert form of a knot with trivial Alexander polynomial. Then we represent this basis by simple closed curves such that for all pairs of curves the geometric intersection number equals the algebraic intersection number, and choose a curve L that separates the curves that represent the last $2g - \deg \Delta_K$ elements of this basis. Thus, N is a Seifert matrix for L and, therefore, L has trivial Alexander polynomial. We note that, if $\Delta_K = 1$, then L is parallel to K and this proof essentially reduces to the proof of [9, Lemma 4.2]; see [Remark 7](#).

In order to provide a detailed proof of [Proposition 2](#), we recall some facts about Seifert matrices and bilinear forms. By choosing a basis for $H_1(S, \mathbb{Z})$, ie by identifying $H_1(S, \mathbb{Z})$ with \mathbb{Z}^{2g} , the Seifert form becomes a bilinear form on \mathbb{Z}^{2g} , which

common divisor of the entries of the last row of $M_{2g} - M_{2g}^T$, which is equal to the last row of M_{2g} , is 1. Therefore, we can change basis again (by performing elementary column operations on M_{2g} simulating the Euclidean algorithm that yields the greatest common divisor of the last row) such that the corresponding Seifert matrix takes the form

$$\left[\begin{array}{ccc|cc} & & & 0 & \\ & & & w_1 & \vdots \\ & & & 0 & \\ \hline & & v_1^T & x & 0 \\ 0 & \cdots & 0 & 1 & 0 \end{array} \right],$$

where M_{2g-2} is a $(2g - 2) \times (2g - 2)$ matrix, x is an integer, and v_1 and w_1 are column vectors with $2g - 2$ entries. By changing basis once more, we can arrange that $x = 0$ and $v_1 = w_1$. The statement of Lemma 6 follows by induction on g , and $2d = \deg \Delta_K$ is immediate from the fact that calculating Δ_K using a Seifert matrix of the form (1) yields

$$\Delta_K = \det\left(M_{2d} \sqrt{t} - M_{2d}^T \frac{1}{\sqrt{t}}\right),$$

which has degree $2d$ since $\det(M_{2d})$ is nonzero. □

Next, we establish that there is also a symplectic (with respect to I) basis for $H_1(S, \mathbb{Z})$ for which the corresponding Seifert matrix is of the form (1); in fact, this follows from a version of Witt’s theorem. Let B be a basis as provided by Lemma 6, and let M_{2g} be the corresponding Seifert matrix. We write $H_1(S, \mathbb{Z}) = V_1 \oplus V_2$, where V_1 denotes the subgroup spanned by the first $2d$ elements of B and V_2 denotes the subgroup spanned by the other elements of B . Denote the lower right square of size $2g - 2d$ of M_{2g} by N_{2g-2d} . By (1), we have that

$$M_{2g} - M_{2g}^T = \begin{bmatrix} M_{2d} - M_{2d}^T & 0 \\ 0 & N_{2g-2d} - N_{2g-2d}^T \end{bmatrix},$$

where $N_{2g-2d} - N_{2g-2d}^T$ equals the standard symplectic form on \mathbb{Z}^{2g-2d} . Since $M_{2d} - M_{2d}^T$ is invertible (which follows from $M_{2g} - M_{2g}^T$ being invertible), there is a symplectic (with respect to the restriction of I) basis B_{V_1} for V_1 . Let $B_{\text{symp}} = (a_1, b_1, \dots, a_g, b_g)$ denote the basis for $H_1(S, \mathbb{Z})$ obtained by replacing the first $2d$ elements of B by B_{V_1} . By construction, B_{symp} is symplectic. The corresponding Seifert matrix M_{symp} is of the form (1) since M_{symp} is obtained from M_{2g} by column (row) operations that involve only the first $2d$ columns (rows); in particular, N_{2g-2d} remains unchanged.

Since B_{sympl} is a symplectic basis for I , it can be realized geometrically; ie for all $1 \leq i \leq g$, there exist simple closed curves α_i and β_i in S representing the classes a_i and b_i , respectively, such that α_i intersects β_i once transversally and no other intersections occur; see eg Farb and Margalit [5, Theorem 6.4]. Let L be any simple closed curve in S separating the curves

$$K, \alpha_1, \beta_1, \dots, \alpha_d, \beta_d \quad \text{from} \quad \alpha_{d+1}, \beta_{d+1}, \dots, \alpha_g, \beta_g,$$

and denote the component of $S \setminus L$ containing $\alpha_{d+1}, \beta_{d+1}, \dots, \alpha_g, \beta_g$ by C . The existence of such a curve L is evident since

$$S \setminus \{K \cup \alpha_1 \cup \beta_1 \cup \dots \cup \alpha_g \cup \beta_g\}$$

is a $(g+1)$ -punctured sphere. The surface C has genus $g - d$ and is a Seifert surface for L . Furthermore, the Seifert matrix corresponding to the basis

$$([\alpha_{d+1}], [\beta_{d+1}], \dots, [\alpha_g], [\beta_g])$$

for $H_1(C, \mathbb{Z})$ is N_{2g-2d} . Therefore, we have

$$\Delta_L = \det \left(N_{2g-2d} \sqrt{t} - N_{2g-2d}^T \frac{1}{\sqrt{t}} \right) \stackrel{(1)}{=} 1. \quad \square$$

Lukas Lewark pointed out Lemma 6. Originally, following the arguments in [6, Lemma 2] and [9, Lemma 4.2], we used changes of basis and S -equivalences to obtain a Seifert matrix of the form (1), which only yields the following weaker version of Proposition 2: every Seifert surface can be stabilized to a Seifert surface that contains a knot with the properties described in Proposition 2. We note that this version still suffices to establish Theorem 1.

The author greatly profited from the nice presentation of Freedman’s result by Garoufalidis and Teichner [9, Appendix], where smooth S^3 - and B^4 -arguments are clearly separated from the application of Freedman’s machinery. In fact, before discovering Proposition 2, which allows one to reduce Theorem 1 to a single application of the fact that knots with trivial Alexander polynomial are topologically slice, our proof of Theorem 1 closely followed the argument in [9, Appendix]. The following remark is related to their presentation:

Remark 7 In the case when $\Delta_K = 1$, the proof of Proposition 2 reduces to the following slight improvement of a lemma [9, Lemma 4.2, first part] by Garoufalidis and Teichner. If a knot has trivial Alexander polynomial, then every Seifert surface has a trivial Alexander basis in the language of [9, Definition 4.1]. This follows by considering the basis $(b_1, \dots, b_g, a_1, \dots, a_g)$ instead of B_{sympl} .




Figure 1: A quasipositive Seifert surface S (gray) for the knot $12n750$ (black) containing two simple closed curves α (red, left) and β (blue, right) such that a neighborhood of $\alpha \cup \beta$ bounds a knot L with trivial Alexander polynomial.

4 Explicit example: the knot $12n750$ and its genera

For the knot $K = 12n750$, which is the closure of the 3–braid

$$(2) \quad aaaba^{-1}baaaba^{-1}b,^2$$

we exhibit the curve L from [Proposition 2](#) explicitly. Let S be the Seifert surface of K depicted in [Figure 1](#). The Seifert surface S has genus 3 and it realizes the genus and the smooth slice genus of K since it is quasipositive [[14](#), slice-Bennequin inequality]; in fact, S is the quasipositive surface canonically associated with the strongly quasipositive braid word given in (2); compare [[11](#); [13](#)]. Let α and β be the two once-intersecting simple closed curves depicted in [Figure 1](#). A neighborhood of their union is a one-holed torus T with a boundary curve $L = \partial T$ that has trivial Alexander polynomial. The latter follows since the Seifert form on T with respect to the basis given by the homology classes of α and β is $\begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix}$. Now, (as in the proof of [Theorem 1](#)) one can modify S by replacing T by a locally flat disc in B^4 to find a locally flat surface S^{top} of genus 2 in B^4 with boundary K . Calculating the signature of K (it is -4) shows that S^{top} realizes $g_4^{\text{top}}(K) = 2$.

References

- [1] **J W Alexander**, *Topological invariants of knots and links*, Trans. Amer. Math. Soc. 30 (1928) 275–306 [MR1501429](#)
- [2] **M Borodzik, S Friedl**, *On the algebraic unknotting number*, Trans. London Math. Soc. 1 (2014) 57–84 [MR3296484](#)

²Here, a and b denote the standard generators of the 3-strand braid group corresponding to a positive crossing of the first two and the last two strands, respectively.

- [3] **M Borodzik, S Friedl**, *The unknotting number and classical invariants, I*, *Algebr. Geom. Topol.* 15 (2015) 85–135 [MR3325733](#)
- [4] **J C Cha, C Livingston**, *KnotInfo: Table of knot invariants* (2015) Available at <http://www.indiana.edu/~knotinfo>
- [5] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Mathematical Series 49, Princeton Univ. Press (2012) [MR2850125](#)
- [6] **M H Freedman**, *A surgery sequence in dimension four; the relations with knot concordance*, *Invent. Math.* 68 (1982) 195–226 [MR666159](#)
- [7] **M H Freedman**, *The topology of four-dimensional manifolds*, *J. Differential Geom.* 17 (1982) 357–453 [MR679066](#)
- [8] **M H Freedman, F Quinn**, *Topology of 4-manifolds*, Princeton Mathematical Series 39, Princeton Univ. Press (1990) [MR1201584](#)
- [9] **S Garoufalidis, P Teichner**, *On knots with trivial Alexander polynomial*, *J. Differential Geom.* 67 (2004) 167–193 [MR2153483](#)
- [10] **L H Kauffman, L R Taylor**, *Signature of links*, *Trans. Amer. Math. Soc.* 216 (1976) 351–365 [MR0388373](#)
- [11] **L Rudolph**, *Constructions of quasipositive knots and links, I*, from: “Knots, braids and singularities”, (C Weber, editor), *Monogr. Enseign. Math.* 31, Enseignement Math., Geneva (1983) 233–245 [MR728589](#)
- [12] **L Rudolph**, *Some topologically locally-flat surfaces in the complex projective plane*, *Comment. Math. Helv.* 59 (1984) 592–599 [MR780078](#)
- [13] **L Rudolph**, *Quasipositivity and new knot invariants*, *Rev. Mat. Univ. Complut. Madrid* 2 (1989) 85–109 [MR1012106](#)
- [14] **L Rudolph**, *Quasipositivity as an obstruction to sliceness*, *Bull. Amer. Math. Soc.* 29 (1993) 51–59 [MR1193540](#)
- [15] **H F Trotter**, *Homology of group systems with applications to knot theory*, *Ann. of Math.* 76 (1962) 464–498 [MR0143201](#)

Department of Mathematics, Boston College
Maloney Hall, Chestnut Hill, MA 02467, United States

peter.feller@math.ch

Proposed: Dmitri Burago
Seconded: Ciprian Manolescu, Ronald Stern

Received: 13 April 2015
Revised: 4 September 2015

