

involve

a journal of mathematics

Induced subgraphs of Johnson graphs

Ramin Naimi and Jeffrey Shaw

 mathematical sciences publishers

2012

vol. 5, no. 1

Induced subgraphs of Johnson graphs

Ramin Naimi and Jeffrey Shaw

(Communicated by Jerrold Griggs)

The Johnson graph $J(n, N)$ is defined as the graph whose vertices are the n -subsets of the set $\{1, 2, \dots, N\}$, where two vertices are adjacent if they share exactly $n - 1$ elements. Unlike Johnson graphs, induced subgraphs of Johnson graphs (JIS for short) do not seem to have been studied before. We give some necessary conditions and some sufficient conditions for a graph to be JIS, including: in a JIS graph, any two maximal cliques share at most two vertices; all trees, cycles, and complete graphs are JIS; disjoint unions and Cartesian products of JIS graphs are JIS; every JIS graph of order n is an induced subgraph of $J(m, 2n)$ for some $m \leq n$. This last result gives an algorithm for deciding if a graph is JIS. We also show that all JIS graphs are edge move distance graphs, but not vice versa.

1. Introduction

We work with finite, simple graphs. Let $F = \{S_1, \dots, S_m\}$ be a family of finite sets. The *intersection graph* of F , denoted $\Omega(F)$, is the graph whose vertices are the elements of F , where two vertices S_i and S_j , $i \neq j$, are adjacent if they share at least one element. More generally, for a fixed positive integer p , the *p -intersection graph* of F , denoted $\Omega_p(F)$, is the graph whose vertices are the elements of F , where two vertices are adjacent if they share at least p elements. (Thus $\Omega_p(F)$ is a subgraph of $\Omega_1(F) = \Omega(F)$.) McKee and McMorris [1999] give an extensive and excellent survey of intersection graphs, which also includes a section on p -intersection graphs. Here we narrow attention to p -intersection graphs of families of $(p + 1)$ -sets, so that two vertices S_i and S_j are adjacent if $|S_i \cap S_j| = |S_i| - 1 = |S_j| - 1$, i.e., S_i and S_j differ by exactly one element.

Another way to view these graphs is as induced subgraphs of Johnson graphs. Given positive natural numbers $n \leq N$, the *Johnson graph* $J(n, N)$ is defined as the graph whose vertices are the n -subsets of the set $\{1, 2, \dots, N\}$, where two vertices are adjacent if they share exactly $n - 1$ elements. Hence a graph G is isomorphic

MSC2000: 05C62.

Keywords: Johnson graph, intersection graph, distance graph.

to an induced subgraph of a Johnson graph if and only if it is possible to assign, for some fixed n , an n -set S_v to each vertex v of G such that distinct vertices have distinct corresponding sets, and vertices v and w are adjacent if and only if S_v and S_w share exactly $n - 1$ elements. When this happens, we say the family of n -sets $F = \{S_v : v \in V(G)\}$ realizes G as an induced subgraph of a Johnson graph, which we abbreviate by saying G is JIS. Thus, F realizes G as a JIS graph if and only if G is isomorphic to $\Omega_{n-1}(F)$, which in turn is isomorphic to an induced subgraph of $J(n, N)$, where $N = |\bigcup_{S \in F} S|$.

Although there is a considerable amount of literature written on Johnson graphs, we have not been able to find any on their induced subgraphs. It would be desirable to obtain “nice” necessary and sufficient conditions for when a graph is JIS. In this paper, we only give some necessary conditions and some sufficient conditions.

A *clique* in a graph G is a complete subgraph of G . A clique L in G is called a *maximal clique*, or a *maxclique* for short, if there is no larger clique $L' \subseteq G$ that contains L . In [Section 2](#) we describe how the maxcliques of a graph play a role in whether or not it is JIS. In particular, [Proposition 2\(1\)](#) states that any two distinct maxcliques in a JIS graph can share at most two vertices. It follows, for example, that the graph “ K_5 minus one edge” is not JIS, since it contains two maximal 4-cliques that share three vertices.

The conditions given in [Section 2](#) are necessary, but not sufficient, for a graph to be JIS. In [Section 3](#) we show that the complete bipartite graph $K_{2,3}$, as well as a few other graphs, satisfy all these necessary conditions but are not JIS. In [Section 3](#) we also give some sufficient conditions for a graph to be JIS, including the following:

- All complete graphs and all cycles are JIS.
- A graph is JIS if and only if all its connected components are JIS.
- The Cartesian product of two JIS graphs is JIS.

Despite not having a “nice” characterization of JIS graphs, for any graph G the question “Is G JIS?” is decidable; this follows from [Theorem 10](#), which says that every JIS graph of order n is isomorphic, for some $m \leq n$, to an induced subgraph of the Johnson graph $J(m, 2n)$. In other words, every JIS graph of order n can, for some $m \leq n$, be realized by m -subsets of $\{1, 2, \dots, 2n\}$. This gives us a simple (albeit slow) algorithm for determining if a graph G is JIS: Do an exhaustive search among all n -families of m -subsets of $\{1, \dots, 2n\}$, where n is the order of G and $m \leq n$, to see if any of them realizes G as a JIS graph.

The p -intersection number of a graph G is defined as the smallest k such that G is isomorphic to the p -intersection graph of a family of subsets of $\{1, \dots, k\}$ ([\[McKee and McMorris 1999\]](#), p. 91). Thus, an immediate corollary of [Theorem 10](#) is that every JIS graph of order n has, for some $m \leq n$, $(m - 1)$ -intersection number at most $2n$.

In the final section of this paper we discuss edge move distance graphs and their relationship to JIS graphs.

2. Maxcliques in JIS Graphs

Given n -sets S_1, \dots, S_k with $n \geq 1$ and $k \geq 2$, we say they share an *immediate subset* if $|\bigcap_{i=1}^k S_i| = n-1$. Similarly, S_1, \dots, S_k share an *immediate superset* if $|\bigcup_{i=1}^k S_i| = n+1$. Observe that for $k=2$, S_1 and S_2 share an immediate subset if and only if they share an immediate superset: $|S_1 \cup S_2| = |S_1| + |S_2| - |S_1 \cap S_2| = 2n - |S_1 \cap S_2|$; hence $|S_1 \cup S_2| = n+1$ if and only if $|S_1 \cap S_2| = n-1$. We begin with the following elementary result on realizations of complete graphs as JIS graphs.

Lemma 1. *Let S_1, \dots, S_k be n -sets that pairwise share an immediate subset, where $n \geq 1$ and $k \geq 3$. Then S_1, \dots, S_k share an immediate subset or an immediate superset, but not both.*

Proof. We first show that for $k \geq 3$, if S_1, \dots, S_k share an immediate subset, then they do not share an immediate superset. Suppose $T = S_1 \cap \dots \cap S_k$ has $n-1$ elements. Then, for each i , $S_i \setminus T$ has exactly one element, x_i . For all $j \neq i$, $x_i \notin S_j$ since $S_i \neq S_j$. It follows that $S_1 \cup \dots \cup S_k$ has at least $n-1+k \geq n+2$ elements, since $k \geq 3$. Thus S_1, \dots, S_k do not share an immediate superset.

Now suppose S_1, \dots, S_k pairwise share an immediate subset. We use induction on k to prove that they share an immediate subset or an immediate superset.

Assume $k=3$. Let $T = S_1 \cap S_2$. If $T \subset S_3$, then $|S_1 \cap S_2 \cap S_3| = |T| = n-1$, and we're done. So assume $T \not\subset S_3$. Note that $|S_1 \setminus T| = |S_2 \setminus T| = 1$. Hence, for S_3 to share $n-1$ elements with each of S_1 and S_2 , it must contain an $(n-2)$ -subset of T , as well as $S_1 \setminus T$ and $S_2 \setminus T$, and no other elements. It follows that $|S_1 \cup S_2 \cup S_3| = n+1$, as desired.

Now assume $k \geq 4$. Then, by our induction hypothesis, S_1, \dots, S_{k-1} share an immediate subset or an immediate superset; and similarly for S_2, \dots, S_k . We have four cases:

Case 1: S_1, \dots, S_{k-1} share an immediate subset and S_2, \dots, S_k share an immediate subset. Then S_1, \dots, S_k share $S_2 \cap S_3$ as an immediate subset.

Case 2: S_1, \dots, S_{k-1} share an immediate superset and S_2, \dots, S_k share an immediate superset. Then S_1, \dots, S_k share $S_2 \cup S_3$ as an immediate superset.

Case 3: S_1, \dots, S_{k-1} share an immediate subset and S_2, \dots, S_k share an immediate superset. Let $T = S_1 \cap \dots \cap S_{k-1}$. Then, for $1 \leq i \leq k-1$, $S_i \setminus T$ has exactly one element, x_i ; and, for $1 \leq j \leq k-1$ with $j \neq i$, $x_i \notin S_j$ since $S_i \neq S_j$. Since $|S_2 \cup \dots \cup S_k| = n+1 = |S_2 \cup S_3|$, S_k is a proper subset of $S_2 \cup S_3 = T \cup \{x_2, x_3\}$. And since S_2, S_3, S_k share an immediate superset, they do not share an immediate subset; hence $T \not\subset S_k$. This implies that $x_2, x_3 \in S_k$ since S_k has n elements and

$T \cup \{x_2, x_3\}$ has $n + 1$ elements. But $x_2, x_3 \notin S_1$, so $|S_1 \cap S_k| < n - 1$, which contradicts the hypothesis of the lemma.

Case 4: S_1, \dots, S_{k-1} share an immediate superset and S_2, \dots, S_k share an immediate subset. This case is similar to Case 3. \square

We now use [Lemma 1](#) to establish restrictions on how maxcliques in a JIS graph can intersect or connect to each other by edges.

Proposition 2. *Suppose G is JIS and L and L' are distinct maxcliques in G .*

- (1) L and L' share at most two vertices.
- (2) If L and L' share exactly two vertices, then no vertex in $V(L) \setminus V(L')$ is adjacent to a vertex in $V(L') \setminus V(L)$.
- (3) If L and L' share exactly one vertex, then each vertex in either of the two sets $V(L) \setminus V(L')$ and $V(L') \setminus V(L)$ is adjacent to at most one vertex in the other set.

Proof. Let $\{S_v : v \in V(G)\}$ be a family of n -sets that realizes G as a JIS graph.

(1) Suppose towards contradiction that L and L' are distinct maxcliques that share three (or more) vertices, u, v , and w . Let x be a vertex of L not in L' , and x' a vertex of L' not in L ; x and x' exist since L and L' are distinct and maximal. Then, by [Lemma 1](#), the sets S_x, S_u, S_v , and S_w share an immediate subset or an immediate superset. Similarly for $S_{x'}, S_u, S_v$, and S_w . But S_u, S_v , and S_w cannot share both an immediate subset and an immediate superset. It follows that S_x and $S_{x'}$ share an immediate subset or an immediate superset, which implies that x and x' are adjacent. Hence every vertex of L is adjacent to every vertex of L' , but this contradicts the assumption that L is a maxclique in G .

(2) Let L and L' be distinct maxcliques that share exactly two vertices, v and w . Suppose towards contradiction that there exist adjacent vertices $x \in V(L) \setminus V(L')$ and $x' \in V(L') \setminus V(L)$. Then the induced subgraph of G containing $\{x, x', v, w\}$ is a 4-clique. Let L'' be the maxclique that contains this 4-clique. Then L'' is distinct from L and shares at least three vertices with it. This contradicts (1).

(3) The proof is similar to the proof of (2). Let L and L' be distinct maxcliques that share exactly one vertex, v . Suppose towards contradiction that there exist vertices $x \in V(L) \setminus V(L')$ and $x', y' \in V(L') \setminus V(L)$ with x adjacent to x' and y' . Then the induced subgraph of G containing $\{x, x', y', v\}$ is a 4-clique, and the maxclique that contains this 4-clique is distinct from L' and shares at least three vertices with it. This contradicts (1). \square

Proposition 3. *Suppose L_1, \dots, L_k , where k is odd and at least 3, are distinct maxcliques in a graph G such that L_i shares exactly two vertices with L_{i+1} for $1 \leq i \leq k - 1$, and L_k shares exactly two vertices with L_1 ; then G is not JIS.*

Proof. In the following, L_{i+1} refers to L_1 whenever $i = k$. Suppose towards contradiction that G is realized as a JIS graph by a family of n -sets. Note that each L_i has at least three vertices, since otherwise it would not be distinct from L_{i+1} . Hence, by [Lemma 1](#), we can label each L_i as either “sub” or “super” according to whether the n -sets assigned to its vertices share an immediate subset or an immediate superset. Then, since k is odd, there exists a j such that L_j and L_{j+1} have the same label. Now, L_j and L_{j+1} share two vertices; therefore the n -sets assigned to their vertices must all share the same immediate subset or immediate superset, which makes all vertices in L_j adjacent to those in L_{j+1} , giving a contradiction. \square

An equivalent way of stating the above result is: One can label every maxclique in a JIS graph with a $+$ or $-$ (or any two symbols) in such a way that any two maxcliques that share two vertices have distinct labels.

3. Miscellaneous JIS and non-JIS graphs

In this section we give some sufficient conditions for when a graph is JIS. We also describe some graphs that satisfy all the conditions listed in the results of the previous section as necessary for a graph to be JIS, but are not JIS.

Proposition 4. *All complete graphs and all cycles are JIS.*

Proof. For each n , K_n is realized as a JIS graph by the 1-sets $\{1\}, \{2\}, \dots, \{n\}$. For each $n \geq 3$, the n -cycle is realized as a JIS graph by the 2-sets $\{1, 2\}, \{2, 3\}, \dots, \{n-1, n\}, \{n, 1\}$. \square

We define the n -core of a graph G as the graph obtained by recursively removing all vertices of degree less than n until there are none left.

Proposition 5. *A graph is JIS if and only if its 2-core is JIS.*

Proof. Suppose G is obtained from a graph G' by removing exactly one vertex, w , which has degree 0 or 1. By induction, it is enough to show that G is JIS if and only if G' is JIS. Clearly, if G' is JIS, then so is G , since any induced subgraph of a JIS graph is JIS. To prove the converse, suppose G is JIS. Let $\{S_x : x \in V(G)\}$ be n -sets that realize G as a JIS graph. Pick distinct a and b that are not in any of the sets S_x . For each $x \in V(G)$, let $S'_x = S_x \cup \{a\}$. Let $S'_w = S_v \cup \{b\}$, where $v \in V(G')$ is arbitrary if w has degree 0, and v is adjacent to w if w has degree 1. Then $\{S'_x : x \in V(G')\}$ are $(n+1)$ -sets that realize G' as a JIS graph, as desired. \square

It follows as a trivial corollary that all trees are JIS.

Proposition 6. *A graph is JIS if and only if all its connected components are JIS.*

Proof. One direction is trivial: every induced subgraph of a JIS graph, and in particular every connected component of it, is JIS. We prove the converse by induction on the number of components of G .

Base step: Suppose that G has two components, G_i , $i = 1, 2$, each realized as a JIS graph by a family of sets F_i . We can assume without loss of generality that each set in F_1 is disjoint from each set in F_2 .

We would like each set in F_1 to have the same size as each set in F_2 , in order to obtain $F_1 \cup F_2$ as a family that realizes G as a JIS graph. If this is not already so, we proceed as follows. Let m_i denote the number of elements in each set in F_i . We can assume $n_1 > n_2$. Now add the first $n_1 - n_2$ elements of the first set in F_1 to every set in F_2 .

Once the sets in the two families all have the same size, we must make sure that sets corresponding to vertices in different components of G do not share immediate subsets. This will automatically be true for sets that had two or more elements before any extra elements were added to them (since we started with the sets in F_1 disjoint from those in F_2), but not for singletons. We remedy this by adding, for each i , an element e_i to every set in F_i , where e_1 and e_2 are distinct elements not already in any set in any F_i . It is now easy to verify that $F_1 \cup F_2$ realizes G as a JIS graph.

The inductive step follows trivially from the base step. \square

Proposition 7. *The Cartesian product of two JIS graphs is JIS.*

Proof. Let G and G' be JIS graphs that are realized, respectively, by sets $\{S_x : x \in V(G)\}$ and $\{S'_{x'} : x' \in V(G')\}$. We can assume without loss of generality that every S_x is disjoint from every $S'_{x'}$.

For each vertex $v = (x, x') \in V(G \times G')$, let $T_v = S_x \cup S'_{x'}$. By definition, two vertices $v = (x, x')$ and $w = (y, y')$ of $G \times G'$ are adjacent if and only if $x = x'$ and y is adjacent to y' or $y = y'$ and x is adjacent to x' . Thus, T_v and T_w share an immediate subset if and only if v and w are adjacent. Hence the sets $\{T_v : v \in G \times G'\}$ realize $G \times G'$ as a JIS graph. \square

Proposition 8. *The complete bipartite graph $K_{2,3}$ is not JIS.*

Proof. Label the two degree-3 vertices of $K_{2,3}$ as v and w , and the three degree-2 vertices as x , y , and z , as in [Figure 1](#). Suppose towards contradiction that there exists a family of n -sets $\{S_u : u \in V(K_{2,3})\}$ that realizes $K_{2,3}$ as a JIS graph. Since v and w have distance two (where *distance* is the number of edges in the shortest




Figure 1. $K_{2,3}$ with labeled vertices.

path joining the two vertices), S_v and S_w must share exactly $n - 2$ elements (this does not work for distance ≥ 3 ; it works only for distance ≤ 2). Let $T = S_v \cap S_w$. Then, since each of x , y , and z is adjacent to both v and w , S_x , S_y , and S_z must each contain T as a subset. Therefore, by subtracting T from every S_u , $u \in V(K_{2,3})$, we get a family of 2-sets that realizes $K_{2,3}$. Hence we will assume that every S_u has exactly two elements. It follows that S_v and S_w are disjoint; and S_x , S_y , and S_z are pairwise disjoint and each shares exactly one element with each of S_v and S_w .

So, without loss of generality, $S_v = \{1, 2\}$, and $S_w = \{3, 4\}$. Therefore, again without loss of generality, $S_x = \{1, 3\}$, and $S_y = \{2, 4\}$. And there is nothing left for S_z . \square

The graph $K_{2,3}$ can be thought of as two 4-cycles that share three vertices. So one may wonder whether the graph θ_n consisting of two n -cycles that share $n - 1$ vertices is also not JIS. It turns out that θ_n is not JIS only for $n = 4$ and $n = 5$. The proof that θ_5 is not JIS is very similar to the proof that $K_{2,3}$ is not JIS, and we therefore omit it. The proof that θ_n is JIS for $n \geq 6$ is a straightforward construction, which we also omit.

One may also wonder whether $K_{2,3}$ becomes JIS if an edge is added to it. There are, up to isomorphism, two ways to add an edge to $K_{2,3}$: add an edge that connects the two degree-3 vertices; or add an edge that connects two of the three degree-2 vertices. It turns out that neither of these two graphs is JIS. The proof that the former graph is not JIS follows immediately from [Proposition 3](#). The proof that the latter graph (which we call Δ_2) is not JIS is given below in [Proposition 9](#).

The graphs Δ_i depicted in [Figure 2](#) have the following pattern (ignore the vertex labels and the $+$ and $-$ signs for now; they are used later): Δ_i consists of a chain of i ‘‘consecutively adjacent’’ triangles, plus one vertex which is connected to the two vertices of degree 2 in the triangle chain. It turns out that, like $K_{2,3}$, Δ_2 , Δ_4 , and Δ_6 satisfy the necessary conditions in the results of the previous sections for being JIS, but are not JIS; Δ_3 and Δ_5 , however, are JIS. We prove these claims below, except for Δ_6 : its proof is similar to that of Δ_2 and Δ_4 , but is more tedious, and in our opinion not worth being included here. We did not check which Δ_i are JIS for $i \geq 7$, but, from the pattern for $i \leq 6$, it seems that:

Conjecture. Δ_i is JIS if and only if i is odd.

Proposition 9. (i) The graphs Δ_2 and Δ_4 are not JIS. (ii) The graphs Δ_3 and Δ_5 are JIS.

Remark. As mentioned above, Δ_2 is isomorphic to $K_{2,3}$ plus an edge that connects two of its three degree-2 vertices. Because of this, the proof that $K_{2,3}$ is not JIS can be easily modified to prove that Δ_2 is not JIS. However, we give a different proof below, one that can be naturally extended to also prove that Δ_4 (and Δ_6) is not JIS.

Proof. Label the vertices of Δ_2 as v_1, \dots, v_5 , as in [Figure 2](#). The $+$ and $-$ signs will be explained shortly. Suppose, towards contradiction, that Δ_2 can be realized as a JIS graph by sets S_1, \dots, S_5 (for simplicity, we write S_i instead of S_{v_i}). Each of the two triangles in Δ_2 is a maxclique. Thus, by [Lemma 1](#), S_1, S_2 , and S_3 must share an immediate subset or an immediate superset; similarly for S_2, S_3 , and S_4 . Furthermore, S_1, S_2 , and S_3 share an immediate subset if and only if S_2, S_3 , and S_4 share an immediate superset, because: if S_1, S_2 , and S_3 share an immediate subset and S_2, S_3 , and S_4 also share an immediate subset, then S_1 and S_4 must share $S_2 \cap S_3$ as an immediate subset, but this contradicts the fact that v_1 and v_4 are not adjacent; and if S_1, S_2 , and S_3 share an immediate superset and S_2, S_3 , and S_4 also share an immediate superset, then S_1 and S_4 must share $S_2 \cup S_3$ as an immediate superset, which implies that they also share an immediate subset, again contradicting the fact that v_1 and v_4 are not adjacent.

Thus, without loss of generality, we will assume that S_1, S_2 , and S_3 share an immediate subset. This is indicated in [Figure 2](#) by the $-$ sign; the $+$ signs indicate immediate supersets. So we will assume that $S_1 = \{1, 2, 3, 4\}$, $S_2 = \{1, 2, 3, 5\}$, and $S_3 = \{1, 2, 3, 6\}$; we explain in the next paragraph why there is no loss of generality in assuming that S_i are 4-sets (as opposed to larger sets). To make the notation more compact, we will drop the commas and the braces from each set; e.g., $S_1 = 1234$. Then S_4 must be a 4-subset of $S_2 \cup S_3 = 12356$. Since S_1 and S_4 have no immediate subset, we can without loss of generality assume that $S_4 = 2356$. Now, S_5 must differ by exactly one element from each of S_1 and S_4 . The only possibilities are 1235, 1236, 2345, and 2346. But the first two are equal to S_2 and S_3 respectively; and the last two differ from S_2 and S_3 respectively by exactly one element, which is not allowed since v_5 is adjacent to neither v_2 nor v_3 . Thus we have a contradiction, as desired.

Note that by assuming that all S_i are 4-sets, we ended up with all of them sharing the two elements 2 and 3. If we instead assumed that S_i were n -sets with $n \geq 5$,




Figure 2. Δ_2 , Δ_3 , and Δ_4 , with vertices labeled in Δ_2 and Δ_4 .




Figure 3. Δ_3 (left) and Δ_5 (right) realized as JIS graphs.

the proof would remain the same except that we would end up with all S_i sharing more than two elements. Hence there is no loss of generality in assuming that S_i are 4-sets (in fact, this shows that we could even assume they are 2-sets).

To prove that Δ_4 is not JIS, we start with the same assumptions that $S_1, S_2,$ and S_3 share an immediate subset, $S_2, S_3,$ and S_4 share an immediate superset, and $S_1 = 1234, S_2 = 1235, S_3 = 1236,$ and $S_4 = 2356.$ Now, $S_3, S_4,$ and S_5 must share an immediate subset. So S_5 must contain $S_3 \cap S_4 = 236.$ Since v_5 is adjacent to neither v_1 nor $v_2,$ S_5 can contain neither 1 nor 4 nor 5. Hence, without loss of generality, $S_5 = 2367.$ Continuing, $S_4, S_5,$ and S_6 must share an immediate superset. So S_6 must be a 4-subset of $S_4 \cup S_5 = 23567;$ i.e., we must drop one element from 23567 to get $S_6.$ Dropping 5 or 7 gets us back to S_4 and $S_5;$ hence we must drop 2, 3, or 6. The roles of 2 and 3 have been identical so far; so, without loss of generality, we must drop 2 or 6; so $S_6 = 2357$ or $3567.$ The former is not possible since v_6 and v_2 are not adjacent. And the latter is ruled out by noticing that 3567 differs from $S_1 = 1234$ by three elements, which contradicts the fact that v_6 and v_1 have distance two¹. Thus we have reached a contradiction, as desired.

Part (ii) of the proposition is proved in [Figure 3](#), which shows sets that realize Δ_3 and Δ_5 as JIS graphs. For the sake of compactness, braces and commas are omitted from the sets. □

We end this section with the following definition and question. Let G be a JIS graph, and suppose $F = \{S_u : u \in V(G)\}$ realizes G as a JIS graph. We define the F -distance between two vertices v and w of G to be $d_F(v, w) = |S_v \setminus S_w|.$ It is easy to show this distance function is indeed a metric. The JIS -diameter of G is defined as

$$\max_{v,w \in V(G)} \min_F \{d_F(v, w)\}$$

where the minimum is taken over all families F that realize G as a JIS graph.

Question. Do there exist JIS graphs with arbitrarily large JIS-diameter?

¹Note that $\Delta_4 - v_7$ is JIS, with S_1 and S_6 differing in three elements. We will refer back to this point at the very end of this section.

From the proof of [Proposition 9](#) and the footnote in it, it follows that Δ_4 minus the degree-2 vertex v_7 has JIS-diameter 3: $S_1 = 1234$, $S_2 = 1235$, $S_3 = 1236$, $S_4 = 2356$, $S_5 = 2367$, and $S_6 = 3567$, i.e., v_1 and v_6 have F -distance 3.

4. An algorithm for recognizing JIS graphs

As mentioned in the introduction, the following theorem provides for an algorithm for deciding if a graph is JIS by doing a bounded exhaustive search.

Theorem 10. *Every JIS graph of order n is isomorphic, for some $m \leq n$, to an induced subgraph of the Johnson graph $J(m, 2n)$.*

Proof. Let G be a JIS graph of order n with c connected components.

Case 1. Assume $c = 1$, i.e., G is connected. In this case we will prove a slightly stronger result, which we will use in the proof of Case 2:

G is isomorphic, for some $m \leq n$, to an induced subgraph of $J(m, 2n-1)$.

The case $n = 1$ is trivial; so we assume $n \geq 2$. Since G is connected, there exists an ordering v_1, v_2, \dots, v_n of the vertices of G such that for each $i \geq 2$, v_i is adjacent to at least one of v_1, \dots, v_{i-1} . Since G is JIS, for some $k \geq 1$ there exist k -sets $\{S_1, \dots, S_n\}$ that realize G as a JIS graph, where S_i corresponds to the vertex v_i . Since v_1 and v_2 are adjacent, $|S_1 \cap S_2| = k - 1$. Since v_3 is adjacent to at least one of v_1 and v_2 , $|S_1 \cap S_2 \cap S_3| \geq k - 2$. Continuing this way, we see that $|S_1 \cap \dots \cap S_n| \geq k - (n - 1)$. Let

$$S'_i = S_i \setminus (S_1 \cap \dots \cap S_n)$$

for $1 \leq i \leq n$. Then for all i , $|S'_i| = m$ where $m \leq k - (k - (n - 1)) = n - 1$, and it is easily verified that the family of sets $\{S'_1, \dots, S'_n\}$ realizes G as a JIS graph.

Now, since v_1 and v_2 are adjacent, $|S'_1 \cup S'_2| = m + 1$. Since v_3 is adjacent to at least one of v_1 and v_2 , $|S'_1 \cup S'_2 \cup S'_3| \leq m + 2$. Continuing this way, we see that $|S'_1 \cup \dots \cup S'_n| \leq m + n - 1 \leq 2n - 2$, which implies G is an induced subgraph of $J(m, 2n - 1)$, $m \leq n - 1$. (Note: we proved the inequalities $|S'_1 \cup \dots \cup S'_n| \leq 2n - 2$ and $m \leq n - 1$ only for $n \geq 2$, not for $n = 1$.)

Case 2. Assume $c \geq 2$. Let n_i be the order of the i th component of G . Then, by Case 1 above, for each i there is a family F_i of m_i -sets, $m_i \leq n_i$, that realizes the i th component of G as a JIS graph, such that the union of the sets in F_i has at most $2n_i - 1$ elements. Thus $\bigcup F_i$ has at most $2n - c$ elements.

We can assume $m_1 \geq m_i$ for all i . We can also assume that for all $i \neq j$, every set in the family F_i is disjoint from every set in F_j . To make all sets in all the families have the same size, for each i such that $m_1 > m_i$ we add the first $m_1 - m_i$ elements of the first set in F_1 to every set in F_i . After adding these extra elements, we must make sure that sets corresponding to vertices in different components of G do not

share immediate subsets. This will automatically be true for sets that had two or more elements before the extra elements were added, but not for singletons. We remedy this by adding, for each i , an element e_i to every set in F_i , where e_1, \dots, e_c are distinct elements not already in any set in any F_i . Let $F = \bigcup F_i$. Then G is realized as a JIS graph by F , which is a family of $(m_1 + 1)$ -sets whose union has at most $2n - c + c = 2n$ elements, where $m_1 + 1 \leq n_1 + 1 \leq n$. Thus G is an induced subgraph of $J(m, 2n)$ where $m = m_1 + 1 \leq n$. \square

Remark. It is not difficult to modify the above proof in Case 1 to show that if G is connected, then it is an induced subgraph of $J(n, 2n)$. It would be interesting to see for which graphs the bounds n and $2n$ can be lowered. Note that if G consists of exactly $n \geq 2$ vertices of degree zero, then the bound $2n$ is optimal.

5. Edge move distance graphs and JIS graphs

Since the 1970s many authors have written on various metrics defined on sets of graphs; see, for instance, [Benadé et al. 1991; Chartrand et al. 1997; 1990; Deza and Deza 2009; Johnson 1987; Kaden 1983; Zelinka 1985]. Among them are edge move, edge rotation, edge jump, and edge slide distances. In general, given a metric d on a set of graphs $S = \{G_1, \dots, G_k\}$, the *distance graph* of S , denoted $D_d(S)$, has S as its vertex set, where two vertices G_i and G_j are adjacent if $d(G_i, G_j) = 1$. We will see shortly that distance graphs associated with the edge move metric are closely related to JIS graphs.

An *edge move* on a graph G consists of removing one edge from and adding a new edge to G , without changing its vertex set $V(G)$; i.e., one edge is “moved to a new position.” The *edge move distance* $d_m(G, H)$ between two graphs G and H is defined as the fewest number of edge moves necessary to transform G into H , up to isomorphism. Note that for $d_m(G, H)$ to be defined, G and H must have the same order and the same size. It is easy to verify that d_m is a metric on any set of graphs of given order and size. Given a set S of graphs of the same order and size, the *edge move distance graph* of S , $D_m(S)$, is the graph whose vertices are the elements of S , where two vertices are adjacent if their edge move distance is one. When we say a graph is an edge move distance graph we mean it is isomorphic to one.

The connection between JIS graphs and edge move distance graphs can be seen by focusing on edge sets. Let G and H be graphs of the same order and size, with n edges each. If the edge sets $E(G)$ and $E(H)$ share exactly $n - 1$ elements, then G and H have edge move distance one. Conversely, if G and H have edge move distance one, then their vertices can be labeled such that $E(G)$ and $E(H)$ share exactly $n - 1$ elements. At first glance, this might seem to suggest that a graph is JIS if and only if it is isomorphic to an edge move distance graphs. We will show, however, that only half (one direction) of this statement is true.

Proposition 11. *Every JIS graph is an edge move distance graph.*

Proof. Let G be realized as a JIS graph by a family of n -sets $\{S_v : v \in V(G)\}$. We will construct a graph G_v for each $v \in V(G)$ such that $d_m(G_v, G_w) = 1$ if and only if S_v and S_w share an immediate subset.

We can assume that each S_v consists of positive integers. Let

$$k = 1 + \max\{i \in S_v : v \in V(G)\},$$

and let P be a path of length $2k$. Denote the vertices of P by p_0, p_1, \dots, p_{2k} . For each $v \in V(G)$, we let G_v be the graph consisting of P plus the edges $p_i p_{2k-i}$ for all $i \in S_v$. Then it is easily verified that for $v \neq w$, G_v is not isomorphic to G_w , and $d_m(G_v, G_w) = 1$ if and only if S_v and S_w share an immediate subset. Therefore G is isomorphic to the edge move distance graph $D_m(\{G_v : v \in V(G)\})$. \square

The converse is not true. The reason is that the number of edges shared by the edge sets of two graphs depends on how their vertices are labeled, whereas edge move distance is measured up to graph isomorphism.

Proposition 12. *The graph obtained by removing one edge from the complete graph K_n , where $n \geq 5$, is an edge move distance graph but is not JIS.*

Proof. Fix $n \geq 5$, and let H be the graph obtained by removing one edge from K_n . Then H contains two maximal $(n-1)$ -cliques which share $n-2$ vertices. Hence, by [Proposition 2\(1\)](#), H is not JIS.

To show that H is an edge move distance graph, we construct a set of graphs $S = \{Q_1, Q_2, \dots, Q_n\}$ such that $H \simeq D_m(S)$. For $1 \leq i \leq n$, Q_i has $n+2$ vertices: $V(Q_i) = \{v_1, v_2, \dots, v_{n+2}\}$. For $1 \leq i \leq n-1$, we have

$$E(Q_i) = \{v_k v_{k+1} : 1 \leq k \leq n\} \cup \{v_{n-1} v_{n+1}, v_i v_{n+2}\};$$

and $E(Q_n) = (E(Q_1) \cup \{v_1 v_{n-2}\}) \setminus \{v_{n-2} v_{n-1}\}$.

Then one readily verifies for all $i \neq j$ except when $\{i, j\} = \{n-1, n\}$ that Q_i and Q_j have edge move distance one. Thus H is an edge move distance graph. \square

Figures [4](#) and [5](#) show some of the Q_i in the case $n = 6$.




Figure 4. Q_1 for $n = 6$.




Figure 5. Q_{n-1} (left) and Q_n (right) for $n = 6$.

Acknowledgments

We thank Terry A. McKee of Wright State University for bringing to our attention that the graphs we were studying are related to Johnson graphs. Naimi thanks Caltech for its hospitality while he did part of this work there during his sabbatical leave. Shaw thanks the Undergraduate Research Center of Occidental College for providing support to do this work as part of an undergraduate summer research project. We also thank the (anonymous) referee for helpful suggestions.

References

- [Benadé et al. 1991] G. Benadé, W. Goddard, T. A. McKee, and P. A. Winter, “On distances between isomorphism classes of graphs”, *Math. Bohem.* **116**:2 (1991), 160–169. [MR 92g:05173](#) [Zbl 0753.05067](#)
- [Chartrand et al. 1990] G. Chartrand, W. Goddard, M. A. Henning, L. Lesniak, H. C. Swart, and C. E. Wall, “Which graphs are distance graphs?”, *Ars Combin.* **29**:A (1990), 225–232. [MR 97f:05156](#) [Zbl 0716.05028](#)
- [Chartrand et al. 1997] G. Chartrand, H. Gavlas, H. Hevia, and M. A. Johnson, “Rotation and jump distances between graphs”, *Discuss. Math. Graph Theory* **17**:2 (1997), 285–300. [MR 99f:05084](#) [Zbl 0902.05022](#)
- [Deza and Deza 2009] M. M. Deza and E. Deza, *Encyclopedia of distances*, Springer, Berlin, 2009. [MR 2011b:51001](#) [Zbl 1167.51001](#)
- [Johnson 1987] M. A. Johnson, “An ordering of some metrics defined on the space of graphs”, *Czechoslovak Math. J.* **37**:1 (1987), 75–85. [MR 88d:05158](#) [Zbl 0641.05027](#)
- [Kaden 1983] F. Kaden, “Graph metrics and distance-graphs”, pp. 145–158 in *Graphs and other combinatorial topics* (Prague, 1982), edited by M. Fiedler, Teubner-Texte in Mathematik **59**, Teubner, Leipzig, 1983. [MR 85e:05060](#) [Zbl 0528.05055](#)
- [McKee and McMorris 1999] T. A. McKee and F. R. McMorris, *Topics in intersection graph theory*, SIAM, Philadelphia, 1999. [MR 2000e:05001](#) [Zbl 0945.05003](#)
- [Zelinka 1985] B. Zelinka, “Comparison of various distances between isomorphism classes of graphs”, *Časopis Pěst. Mat.* **110**:3 (1985), 289–293. [MR 87c:05114](#) [Zbl 0579.05056](#)

Received: 2010-08-04

Revised: 2011-07-01

Accepted: 2011-07-09

rnaimi@oxy.edu

Department of Mathematics, Occidental College, 1600
Campus Road, Los Angeles, CA 90041-3314, United States

jeffreyshaw@gmail.com

Department of Mathematics, Occidental College, 1600
Campus Road, Los Angeles, CA 90041-3314, United States

EDITORS

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Pietro Cerone	Victoria University, Australia pietro.cerone@vu.edu.au	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Joseph Gallian	University of Minnesota Duluth, USA kgallian@d.umn.edu	Robert J. Plemmons	Wake Forest University, USA rjplemmons@wfu.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Andrew Granville	Université Montréal, Canada andrew@dms.umontreal.ca	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Sat Gupta	U of North Carolina, Greensboro, USA sgupta@uncg.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Karen Kafadar	University of Colorado, USA karen.kafadar@cudenver.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
David Larson	Texas A&M University, USA larson@math.tamu.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu	Michael E. Zieve	University of Michigan, USA zieve@umich.edu

PRODUCTION

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor

Cover design: ©2008 Alex Scorpan

See inside back cover or <http://msp.berkeley.edu/involve> for submission instructions.

The subscription price for 2012 is US \$105/year for the electronic version, and \$145/year (+\$35 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94704-3840, USA.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.



PUBLISHED BY
mathematical sciences publishers
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Mathematical Sciences Publishers

involve

2012

vol. 5

no. 1

Elliptic curves, eta-quotients and hypergeometric functions	1
DAVID PATHAKJEE, ZEF ROSNBRICK AND EUGENE YOONG	
Trapping light rays aperiodically with mirrors	9
ZACHARY MITCHELL, GREGORY SIMON AND XUEYING ZHAO	
A generalization of modular forms	15
ADAM HAQUE	
Induced subgraphs of Johnson graphs	25
RAMIN NAIMI AND JEFFREY SHAW	
Multiscale adaptively weighted least squares finite element methods for convection-dominated PDEs	39
BRIDGET KRAYNIK, YIFEI SUN AND CHAD R. WESTPHAL	
Diameter, girth and cut vertices of the graph of equivalence classes of zero-divisors	51
BLAKE ALLEN, ERIN MARTIN, ERIC NEW AND DANE SKABELUND	
Total positivity of a shuffle matrix	61
AUDRA MCMILLAN	
Betti numbers of order-preserving graph homomorphisms	67
LAUREN GUERRA AND STEVEN KLEE	
Permutation notations for the exceptional Weyl group F_4	81
PATRICIA CAHN, RUTH HAAS, ALOYSIUS G. HELMINCK, JUAN LI AND JEREMY SCHWARTZ	
Progress towards counting D_5 quintic fields	91
ERIC LARSON AND LARRY ROLEN	
On supersingular elliptic curves and hypergeometric functions	99
KEENAN MONKS	