

A FANO 3-FOLD WITH NON-RATIONAL SINGULARITIES AND A TWO DIMENSIONAL BASIS

BY SHIHOKO ISHII

Introduction

In this paper, the author gives a summary of the paper [I2] and pictures of the Fano 3-folds which appear in it. Here a Fano 3-fold means a normal projective variety of dimension three over \mathbb{C} whose anticanonical sheaf is ample and invertible. During the past fifteen years, there has been big progress in the investigation of a non-singular Fano 3-fold owing to Iskovskih, Mori, Mukai and Shokurov. And it is still developing. On the other hand, in singular Fano 3-folds, progress seems to have started recently. Here we study the structure of a Fano 3-fold with non-rational singularities.

Let Σ be the locus of non-rational singular points of a Fano 3-fold X . As X is normal, $\dim \Sigma \leq 1$. If $\dim \Sigma = 0$, then X is isomorphic to a projective cone over a normal K3-surface or an Abelian surface (Theorem 1A, 1B). The proof of this theorem also works in the case that Σ contains an isolated point. So what we should study next is the case that Σ has pure dimension one. Such a Fano 3-fold is classified in three families according to the maximal basis-dimension of its \mathbb{Q} -factorial terminal modification (Theorem 2, Definition 1). We obtain the fact that a Fano 3-fold with the maximal basis-dimension 2 admits a projective bundle over a non-singular surface as a \mathbb{Q} -factorial terminal modification (Theorem 3). We try to make clear the structure of a Fano 3-fold in this family: what kind of surface occurs as a basis, what kind of projective bundle appears as a \mathbb{Q} -factorial terminal modification and which parts on the projective bundle are contracted in a Fano 3-fold.

The author would like to thank Professors Nakayama and Kei-ichi Watanabe and also other members of Waseda Seminar for their stimulating discussion during the preparation of this article. In particular Nakayama's proof of Proposition 1 helped her very much and also K-i. Watanabe's comment "a Weil divisor on a \mathbb{Q} -factorial terminal singularity is Cohen-Macaulay" was very helpful in the proof of Theorem 2.

1. The case $\dim \Sigma = 0$

THEOREM 1A([I]). *Let X be a Fano 3-fold with $\dim \Sigma = 0$. Then there exist a normal surface S which is either an Abelian surface or a normal K3-surface and an ample invertible sheaf \mathcal{L} on S such that X is the contraction of the negative section of a projective bundle $\mathbf{P}(\mathcal{O}_S \oplus \mathcal{L})$. Here a normal K3-surface implies a normal projective surface with the trivial canonical sheaf and has only rational singularities.*

THEOREM 1B([I]). *Let X be a projective cone over a surface S which is either an*

Abelian or a normal K3-surface. Then X is a Fano 3-fold with $\Sigma = \{\text{the vertex}\}$.

2. Basic structure theorem of \mathbf{Q} -factorial terminal modifications for the case $\dim \Sigma = 1$

THEOREM 2. *Let X be a Fano 3-fold with Σ of pure dimension one. Let $g : Y \rightarrow X$ be a \mathbf{Q} -factorial terminal modification whose existence is proved by Mori ([M]). Denote $K_Y = g^*K_X - \Delta$. Then we have a sequence of projective morphisms: $Y = Y_0 \xrightarrow{\varphi_0} Y_1 \xrightarrow{\varphi_1} Y_2 \dots \rightarrow Y_r \xrightarrow{\varphi_r} Z$, where for each i , φ_i is the contraction of an extremal ray R_i on Y_i such that $R_i \cdot \Delta_i > 0$ (here, $\Delta_0 = \Delta$, and $\Delta_i = (\varphi_{i-1})_*\Delta_{i-1}$). For $i \leq r - 1$, φ_i is a birational contraction of a divisor isomorphic to $F_{a,0}$ ($a \geq 1$) to a non-singular point and φ_r is a fibration to a lower dimensional variety Z .*

DEFINITION 1. The variety Z above is called a basis of X . And each φ_i is called a Δ -extremal contraction. Of course a basis of X is not unique for X . It depends on the choice of a \mathbf{Q} -factorial terminal modification Y and also on the choice of extremal rays R_i 's.

From now on, we devote to study X which has a two dimensional basis Z . In this case, the last contraction $\varphi_r : Y_r \rightarrow Z$ satisfies the assumption of the following proposition. So we can see that it is a \mathbf{P}^1 -bundle over a non-singular surface Z .

PROPOSITION 1 (NAKAYAMA). *Let $\varphi : Y \rightarrow Z$ be a contraction of an extremal ray on a 3-fold Y with at worst \mathbf{Q} -factorial terminal singularities on it to a surface Z . Assume there exists an invertible sheaf on Y whose degree on a general fiber is 1. Then Z is non-singular and Y is a \mathbf{P}^1 -bundle over Z .*

THEOREM 3. *Let X be a Fano 3-fold with one dimensional Σ and a two dimensional basis. Then there exists a \mathbf{Q} -factorial terminal modification $g : Y \rightarrow X$ such that a Δ -extremal contraction $\varphi_0 : Y \rightarrow Z$ gives a \mathbf{P}^1 -bundle over a non-singular surface Z .*

This theorem is proved by applying the following lemma successively.

LEMMA. *Let X be as above and $Y = Y_0 \xrightarrow{\varphi_0} Y_1 \xrightarrow{\varphi_1} Y_2 \dots \rightarrow Y_r \xrightarrow{\varphi_r} Z$ be a sequence of Δ -extremal contractions of \mathbf{Q} -factorial terminal modification Y of X with 2-dimensional basis Z . If $r > 0$, then there is a flop Y'_i of Y_i for each i ($i \leq r - 1$) such that $g' : Y' = Y'_0 \rightarrow X$ is a \mathbf{Q} -factorial terminal modification of X and $Y' = Y'_0 \xrightarrow{\varphi'_0} Y'_1 \xrightarrow{\varphi'_1} Y'_2 \dots \rightarrow Y'_{r-1} \xrightarrow{\varphi'_{r-1}} Z'$ is a sequence of Δ' -extremal contractions with 2-dimensional basis Z' , where Δ' is a \mathbf{Q} -divisor such that $K_{Y'} = g'^*K_X - \Delta'$.*

3. Fano 3-folds which have \mathbf{P}^1 -bundles as \mathbf{Q} -factorial terminal modifications.

Let X be a Fano 3-fold with a 2-dimensional basis. Then, by Theorem 3, we can take a \mathbf{Q} -factorial terminal modification $g : Y \rightarrow X$ such that a Δ -extremal contraction

$\varphi : Y \rightarrow Z$ gives a \mathbf{P}^1 -bundle over a non-singular surface Z . Then we have the following facts:

- (i) $-g^*K_X \cdot \ell = \Delta \cdot \ell = 1$, where ℓ is a fiber of $\varphi : Y \rightarrow Z$.
- (ii) Δ is denoted by $E_0 + \varphi^*(\Delta')$, where E_0 is an irreducible component with $E_0 \cdot \ell = 1$ and $\Delta' \in \text{Pic}(Z)$.

The case $\text{Supp}\Delta$ contains a vertical component

We call an irreducible divisor D in Y a vertical divisor for g , if D is mapped to a point of X by g . We call a singular point an intensive singular point, if it is the image of a vertical component for some \mathbf{Q} -factorial terminal modification.

THEOREM 4A. *Let $X, g : Y \rightarrow X, \Delta$ and $\varphi : Y \rightarrow Z$ be as in the beginning of this section. Assume $\text{Supp}\Delta$ contains a vertical component.*

Then, (i) a vertical component is unique and coincides with E_0 and it is a section of the projection φ ,

(ii) there exists a normal surface Z_0 with at least one non-rational singular point on it whose canonical sheaf is trivial and whose minimal resolution is $h : Z \rightarrow Z_0$ and

(iii) the \mathbf{P}^1 -bundle $\varphi : Y \rightarrow Z$ is a pull back of a \mathbf{P}^1 -bundle $\varphi_0 : Y_0 \rightarrow Z_0$ by h and $g : Y \rightarrow X$ factors as $Y \xrightarrow{h} Y_0 \xrightarrow{g_0} X$, where g_0 is a contraction of the negative section $h(E_0)$.

THEOREM 4B. *Let S be a normal surface with trivial canonical sheaf and at least one non-rational singular point on it. Then an arbitrary projective cone X over S is a Fano 3-fold and Σ is generating lines over a non-rational singular points of S .*

Remark. Normal surfaces with the trivial canonical sheaf and at least one non-rational singular point are studied in [U] among others. The number of non-rational singular points is less than or equal to 2. It is 2, if and only if both of them are simple elliptic singularities [U, Theorem 1].

The case $\text{Supp}\Delta$ contains no vertical component

In the previous case, E_0 is a section of φ . But in this case, it is not necessarily true. First we consider the case that E_0 is a section. Since E_0 is not a vertical component, $g|_{E_0} : E_0 \rightarrow C$ is a fibration to a curve C .

PROPOSITION 2. *The possible triples $(E_0, g|_{E_0}, \Delta')$ are the following:*

- (i) $(\mathbf{P}^1 \times \text{elliptic curve, the first projection } p_1, \varphi)$,
- (ii) $(\text{a rational elliptic surface, the elliptic fibration, } \varphi)$,
- (iii) E_0 is the composite of r -blowing ups $E_0 \xrightarrow{\sigma_r} \dots \xrightarrow{\sigma_1} \mathbf{P}^1 \times \text{elliptic curve}$, where σ_1 is the blow up at a point on the fiber $C = p_1^{-1}(z)$ of a point $z \in \mathbf{P}^1$ and σ_i ($i > 1$) is the blow up at the intersection of the proper transform of C and the exceptional curve of σ_{i-1} . The morphism $g|_{E_0}$ is $p_1 \cdot \sigma_1 \cdot \sigma_2 \dots \sigma_r$ and Δ' = the proper transform of C .

(iv) E_0 is a ruled surface $p : E_0 \rightarrow S$ such that there exist a covering $\pi : S \rightarrow \mathbf{P}^1$ and a member D in $|-K_{E_0}|$ of type $D = (\pi \cdot p)^*(z) + \Delta'$, where $z \in \mathbf{P}^1$ and Δ' is an

effective divisor with $K_{E_0} \cdot C \geq 0$ for every component $C \subset \Delta'$. The morphism $g|_{E_0}$ is πp .

THEOREM 5A. *Let X be a Fano 3-fold with a non-rational singular point but no intensive point. Assume X has a \mathbf{P}^1 -bundle model $\varphi : Y \rightarrow Z$ and E_0 is a section of φ , where $g : Y \rightarrow X$ is the modification and $E_0 < \Delta = g^*K_X - K_Y$. Denote $\Delta = E_0 + \varphi^*\delta$ and $-g^*K_X = E_0 + \varphi^*L$ for divisors δ, L on Z .*

Then $\varphi : Y \rightarrow Z$ is defined by a locally free sheaf \mathcal{E} of rank 2 on Z and satisfies the following conditions:

- (I) *the triple $(E_0, g|_{E_0}, \bar{\delta})$ is as one of (i)~(iv) in Proposition 4,*
- (II) *$L - \delta$ is semi-ample and $(L - \delta) \cdot (L - \delta - K_Z) > 0$,*
- (III-1) *\mathcal{E} is an extension of $\mathcal{N} = \mathcal{O}_Z(-K_Z - \delta - L)$ by \mathcal{O}_Z :*

$$(\mathcal{E}) \quad 0 \rightarrow \mathcal{O}_Z \rightarrow \mathcal{E} \rightarrow \mathcal{N} \rightarrow 0$$

- (III-2) *$\mathcal{E}|_C = \mathcal{O}_C(-L) \oplus \mathcal{O}_C(-L)$ for every component $C < \delta$ and*
- (III-3) *$(\mathcal{O}(L) \otimes \mathcal{E})_y$ is generated by its global sections for each $y \in \delta$.*

THEOREM 5B. *Let Z be a non-singular projective surface, $g_0 : Z \rightarrow \mathbf{P}^1$ be a surjective morphism, L, δ ($\delta \geq 0$) be divisors on Z and \mathcal{E} be a locally free sheaf of rank 2 on Z such that*

- (I) *the triple (Z, g_0, δ) is as one of (i) ~ (iv) of Proposition 2,*
- (II) *$L - \delta$ is semi-ample and $(L - \delta) \cdot (L - \delta - K_Z) > 0$, and*
- (III-1) *\mathcal{E} is an extension of $\mathcal{N} = \mathcal{O}_Z(-K_Z - \delta - L)$ by \mathcal{O}_Z :*

$$(\mathcal{E}) \quad 0 \rightarrow \mathcal{O}_Z \rightarrow \mathcal{E} \rightarrow \mathcal{N} \rightarrow 0$$

- (III-2) *$\mathcal{E}|_C = \mathcal{O}_C(-L) \oplus \mathcal{O}_C(-L)$ for every component $C < \delta$ and*
- (III-3) *$(\mathcal{O}(L) \otimes \mathcal{E})_y$ is generated by its global sections for each $y \in \delta$.*

Let $\varphi : Y \rightarrow Z$ be a \mathbf{P}^1 -bundle defined by \mathcal{E} and E_0 be the section corresponding to the surjection $\mathcal{E} \rightarrow \mathcal{N}$ in (III-1).

*Then $E_0 + \varphi^*L$ is semi-ample and the image X of Y by $g := \Phi_{|m(E_0 + \varphi^*L)|}$ is a Fano 3-fold with $\Sigma \neq \emptyset$ and no intensive singular point and $g : Y \rightarrow X$ is a \mathbf{Q} -factorial terminal modification.*

Now we give an example of a Fano 3-fold with E_0 not a section.

Example. Let Z be the projective plane \mathbf{P}^2 , C and C' be two general curves of degree 3 on Z . Let $\sigma : \tilde{Z} \rightarrow Z$ be the blowing up at 9-distinct points $\{p_1, p_2, \dots, p_9\} = C \cap C'$, then \tilde{Z} becomes an elliptic surface with elliptic fibers $[C], [C']$, where $[C]$ is the proper transform of C on \tilde{Z} . Denote the fiber $\sigma^{-1}(p_i)$ by ℓ_i . Let L be $\sigma^*L_0 + \sum_{i=1}^9 \ell_i$ where L_0 is an ample divisor on Z .

Since $H^1(\tilde{Z}, L - [C]) \simeq \oplus H^1(\ell_i, L - [C]|_{\ell_i}) \simeq \mathbf{C}^{\oplus 9}$, we can take an extension sheaf $\tilde{\mathcal{E}}$ of $\mathcal{O}([C] - L)$ by $\mathcal{O}_{\tilde{Z}}$ such that the restriction $[\tilde{\mathcal{E}}]_{\ell_i} \in H^1(\ell_i, L - [C]|_{\ell_i})$ is not zero for every i ($i = 1, 2, \dots, 9$). Now $0 \rightarrow \mathcal{O}_{\tilde{Z}}|_{\ell_i} \rightarrow \tilde{\mathcal{E}}|_{\ell_i} \rightarrow \mathcal{O}([C] - L)|_{\ell_i} = \mathcal{O}_{\mathbf{P}^1}(2) \rightarrow 0$ does not split, so $\tilde{\mathcal{E}}|_{\ell_i} \simeq \mathcal{O}_{\mathbf{P}^1}(1) \oplus \mathcal{O}_{\mathbf{P}^1}(1)$. Put $\tilde{\mathcal{E}}' = \tilde{\mathcal{E}}(\Sigma \ell_i)$, then $\tilde{\mathcal{E}}'|_{\ell_i}$ is trivial for each i . By Schwarzenberger's Theorem, $\tilde{\mathcal{E}}' = \sigma^*\mathcal{E}$ for some locally free sheaf \mathcal{E} on Z . Let Y be the

projective bundle $\mathbf{P}(\mathcal{E})$ and \tilde{Y} be $\mathbf{P}(\tilde{\mathcal{E}}')$. Then we have the diagram of a fiber product


$$\begin{array}{ccc} \tilde{Y} & \xrightarrow{\sigma} & Y \\ \downarrow \tilde{\varphi} & \square & \varphi \downarrow \\ \tilde{Z} & \xrightarrow{\sigma} & Z \end{array}$$

Let \tilde{E}_0 be the section of $\tilde{\varphi}$ defined by the surjection $\tilde{\mathcal{E}} \rightarrow \mathcal{O}([C] - L)$ and E_0 be the image $\sigma(\tilde{E}_0)$. Then $H = E_0 + \varphi^* L_0$ is a semipositive divisor on Y . The image X of the morphism $\Phi_{|mH|} : Y \rightarrow \mathbf{P}^M$ becomes a Fano 3-fold with $\Sigma \simeq \mathbf{P}^1$ and Y is a \mathbf{Q} -factorial terminal modification of X . It is easy to see that $\Delta = E_0$ and E_0 contains the fibers of φ over $p_1, p_2, \dots, p_9 \in Z$

4. Pictures of Y and X of Theorem 5


(i) In the case the triple is $(\mathbf{P}^1 \times C, \text{the first projection } p_1, \phi)$, where C is an elliptic curve. Then $\Delta = E_0$. If we denote $L = p_1^* \mathcal{O}_{\mathbf{P}^1}(a) \otimes p_2^* B$, then $a \geq 0$ and B is ample.

(i-1) $a > 0$. $g|_{Y-E_0} : Y - E_0 \simeq X - \Sigma$.




(i-2) $a = 0$ and the exact sequence $(\mathcal{E}) : 0 \rightarrow \mathcal{O}_Z \rightarrow \mathcal{E} \rightarrow \mathcal{N} \rightarrow 0$ splits.

$g|_{Y-E_0-E_\infty} : Y - E_0 - E_\infty \simeq X - \Sigma - \Sigma_0$, and $g|_{E_\infty} = p_2$, where Σ_0 is the locus of canonical singularities.




(i-3) $a = 0$ and the exact sequence $(\mathcal{E}) : 0 \rightarrow \mathcal{O}_Z \rightarrow \mathcal{E} \rightarrow \mathcal{N} \rightarrow 0$ does not split. There exists a divisor $\sum_{i=1}^s m_i q_i \in |B|$ such that the restriction $(\mathcal{E})|_f$, splits for each $i, (i = 1, \dots, s)$, where $f_i = p_2^{-1}(q_i)$. For a general fiber $f = p_2^{-1}(q), q \in C, Y_f$ is $\mathbf{P}^1 \times \mathbf{P}^1$ and for $f_i, Y_{f_i} \simeq \mathbf{P}(\mathcal{O} \oplus \mathcal{O}(2))$. $E_0|_{Y_f}$ is an ample section for general f and is the disjoint section from the negative section for $f = f_i$. Denote the negative section of Y_{f_i} by \tilde{f}_i . Then the restriction $g|_{Y-E_0-\cup \tilde{f}_i}$ is an isomorphism, $g|_{E_0} = p_1$, and each \tilde{f}_i is contracted

to a canonical singularity in X .




(ii) The case that the triple $(E_0, g|_{E_0}, \Delta')$ is (rational elliptic surface, the elliptic fibration, φ). Then $E_0 = \Delta$ in this case too. If L is big then the exact sequence (\mathcal{E}) splits and if L is not big $|L|$ gives a fibration $\Phi = \Phi|_{|L|} : Z \rightarrow \mathbf{P}^1$ with a general fiber \mathbf{P}^1 . Let C_i ($i = 1, 2, \dots, r$) be (-2) -curves on Z with $LC_i = 0$ and f_j ($j = 1, \dots, s$) be (-1) -curves on Z with $Lf_j = 0$. Then $E_0|_{Y_f}$ is the section disjoint from the negative section. Denote the negative section of Y_f by \tilde{f}_j . Then the normal bundle of \tilde{f}_j in Y is $\mathcal{O}_{\mathbf{P}^1}(1) \oplus \mathcal{O}_{\mathbf{P}^1}(1)$.


(ii-1) L is big. Then the restriction $g|_{Y-E_0-\cup Y_{C_i}-\cup \tilde{f}_i}$ is an isomorphism, $g|_{Y_{C_i}} : Y_{C_i} \simeq C_i \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is the projection to the second factor and $g(\tilde{f}_j)$ is an isolated canonical singular point for each j . A point of $g(Y_{C_i})$ away from $g(E_0)$ is non-isolated canonical singularities.



(ii-2) L is not big and (\mathcal{E}) splits. Let E_∞ be the section of φ disjoint from E_0 . Then the restriction $g|_{E_0-E_\infty-\cup Y_{C_i}}$ is an isomorphism, $g|_{Y_{C_i}}$ is as above and $g|_{E_\infty} = \Phi$.




(ii-3) L is not big and (\mathcal{E}) does not split. Denote $L = \Phi^* L_0$ for a Cartier divisor L_0 on \mathbf{P}^1 . Then the extension \mathcal{E} of \mathcal{N} corresponds to a non-zero section $\varphi_{\mathcal{E}}$ of $\Gamma(\mathbf{P}^1, L_0 + K_{\mathbf{P}^1})$. Let $\varphi_{\mathcal{E}}$ define a divisor $\sum_{k=1}^d m_k q_k$ ($d \geq 0, m_k > 0$) and ℓ_k $k = 1, \dots, b$ ($0 \leq b \leq d$) be smooth fibers among $\{\Phi^{-1}(q_k)\}$. A component of a singular fiber of Φ is either one of C'_i 's or f'_i 's defined above. For a general fiber $\ell = \Phi^{-1}(q)$ $q \in \mathbf{P}^1$, Y_{ℓ} is $\mathbf{P}^1 \times \mathbf{P}^1$ and for ℓ_k , $Y_{\ell_k} \simeq \mathbf{P}(\mathcal{O} \oplus \mathcal{O}(2))$. $E_0|Y_{\ell}$ is an ample section for general ℓ , while it is the section disjoint from the negative section for $\ell = \ell_k$ ($0 \leq k \leq b$). Denote the negative section of Y_{ℓ_k} by $\tilde{\ell}_k$. Then the restriction $g|_{Y-E_0-\cup_{i=1}^r Y_{C_i}-\cup_{j=1}^s \tilde{f}_j-\cup_{k=1}^b \tilde{\ell}_k}$ is isomorphic, $g|_{Y_{C_i}}$ is the second projection, \tilde{f}'_j 's and $\tilde{\ell}'_k$'s are contracted to canonical singularities in X .




(iii) The case that the triple $(E_0, g|_{E_0}, \Delta')$ is as follows: E_0 is the composite of r -blowing ups $E_0 \xrightarrow{\sigma_r} \dots \xrightarrow{\sigma_1} \mathbf{P}^1 \times \text{elliptic curve}$, where σ_1 is the blow up at a point on the fiber $C = p_1^{-1}(z)$ of a point $z \in \mathbf{P}^1$ and σ_i ($i > 1$) is the blow up at the intersection of the proper transform of C and the exceptional curve of σ_{i-1} . The morphism $g|_{E_0}$ is $p_1 \sigma_1 \sigma_2 \dots \sigma_r$ and $\Delta' =$ the proper transform of C .

Then L is nef and big, with $L\ell_r > 0$ and the exact sequence (\mathcal{E}) splits, where ℓ_i ($i = 1, 2, \dots, r$) are the exceptional curves of σ_i respectively. Let E_{∞} be the section of φ disjoint from E_0 , and ℓ_j $j \in J \subset \{1, 2, \dots, r-1\}$ be the exceptional curves with $L\ell_j = 0$ and C_i $i = 1, 2, \dots, s$ be the curves on E_{∞} with $LC_i = 0$. Then $g|_{Y-E_0-Y_{\Delta'}-\cup_{j \in J} Y_{\ell_j}-\cup_{i=1}^n C_i}$ is an isomorphism, $g|_{Y_{\ell_j}} \simeq \ell_j \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is the projection to the second factor and $g(C_i)$ is an isolated canonical singular point on X for $i = 1, 2, \dots, s$.



(iv) The case that the triple is as in (iv) of Proposition 2. Then the exact sequence (\mathcal{E}) does not split. Let C_i ($i = 1, 2, \dots, r$) be (-2) -curves on Z with $eC_i = LC_i = 0$ and f_j ($j = 1, \dots, s$) be (-1) -curves on Z with $ef_j > 0$ and $Lf_j = 0$. Then we can take the negative section \tilde{f}_j of $Y_{\tilde{f}_j}$ disjoint from E_0 . Then $g|_{Y-\Delta-\cup_{i=1}^r Y_{C_i}-\cup_{j=1}^s \tilde{f}_j}$ is an isomorphism, $g|_{Y_C} : Y_C \simeq C \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is the projection to the second factor for a component $C < \Delta$ and for $C = C_i$ ($i = 1, \dots, r$) and $g(\tilde{f}_j)$ is an isolated canonical singularity for $j = 1, \dots, s$.



, where is the fiber of a point in Σ

REFERENCES

- [I] S. Ishii: *Quasi-Gorenstein Fano-3-folds with isolated non-rational loci*. *Compositio Math.* 77 (1991) 335-341.
- [I2] S. Ishii: *On Fano 3-folds with non-rational singularities and two dimensional base*. to appear in *Abh. Math. Sem. Univ. Hamburg*.
- [M] S. Mori: *Flip theorem and the existence of minimal models for 3-folds*. *J. Amer. Math. Soc.* 1 (1988) 117-253.
- [U] Y. Umezumi: *On normal projective surfaces with trivial dualizing sheaf*. *Tokyo J. Math.* 4 (1981) 343-354.

DEPARTMENT OF MATHEMATICS
 TOKYO INSTITUTE OF TECHNOLOGY
 OH-OKAYAMA, MEGURO
 152 TOKYO, JAPAN
 e-mail: shihoko@math.titech.ac.jp