

# Local time and Tanaka formula for a Volterra-type multifractional Gaussian process

BRAHIM BOUFOUSSI<sup>1</sup>, MARCO DOZZI<sup>2,\*</sup> and RENAUD MARTY<sup>2,\*\*</sup>

<sup>1</sup>*Faculté des Sciences Semlalia, Department of Mathematics, Cadi Ayyad University, BP 2390 Marrakech, Morocco. E-mail: boufoussi@ucam.ac.ma*

<sup>2</sup>*Institut Elie Cartan de Nancy, Nancy-Université, CNRS, INRIA, B.P. 239, F-54506 Vandoeuvre-lès-Nancy Cedex, France. E-mail: \*marco.dozzi@iecn.u-nancy.fr; \*\*renaud.marty@iecn.u-nancy.fr*

The stochastic calculus for Gaussian processes is applied to obtain a Tanaka formula for a Volterra-type multifractional Gaussian process. The existence and regularity properties of the local time of this process are obtained by means of Berman's Fourier analytic approach.

*Keywords:* Gaussian processes; local nondeterminism; local time; multifractional processes; Tanaka formula

## 1. Introduction

Several types of multifractional Gaussian processes have been studied, including the processes introduced by Lévy-Véhel and Peltier [12] and Benassi, Jaffard and Roux [6], known as the moving average and harmonizable versions of multifractional Brownian motion, and the processes introduced by Benassi, Cohen and Istas [2] and by Surgailis [16]. These processes are usually defined by replacing, in certain representations of fractional Brownian motion (fBm), the Hurst parameter  $H$  by a Hurst function  $h$ , that is, a real function of the time parameter with values in  $(0,1)$ . Generalizations to stable processes have also been studied [10,14,15]. The stochastic properties and the regularity of the trajectories of these processes can be characterized in the same terms as for fBm, in particular, by the property of local asymptotic self-similarity or by the pointwise Hölder exponent, which is constant and equal to  $H$  for fBm, but which varies in time following  $h$  in the multifractional case.

The aim of this article is to study a Volterra-type multifractional Gaussian process  $B_h$  which fits into the framework of stochastic calculus, and its local time. In fact, the stochastic calculus, including the Itô formula and stochastic differential equations, is now well established for fBm and stochastic integrals have been defined in the Malliavin sense or by means of Wick products. The stochastic calculus for multifractional Gaussian processes has not yet been developed explicitly, but important elements, including a divergence integral and an Itô formula, have been proven in [1] for a class of Gaussian processes admitting a kernel representation with respect to Brownian motion. The process  $B_h$  we study in this article is defined by allowing for a Hurst function in the kernel representation of fBm and it belongs to the class of Gaussian processes studied in [1].

The Itô formula in [1] is applied to obtain a representation of the local time for which, as is usually the case in Tanaka-like formulas, the occupation measure is not the Lebesgue measure, but the quadratic variation process. However, contrary to fBm, the interpretation in our case is more delicate because the quadratic variation process is not necessarily increasing (but of bounded variation). We compare this local time to the local time with respect to the Lebesgue measure, which we obtain, together with the regularity properties of its trajectories, by the Fourier analytic approach initiated by Berman [3] based on the notion of local non-determinism (LND).

It is well known that the cases where the Hurst parameter or the Hurst function is  $<1/2$  (resp.,  $>1/2$ ) have to be treated separately (see [1] and [11]). We are interested in the case where the Hurst function  $h$  takes values in  $(1/2, 1)$ . In fact, the Volterra-type process we study here is defined only for this case, which is appropriate for long memory applications [13]. The article is organized as follows. In Section 2, stochastic properties and the regularity of the trajectories of  $B_h$  are proved. The form of the covariance function (Proposition 2) shows, in particular, that  $B_h$  differs from the harmonizable and moving average multifractional Brownian motions. The continuity of the trajectories of  $B_h$  is obtained by classical criteria from estimates for the second order moments of the increments (Proposition 3). The lass property is proved in Proposition 5; it implies that the pointwise Hölder exponent of  $B_h$  is equal to  $h$  (Proposition 6). In Section 3, it is shown that  $B_h$  satisfies the property labeled (K4) in [1] and that the quadratic variation of  $B_h$  is of bounded variation. Therefore, the divergence integral and the Itô formula developed in [1] (recalled briefly in this section) hold for  $B_h$ . Section 4 is devoted to the local time of  $B_h$ . Its existence and square integrability with respect to the space variable follow from a classical criterion due to Berman [3] (Proposition 11). A Tanaka-type formula with the divergence integral of Section 3 is then given (Theorem 12), where the occupation measure of the local time is the quadratic variation of  $B_h$ . Finally, the local time with respect to the Lebesgue measure and its regularity in the space and time variables are given in Theorem 16. The proofs are based on the LND property, which is shown to hold for  $B_h$  in Proposition 14.

## 2. A Volterra-type multifractional Gaussian process

It is well known that the fractional Brownian motion  $B_H$  with (fixed) Hurst parameter  $H \in (1/2, 1)$  can be represented for any  $t \geq 0$  as

$$B_H(t) = \int_0^t K_H(t, u) W(du),$$

where

$$K_H(t, u) = u^{1/2-H} \int_u^t (y-u)^{H-3/2} y^{H-1/2} dy \tag{1}$$

and  $W(dy)$  is a Gaussian measure. Let  $a$  and  $b$  be two real numbers satisfying  $1/2 < a < b < 1$ . Throughout the paper, we consider a function  $h: \mathbb{R} \rightarrow [a, b]$ . We assume that this function is  $\beta$ -Hölder with  $\sup h < \beta$ . Define the centered Gaussian process  $B_h = \{B_h(t), t \geq 0\}$  by

$$B_h(t) = B_{h(t)}(t) = \int_0^t K_{h(t)}(t, u) W(du),$$

where

$$K_{h(t)}(t, u) = u^{1/2-h(t)} \int_u^t (y-u)^{h(t)-3/2} y^{h(t)-1/2} dy. \tag{2}$$

Before establishing properties of  $B_h$ , we state a lemma regarding an estimate on  $K_H$  that we use throughout the paper.

**Lemma 1.** *For every  $T > 0$ , there exists a function  $\Phi_T \in L^2((0, T], \mathbb{R}_+)$  such that for every  $s \in (0, T]$ ,*

$$\sup_{\lambda \in [a, b], t \in (0, T]} \left| \frac{\partial}{\partial \lambda} K_\lambda(t, s) \right| \leq \Phi_T(s).$$

**Proof.** We have

$$\begin{aligned} \frac{\partial}{\partial \lambda} K_\lambda(t, s) &= (-\log s) s^{1/2-\lambda} \int_s^t (y-s)^{\lambda-3/2} y^{\lambda-1/2} dy \\ &\quad + s^{1/2-\lambda} \int_s^t (y-s)^{\lambda-3/2} y^{\lambda-1/2} (\log(y-s) + \log y) dy. \end{aligned}$$

Then, for every  $T > 0$ , there exists a constant  $C_{a,b,T}$  such that for every  $\lambda \in [a, b]$  and  $s \in (0, T]$ ,

$$\left| \frac{\partial}{\partial \lambda} K_\lambda(t, s) \right| \leq C_{a,b,T} (1 \vee |\log s|) s^{1/2-b} =: \Phi_T(s),$$

which concludes the proof. □

The following proposition gives the covariance of this process.

**Proposition 2.** *Let  $X = \{X(t, \lambda), t \geq 0, \lambda \in (1/2, 1)\}$  be the two-parameter process given by  $X(t, \lambda) = \int_0^t K_\lambda(t, u) W(du)$ . Then*

$$\mathbb{E}[X(t, \lambda) X(s, \lambda')] = \int_0^t dy \int_0^s dz \tilde{\beta}(y, z, \lambda, \lambda') |y-z|^{\lambda+\lambda'-2} \left(\frac{y}{z}\right)^{\lambda-\lambda'},$$

where

$$\tilde{\beta}(y, z, \lambda, \lambda') = \beta(2-\lambda-\lambda', \lambda'-1/2) 1_{\{y>z\}} + \beta(2-\lambda-\lambda', \lambda-1/2) 1_{\{y<z\}}$$

and  $\beta(a, b)$  ( $a, b > 0$ ) is the beta function. In particular,

$$\mathbb{E}[X(t, \lambda)^2] = \int_0^t dy \int_0^t dz \beta(2-2\lambda, \lambda-1/2) |y-z|^{2\lambda-2} = \frac{t^{2\lambda}}{c_\lambda^2}$$

with

$$c_\lambda^2 = \frac{2\pi\lambda(\lambda-1/2)^3}{\Gamma(2-2\lambda)\Gamma(\lambda+1/2)^2 \sin(\pi(\lambda-1/2))}. \tag{3}$$

**Proof.** We have

$$\begin{aligned} &\mathbb{E}[X(t, \lambda)X(s, \lambda')] \\ &= \int_0^{t \wedge s} K_\lambda(t, u)K_{\lambda'}(s, u) \, du \\ &= \int_0^{t \wedge s} u^{1-\lambda-\lambda'} \left( \int_u^t (y-u)^{\lambda-3/2} y^{\lambda-1/2} \, dy \right) \left( \int_u^s (z-u)^{\lambda'-3/2} z^{\lambda'-1/2} \, dz \right) \, du \\ &= \int_0^t dy \int_0^s dz y^{\lambda-1/2} z^{\lambda'-1/2} \int_0^{y \wedge z} u^{1-\lambda-\lambda'} (y-u)^{\lambda-3/2} (z-u)^{\lambda'-3/2} \, du. \end{aligned}$$

We fix  $y > z$  and calculate the following integral by making the successive substitutions  $u = vz$ ,  $w = (y - vz)/(1 - v)$ ,  $t = w/y$  and  $s = 1/t$ :

$$\begin{aligned} &\int_0^z u^{1-\lambda-\lambda'} (y-u)^{\lambda-3/2} (z-u)^{\lambda'-3/2} \, du \\ &= z^{1/2-\lambda} \int_0^1 \lambda' (y-vz)^{\lambda-3/2} (1-v)^{\lambda'-3/2} \, dv \\ &= z^{1/2-\lambda} (y-z)^{\lambda+\lambda'-2} \int_y^{+\infty} (w-y)^{1-\lambda-\lambda'} w^{\lambda-3/2} \, dw \\ &= z^{1/2-\lambda} y^{1/2-\lambda'} (y-z)^{\lambda+\lambda'-2} \int_1^{+\infty} (t-1)^{1-\lambda-\lambda'} t^{\lambda-3/2} \, dt \\ &= z^{1/2-\lambda} y^{1/2-\lambda'} (y-z)^{\lambda+\lambda'-2} \beta(2-\lambda-\lambda', \lambda'-1/2). \end{aligned}$$

In the same way, for  $y < z$ , we obtain

$$\begin{aligned} &\int_0^y u^{1-\lambda-\lambda'} (y-u)^{\lambda-3/2} (z-u)^{\lambda'-3/2} \, du \\ &= z^{1/2-\lambda} y^{1/2-\lambda'} (z-y)^{\lambda+\lambda'-2} \beta(2-\lambda-\lambda', \lambda-1/2). \end{aligned}$$

This concludes the proof. □

In the sequel, we need the estimates which we establish in the following proposition.

**Proposition 3.** *The process  $X$  satisfies the following estimates:*

(a) for all  $s$  and  $t \geq 0$

$$\mathbb{E}[(X(t, \lambda) - X(s, \lambda))^2] = c_\lambda^{-2} |t - s|^{2\lambda}; \tag{4}$$

(b) for every  $T > 0$ , there exists a constant  $C_T > 0$  such that for every  $t \in [0, T]$  and every  $\lambda$  and  $\lambda' \in [a, b]$ ,

$$\mathbb{E}[(X(t, \lambda) - X(t, \lambda'))^2] \leq C_T |\lambda - \lambda'|^2. \tag{5}$$

**Proof.** (a) For every  $\lambda \in [a, b]$ , the process  $X(\cdot, \lambda) : t \mapsto X(t, \lambda)$  is a fractional Brownian motion with variance  $c_\lambda^{-2}$ , so we deduce (4).

(b) We have

$$\mathbb{E}[(X(t, \lambda) - X(t, \lambda'))^2] = \int_0^t (K_\lambda(t, u) - K_{\lambda'}(t, u))^2 du.$$

There exists  $\xi = \xi(\lambda, \lambda') \in [\min\{\lambda, \lambda'\}, \max\{\lambda, \lambda'\}]$  such that

$$K_\lambda(t, u) - K_{\lambda'}(t, u) = (\lambda - \lambda') \left. \frac{\partial}{\partial \lambda} K_\lambda(t, u) \right|_{\lambda=\xi(\lambda, \lambda')}.$$

Then, thanks to Lemma 1, for every  $t, \lambda$  and  $\lambda'$ , we obtain

$$\mathbb{E}[(X(t, \lambda) - X(t, \lambda'))^2] \leq |\lambda - \lambda'|^2 \int_0^T (\Phi_T(u))^2 du,$$

which concludes the proof. □

From the last proposition, we can deduce the continuity of  $B_h$ .

**Corollary 4.** *The process  $B_h$  defined above has continuous trajectories.*

**Proof.** We deduce from Proposition 3 and the assumption  $\sup h < \beta$  that for every  $s$  and  $t$  in a compact interval  $[0, T]$  such that  $|t - s| < 1$ , we have

$$\begin{aligned} \mathbb{E}[(B_h(t) - B_h(s))^2] &= \mathbb{E}[(X(t, h(t)) - X(s, h(s)))^2] \\ &\leq 2\mathbb{E}[(X(t, h(t)) - X(t, h(s)))^2] \\ &\quad + 2\mathbb{E}[(X(t, h(s)) - X(s, h(s)))^2] \\ &\leq 2C_T |h(t) - h(s)|^2 + 2c_{h(s)}^{-2} |t - s|^{2h(s)} \\ &\leq 2C_T |t - s|^{2\beta} + 2 \sup_{\lambda \in [a, b]} (c_\lambda^{-2}) |t - s|^{2\alpha}. \end{aligned}$$

Then  $\mathbb{E}[(B_h(t) - B_h(s))^2]/|t - s|^{2\alpha}$  is bounded and since  $B_h$  is Gaussian, we can deduce from [5] its continuity. □

We now deal with the local self-similarity property of  $B_h$ .

**Proposition 5.** *The process  $B_h$  is locally self-similar. More precisely, for every  $t$ , we have the following convergence in distribution:*

$$\lim_{\varepsilon \rightarrow 0} \left( \frac{B_h(t + \varepsilon u) - B_h(t)}{\varepsilon^{h(t)}} \right)_{u \geq 0} = (c_{h(t)}^{-1} B_{h(t)}(u))_{u \geq 0},$$

where  $\lim_{\varepsilon \rightarrow 0}$  stands for the limit in distribution in the space of continuous functions endowed with the uniform norm on every compact set.

**Proof.** Let us start by proving the convergence of the finite-dimensional distribution. Because  $B_h$  is Gaussian, it suffices to show the convergence of the second order moments. We then can write, for every  $u$  and  $v$ ,

$$\begin{aligned} & \frac{1}{\varepsilon^{2h(t)}} \mathbb{E}[(B_h(t + \varepsilon u) - B_h(t))(B_h(t + \varepsilon v) - B_h(t))] \\ &= \frac{1}{\varepsilon^{2h(t)}} (I_1(\varepsilon) + I_2(\varepsilon) + I_3(\varepsilon) + I_4(\varepsilon)), \end{aligned}$$

where

$$\begin{aligned} I_1(\varepsilon) &= \mathbb{E}[(X(t + \varepsilon u, h(t)) - X(t, h(t)))(X(t + \varepsilon v, h(t)) - X(t, h(t)))], \\ I_2(\varepsilon) &= \mathbb{E}[(X(t + \varepsilon u, h(t + \varepsilon u)) - X(t + \varepsilon u, h(t)))(X(t + \varepsilon v, h(t)) - X(t, h(t)))], \\ I_3(\varepsilon) &= \mathbb{E}[(X(t + \varepsilon u, h(t)) - X(t, h(t)))(X(t + \varepsilon v, h(t + \varepsilon v)) - X(t + \varepsilon v, h(t)))], \\ I_4(\varepsilon) &= \mathbb{E}[(X(t + \varepsilon u, h(t + \varepsilon u)) - X(t + \varepsilon u, h(t)))(X(t + \varepsilon v, h(t + \varepsilon v)) - X(t + \varepsilon v, h(t)))]. \end{aligned}$$

By the self-similarity of the fractional Brownian motion and the stationarity of its increments, we obtain

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^{2h(t)}} I_1(\varepsilon) = \frac{1}{2c_{h(t)}^2} (|u|^{2h(t)} + |v|^{2h(t)} - |u - v|^{2h(t)}). \quad (6)$$

Then, because of the Cauchy–Schwarz inequality, it is enough to prove

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^{2h(t)}} \mathbb{E}[(X(t + \varepsilon u, h(t + \varepsilon u)) - X(t + \varepsilon u, h(t)))^2] = 0 \quad (7)$$

to get

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^{2h(t)}} (I_2(\varepsilon) + I_3(\varepsilon) + I_4(\varepsilon)) = 0. \quad (8)$$

Thus, using Lemma 1, we obtain

$$\begin{aligned} & \frac{1}{\varepsilon^{2h(t)}} \mathbb{E}[(X(t + \varepsilon u, h(t + \varepsilon u)) - X(t + \varepsilon u, h(t)))^2] \\ &= \frac{1}{\varepsilon^{2h(t)}} \int_0^{t+\varepsilon u} (K_{h(t+\varepsilon u)}(t + \varepsilon u, s) - K_{h(t)}(t + \varepsilon u, s))^2 ds \\ &\leq \frac{(h(t + \varepsilon u) - h(t))^2}{\varepsilon^{2h(t)}} \int_0^{t+\varepsilon u} \sup_H \left| \frac{\partial}{\partial H} K_H(t + \varepsilon u, s) \right|^2 ds \\ &\leq C \varepsilon^{2\beta - 2h(t)} \int_0^{t+u} |\Phi_{t+u}(s)|^2 ds \xrightarrow{\varepsilon \rightarrow 0} 0. \end{aligned}$$

It now remains to prove tightness in the space of continuous functions endowed with the uniform norm. We also consider  $T > 0$  such that  $t, t + \varepsilon u$  and  $t + \varepsilon v \in [0, T]$  for all  $\varepsilon$ . Performing similar calculations as in the proof of Corollary 4, we get that there exist  $C_T > 0$  such that

$$\begin{aligned} & \mathbb{E} \left[ \left( \frac{B_h(t + \varepsilon u) - B_h(t)}{\varepsilon^{h(t)}} - \frac{B_h(t + \varepsilon v) - B_h(t)}{\varepsilon^{h(t)}} \right)^2 \right] \\ &= \frac{1}{\varepsilon^{2h(t)}} \mathbb{E} \left[ (B_h(t + \varepsilon u) - B_h(t + \varepsilon v))^2 \right] \\ &\leq \frac{C_T}{\varepsilon^{2h(t)}} |\varepsilon u - \varepsilon v|^{2h(t+\varepsilon v)} \\ &= C_T \varepsilon^{2(h(t+\varepsilon v)-h(t))} |u - v|^{2h(t+\varepsilon v)}. \end{aligned}$$

Since  $h$  is  $\beta$ -Hölder, it follows that  $\varepsilon^{2(h(t+\varepsilon v)-h(t))}$  is uniformly bounded. Moreover,  $h(t + \varepsilon v) \geq a$ , thus

$$\mathbb{E} \left[ \left( \frac{B_h(t + \varepsilon u) - B_h(t)}{\varepsilon^{h(t)}} - \frac{B_h(t + \varepsilon v) - B_h(t)}{\varepsilon^{h(t)}} \right)^2 \right] \leq C_T |u - v|^{2a}.$$

This completes the proof. □

It is classical to deduce pointwise Hölder continuity from local self-similarity [2]. We recall that the pointwise Hölder continuity of a function  $f$  is characterized by the pointwise Hölder exponent  $\alpha_f(t_0)$  defined at each point  $t_0$  as

$$\alpha_f(t_0) = \sup \left\{ \alpha > 0 : \lim_{t \rightarrow t_0} \frac{|f(t) - f(t_0)|}{|t - t_0|^\alpha} = 0 \right\}.$$

**Proposition 6.** *For every  $t_0 \in \mathbb{R}_+$ , the pointwise Hölder exponent  $\alpha_{B_h}(t_0)$  of  $B_h$  is almost surely equal to  $h(t_0)$ .*

**Proof.** We deduce from Proposition 5 and [2] that  $\alpha_{B_h}(t_0) \leq h(t_0)$ . We now prove that  $\alpha_{B_h}(t_0) \geq h(t_0)$ . By Proposition 3, there exists  $C > 0$  such that for every  $\varepsilon > 0$  and every  $s$  and  $t \in [t_0 - \varepsilon, t_0 + \varepsilon]$  such that  $|t - s| < 1$ , we have

$$\mathbb{E} \left[ (B_h(t) - B_h(s))^2 \right] \leq C |t - s|^{2 \inf_{[t_0 - \varepsilon, t_0 + \varepsilon]} h}.$$

By using the fact that  $B_h$  is Gaussian and applying Kolmogorov’s theorem [5], we get that  $\lim_{t \rightarrow t_0} |B_h(t) - B_h(t_0)|/|t - t_0|^\alpha = 0$  for every  $\alpha < \inf_{[t_0 - \varepsilon, t_0 + \varepsilon]} h$ . This holds for every  $\varepsilon > 0$ , so by continuity of  $h$ , we have  $\lim_{t \rightarrow t_0} |B_h(t) - B_h(t_0)|/|t - t_0|^\alpha = 0$  for every  $\alpha < h(t_0)$ . We can deduce that  $\alpha_{B_h}(t_0) \geq h(t_0)$  and hence  $\alpha_{B_h}(t_0) = h(t_0)$ . □

### 3. Stochastic calculus for $B_h$

The aim of this section is to apply the stochastic calculus developed by Alòs, Mazet and Nualart in [1] to get a stochastic integral for  $B_h$  and an Itô formula. We recall that in [1], the following hypothesis, called (K4), appears for regular kernels:

- **Hypothesis (K4).** For all  $s \in [0, T]$ ,  $K(\cdot, s)$  has bounded variation on the interval  $(s, T]$  and  $\int_0^T |K|((s, T], s)^2 ds < \infty$  (where, for all  $s \in [0, T]$ ,  $|K|((s, T], s)$  denotes the total variation of  $K(\cdot, s)$  on  $(s, T]$ ).

**Lemma 7.** *Suppose that  $h$  is of bounded variation on  $(s, T]$  for all  $s \in [0, T]$ . Then (K4) holds for  $(t, s) \mapsto K_{h(t)}(t, s)$  defined by (2).*

**Proof.** Let

$$\text{Var}_{(s, T]}^n(\cdot, s) = \sup_{t_0=s < t_1 < \dots < t_n=T} \sum_{i=1}^n |K_{h(t_i)}(t_i, s) - K_{h(t_{i-1})}(t_{i-1}, s)|$$

and suppose, without restriction of generality, that  $T = 1$ . Then

$$|K_{h(t_i)}(t_i, s) - K_{h(t_{i-1})}(t_{i-1}, s)| \leq I_1(i) + I_2(i),$$

where

$$I_1(i) = |K_{h(t_i)}(t_i, s) - K_{h(t_i)}(t_{i-1}, s)|$$

and

$$I_2(i) = |K_{h(t_i)}(t_{i-1}, s) - K_{h(t_{i-1})}(t_{i-1}, s)|.$$

We have

$$\begin{aligned} I_1(i) &= s^{1/2-h(t_i)} \int_{t_{i-1}}^{t_i} (y-s)^{h(t_i)-3/2} y^{h(t_i)-1/2} dy \\ &\leq s^{1/2-b} \int_{t_{i-1}}^{t_i} (y-s)^{a-3/2} y^{a-1/2} dy. \end{aligned}$$

Therefore,

$$\begin{aligned} &\sum_{i=1}^n |K_{h(t_i)}(t_i, s) - K_{h(t_i)}(t_{i-1}, s)| \\ &\leq s^{1/2-b} \int_s^1 (y-s)^{a-3/2} y^{a-1/2} dy \\ &\leq C(a) s^{1/2-b}. \end{aligned} \tag{9}$$



Regarding  $I_2$ , we have

$$\begin{aligned} & \sum_{i=1}^n |K_{h(t_i)}(t_{i-1}, s) - K_{h(t_{i-1})}(t_{i-1}, s)| \\ & \leq \sum_{i=1}^n |h(t_i) - h(t_{i-1})| \sup_{s \leq t \leq 1, a \leq \lambda \leq b} \left| \frac{\partial}{\partial \lambda} K_\lambda(t, s) \right| \\ & \leq \operatorname{Var}_{[0, T]} h \sup_{s \leq t \leq 1, a \leq \lambda \leq b} \left| \frac{\partial}{\partial \lambda} K_\lambda(t, s) \right|. \end{aligned}$$

The proof of Lemma 1 implies that  $\operatorname{Var}_{(s, T]}^n(\cdot, s) \leq C(1 \vee |\log s|)s^{1/2-b}$ , where the constant  $C > 0$  depends on  $h$  (but not on  $n$ ). This implies that (K4) holds for  $K_h$ .  $\square$

**Remark 8.** Since  $h$  is also supposed  $\beta$ -Hölder continuous for some  $\beta \leq 1$ , we will hereafter suppose that  $h$  is Lipschitz continuous. Note that  $\lim_{t \searrow u} K_{h(t)}(t, u) = 0$  for all  $u > 0$ .

For simplicity, we write  $K$  instead of  $K_h$ , but keep in mind that  $(t, s) \rightarrow K(t, s)$  means  $(t, s) \rightarrow K_{h(t)}(t, s)$  and that differentials with respect to  $t$  also act via  $h$ .

In the sequel, we need the following proposition regarding the variance of  $B_h$ .

**Proposition 9.** *The variance  $s \mapsto R_s = \mathbb{E}[B_h(s)^2]$  is of bounded variation on  $(0, T]$ .*

**Proof.** We have

$$\begin{aligned} \sum_{i=1}^n |R_{s_{i+1}} - R_{s_i}| &= \sum_{i=1}^n \left| \int_0^{s_{i+1}} K(s_{i+1}, r)^2 \, dr - \int_0^{s_i} K(s_i, r)^2 \, dr \right| \\ &\leq \sum_{i=1}^n \int_0^{s_{i+1}} K(s_{i+1}, r)^2 \, dr + \sum_{i=1}^n \left| \int_0^{s_i} [K(s_{i+1}, r)^2 - K(s_i, r)^2] \, dr \right|. \end{aligned}$$

The functions  $|K(s, r)1_{[0, s]}(r)|$  are bounded by the square-integrable function  $k(r) = |K(T, r)| + |K((r, T], r)|$ . Hence, the first term above is upper bounded by  $\int_0^T k(r)^2 \, dr$ . The second term is upper bounded by

$$\begin{aligned} & \sum_{i=1}^n \int_0^{s_i} |K((s_i, s_{i+1}], r)|K(s_{i+1}, r) + K(s_i, r)| \, dr \\ & \leq 2 \int_0^T k(r)|K((r, T], r)| \, dr < \infty. \end{aligned}$$

This concludes the proof.  $\square$

For any  $f \in L^2([0, T])$ , let  $Kf$  be defined by  $(Kf)(t) = \int_0^t K(t, s)f(s) \, ds$ . Let  $\mathcal{E}$  be the set of step functions on  $[0, T]$  and let the operator  $K^*$  be defined on  $\mathcal{E}$  by  $(K^*\varphi)(s) =$

$\int_s^T \varphi(t)K(dt, s)$ . Then  $K^*$  is the adjoint of  $K$ . In fact, for  $a_i \in \mathbb{R}$ ,  $0 = s_1 < s_2 < \dots < s_{n+1} = T$ ,  $\varphi = \sum_{i=1}^n a_i 1_{(s_i, s_{i+1}]}(s)$  and  $f \in L^2([0, T])$ , we write

$$(K^*\varphi)(s) = \sum_{i=1}^n 1_{(s_i, s_{i+1}]}(s) \sum_{j=i}^n a_j (K(s_{j+1}, s) - K(s_j, s))$$

and

$$\begin{aligned} \int_0^T (K^*\varphi)(s)f(s) ds &= \sum_{j=1}^n a_j \int_0^T \sum_{i=1}^j 1_{(s_i, s_{i+1}]}(s) (K(s_{j+1}, s) - K(s_j, s))f(s) ds \\ &= \sum_{j=1}^n a_j [(Kf)(s_{j+1}) - (Kf)(s_j)] = \int_0^T \varphi(t)(Kf)(dt). \end{aligned} \tag{10}$$

As usual, the reproducing kernel Hilbert space (RKHS)  $\mathcal{H}$  is defined as the closure of the linear span of the indicator functions  $\{1_{[0,t]}, t \in [0, T]\}$  with respect to the scalar product  $\langle 1_{[0,t]}, 1_{[0,s]} \rangle_{\mathcal{H}} = \mathbb{E}[B_{h(t)}(t)B_{h(s)}(s)] \equiv R(t, s)$ . By replacing  $f(s) ds$  in (10) by  $W(ds)$ , we have, by (1),

$$B_h(\varphi) \equiv \int_0^T \varphi(t)B_h(dt) = \int_0^T (K^*\varphi)(s)W(ds).$$

Therefore,

$$\|\varphi\|_{\mathcal{H}}^2 = \mathbb{E}[B_h(\varphi)^2] = \|K^*\varphi\|_{L^2([0,T])}^2 \leq \int_0^T \left[ \int_0^T |\varphi(t)||K|(dt, s) \right]^2 ds =: \|\varphi\|_K^2.$$

Let us denote by  $\mathcal{H}_K$  the completion of  $\mathcal{E}$  with respect to the  $\|\cdot\|_K$ -norm. Then  $\mathcal{H}_K$  is continuously embedded in  $\mathcal{H}$ .

In order to define the stochastic integral with respect to  $B_h$ , let us denote by  $\mathcal{S}$  the set of smooth cylindrical random variables of the form  $F = f(B_h(\varphi_1), B_h(\varphi_2), \dots, B_h(\varphi_n))$ , where  $n \geq 1$ ,  $f \in C_b^\infty(\mathbb{R}^n)$  ( $f$  and all its derivatives are bounded) and  $\varphi_1, \varphi_2, \dots, \varphi_n \in \mathcal{H}$ . Let us also denote by  $\mathbb{D}^{1,2}(\mathcal{H}_K)$  the closure of  $\{F \in \mathcal{S} : F \in L^2(\Omega, \mathcal{H}_K), DF \in L^2(\Omega \times \mathcal{H}_K, \mathcal{H}_K)\}$ . Then  $\mathbb{D}^{1,2}(\mathcal{H}_K)$  is included in the domain  $\text{Dom}(\delta^{B_h})$  of the divergence operator of  $B_h$ , and the integral of  $u \in \text{Dom}(\delta^{B_h})$  with respect to  $B_h$  is given by

$$\delta^{B_h}(u) \equiv \int_0^T u \delta B_h = \int_0^T (K^*u) \delta W \equiv \int_0^T \left[ \int_s^T u(r)K(dr, s) \right] \delta W(s),$$

where the last two integrals are the divergence integrals with respect to Brownian motion. Let us recall that the integral  $\delta^{B_h}(u)$  is defined, for any  $u \in L^2(\Omega, \mathcal{H})$ , as the unique element in  $L^2(\Omega)$  which satisfies the duality relationship  $\mathbb{E}(\delta^{B_h}(u)F) = \mathbb{E}\langle DF, u \rangle_{\mathcal{H}}$  for all  $F \in \mathcal{S}$ .

The following Itô formula, due to Alòs, Mazet and Nualart [1], will be applied in the next section. Let  $F$  be a function belonging to the class  $C^2(\mathbb{R})$  and satisfying the condition

$$\max\{|F(x)|, |F'(x)|, |F''(x)|\} \leq ce^{\lambda|x|^2},$$

where  $c$  and  $\lambda$  are positive constants such that  $\lambda < \frac{1}{4}(\sup_{0 \leq t \leq T} R_t)^{-1}$ . This implies that the process  $F'(B_h)$  belongs to  $\mathbb{D}^{1,2}(\mathcal{H}_K)$ . The integral  $\int_0^t F'(B_h(s)) \delta B_h(s)$  is therefore well defined for all  $t \in [0, T]$  and the following Itô-type formula holds ([1], Theorem 2):

$$F(B_h(t)) = F(0) + \int_0^t F'(B_h(s)) \delta B_h(s) + \frac{1}{2} \int_0^t F''(B_h(s)) dR(s). \tag{11}$$

### 4. Local time and Tanaka formula for $B_h$

First, we prove, by means of a criterion for Gaussian processes due to Berman, that  $B_h$  has a local time with respect to the Lebesgue measure. We then derive a Tanaka-type formula from the Itô formula of Section 3 and show that  $B_h$  satisfies the LND property. This implies continuity and Hölder regularities in space and in time of the trajectories of the local time.

**Definition 10.** For any Borel set  $C \subset \mathbb{R}_+$ , the occupation measure  $m_C$  of  $B_h$  on  $C$  is defined, for all Borel sets  $A \subset \mathbb{R}$ , by  $m_C(A) = \lambda\{t \in C, B_{h(t)}(t) \in A\}$ , where  $\lambda$  is the Lebesgue measure on  $\mathbb{R}_+$ . If  $m_C$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}$ , the local time (or occupation density) of  $B_h$  on  $C$  is defined as the Radon–Nikodym derivative of  $m_C$  and will be denoted by  $\{L(C, x), x \in \mathbb{R}\}$ . Sometimes we write  $L(t, x)$  instead of  $L([0, t], x)$ .

This definition implies that the local time of  $B_h$  satisfies the occupation density formula

$$\int_C g(t, B_h(t)) dt = \int_{C \times \mathbb{R}} g(t, x) L(dt, x) dx \tag{12}$$

for all continuous functions with compact support  $g : C \times \mathbb{R} \rightarrow \mathbb{R}$ . If  $g$  does not depend explicitly on  $t$ , then we get the more classical occupation density formula where the right-hand side of (12) is replaced by  $\int_{\mathbb{R}} g(x) L(C, x) dx$ .

**Proposition 11.** The local time of  $B_h$  exists  $P$ -a.s. on any interval  $[0, T]$  and is a square-integrable function of  $x$ .

**Proof.** By [3], for any continuous and zero-mean Gaussian process  $\{X_t, t \in [0, T]\}$  with bounded covariance function, the condition

$$\int_0^T \int_0^T \frac{ds dt}{\sqrt{\mathbb{E}[|X_t - X_s|^2]}} < +\infty \tag{13}$$

is sufficient for the local time of  $X$  to exist and to be a square-integrable function of  $x$ . For  $(s, t)$  away from the diagonal, we write, for  $s < t$ ,

$$\mathbb{E}[|B_h(t) - B_h(s)|^2] = \int_0^s (K_{h(t)}(t, u) - K_{h(s)}(s, u))^2 du + \int_s^t K_{h(t)}(t, u)^2 du$$

and deduce from (1) that the second term stays strictly positive as  $(s, t)$  varies in  $[0, t - \varepsilon] \times [0, T]$ . If  $(s, t)$  is close to the diagonal, say  $0 \leq t - s < \varepsilon$ , then, by Proposition 5,

$$\mathbb{E}[|B_h(t) - B_h(s)|^2] \sim c_{h(t)}^{-2}(t - s)^{2h(t)}$$

as  $s \nearrow t$  and a direct calculation shows that (13) is satisfied. □

Let us now derive a Tanaka-type formula for  $B_h$ . Since the last term in the Itô formula (11) is an integral with respect to the variance function  $R$  and since  $R$  is not, in general, increasing, but only of finite variation, the trajectorial representation of the local time is more delicate than for Brownian motion or for fractional Brownian motion. In fact, on the time intervals where  $R$  is (strictly) increasing or decreasing, this formula gives an occupation density  $\widehat{L}$  related to  $L$ .

**Theorem 12.** *Suppose that  $h$  is continuously differentiable. Then, for all  $a \in \mathbb{R}$ ,*

$$|B_h(t) - a| - |B_h(s) - a| = \int_s^t \text{sign}(B_h(u) - a) \delta B_h(u) + \widehat{L}([s, t], a)$$

*P*-a.s., where  $\widehat{L}([s, t], a) = \int_s^t R'(u)L(du, a)$  and  $L$  is the local time with respect to the Lebesgue measure of  $B_h$ . On the time intervals  $[s, t]$  where  $R$  is strictly increasing (resp., strictly decreasing),  $\widehat{L}$  (resp.,  $-\widehat{L}$ ) is the (positive) occupation density of  $B_h$  with respect to the measure induced by  $R$ .

**Remark 13.** (a) No information on the local time of  $B_h$  can be deduced from the above formula if  $R'(u) = 0$  on the interval  $[s, t]$ . In fact,

$$R'(u) = \frac{d}{du} \mathbb{E}[B_{h(u)}^2(u)] = \frac{d}{du} (u^{2h(u)}) = 2 \left( h'(u) \log u + \frac{h(u)}{u} \right) u^{2h(u)}$$

and  $R'(u) = 0$  if  $h(u) = 1/\log u \in (1/2, 1)$  on an interval  $(s, t)$ . In this case,  $\widehat{L}([s, t], a) = 0$  for all  $a$ .

(b) Tanaka formulas for fractional Brownian motion have been shown by several authors. We refer to the survey [8] for references.

**Proof of Theorem 12.** For  $\varepsilon > 0$ , let  $p_\varepsilon(x) = (2\pi\varepsilon)^{-1/2} \exp(-x^2/(2\varepsilon))$ . We apply the Itô formula of the previous section to the function

$$F_\varepsilon(x) = \int_0^x F'_\varepsilon(y) dy,$$

where

$$F'_\varepsilon(x) = 2 \int_{-\infty}^x p_\varepsilon(y) dy - 1.$$

$F'_\varepsilon(x)$  then converges to  $\text{sign}(x)$  and  $F_\varepsilon(x)$  converges to  $|x|$  as  $\varepsilon \rightarrow 0$ . By (11), for  $\varepsilon > 0$  fixed,

$$F_\varepsilon(B_h(t) - a) = F_\varepsilon(-a) + \int_0^t \int_s^t F'_\varepsilon(B_h(r) - a) K(dr, s) \delta W_s + \int_0^t p_\varepsilon(B_h(s) - a) dR(s). \tag{14}$$

Note that, by (K4), the process  $\{F'_\varepsilon(B_h(r) - a), r \in [0, t]\}$  belongs to  $L^2(\Omega, \mathcal{H}_K)$  and therefore to  $\text{Dom}(\delta^{B_h})$ . Or, equivalently, the process  $\{\int_s^t F'_\varepsilon(B_h(r) - a) K_h(dr, s), s \in [0, t]\}$  belongs to  $\text{Dom}(\delta^W)$ . Clearly,  $F_\varepsilon(B_h(t) - a)$  converges to  $|B_h(t) - a|$  in  $L^2(\Omega)$  and  $F_\varepsilon(-a)$  converges to  $|a|$  if  $\varepsilon \rightarrow 0$ . Moreover, the process  $\{F'_\varepsilon(B_h(r) - a), r \in [0, t]\}$  converges, as  $\varepsilon \rightarrow 0$ , to  $\{\text{sign}(B_h(r) - a), r \in [0, t]\}$  in  $L^2(\Omega, \mathcal{H}_K)$ . In fact, by (K4),

$$\mathbb{E} \left[ \int_0^t \left( \int_s^t |F'_\varepsilon(B_h(r) - a) - \text{sign}(B_h(r) - a)| |K_h|(dr, s) \right)^2 ds \right]$$

is upper bounded independently of  $\varepsilon$  and we may apply the dominated convergence theorem ([9], Lemma 1). Therefore, the last term in (14) converges in  $L^2(\Omega)$ . Let us denote the limit by  $\Lambda_t^a$ . Therefore, for any continuous function  $f$  with compact support in  $\mathbb{R}$ ,

$$\int \left( \int_0^t p_\varepsilon(B_h(s) - a) dR(s) \right) f(a) da \tag{15}$$

converges in  $L^1(\Omega)$  to  $\int \Lambda_t^a f(a) da$ . In fact, the dominated convergence theorem applies because

$$\int_0^t \mathbb{E}[p_\varepsilon(B_h(s) - a)] dR(s) = \int_0^t p_{R(s)+\varepsilon}(a) dR(s) \leq \int_0^t s^{-b} R'(s) ds < \infty.$$

But (15) also converges to  $\int_0^t f(B_h(s)) dR(s) = \int_0^t f(B_h(s)) R'(s) ds$ , where we use the fact that  $R$  is differentiable if  $h$  is differentiable. Hence,

$$\int_0^t f(B_h(s)) R'(s) ds = \int \Lambda_t^a f(a) da.$$

By the occupation density formula (12) applied to  $g(s, x) = f(x)R'(s)$ , we get  $\Lambda_t^a = \int_0^t R'(s) L(ds, a) = \widehat{L}(t, a)$  for  $\lambda$ -a.e.  $a$ ,  $P$ -a.s. We can extend this to all  $a \in \mathbb{R}$  since there exists a jointly continuous version of  $L$ , and therefore of  $\widehat{L}$ , as will be shown next by Berman’s methods, which are independent of the Tanaka formula.  $\square$

We now state regularity properties in time and space of the trajectories of  $L$ . The regularity properties of  $\widehat{L}$  follow easily since  $\widehat{L}(t, x) = \int_0^t R'(s) L(ds, x)$ . In order to show the existence of a jointly continuous version of  $L$ , we recall the hypotheses introduced in [7] and show that they are satisfied for  $B_h$ . We recall them in terms of any real-valued separable random process  $\{X(t), t \in [0, T]\}$  with Borel sample functions.

- **Hypothesis**  $(\mathbb{A})$ . There exist numbers  $\rho_0 > 0$  and  $H \in (0, 1)$  and a positive function  $\psi \in L^1(\mathbb{R})$  such that for all  $\lambda \in \mathbb{R}$  and  $t, s \in [0, T]$ ,  $0 < |t - s| < \rho_0$ , we have

$$\left| \mathbb{E} \left[ \exp \left( i \lambda \frac{X(t) - X(s)}{|t - s|^H} \right) \right] \right| \leq \psi(\lambda).$$

- **Hypothesis**  $(\mathbb{A}_m)$ . There exist positive constants  $\delta$  and  $c$  (both eventually depending on  $m \geq 2$ ) such that for all  $t_1, t_2, \dots, t_m$  with  $0 =: t_0 < t_1 < \dots < t_m \leq T$  and  $|t_m - t_1| < \delta$ , we have

$$\left| \mathbb{E} \left[ \exp \left( i \sum_{j=1}^m u_j (X(t_j) - X(t_{j-1})) \right) \right] \right| \leq \prod_{j=1}^m \left| \mathbb{E} \left[ \exp( i c u_j (X(t_j) - X(t_{j-1}))) \right] \right|$$

for all  $u_1, u_2, \dots, u_m \in \mathbb{R}$ .

Hypothesis  $(\mathbb{A})$  is evidently satisfied for self-similar processes with stationary increments: here,  $\psi(\lambda) = |\mathbb{E}[e^{i\lambda X(1)}]|$ . Hypothesis  $(\mathbb{A})$  is also closely related to asymptotic self-similarity and, in fact, it holds for a fairly large class of processes (see Proposition 15).

If  $X$  has independent increments,  $(\mathbb{A}_m)$  is trivially true for all  $m \geq 2$ . When the left- and right-hand sides of the inequality are applied to the characteristic function of a Gaussian process  $X$ , we get the condition which is known in the literature as *local non-determinism* (LND). Loosely speaking, LND says how much dependence is allowed for the increments of the process if the local time has certain regularity properties. As a general rule, the trajectories of local time become more regular if the trajectories of the process become less regular.

**Proposition 14.** *For all  $T > 0$ , the process  $\{B_h(t), t \in [0, T]\}$  satisfies hypotheses  $(\mathbb{A})$  and  $(\mathbb{A}_m)$  for all  $m \geq 2$  with  $H = \max_{0 \leq t \leq T} h(t)$ .*

**Proof.** Let us start by showing  $(\mathbb{A})$ . By similar calculations as in the proof of Proposition 5, for every  $t \in [0, T]$ ,

$$\left| \mathbb{E} \left[ e^{-2h(t)} (B_h(t + \varepsilon) - B_h(t))^2 \right] - \frac{1}{c_h^2(t)} \right| \leq \varepsilon^{\beta - \sup_{[0, T]} h} \int_0^{T+1} (\Phi_{T+1}(s))^2 ds.$$

We can now conclude by applying Proposition 15.

We now prove  $(\mathbb{A}_m)$  for all  $m \geq 2$ . We show that  $B_h$  satisfies the LND property as introduced by Berman for Gaussian processes. For simplicity, we write  $B$  instead of  $B_h$  in this proof. Let  $t_1 < t_2 < \dots < t_m$  and let  $\mathcal{V}_m$  be the relative conditioning error given by

$$\mathcal{V}_m = \frac{\text{Var}[B(t_m) - B(t_{m-1}) | B(t_1), \dots, B(t_{m-1})]}{\text{Var}[B(t_m) - B(t_{m-1})]}.$$

The Gaussian process is said to be LND if

$$\liminf_{\substack{c \searrow 0^+ \\ 0 < t_m - t_1 \leq c}} \mathcal{V}_m > 0. \tag{16}$$

This condition means that a small increment of the process is not completely predictable on the basis of a finite number of observations from the immediate past. For Gaussian processes, Berman [4] has proven that (16) implies  $(\mathbb{A}_m)$ . More precisely, he has shown that if  $X$  satisfies (16) for all  $m \geq 2$ , then there exist constants  $C_m > 0$  and  $\delta_m > 0$  such that, for all  $u_1, u_2, \dots, u_m \in \mathbb{R}$ ,

$$\text{Var} \left[ \sum_{j=1}^m u_j [B(t_j) - B(t_{j-1})] \right] \geq C_m \sum_{j=1}^m u_j^2 \text{Var}[B(t_j) - B(t_{j-1})],$$

where  $t_0 = 0$  and  $t_1 < \dots < t_m$  are different and lie in an interval of length at most  $\delta_m$ . This implies  $(\mathbb{A}_m)$ . In order to prove that  $B$  verifies (16), fix  $m \geq 2$  and let  $0 < t_1 < t_2 < \dots < t_m < T$ . By Proposition 3,

$$\text{Var}[B(t_m) - B(t_{m-1})] \leq C_{h(T)}(t_m - t_{m-1})^{2h(t_m)}, \tag{17}$$

where  $C_{h(T)}$  is a constant depending on  $T$ . We now study the conditional variance. Because of the definition of  $B_h$  in terms of  $W$ , we have the inclusion of  $\sigma$ -algebra

$$\sigma\{B(t_1), \dots, B(t_{m-1})\} \subset \sigma\{W(t), t \in [0, t_{m-1}]\}.$$

We then get

$$\begin{aligned} &\text{Var}[B(t_m) - B(t_{m-1}) | B(t_1), \dots, B(t_{m-1})] \\ &\geq \text{Var}[B(t_m) - B(t_{m-1}) | W(t), t \in [0, t_{m-1}]] \\ &= \text{Var}[B(t_m) | W(t), t \in [0, t_{m-1}]] \\ &= \text{Var} \left[ \int_{t_{m-1}}^{t_m} K_{h(t_m)}(t_m, u) W(du) \right] \\ &= \int_{t_{m-1}}^{t_m} (K_{h(t_m)}(t_m, u))^2 du. \end{aligned} \tag{18}$$

We simplify the notation by letting  $t := t_m$ ,  $s := t_{m-1}$  and  $H := h(t)$  and considering  $\int_s^t (K_H(t, u))^2 du$ . We recall that we have the relation

$$\int_s^t (K_H(t, u))^2 du = \mathbb{E}[(B_H(t) - B_H(s))^2] - \int_0^s (K_H(t, u) - K_H(s, u))^2 du \tag{19}$$

so we study  $\int_0^s (K_H(t, u) - K_H(s, u))^2 du$ . We make the same substitutions as in the proof of Proposition 2 and thereby obtain

$$\begin{aligned} &\int_0^s (K_H(t, u) - K_H(s, u))^2 du \\ &= \int_0^s u^{1-2H} \left( \int_s^t (y - u)^{H-3/2} y^{H-1/2} dy \right)^2 du \end{aligned}$$

$$\begin{aligned}
 &= \int_s^t dy \int_s^t dz y^{H-1/2} z^{H-1/2} \int_0^{y \wedge z} u^{1-2H} (y-u)^{H-3/2} (z-u)^{H-3/2} du \\
 &= \int_s^t dy \int_s^t dz |y-z|^{2H-2} \int_1^{\Psi_1(s,y,z)} (v-1)^{1-2H} v^{H-3/2} dv,
 \end{aligned}$$

where

$$\Psi_1(s, y, z) = \frac{yz - zs}{yz - ys} 1_{y>z} + \frac{yz - ys}{yz - zs} 1_{y \leq z}.$$

Then, using (19) and making the substitutions  $(y, z) \rightarrow (y + s, z + s)$  and  $(y, z) \rightarrow ((t - s)y, (t - s)z)$ , we get

$$\int_s^t (K_H(t, u))^2 du = (t - s)^{2H} \int_0^1 dy \int_0^1 dz |y - z|^{2H-2} \int_{\Psi_2(s,t,y,z)}^\infty (v - 1)^{1-2H} v^{H-3/2} dv,$$

where

$$\Psi_2(s, t, y, z) = \frac{(t - s)yz + ys}{(t - s)yz + zs} 1_{y>z} + \frac{(t - s)yz + zs}{(t - s)yz + ys} 1_{y \leq z}.$$

We can therefore deduce that

$$\liminf_{(t-s) \rightarrow 0} (t - s)^{-2H} \int_s^t (K_H(t, u))^2 du \geq \mathcal{I}, \tag{20}$$

where

$$\mathcal{I} = \int_0^1 dy \int_0^1 dz |y - z|^{2b-2} \int_{\Psi_3(y,z)}^\infty (v - 1)^{1-2b} v^{a-3/2} dv > 0$$

with

$$\Psi_3(y, z) = \sup \left\{ \frac{y}{z}, \frac{z}{y}, 2 \right\}.$$

Combining (17), (18) and (20), we then get

$$\liminf_{\substack{c \searrow 0^+ \\ 0 < t_m - t_1 \leq c}} \mathcal{V}_m \geq \frac{\mathcal{I}}{C_h(T)} > 0, \tag{21}$$

which concludes the proof. □

The beginning of the proof of Proposition 14 shows that the hypothesis  $(\mathbb{A})$  holds for a much larger class of processes. In fact, we have only used the assumption of the following proposition.

**Proposition 15.** *Let  $\{X_t, t \in [0, T]\}$  be a centered Gaussian process. Suppose that for some positive continuous functions  $f : [0, T] \rightarrow (0, 1)$  and  $g : [0, T] \rightarrow (0, \infty)$ ,*

$$\mathbb{E}[\varepsilon^{-2f(t)} (X(t + \varepsilon) - X(t))^2] \xrightarrow{\varepsilon \rightarrow 0} g(t) \tag{22}$$



uniformly in  $t$ . Hypothesis (A) then holds with  $H = \sup f$ .

**Proof.** Let us fix  $H = \sup f$ . Because  $X$  is Gaussian and centered, we have

$$\mathbb{E} \left[ \exp \left( i\lambda \frac{X(t + \varepsilon) - X(t)}{\varepsilon^H} \right) \right] = \exp \left( -\frac{\lambda^2}{2} \mathbb{E} \left[ \left( \frac{X(t + \varepsilon) - X(t)}{\varepsilon^H} \right)^2 \right] \right). \tag{23}$$

Because of (22), there exists  $\varepsilon_0$  such that for every  $\varepsilon$  satisfying  $|\varepsilon| < \varepsilon_0$  and for every  $t$ , we have

$$\mathbb{E} \left[ \left( \frac{X(t + \varepsilon) - X(t)}{\varepsilon^{f(t)}} \right)^2 \right] \geq \frac{c}{2}, \tag{24}$$

where  $c = \inf_{[0, T]} g$ . Besides,  $\varepsilon^{2f(t)-2H} \geq 1$  and thus

$$\mathbb{E} \left[ \left( \frac{X(t + \varepsilon) - X(t)}{\varepsilon^H} \right)^2 \right] = \varepsilon^{2f(t)-2H} \mathbb{E} \left[ \left( \frac{X(t + \varepsilon) - X(t)}{\varepsilon^{f(t)}} \right)^2 \right] \geq \frac{c}{2}. \tag{25}$$

Combining (23) and (25), we then get, for every  $\lambda$ , and  $t$  and  $s$  satisfying  $|t - s| < \varepsilon_0$ , that

$$\left| \mathbb{E} \left[ \exp \left( i\lambda \frac{X(t + \varepsilon) - X(t)}{\varepsilon^H} \right) \right] \right| \leq \exp \left( -\frac{\lambda^2 c}{4} \right).$$

We then choose  $\psi(\lambda) = \exp(-\lambda^2 c/4)$  to conclude the proof. □

Let us now state a regularity result for the trajectories of  $L$ . The following theorem states that  $L(t, x)$  is Hölder continuous in  $t$  of order  $1 - H$  and Hölder continuous in  $x$  of order  $\frac{1-H}{2H}$ , where  $H$  is the constant appearing in (A). For the proofs, we refer to [7], where it is shown that these regularities hold for any process starting at zero and satisfying (A) and (A<sub>m</sub>) for all  $m \geq 2$ . Here, the function  $\psi$  in (A) satisfies

$$\int_{|u| \geq 1} |u|^{(1-H)/H} \psi(u) \, du < \infty.$$

**Theorem 16.**  $\{B_h(t), t \in [0, T]\}$  has a jointly continuous local time  $\{L(t, x), (t, x) \in [0, T] \times \mathbb{R}\}$ . Moreover, for any compact set  $K \subset \mathbb{R}$  and any interval  $I \subset [0, T]$  with length less than  $\rho_0$  (the constant appearing in (A)):

- (i) if  $0 < \xi < (1 - H)/2H$ , then  $|L(I, x) - L(I, y)| \leq \eta |x - y|^\xi$  for all  $x, y \in K$ ;
- (ii) if  $0 < \delta < 1 - H$ , then  $\sup_{x \in K} L(I, x) \leq \eta |I|^\delta$ , where  $\eta$  is a random variable, almost surely positive and finite, and  $|I|$  is the length of  $I$ .

### Acknowledgements

The authors would like to thank the referee for his/her comments which resulted in a new proof of Proposition 14. This work was partly supported by the program Volubilis ‘‘Action Intégrée MA/06/142’’ and by the Commission of the European Communities Grant PIRSES-GA-2008-230804 within the program ‘Marie Curie Actions’.

## References

- [1] Alòs, E., Mazet, O. and Nualart, D. (2001). Stochastic calculus with respect to Gaussian processes. *Ann. Probab.* **29** 766–801. [MR1849177](#)
- [2] Benassi, A., Cohen, S. and Istas, J. (1998). Identifying the multifractional function of a Gaussian process. *Statist. Probab. Lett.* **39** 337–345. [MR1646220](#)
- [3] Berman, S.M. (1969). Local times and sample function properties of stationary Gaussian process. *Trans. Amer. Math. Soc.* **137** 277–299. [MR0239652](#)
- [4] Berman, S.M. (1973). Local nondeterminism and local times of Gaussian processes. *Indiana Univ. Math. J.* **23** 69–94. [MR0317397](#)
- [5] Billingsley, P. (1968). *Convergence of Probability Measures*. New York: Wiley. [MR0233396](#)
- [6] Benassi, A., Jaffard, S. and Roux, D. (1997). Elliptic Gaussian random processes. *Rev. Mat. Iberoamericana* **13** 19–90. [MR1462329](#)
- [7] Boufoussi, B., Dozzi, M. and Guerbaz, R. (2008). Path properties of a class of locally asymptotically self similar processes. *Electron. J. Probab.* **13** 898–921. [MR2413288](#)
- [8] Coutin, L. (2007). An introduction to (stochastic) calculus with respect to fractional Brownian motion. In *Séminaire de Probabilités XL. Lecture Notes in Math.* **1899** 3–65. Berlin: Springer. [MR2408998](#)
- [9] Coutin, L., Nualart, D. and Tudor C.A. (2001). Tanaka formula for the fractional Brownian motion. *Stochastic Process. Appl.* **94** 301–315. [MR1840834](#)
- [10] Lacaux, C. (2004). Real harmonizable multifractional Lévy motions. *Ann. Inst. H. Poincaré Probab. Statist.* **40** 259–277. [MR2060453](#)
- [11] Nualart, D. (2006). *The Malliavin Calculus and Related Topics*. Berlin: Springer. [MR2200233](#)
- [12] Peltier, R.F and Lévy Véhel, J. (1995). Multifractional Brownian motion: Definition and preliminary results. Technical Report RR-2645, INRIA.
- [13] Samorodnitsky, G. and Taqqu, M.S. (1994). *Stable Non-Gaussian Random Processes*. New York: Chapman and Hall. [MR1280932](#)
- [14] Stoev, S. and Taqqu, M.S. (2004). Stochastic properties of the linear multifractional stable motion. *Adv. Appl. Probab.* **36** 1085–1115. [MR2119856](#)
- [15] Stoev, S. and Taqqu, M.S. (2005). Path properties of the linear multifractional stable motion. *Fractals* **13** 157–178. [MR2151096](#)
- [16] Surgailis, D. (2008). Nonhomogenous fractional integration and multifractional processes. *Stochastic Process. Appl.* **118** 171–198. [MR2376898](#)

Received May 2009 and revised January 2010