

**ONE-TO-ONE CONTINUOUS EXTENSIONS
OF ANALYTIC FUNCTIONS**

Ramesh Garimella

Northwest Missouri State University

Let U be the open unit disc in the complex plane. Let $H(U)$ stand for the space of functions analytic on U . Let

$$A = \{g \in H(U) : g'(0) \neq 0\} .$$

For $g \in A$,

$$g(z) = \sum_{n=0}^{\infty} a_n z^n ,$$

following the lead of Walter Rudin (see [1] problem E3325 p.445), we say g has the property P_t if

$$\sum_{n=2}^{\infty} |a_n| n \leq t .$$

In this short note we prove the following result which is an extension of the problem E3325 of [1].

Theorem: Let $g \in A$ have the property P_t for some $t > 0$. Then g is one-to-one and admits a one-to-one continuous extension to the closed unit disc if $t \leq |g'(0)|$. First we prove a lemma.

Lemma: Assume $g \in A$ and has the property P_t for some $t > 0$.

Then g has a continuous extension to the closure of U .

Proof: Let

$$f(z) = (g(z) - g(0))(g'(0))^{-1} .$$

Obviously $f \in A$, $f(0) = 0$ and $f'(0) = 1$ and has the property P_s

where $s = t|g'(0)|^{-1}$. Let

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n .$$

Since f has the property P_s , the series

$$z + \sum_{n=2}^{\infty} a_n z^n$$

is absolutely convergent for every z in the unit circle. Now we show that f is continuous on the closed unit disc. Let z, w be in

the closed unit disc. We have,

$$\begin{aligned}
|f(z) - f(w)| &= \left| (z - w) + \sum_{n=2}^{\infty} a_n (z^n - w^n) \right| \\
&= |z - w| \left| 1 + \sum_{n=2}^{\infty} a_n (z^{n-1} + z^{n-2}w + \dots \right. \\
&\quad \left. + zw^{n-2} + w^{n-1}) \right| \\
&\leq |z - w| \left[1 + \sum_{n=2}^{\infty} |a_n| (|z|^{n-1} + |z|^{n-2}|w| + \dots + |w|^{n-1}) \right] \\
&\leq |z - w| \left[1 + \sum_{n=2}^{\infty} n|a_n| \right] \\
&\leq |z - w|(1 + s) .
\end{aligned}$$

The next-to-last inequality is true since $|z| \leq 1$, and $|w| \leq 1$. From the above it follows that f has a continuous extension to the closed unit disc. Hence, g has a continuous extension to the closed unit disc. Q.E.D.

Proof of the Theorem : Let

$$f = (g(z) - g(0))^{-1}(g'(0))^{-1} .$$

Then $f \in A$, $f(0) = 0$, $f'(0) = 1$ and has the property P_t where $t \leq 1$. Since f has continuous extension to the closed unit disc (by the lemma), it is enough to show that f is one-to-one and the extension of f is one-to-one on the closed unit disc. Let

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

For z, w in the open unit disc, since

$$\begin{aligned} |f(z) - f(w)| &= \left| (z - w) + \sum_{n=2}^{\infty} a_n (z^n - w^n) \right| \\ &= |z - w| \left| 1 + \sum_{n=2}^{\infty} a_n (z^{n-1} + z^{n-2}w + \dots + w^{n-1}) \right|, \end{aligned}$$

and since

$$\left| \sum_{n=2}^{\infty} a_n (z^{n-1} + z^{n-2}w + \dots + w^{n-1}) \right| < \sum_{n=2}^{\infty} n|a_n| \leq t \leq 1$$

it follows that f is one-to-one on the open unit disc. Also from the above it follows that $f(z) \neq f(w)$ if one of z, w is in the open unit disc and the other on the unit circle. Now we show that f is one-to-one on the unit circle. By the lemma, f has continuous extension

to the closed unit disc. Let $f(U) = \Omega$. If possible assume that z, w are distinct points on the unit circle such that $f(z) = f(w)$. Let $z_n = (1 - n^{-1})z$, $w_n = (1 - n^{-1})w$. Since f is continuous on the closed unit disc, $f(z_n) \rightarrow f(z)$ and $f(w_n) \rightarrow f(w)$. Since f is a homeomorphism of U onto Ω , $f(z) (= f(w))$ is a boundary point of Ω . Now for each $n \geq 1$, let

$$s_n = \begin{cases} f(z_n) & \text{if } n \text{ is even} \\ f(w_n) & \text{if } n \text{ is odd.} \end{cases}$$

Clearly s_n is a sequence in Ω converging to the boundary point of Ω . Since f^{-1} from Ω onto U is a homeomorphism, the sequence $f^{-1}(s_n)$ must converge to a point of the unit circle. This is a contradiction because by definition of s_n , the sequence $f^{-1}(s_n)$ has two subsequences converging to two different limits. Hence f is one-to-one on the unit circle. It follows that g is one-to-one in the unit disc and has a one-to-one continuous extension to the closed unit disc. Q.E.D.

Remark: For any $t > 1$ there exists a function $g \in A$ having the property P_t and fails to be one-to-one (refer to (c) of problem

E3325 of [1]). Let $t > 1$. Write $t = 1 + c$. Define

$$g(z) = z - \frac{1}{2}z^2 + \frac{\alpha}{3}z^3 + \frac{\beta}{4}z^4$$

where

$$\alpha = \frac{-72 - 3c}{5} \text{ and } \beta = \frac{8c + 72}{5} .$$

Since

$$\sum_{n=2}^4 n|a_n| = 1 + \alpha + \beta = 1 + c = t ,$$

g has the property P_t . Clearly $g \in A$. Now by the choice of α, β it is easy to verify that $g(1/2) = 0$. Also $g(0) = 0$. Hence g is not one-to-one on the unit disc.

References

1. *The American Mathematical Monthly*, Vol. 96, No. 5, May 1989.