

## A NEW GENERALIZATION OF SHANIN'S NOTION of $R_0$ TOPOLOGICAL SPACES

M. Caldas, E. Hatir, and S. Jafari

**Abstract.** In this paper, we introduce and investigate some weak separation axioms by using the notions of  $(\Lambda, \alpha)$ -open sets and the  $(\Lambda, \alpha)$ -closure operator.

**1. Introduction.** In 1943, N. A. Shanin [8] offered a new weak separation axiom called  $R_0$  to the world of general topology. In 1961, A. S. Davis [2] rediscovered this axiom and established some properties of topological spaces endowed with it. Later on several topologists further investigated  $R_0$  topological spaces [3,4,5,6]. The notion of  $\alpha$ -open set was introduced by O. Njåstad [7] in 1965. Since then it has been investigated in different respects in the literature. Quite recently Caldas, et. al. [1] introduced and studied the notions of  $(\Lambda, \alpha)$ -open sets,  $(\Lambda, \alpha)$ -closed sets, and the  $(\Lambda, \alpha)$ -closure operator. This paper deals with some new low separation axioms by utilizing  $(\Lambda, \alpha)$ -open and  $(\Lambda, \alpha)$ -closed sets. There is no doubt that low separation axioms play a very important role in general topology. Indeed, there are lots of research papers which deal with different low separation axioms and also many topologists worldwide are doing research in this area. It is the aim of this paper to offer some new types of low separation axioms by using  $(\Lambda, \alpha)$ -open sets and  $(\Lambda, \alpha)$ -closure operator.

In this paper, by  $(X, \tau)$  and  $(Y, \sigma)$  (or  $X$  and  $Y$ ) we always mean topological spaces. Let  $A$  be a subset of  $X$ . The subset  $A$  of the topological space  $(X, \tau)$  is called  $\alpha$ -open (originally called  $\alpha$ -sets) [7] if  $A \subseteq \text{Int}(\text{Cl}(\text{Int}(A)))$ . The complement of a  $\alpha$ -open set is called  $\alpha$ -closed. By  $\alpha\text{O}(X, \tau)$  (respectively  $\alpha\text{C}(X, \tau)$ ), we denote the family of all  $\alpha$ -open (respectively  $\alpha$ -closed) sets of  $(X, \tau)$ . Observe that  $\alpha$ -open sets form a topology, and also  $\alpha$ -openness does not imply openness in the underlying topology. Let  $C$  denote the standard "middle thirds" Cantor set in the unit interval  $[0, 1]$  with the standard topology, and let  $x \in C$ . Then,  $[0, 1] - \{C - \{x\}\}$  would be an  $\alpha$ -open set. The intersection of all  $\alpha$ -closed sets containing  $A$  is called the  $\alpha$ -closure of  $A$ , denoted by  $\text{Cl}_\alpha(A)$ . A subset  $A$  is also  $\alpha$ -closed if and only if  $A = \text{Cl}_\alpha(A)$ . A set  $U$  in a topological space  $(X, \tau)$  is a  $\alpha$ -neighborhood [7] of a point  $x$  if  $U$  contains an  $\alpha$ -open set  $V$  such that  $x \in V$ .

**Remark 1.1.**

- (i) It is shown in [7] that  $\alpha O(X, \tau)$  is a topology on  $X$  and  $\tau \subset \alpha O(X, \tau)$ .
- (ii) Clearly open sets are  $\alpha$ -open but one easily finds in the real line with the usual topology  $\alpha$ -open sets that are not open.

**Lemma 1.2.** Let  $A$  be a subset of a topological space  $(X, \tau)$ .

- (1)  $Cl_\alpha(A) = \cap \{F \in \alpha C(X, \tau) \mid A \subset F\}$ .
- (2)  $Cl_\alpha(A)$  is  $\alpha$ -closed, that is  $Cl_\alpha(Cl_\alpha(A)) = Cl_\alpha(A)$ .

**Definition 1.** Let  $A$  be a subset of a topological space  $(X, \tau)$ . A subset  $\Lambda_\alpha(A)$  is defined as follows:  $\Lambda_\alpha(A) = \cap \{O \in \alpha O(X, \tau) \mid A \subset O\}$  and  $A$  is called a  $\Lambda_\alpha$ -set if  $A = \Lambda_\alpha(A)$  [1].

**Definition 2.** Let  $A$  be a subset of a topological space  $(X, \tau)$ .

- (i)  $A$  is called a  $(\Lambda, \alpha)$ -closed set [1] if  $A = T \cap C$ , where  $T$  is a  $\Lambda_\alpha$ -set and  $C$  is an  $\alpha$ -closed set. The complement of a  $(\Lambda, \alpha)$ -closed set is called  $(\Lambda, \alpha)$ -open. We denoted the collection of all  $(\Lambda, \alpha)$ -open sets (respectively  $(\Lambda, \alpha)$ -closed sets) by  $\Lambda_\alpha O(X, \tau)$  (respectively by  $\Lambda_\alpha C(X, \tau)$ ).
- (ii) A point  $x \in X$  is called a  $(\Lambda, \alpha)$ -cluster point of  $A$  [1] if for every  $(\Lambda, \alpha)$ -open set  $U$  of  $X$  containing  $x$  we have  $A \cap U \neq \emptyset$ . The set of all  $(\Lambda, \alpha)$ -cluster points of  $A$  is called the  $(\Lambda, \alpha)$ -closure of  $A$  and is denoted by  $A^{(\Lambda, \alpha)}$ .

**Lemma 1.3** Let  $A$  and  $B$  be subsets of a topological space  $(X, \tau)$ . For the  $(\Lambda, \alpha)$ -closure, the following properties hold [1].

- (1)  $A \subset A^{(\Lambda, \alpha)}$ .
- (2)  $A^{(\Lambda, \alpha)} = \cap \{F \mid A \subset F \text{ and } F \text{ is } (\Lambda, \alpha)\text{-closed}\}$ .
- (3) If  $A \subset B$ , then  $A^{(\Lambda, \alpha)} \subset B^{(\Lambda, \alpha)}$ .
- (4)  $A$  is  $(\Lambda, \alpha)$ -closed if and only if  $A = A^{(\Lambda, \alpha)}$ .
- (5)  $A^{(\Lambda, \alpha)}$  is  $(\Lambda, \alpha)$ -closed.

**2. Sober  $\Lambda_\alpha$ - $R_0$  Topological Spaces.** A. S. Davis [2] introduced the notion of  $R_0$ -axiom which in some aspects is more natural than the  $T_0$ -axiom. In this section we introduce the concept of sober  $\Lambda_\alpha$ - $R_0$  topological space and we show that sober  $\Lambda_\alpha$ - $R_0$  and  $R_0$  are independent of each other.

**Lemma 2.1.** Let  $A$  be a subset of a space  $X$ . Then the following hold.

- (1) If  $A$  is  $\alpha$ -closed, then  $A$  is  $(\Lambda, \alpha)$ -closed.
- (2) If  $A$  is  $(\Lambda, \alpha)$ -closed, then  $A = \Lambda_\alpha(A) \cap A^{(\Lambda, \alpha)}$ .
- (3) If  $A_i$  is  $(\Lambda, \alpha)$ -closed for each  $i \in I$ , then  $\cap_{i \in I} A_i$  is  $(\Lambda, \alpha)$ -closed.
- (4) If  $A_i$  is  $(\Lambda, \alpha)$ -open for each  $i \in I$ , then  $\cup_{i \in I} A_i$  is  $(\Lambda, \alpha)$ -open.

Proof. (1) It is sufficient to observe that  $A = X \cap A$  where the whole set  $X$  is a  $\Lambda_\alpha$ -set.

(2) Let  $A$  be  $(\Lambda, \alpha)$ -closed, then there exists a  $\Lambda_\alpha$ -set  $T$  and a  $\alpha$ -closed set  $C$  such that  $A = T \cap C$ . By  $A \subset T$ , we have  $A \subset \Lambda_\alpha(A) \subset \Lambda_\alpha(T) = T$ , and also by  $A \subset C$ ,  $A \subset A^{(\Lambda, \alpha)} \subset C^{(\Lambda, \alpha)} = C$ . Now,  $A \subset \Lambda_\alpha(A) \cap A^{(\Lambda, \alpha)} \subset T \cap C = A$ . Hence,  $A = \Lambda_\alpha(A) \cap A^{(\Lambda, \alpha)}$ .

(3) Suppose that  $A_i$  is  $(\Lambda, \alpha)$ -closed for each  $i \in I$ . Then, for each  $i \in I$  there exists a  $\Lambda_\alpha$ -set  $T_i$  and a  $\alpha$ -closed set  $C_i$  such that  $A_i = T_i \cap C_i$ . Now,  $\bigcap_{i \in I} A_i = \bigcap_{i \in I} (T_i \cap C_i) = (\bigcap_{i \in I} T_i) \cap (\bigcap_{i \in I} C_i)$ . By Lemma 2.4 of [1],  $\bigcap_{i \in I} T_i$  is a  $\Lambda_\alpha$ -set and  $\bigcap_{i \in I} C_i$  is  $\alpha$ -closed. This shows that  $\bigcap_{i \in I} A_i$  is  $(\Lambda, \alpha)$ -closed.

(4) Follows from (3).

Definition 3. Let  $(X, \tau)$  be a topological space,  $A \subset X$  and  $x \in X$ . Then

- (i) The  $\Lambda_\alpha$ -kernel of  $A$ , denoted by  $\Lambda_\alpha Ker(A)$ , is defined to be the set  $\Lambda_\alpha Ker(A) = \bigcap \{G \in \Lambda_\alpha O(X, \tau) \mid A \subset G\}$ .
- (ii)  $\langle x \rangle = \{x\}^{(\Lambda, \alpha)} \cap \Lambda_\alpha Ker(\{x\})$ .

Lemma 2.2. Let  $(X, \tau)$  be a topological space and  $x \in X$ . Then

$$\Lambda_\alpha Ker(A) = \{x \in X \mid \{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset\}.$$

Proof. Let  $x \in \Lambda_\alpha Ker(A)$  and suppose  $\{x\}^{(\Lambda, \alpha)} \cap A = \emptyset$ . Hence,  $x \notin X - \{x\}^{(\Lambda, \alpha)}$  which is a  $(\Lambda, \alpha)$ -open set containing  $A$ . This is absurd, since  $x \in \Lambda_\alpha Ker(A)$ . Consequently,  $\{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset$ . Next, let  $x$  such that  $\{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset$  and suppose that  $x \notin \Lambda_\alpha Ker(A)$ . Then, there exists a  $(\Lambda, \alpha)$ -open set  $D$  containing  $A$  and  $x \notin D$ . Let  $y \in \{x\}^{(\Lambda, \alpha)} \cap A$ . Hence,  $D$  is a  $(\Lambda, \alpha)$ -neighborhood of  $y$  which does not contain  $x$ . But this contradicts  $x \in Ker_\alpha(A)$  and the claim follows.

Lemma 2.3. If  $A, B \subset X$ , then

- (1)  $A \subset B$  implies  $\Lambda_\alpha Ker(A) \subset \Lambda_\alpha Ker(B)$ .
- (2)  $\Lambda_\alpha Ker(\Lambda_\alpha Ker(A)) = \Lambda_\alpha Ker(A)$ .

Lemma 2.4. Let  $(X, \tau)$  be a topological space and  $x, y \in X$ . Then  $y \in \Lambda_\alpha Ker(\{x\})$  if and only if  $x \in \{y\}^{(\Lambda, \alpha)}$ .

Proof. Let  $y \notin \Lambda_\alpha Ker(\{x\})$ . Then there exists a  $(\Lambda, \alpha)$ -open set  $V$  containing  $x$  such that  $y \notin V$ . Hence,  $x \notin \{y\}^{(\Lambda, \alpha)}$ . The converse is similarly shown.

A subset  $B_x$  of a topological space  $X$  is said to be  $(\Lambda, \alpha)$ -neighborhood of a point  $x \in X$  if there exists a  $(\Lambda, \alpha)$ -open set  $U$  such that  $x \in U \subset B_x$ .

Lemma 2.5. A subset of a topological space  $X$  is  $(\Lambda, \alpha)$ -open in  $X$  if and only if it contains a  $(\Lambda, \alpha)$ -neighborhood of each of its points.

**Proposition 2.6.** If  $(X, \tau)$  is a topological space and  $A \subset X$ . Then

- (1)  $\Lambda_\alpha Ker(A) = \{x \in X / \{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset\}$ .
- (2) For each  $x \in X$ ,  $\Lambda_\alpha Ker(\langle x \rangle) = \Lambda_\alpha Ker(\{x\})$ .
- (3) For each  $x \in X$ ,  $\{\langle x \rangle\}^{(\Lambda, \alpha)} = \{x\}^{(\Lambda, \alpha)}$ .
- (4) For each  $(\Lambda, \alpha)$ -open set  $U \subset X$ , if  $x \in U$  then  $\langle x \rangle \subset U$ .
- (5) For each  $(\Lambda, \alpha)$ -closed set  $F \subset X$ , if  $x \in F$  then  $\langle x \rangle \subset F$ .

**Proof.** (1) Let  $x \in \Lambda_\alpha Ker(A)$  and suppose  $\{x\}^{(\Lambda, \alpha)} \cap A = \emptyset$ . Then,  $x \notin X \setminus \{x\}^{(\Lambda, \alpha)}$  which is a  $(\Lambda, \alpha)$ -open set containing  $A$ . This is impossible, since  $x \in \Lambda_\alpha Ker(A)$ . Consequently,  $\{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset$ . Next, let  $x \in X$  such that  $\{x\}^{(\Lambda, \alpha)} \cap A \neq \emptyset$  and suppose that  $x \notin \Lambda_\alpha Ker(A)$ . Then, there exists a  $(\Lambda, \alpha)$ -open set  $U$  containing  $A$  and  $x \notin U$ . Let  $y \in \{x\}^{(\Lambda, \alpha)} \cap A$ . Hence,  $U$  is a  $(\Lambda, \alpha)$ -neighborhood of  $y$  which does not contain  $x$ . But this contradicts  $x \in \Lambda_\alpha Ker(A)$ .

(2) Follows easily from Definition 2 and Lemma 2.4.

(3) The proof is quite similar to that of (2).

(4) Since  $x \in U$  and  $U$  is a  $(\Lambda, \alpha)$ -open set, we have that  $\Lambda_\alpha Ker(\{x\}) \subset U$ . Hence,  $\langle x \rangle \subset U$ .

(5) Since  $x \in F$  and  $F$  is a  $(\Lambda, \alpha)$ -closed set, we have that  $\langle x \rangle = \{x\}^{(\Lambda, \alpha)} \cap \Lambda_\alpha Ker(\{x\}) \subset \{x\}^{(\Lambda, \alpha)} \subset F$ .

**Lemma 2.7.** The following statements are equivalent for any points  $x$  and  $y$  in a topological space  $(X, \tau)$ .

- (1)  $\Lambda_\alpha Ker(\{x\}) \neq \Lambda_\alpha Ker(\{y\})$ .
- (2)  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ .

**Proof.** (1)  $\rightarrow$  (2): Suppose that  $\Lambda_\alpha Ker(\{x\}) \neq \Lambda_\alpha Ker(\{y\})$ , then there exists a point  $z$  in  $X$  such that  $z \in \Lambda_\alpha Ker(\{x\})$  and  $z \notin \Lambda_\alpha Ker(\{y\})$ . From  $z \in \Lambda_\alpha Ker(\{x\})$  it follows that  $\{x\} \cap \{z\}^{(\Lambda, \alpha)} \neq \emptyset$  which implies  $x \in \{z\}^{(\Lambda, \alpha)}$ . By  $z \notin \Lambda_\alpha Ker(\{y\})$ , we have  $\{y\} \cap \{z\}^{(\Lambda, \alpha)} = \emptyset$ . Since  $x \in \{z\}^{(\Lambda, \alpha)}$ ,  $\{x\}^{(\Lambda, \alpha)} \subset \{z\}^{(\Lambda, \alpha)}$  and  $\{y\} \cap \{x\}^{(\Lambda, \alpha)} = \emptyset$ . Therefore, it follows that  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ . Now  $\Lambda_\alpha Ker(\{x\}) \neq \Lambda_\alpha Ker(\{y\})$  implies that  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ .

(2)  $\rightarrow$  (1): Suppose that  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ . Then there exists a point  $z$  in  $X$  such that  $z \in \{x\}^{(\Lambda, \alpha)}$  and  $z \notin \{y\}^{(\Lambda, \alpha)}$ . It follows that there exists a  $(\Lambda, \alpha)$ -open set containing  $z$  and therefore  $x$  but not  $y$ , namely,  $y \notin \Lambda_\alpha Ker(\{x\})$  and thus,  $\Lambda_\alpha Ker(\{x\}) \neq \Lambda_\alpha Ker(\{y\})$ .

**Definition 4.** A topological space  $(X, \tau)$  is said to be sober  $\Lambda_\alpha$ - $R_0$  if  $\bigcap_{x \in X} \{x\}^{(\Lambda, \alpha)} = \emptyset$ .

**Theorem 2.8.** A topological space  $(X, \tau)$  is sober  $\Lambda_\alpha$ - $R_0$  if and only if  $\Lambda_\alpha \text{Ker}(\{x\}) \neq X$  for every  $x \in X$ .

**Proof.** Suppose that the space  $(X, \tau)$  is sober  $\Lambda_\alpha$ - $R_0$ . Assume that there is a point  $y$  in  $X$  such that  $\Lambda_\alpha \text{Ker}(\{y\}) = X$ . Then  $y \notin O$  which  $O$  is some proper  $(\Lambda, \alpha)$ -open subset of  $X$ . This implies that  $y \in \bigcap_{x \in X} \{x\}^{(\Lambda, \alpha)}$ . But this is a contradiction. Now assume that  $\Lambda_\alpha \text{Ker}(\{x\}) \neq X$  for every  $x \in X$ . If there exists a point  $y$  in  $X$  such that  $y \in \bigcap_{x \in X} \{x\}^{(\Lambda, \alpha)}$ , then every  $(\Lambda, \alpha)$ -open set containing  $y$  must contain every point of  $X$ . This implies that the space  $X$  is the unique  $(\Lambda, \alpha)$ -open set containing  $y$ . Hence,  $\Lambda_\alpha \text{Ker}(\{y\}) = X$  which is a contradiction. Therefore,  $(X, \tau)$  is sober  $\Lambda_\alpha$ - $R_0$ .

Recall that a topological space  $(X, \tau)$  is said to be  $R_0$  [2] if every open set contains the closure of each of its singletons.

**Example 2.9.** Let  $(X, \tau)$  be a topological space such that  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, \{a\}, X\}$ . Observe that  $(X, \tau)$  is sober  $\Lambda_\alpha$ - $R_0$ , but it is not  $R_0$ .

**Example 2.10.** Let  $(X, \tau)$  with  $\tau = \{\emptyset, X\}$ . Clearly  $(X, \tau)$  is not sober  $\Lambda_\alpha$ - $R_0$ , but it is  $R_0$ .

Examples 2.9 and 2.10 show that sober  $\Lambda_\alpha$ - $R_0$  and  $R_0$  are independent of each other.

**Definition 5.** A function  $f: X \rightarrow Y$  is always called  $\Lambda_\alpha$ -closed if the image of every  $(\Lambda, \alpha)$ -closed subset of  $X$  is  $(\Lambda, \alpha)$ -closed in  $Y$ .

**Theorem 2.11.** If  $f: X \rightarrow Y$  is an injective always  $\Lambda_\alpha$ -closed function and  $X$  is sober  $\Lambda_\alpha$ - $R_0$ , then  $Y$  is sober  $\alpha$ - $R_0$ .

**Proof.** Straightforward.

**Theorem 2.12.** If the topological space  $X$  is sober  $\Lambda_\alpha$ - $R_0$  and  $Y$  is any topological space, then the product  $X \times Y$  is sober  $\Lambda_\alpha$ - $R_0$ .

**Proof.** By showing that  $\bigcap_{(x,y) \in X \times Y} \{x, y\}^{(\Lambda, \alpha)} = \emptyset$  we are done. We have  $\bigcap_{(x,y) \in X \times Y} \{x, y\}^{(\Lambda, \alpha)} \subseteq \bigcap_{(x,y) \in X \times Y} \{x\}^{(\Lambda, \alpha)} \times \{y\}^{(\Lambda, \alpha)} = \bigcap_{x \in X} \{x\}^{(\Lambda, \alpha)} \times \bigcap_{y \in Y} \{y\}^{(\Lambda, \alpha)} \subseteq \emptyset \times Y = \emptyset$ .

**3.  $\Lambda_\alpha$ - $R_0$  Topological Spaces.** In this section  $(\Lambda, \alpha)$ -open sets and  $(\Lambda, \alpha)$ -closure operator are used to define a new separation axiom analogous to  $R_0$ - axiom and we obtain several characterizations of it.

**Definition 6.** A topological space  $(X, \tau)$  is said to be a  $\Lambda_\alpha$ - $R_0$  space if every  $(\Lambda, \alpha)$ -open set contains the  $(\Lambda, \alpha)$ -closure of each of its singletons.

The next result will give the characterization of the  $\Lambda_\alpha$ - $R_0$  space in terms of the  $(\Lambda, \alpha)$ -closure of points.

**Theorem 3.1.** For a topological space  $(X, \tau)$ , the following properties are equivalent.

- (1)  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.
- (2) For any  $F \in \Lambda_\alpha C(X, \tau)$ ,  $x \notin F$  implies that there exist  $U \in \Lambda_\alpha O(X, \tau)$ , such that  $F \subset U$  and  $x \notin U$ .
- (3) For any  $F \in \Lambda_\alpha C(X, \tau)$ ,  $x \notin F$  implies  $F \cap \{x\}^{(\Lambda, \alpha)} = \emptyset$ .
- (4) For any distinct points  $x$  and  $y$  of  $X$ , either  $\{x\}^{(\Lambda, \alpha)} = \{y\}^{(\Lambda, \alpha)}$  or  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$ .

**Proof.** (1)  $\rightarrow$  (2): Let  $F \in \Lambda_\alpha C(X, \tau)$  and  $x \notin F$ . Then by (1)  $\{x\}^{(\Lambda, \alpha)} \subset X \setminus F$ . Set  $U = X \setminus \{x\}^{(\Lambda, \alpha)}$ , then  $U \in \Lambda_\alpha O(X, \tau)$ ,  $F \subset U$  and  $x \notin U$ .

(2)  $\rightarrow$  (3): Let  $F \in \Lambda_\alpha C(X, \tau)$  and  $x \notin F$ . There exists  $U \in \Lambda_\alpha O(X, \tau)$  such that  $F \subset U$  and  $x \notin U$ . Since  $U \in \Lambda_\alpha O(X, \tau)$ ,  $U \cap \{x\}^{(\Lambda, \alpha)} = \emptyset$  and  $F \cap \{x\}^{(\Lambda, \alpha)} = \emptyset$ .

(3)  $\rightarrow$  (4): Assume that  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$  for distinct points  $x, y \in X$ . There exists  $z \in \{x\}^{(\Lambda, \alpha)}$  such that  $z \notin \{y\}^{(\Lambda, \alpha)}$  (or  $z \in \{y\}^{(\Lambda, \alpha)}$  such that  $z \notin \{x\}^{(\Lambda, \alpha)}$ ). There exists  $V \in \Lambda_\alpha O(X, \tau)$  such that  $y \notin V$  and  $z \in V$ ; hence  $x \in V$ . Therefore, we have  $x \notin \{y\}^{(\Lambda, \alpha)}$ . By (3), we obtain  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$ . The proof for otherwise case is similar.

(4)  $\rightarrow$  (1): Let  $V \in \Lambda_\alpha O(X, \tau)$  and  $x \in V$ . For each  $y \notin V$ ,  $x \neq y$  and  $x \notin \{y\}^{(\Lambda, \alpha)}$ . This shows that  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ . By (4),  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$  for each  $y \in X \setminus V$  and hence,

$$\{x\}^{(\Lambda, \alpha)} \cap \left( \bigcup_{y \in X \setminus V} \{y\}^{(\Lambda, \alpha)} \right) = \emptyset.$$

On the other hand, since  $V \in \Lambda_\alpha O(X, \tau)$  and  $y \in X \setminus V$ , we have  $\{y\}^{(\Lambda, \alpha)} \subset X \setminus V$ . Therefore,

$$X \setminus V = \bigcup_{y \in X \setminus V} \{y\}^{(\Lambda, \alpha)}.$$

Therefore, we obtain  $(X \setminus V) \cap \{x\}^{(\Lambda, \alpha)} = \emptyset$  and  $\{x\}^{(\Lambda, \alpha)} \subset V$ . Hence,  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.

**Corollary 3.2.** A topological space  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space if and only if for any  $x$  and  $y$  in  $X$ ,  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$  implies  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$ .

**Proposition 3.3.** A topological space  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_0$  if and only if for any points  $x$  and  $y$  and  $X$ ,  $\Lambda_\alpha \text{Ker}(\{x\}) \neq \Lambda_\alpha \text{Ker}(\{y\})$  implies  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\}) = \emptyset$ .

**Proof.** Suppose that  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space. Thus by Lemma 2.7, for any points  $x$  and  $y$  in  $X$  if  $\Lambda_\alpha \text{Ker}(\{x\}) \neq \Lambda_\alpha \text{Ker}(\{y\})$  then  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ . Now we prove that  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\}) = \emptyset$ . Assume that  $z \in \Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\})$ . By  $z \in \Lambda_\alpha \text{Ker}(\{x\})$ , it follows that  $x \in \{z\}^{(\Lambda, \alpha)}$ . Since  $x \in \{x\}^{(\Lambda, \alpha)}$ , by Corollary 3.2  $\{x\}^{(\Lambda, \alpha)} = \{z\}^{(\Lambda, \alpha)}$ . Similarly, we have  $\{y\}^{(\Lambda, \alpha)} = \{z\}^{(\Lambda, \alpha)} = \{x\}^{(\Lambda, \alpha)}$ . This is a contradiction. Therefore, we have  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\}) = \emptyset$ .

Conversely, let  $(X, \tau)$  be a topological space such that for any points  $x$  and  $y$  in  $X$ ,  $\Lambda_\alpha \text{Ker}(\{x\}) \neq \Lambda_\alpha \text{Ker}(\{y\})$  implies  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\}) = \emptyset$ . If  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ , then by Lemma 2.7,  $\Lambda_\alpha \text{Ker}(\{x\}) \neq \Lambda_\alpha \text{Ker}(\{y\})$ . Therefore,  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{y\}) = \emptyset$  which implies  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$ . Because  $z \in \{x\}^{(\Lambda, \alpha)}$  implies that  $x \in \Lambda_\alpha \text{Ker}(\{z\})$ ,  $\Lambda_\alpha \text{Ker}(\{x\}) \cap \Lambda_\alpha \text{Ker}(\{z\}) \neq \emptyset$ . By hypothesis, we therefore have  $\Lambda_\alpha \text{Ker}(\{x\}) = \Lambda_\alpha \text{Ker}(\{z\})$ . Then  $z \in \{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)}$  implies that  $\{x\}^{(\Lambda, \alpha)} = \{z\}^{(\Lambda, \alpha)} = \{y\}^{(\Lambda, \alpha)}$ . This is a contradiction. Therefore,  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} = \emptyset$  and by Corollary 3.2,  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.

**Proposition 3.4.** For a topological space  $(X, \tau)$ , the following properties are equivalent.

- (1)  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.
- (2) For any nonempty set  $A$  and  $G \in \Lambda_\alpha O(X, \tau)$  such that  $A \cap G \neq \emptyset$ , there exists  $F \in \Lambda_\alpha C(X, \tau)$  such that  $A \cap F \neq \emptyset$  and  $F \subset G$ .
- (3) Any  $G \in \Lambda_\alpha O(X, \tau)$ ,  $G = \cup\{F \in \Lambda_\alpha C(X, \tau) \mid F \subset G\}$ .
- (4) Any  $F \in \Lambda_\alpha C(X, \tau)$ ,  $F = \cap\{G \in \Lambda_\alpha O(X, \tau) \mid F \subset G\}$  (i.e.,  $F = \Lambda_\alpha \text{Ker}(F)$ ).
- (5) For any  $x \in X$ ,  $\{x\}^{(\Lambda, \alpha)} \subset \Lambda_\alpha \text{Ker}(\{x\})$ .

**Proof.** (1)  $\rightarrow$  (2): Let  $A$  be a nonempty set of  $X$  and  $G \in \Lambda_\alpha O(X, \tau)$  such that  $A \cap G \neq \emptyset$ . There exists  $x \in A \cap G$ . Since  $x \in G \in \Lambda_\alpha O(X, \tau)$ ,  $\{x\}^{(\Lambda, \alpha)} \subset G$ . Set  $F = \{x\}^{(\Lambda, \alpha)}$ , then  $F \in \Lambda_\alpha C(X, \tau)$ ,  $F \subset G$  and  $A \cap F \neq \emptyset$ .

(2)  $\rightarrow$  (3): Let  $G \in \Lambda_\alpha O(X, \tau)$ , then  $G \supset \cup\{F \in \Lambda_\alpha C(X, \tau) \mid F \subset G\}$ . Let  $x$  be any point of  $G$ . There exists  $F \in \Lambda_\alpha C(X, \tau)$  such that  $x \in F$  and  $F \subset G$ . Therefore, we have  $x \in F \subset \cup\{F \in \Lambda_\alpha C(X, \tau) \mid F \subset G\}$  and hence,  $G = \cup\{F \in \Lambda_\alpha C(X, \tau) \mid F \subset G\}$ .

(3)  $\rightarrow$  (4): This is obvious.

(4)  $\rightarrow$  (5): It follows from (4) and Lemma 1.3.

(5)  $\rightarrow$  (1): Let  $G \in \Lambda_\alpha O(X, \tau)$  and  $x \in G$ . Let  $y \in \Lambda_\alpha Ker(\{x\})$ , then  $x \in \{y\}^{(\Lambda, \alpha)}$  and  $y \in G$ . This implies that  $\Lambda_\alpha Ker(\{x\}) \subset G$ . Therefore, we obtain  $x \in \{x\}^{(\Lambda, \alpha)} \subset \Lambda_\alpha Ker(\{x\}) \subset G$ . This shows that  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.

Corollary 3.5. For a topological space  $(X, \tau)$ , the following properties are equivalent.

- (1)  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.
- (2)  $\{x\}^{(\Lambda, \alpha)} = \Lambda_\alpha Ker(\{x\})$  for all  $x \in X$ .

Proof. (1)  $\rightarrow$  (2): Suppose that  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space. By Proposition 3.4,  $\{x\}^{(\Lambda, \alpha)} \subset \Lambda_\alpha Ker(\{x\})$  for each  $x \in X$ . Let  $y \in \Lambda_\alpha Ker(\{x\})$ , then  $\{x\}^{(\Lambda, \alpha)} \cap \{y\}^{(\Lambda, \alpha)} \neq \emptyset$ . By Corollary 3.2 we obtain  $\{x\}^{(\Lambda, \alpha)} = \{y\}^{(\Lambda, \alpha)}$ . Therefore,  $y \in \{x\}^{(\Lambda, \alpha)}$  and hence,  $\Lambda_\alpha Ker(\{x\}) \subset \{x\}^{(\Lambda, \alpha)}$ . This shows that  $\{x\}^{(\Lambda, \alpha)} = \Lambda_\alpha Ker(\{x\})$ .

(2)  $\rightarrow$  (1): Proposition 3.4.

Corollary 3.6. Let  $X$  be a  $\Lambda_\alpha$ - $R_0$  topological space. For any  $x \in X$  if  $\langle x \rangle = \{x\}$ , then  $\{x\}^{(\Lambda, \alpha)} = \{x\}$ .

Proof. It is a consequence of Corollary 3.5.

Proposition 3.7. For a topological space  $(X, \tau)$ , the following properties are equivalent.

- (1)  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.
- (2)  $x \in \{y\}^{(\Lambda, \alpha)}$  if and only if  $y \in \{x\}^{(\Lambda, \alpha)}$ .

Proof. (1)  $\rightarrow$  (2): Let  $X$  be  $\Lambda_\alpha$ - $R_0$ . Let  $x \in \{y\}^{(\Lambda, \alpha)}$  and  $U$  be any  $(\Lambda, \alpha)$ -open set such that  $y \in U$ . Hence,  $\Lambda_\alpha Ker(\{y\}) \subset U$ . Since  $x \in \{y\}^{(\Lambda, \alpha)}$  and  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_0$ , by Corollary 3.5,  $x \in \Lambda_\alpha Ker(\{y\}) \subset U$ . Therefore, every  $(\Lambda, \alpha)$ -open set which contains  $y$  contains  $x$ . Hence,  $y \in \{x\}^{(\Lambda, \alpha)}$ .

(2)  $\rightarrow$  (1): Let  $U$  be a  $(\Lambda, \alpha)$ -open set and  $x \in U$ . If  $y \notin U$ , then  $x \notin \{y\}^{(\Lambda, \alpha)}$  and hence,  $y \notin \{x\}^{(\Lambda, \alpha)}$ . This implies that  $\{x\}^{(\Lambda, \alpha)} \subset U$ . Hence,  $(X, \tau)$  is a  $\Lambda_\alpha$ - $R_0$  space.

Proposition 3.8. For a topological space  $(X, \tau)$ , the following properties are equivalent.

- (1)  $X$  is a  $\Lambda_\alpha$ - $R_0$  space.
- (2)  $\langle x \rangle = \{x\}^{(\Lambda, \alpha)}$  for each  $x \in X$ .
- (3)  $\langle x \rangle$  is  $(\Lambda, \alpha)$ -closed for each  $x \in X$ .

Proof. (1)  $\rightarrow$  (2): By Corollary 3.5,  $\{x\}^{(\Lambda, \alpha)} = \Lambda_\alpha Ker(\{x\})$  for each  $x \in X$ . Hence,  $\{x\}^{(\Lambda, \alpha)} = \{x\}^{(\Lambda, \alpha)} \cap \Lambda_\alpha Ker(\{x\}) = \langle x \rangle$ .



(2)  $\rightarrow$  (1): Since  $\{x\}^{(\Lambda, \alpha)} = \langle x \rangle$  for each  $x \in X$ , we have  $\{x\}^{(\Lambda, \alpha)} \subset \Lambda_\alpha \text{Ker}(\{x\})$ . By Proposition 3.4(5),  $X$  is  $\Lambda_\alpha$ - $R_0$ .

(2)  $\longleftarrow$  (3): It is consequence of Lemma 2.1.

**4.  $\Lambda_\alpha$ - $R_1$  Topological Spaces and a Question.** A. S. Davis [2] also introduced the notion of  $R_1$  topological spaces which is strictly weaker than  $T_2$ . Now we offer a new generalization of  $R_1$  by utilizing the notions of  $(\Lambda, \alpha)$ -open sets and  $(\Lambda, \alpha)$ -closure operator.

**Definition 7.** A topological space  $(X, \tau)$  is said to be  $\Lambda_\alpha$ - $R_1$  if for  $x, y$  in  $X$  with  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$ , there exists disjoint  $(\Lambda, \alpha)$ -open sets  $U$  and  $V$  such that  $\{x\}^{(\Lambda, \alpha)}$  is a subset of  $U$  and  $\{y\}^{(\Lambda, \alpha)}$  is a subset of  $V$ .

**Proposition 4.1.** If  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_1$ , then  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_0$ .

**Proof.** Let  $U$  be  $(\Lambda, \alpha)$ -open and let  $x \in U$ . If  $y \notin U$ , then since  $x \notin \{y\}^{(\Lambda, \alpha)}$ ,  $\{x\}^{(\Lambda, \alpha)} \neq \{y\}^{(\Lambda, \alpha)}$  and there exists a  $(\Lambda, \alpha)$ -open  $V_y$  such that  $\{y\}^{(\Lambda, \alpha)} \subset V_y$  and  $x \notin V_y$ , which implies  $y \notin \{x\}^{(\Lambda, \alpha)}$ . Thus,  $\{x\}^{(\Lambda, \alpha)} \subset U$ . Hence,  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_0$ .

Recall that a topological space  $(X, \tau)$  is said to be  $R_1$  [2] if for  $x, y$  in  $X$  with  $Cl(\{x\}) \neq Cl(\{y\})$ , there exists disjoint open sets  $U$  and  $V$  such that  $Cl(\{x\})$  is a subset of  $U$  and  $Cl(\{y\})$  is a subset of  $V$ .

**Example 4.2.** Let  $(X, \tau)$  be a topological space such that  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, \{a\}, X\}$ . Clearly, the family of all  $(\Lambda, \alpha)$ -closed sets consists of

$\{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, X\}$ . We have the following for the space  $(X, \tau)$ .

- 1)  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_0$ , but it is not  $R_0$ .
- 2)  $(X, \tau)$  is  $\Lambda_\alpha$ - $R_1$ , but it is not  $R_1$ .

**Question.** Characterize  $\Lambda_\alpha$ - $R_1$  spaces. Is there any example showing that a topological space is  $\Lambda_\alpha$ - $R_0$  but not  $\Lambda_\alpha$ - $R_1$ ?

**Acknowledgment.** The authors are grateful to the referee for his remarkable work which improved the quality of this paper.

#### References

1. M. Caldas, D. N. Georgiou, and S. Jafari, "Study of  $\alpha$ -open Sets and the Related Notions in Topological Spaces (submitted).
2. A. S. Davis, "Indexed Systems of Neighborhoods for General Topological Spaces," *American Mathematical Monthly*, 68 (1961), 886–893.
3. K. K. Dube, "A Note on  $R_0$  Topological Spaces," *Math. Vesnik*, 11 (1974), 203–208.

4. D. W. Hall, S. K. Murphy, and J. Rozycki, "On Spaces Which are Essentially  $T_1$ ," *J. Austr. Math. Soc.*, 12 (1974), 451–455.
5. H. Herlich, "A Concept of Nearness," *Gen. Topol. Appl.*, 4 (1974), 191–212.
6. S. A. Naimpally, "On  $R_0$ -Topological Spaces," *Ann. Univ. Sci. Budapest Eotvos, Sect. Math.* 10 (1967), 53–54.
7. O. Njåstad, "On Some Classes of Nearly Open Sets," *Pacific Journal of Mathematics*, 15 (1965), 961–970.
8. N. A. Shanin, "On Separation in Topological Spaces," *Dokl. Akad. Nauk. SSSR*, 38 (1943), 110–113.

Miguel Caldas  
Departamento de Matematica Aplicada  
Universidade Federal Fluminense  
Rua Mario Santos Braga, s/n  
24020-140, Niteroi, RJ Brasil  
email: gmamccs@vm.uff.br

Esref Hatir  
Department of Mathematics  
Education Faculty  
Selcuk University, 42090  
Konya, Turkey  
email: ehatir@selcuk.edu.tr

Saeid Jafari  
Department of Mathematics and Physics  
Roskilde University, Postbox 260, 4000  
Roskilde, Denmark  
email: sjafari@ruc.dk