

94. On Osima's Blocks of Group Characters

By Kenzo IIZUKA

Department of Mathematics, Kumamoto University, Kumamoto, Japan

(Comm. by Z. SUETUNA, M.J.A., July 12, 1960)

Let \mathcal{G} be a group of finite order g and p be a fixed rational prime. M. Osima, in his earlier paper [4], introduced a concept of blocks of characters with regard to a subgroup \mathfrak{H} of \mathcal{G} (" \mathfrak{H} -blocks"). Let \mathfrak{H}_0 be the maximal normal subgroup of \mathcal{G} contained in \mathfrak{H} . It is well known that the irreducible characters¹⁾ $\phi_1, \phi_2, \dots, \phi_k$ of \mathfrak{H}_0 are distributed into the classes $\mathfrak{B}_1, \mathfrak{B}_2, \dots, \mathfrak{B}_s$ of associated characters in \mathcal{G} . If $\mathfrak{B}'_1, \mathfrak{B}'_2, \dots, \mathfrak{B}'_r$ are the classes of associated irreducible characters of \mathfrak{H}_0 in \mathfrak{H} , then each class \mathfrak{B}_σ is a collection of classes \mathfrak{B}'_ρ . Let $\chi_1, \chi_2, \dots, \chi_n$ be the irreducible characters of \mathcal{G} and $\theta_1, \theta_2, \dots, \theta_h$ be those of \mathfrak{H} . As is well known, there corresponds to each character χ_i exactly one class \mathfrak{B}_σ such that

$$\chi_i(H_0) = s_{i\sigma} \sum_{\phi_\mu \in \mathfrak{B}_\sigma} \phi_\mu(H_0) \quad (H_0 \in \mathfrak{H}_0)$$

where $s_{i\sigma}$ is a positive rational integer. If a class \mathfrak{B}_σ corresponds to a character χ_i in this sense, we say that χ_i belongs to \mathfrak{B}_σ by counting χ_i in \mathfrak{B}_σ . We also say that θ_λ belongs to \mathfrak{B}_σ if θ_λ belongs to \mathfrak{B}'_ρ contained in \mathfrak{B}_σ . Then the classes \mathfrak{B}_σ are the \mathfrak{H} -blocks of \mathcal{G} in Osima's sense. From the definition, we see that χ_i and χ_j belong to the same \mathfrak{H} -block of \mathcal{G} if and only if $\chi_i(H_0)/\chi_i(1) = \chi_j(H_0)/\chi_j(1)$ for all elements H_0 of \mathfrak{H}_0 [4], where 1 denotes the identity of \mathcal{G} .

In the following, "block" of a group will always mean block with regard to a p -Sylow subgroup of the group. While Brauer's blocks for a rational prime q will be referred always as q -blocks. The purpose of this paper is to consider a connection between blocks of \mathcal{G} and the blocks of the normalizer $\mathfrak{N}(R)$ of a p -regular element R in \mathcal{G} .

The author wishes to thank Prof. M. Osima for several helpful suggestions.

1. Let \mathfrak{P} be a p -Sylow subgroup of \mathcal{G} and \mathfrak{P}_0 be the maximal normal p -subgroup of \mathcal{G} . We shall denote by $\mathfrak{B}_1, \mathfrak{B}_2, \dots, \mathfrak{B}_s$ the blocks of \mathcal{G} with regard to \mathfrak{P} . For each \mathfrak{B}_σ we set

$$(1.1) \quad A_\sigma = \sum_{\chi_i \in \mathfrak{B}_\sigma} e_i,$$

where e_i is the primitive idempotent of the center Z of the group ring of \mathcal{G} over the field Ω of g -th roots of unity which belongs to χ_i . Let K_1, K_2, \dots, K_n be the classes of conjugate elements in \mathcal{G} and $G_1,$

1) The term "irreducible character" will always mean absolutely irreducible ordinary character.

G_2, \dots, G_n be a complete system of representatives for the classes. If we interpret each class K_ν as the sum of all its elements, then we may write

$$(1.2) \quad \Delta_\sigma = \sum_\nu a_\nu^\sigma K_\nu,$$

where $a_\nu^\sigma = \frac{1}{g} \sum_{\chi_i \in \mathfrak{B}_\sigma} \chi_i(1) \bar{\chi}_i(G_\nu)$.²⁾ By Frobenius' theorem on induced characters, we have the following:

Lemma 1. 1) $a_\nu^\sigma = 0$ for all classes K_ν , which are not contained in \mathfrak{B}_0 . 2) All $p^{\alpha_\nu} a_\nu^\sigma$ are algebraic integers, where p^{α_ν} is the order of \mathfrak{B}_0 .

The converse of this lemma also holds in the following form: If, for a set \mathfrak{B} of characters χ_i , the idempotent $\Delta = \sum_{\chi_i \in \mathfrak{B}} e_i$ of Z is expressed as a linear combination of classes K_ν , contained in \mathfrak{B}_0 , then \mathfrak{B} is a collection of blocks \mathfrak{B}_ν of \mathfrak{G} .

2. Let q be an arbitrarily fixed rational prime, different from p , and Q be an arbitrarily given element of \mathfrak{G} whose order is a power of q . It follows from Lemma 1 that each block \mathfrak{B}_ν of \mathfrak{G} is a collection of q -blocks B_τ of \mathfrak{G} . Let $B^{(\tau)}(Q)$ be the collection of q -blocks of the normalizer $\mathfrak{N}(Q)$ of Q in \mathfrak{G} which determine a q -block B_τ of \mathfrak{G} . We set $\mathfrak{B}^{(\sigma)}(Q) = \bigcup_{B_\tau \subseteq \mathfrak{B}_\sigma} B^{(\tau)}(Q)$. By (4.16) in [2]³⁾ and Lemma 1, we have the following:

Lemma 2. Each $\mathfrak{B}^{(\sigma)}(Q)$ is a collection of blocks $\hat{\mathfrak{B}}_p$ of $\mathfrak{N}(Q)$.

Let now R be a p -regular element of \mathfrak{G} whose order is a product of powers of distinct rational primes q_1, q_2, \dots, q_r . As is well known, R is decomposed uniquely into

$$(2.1) \quad R = Q_1 Q_2 \cdots Q_r \quad (Q_i Q_j = Q_j Q_i),$$

where Q_i is the q_i -factor of R . Let a block \mathfrak{B}_ν of \mathfrak{G} be given arbitrarily. First, applying Lemma 2 for $Q = Q_1$, \mathfrak{B}_ν and \mathfrak{G} , we have a collection $\mathfrak{B}^{(\sigma)}(Q_1)$ of blocks of $\mathfrak{N}(Q_1)$. Secondly, working similarly for $Q = Q_2$, $\mathfrak{B}_\nu = \mathfrak{B}^{(\sigma)}(Q_1)$ and $\mathfrak{G} = \mathfrak{N}(Q_1)$, we have a collection $\mathfrak{B}^{(\sigma)}(Q_1, Q_2)$ of blocks of $\mathfrak{N}(Q_1 Q_2)$. Continuing this process, we have finally a collection $\tilde{\mathfrak{B}}^{(\sigma)} = \mathfrak{B}^{(\sigma)}(Q_1, Q_2, \dots, Q_r)$ of blocks $\tilde{\mathfrak{B}}_p$ of $\tilde{\mathfrak{G}} = \mathfrak{N}(R)$. If a block $\tilde{\mathfrak{B}}_p$ of $\tilde{\mathfrak{G}}$ belongs to the collection $\tilde{\mathfrak{B}}^{(\sigma)}$, we say that the block \mathfrak{B}_ν of \mathfrak{G} is determined by the block $\tilde{\mathfrak{B}}_p$ of $\tilde{\mathfrak{G}}$. It follows from Theorem 1 in §3 that $\tilde{\mathfrak{B}}^{(\sigma)}$ is independent of the order of Q_1, Q_2, \dots, Q_r .

3. Let R be a p -regular element of \mathfrak{G} and $S(R)$ be the p -regular section ("Oberklasse")⁴⁾ of R in \mathfrak{G} , i.e. the set of all elements of \mathfrak{G}

2) If α is a complex number, we denote by $\bar{\alpha}$ the conjugate complex number of α .

3) Cf. also [7, p. 181].

4) Cf. [9].

whose p -regular factors are conjugate to R in \mathfrak{G} . Let $\tilde{K}_1, \tilde{K}_2, \dots, \tilde{K}_v$ be the classes of conjugate elements in $\tilde{\mathfrak{G}} = \mathfrak{N}(R)$ whose orders are powers of p . We may assume that the maximal normal p -subgroup $\tilde{\mathfrak{F}}_0$ of $\tilde{\mathfrak{G}}$ is the union of the first u classes \tilde{K}_α . We may also assume that $K_\alpha \cong RK_\alpha$, $\alpha = 1, 2, \dots, v$; $S(R)$ is the union of K_1, K_2, \dots, K_v . We denote by $S_0(R)$ the union of K_1, K_2, \dots, K_u and denote by $S_1(R)$ the union of $K_{u+1}, K_{u+2}, \dots, K_v$; $S(R) = S_0(R) \cup S_1(R)$.

Let $\tilde{\mathfrak{B}}_1, \tilde{\mathfrak{B}}_2, \dots, \tilde{\mathfrak{B}}_s$ be the blocks of $\tilde{\mathfrak{G}}$ with regard to a p -Sylow subgroup $\tilde{\mathfrak{P}}$ of $\tilde{\mathfrak{G}}$. Let, for each block $\tilde{\mathfrak{B}}_\sigma$ of $\tilde{\mathfrak{G}}$, Δ_σ be given by (1.1). Similarly, for each block $\tilde{\mathfrak{B}}_\rho$ of $\tilde{\mathfrak{G}}$, we define an idempotent $\tilde{\Delta}_\rho$ of the center \tilde{Z} of the group ring of $\tilde{\mathfrak{G}}$ over Ω . We set $\tilde{\Delta}^{(\sigma)} = \sum_{\tilde{\mathfrak{B}}_\rho \subseteq \tilde{\mathfrak{B}}^{(\sigma)}} \tilde{\Delta}_\rho$, where $\tilde{\mathfrak{B}}^{(\sigma)}$ is the collection of blocks $\tilde{\mathfrak{B}}_\rho$ of $\tilde{\mathfrak{G}}$ which determine $\tilde{\mathfrak{B}}_\sigma$, and set

$$(3.1) \quad K_\mu \Delta_\sigma = \sum_{\nu=1}^n \alpha_{\mu\nu}^\sigma K_\nu \quad (\mu = 1, 2, \dots, n).$$

Then, by Theorem 2 in [3] and Lemmas 1 and 2, we obtain the following:

Theorem 1. *For $\alpha = 1, 2, \dots, u$, we have*

$$K_\alpha \Delta_\sigma = \sum_{\beta=1}^u \alpha_{\alpha\beta}^\sigma K_\beta$$

and

$$\tilde{K}_\alpha \tilde{\Delta}^{(\sigma')} = \sum_{\beta=1}^u \alpha_{\alpha\beta}^\sigma \tilde{K}_\beta.$$

For $\alpha = u+1, u+2, \dots, v$, we have

$$K_\alpha \Delta_\sigma = \sum_{\beta=u+1}^v \alpha_{\alpha\beta}^\sigma K_\beta$$

and

$$\tilde{K}_\alpha \tilde{\Delta}^{(\sigma')} = \sum_{\beta=u+1}^v \alpha_{\alpha\beta}^\sigma \tilde{K}_\beta.$$

Let $\tilde{\chi}_1, \tilde{\chi}_2, \dots, \tilde{\chi}_n$ be the irreducible characters of $\tilde{\mathfrak{G}}$ and $\tilde{\theta}_1, \tilde{\theta}_2, \dots, \tilde{\theta}_{\tilde{h}}$ be those of $\tilde{\mathfrak{P}}$. We set

$$(3.2) \quad \tilde{\chi}_j(P) = \sum_{\lambda=1}^{\tilde{h}} \tilde{r}_{j\lambda} \tilde{\theta}_\lambda(P) \quad (P \in \tilde{\mathfrak{P}})$$

and

$$(3.3) \quad \chi_i(RP) = \sum_{\lambda=1}^{\tilde{h}} r_{i\lambda}^R \tilde{\theta}_\lambda(P) \quad (P \in \tilde{\mathfrak{P}})$$

[5, 6]. Setting

$$(3.4) \quad \tilde{w}_{\lambda\mu} = \sum_{j=1}^{\tilde{n}} \tilde{r}_{j\lambda} \tilde{r}_{j\mu} \quad (\lambda, \mu = 1, 2, \dots, \tilde{h}),$$

by Theorem 1 we have

$$(3.5) \quad \sum_{\lambda_i \in \tilde{\mathfrak{B}}_\sigma} r_{i\lambda}^R \tilde{r}_{i\mu}^R = \begin{cases} \tilde{w}_{\lambda\mu} & (\tilde{\theta}_\lambda, \tilde{\theta}_\mu \in \tilde{\mathfrak{B}}_\rho \subseteq \tilde{\mathfrak{B}}^{(\sigma')}, \\ 0 & (\text{elsewhere}). \end{cases}$$

In particular,

$$\sum_{\chi_i \in \mathfrak{B}_\sigma} r_{i\lambda}^R \bar{r}_{i\lambda}^R = 0 \quad (\tilde{\theta}_\lambda \notin \tilde{\mathfrak{B}}^{(\sigma)}),$$

hence

$$r_{i\lambda}^R = 0 \quad (\chi_i \in \mathfrak{B}_\sigma, \tilde{\theta}_\lambda \notin \tilde{\mathfrak{B}}^{(\sigma)}).$$

Thus we obtain the following theorem.⁵⁾

Theorem 2. *If an irreducible character $\tilde{\theta}_\lambda$ of $\tilde{\mathfrak{F}}$ belongs to a block $\tilde{\mathfrak{B}}_\rho$ of $\tilde{\mathfrak{G}}$, then $r_{i\lambda}^R$ can be different from zero only for irreducible characters χ_i of \mathfrak{G} which belong to the block \mathfrak{B}_σ of \mathfrak{G} determined by the block $\tilde{\mathfrak{B}}_\rho$ of $\tilde{\mathfrak{G}}$.*

By Theorem 1, we also have the following refinements of some of the orthogonality relations for group characters.

Theorem 3. 1) *If two elements L and M of \mathfrak{G} belong to different p -regular sections of \mathfrak{G} , then*

$$(3.6) \quad \sum_{\chi_i \in \mathfrak{B}_\sigma} \chi_i(L) \bar{\chi}_i(M) = 0$$

for each block \mathfrak{B}_σ of \mathfrak{G} [8].⁶⁾

2) *If L and M belong to the same p -regular section $S(R)$ of \mathfrak{G} and if exactly one of the p -factors of them belongs to the maximal normal p -subgroup of $\mathfrak{R}(R)$, then (3.6) also holds for each block \mathfrak{B}_σ of \mathfrak{G} .*

Theorem 4. *If χ_i and χ_j are two irreducible characters of \mathfrak{G} which belong to different blocks of \mathfrak{G} , then*

$$\sum_{G \in S_0(R)} \chi_i(G) \bar{\chi}_j(G) = 0$$

and

$$\sum_{G \in S_1(R)} \chi_i(G) \bar{\chi}_j(G) = 0$$

for each p -regular section $S(R)$ of \mathfrak{G} .⁷⁾

References

- [1] R. Brauer: On blocks of characters of groups of finite order II, Proc. Nat. Acad. Sci. U. S. A., **32**, 215-219 (1946).
- [2] —: Zur Darstellungstheorie der Gruppen endlicher Ordnung II, Math. Zeitschr., **72**, 25-46 (1959).
- [3] K. Iizuka: On Brauer's theorem on sections in the theory of blocks of group characters, forthcoming.
- [4] M. Osima: On the representations of groups of finite order, Math. Jour. Okayama Univ., **1**, 33-61 (1952).

5) Prof. M. Osima has pointed out the fact that the theorem follows also from Theorem 1 in [1].

6) We have a refinement of this result, which is a dual theorem of Theorem 2 in [1].

7) This theorem is an improvement of a result in [8].

- [5] M. Osima: On the induced characters of a group, Proc. Japan Acad., **28**, 243-248 (1952).
- [6] —: On the induced characters of groups of finite order, Math. Jour. Okayama Univ., **3**, 47-64 (1953).
- [7] —: Notes on blocks of group characters, *ibid.*, **4**, 175-188 (1955).
- [8] —: On some properties of group characters II, forthcoming.
- [9] P. Roquette: Arithmetische Untersuchung des Charakterrings einer endlichen Gruppe, Crelles Jour., **190**, 148-168 (1952).