

On the homology of Torelli groups and Torelli spaces

By Toshiyuki AKITA

Department of Applied Mathematics, Fukuoka University, 8-19-1 Nanakuma, Jonan-ku, Fukuoka 814-0180

(Communicated by Heisuke HIRONAKA, M.J.A., Feb. 12, 1999)

The purpose of the present note is to announce our recent results concerning of the rational homology of Torelli groups, Torelli spaces, and certain subgroups of Torelli groups. The detail will appear elsewhere. Let Σ_g be a closed orientable surface of genus $g \geq 2$. Let $\Gamma_{g,r}^n$ be the mapping class group of Σ_g relative to n marked points and r embedded disks. Namely it is the group of connected components of all the orientation-preserving diffeomorphisms of Σ_g which fix marked points and embedded disks pointwise. The action of $\Gamma_{g,r}^n$ on the first integral homology of Σ_g induces a surjective homomorphism

$$\Gamma_{g,r}^n \rightarrow Sp(2g, \mathbf{Z}),$$

where $Sp(2g, \mathbf{Z})$ is the Siegel modular group of degree g . The *Torelli group* $\mathcal{I}_{g,r}^n$ is defined to be its kernel so that we have an extension

$$1 \rightarrow \mathcal{I}_{g,r}^n \rightarrow \Gamma_{g,r}^n \rightarrow Sp(2g, \mathbf{Z}) \rightarrow 1.$$

Henceforth we omit the decorations n and r when they are zero (e.g. $\mathcal{I}_{g,0}^n = \mathcal{I}_g^n$ and $\mathcal{I}_{g,r}^0 = \mathcal{I}_{g,r}$). The Torelli groups are related to characteristic classes of surface bundles as well as the Casson invariants and the finite type invariants of homology 3-spheres (see [2, 13, 14, 15] for instance).

Let $\mathcal{T}_{g,r}^n$ be the Teichmüller space of genus g with $n+r$ marked points and r tangent vectors. Namely it is the space of all conformal structures on Σ_g up to isotopies that fix $n+r$ marked points $\{p_1, \dots, p_{n+r}\}$ pointwise and act trivially on the tangent space $T_{p_i} \Sigma_g$ for $n+1 \leq i \leq n+r$. It is a complex manifold of dimension $3g-3+n+2r$. The mapping class group $\Gamma_{g,r}^n$ acts on $\mathcal{T}_{g,r}^n$ properly discontinuously and the quotient space $\mathcal{T}_{g,r}^n / \Gamma_{g,r}^n$ is identified with the moduli space of curves of genus g with $n+r$ marked points and r tangent vectors. The Torelli group $\mathcal{I}_{g,r}^n$ is torsion-free and hence it acts on $\mathcal{T}_{g,r}^n$ freely so that the quotient space $T_{g,r}^n = \mathcal{T}_{g,r}^n / \mathcal{I}_{g,r}^n$ is

a complex manifold which is called the *Torelli space*. The Torelli space admits a moduli interpretation as well (see [3, 5]). Moreover, since $\mathcal{T}_{g,r}^n$ is contractible, $T_{g,r}^n$ is the classifying space of $\mathcal{I}_{g,r}^n$ so that there is a canonical isomorphism

$$H_*(\mathcal{I}_{g,r}^n, \mathbf{Z}) \cong H_*(T_{g,r}^n, \mathbf{Z}).$$

Torelli spaces are related to the study of the moduli spaces of curves with or without level structure (see [3, 4, 5] for instance).

Little is known about the structure of the homology of Torelli groups and Torelli spaces apart from D. Johnson's several fundamental results obtained in a series of papers [6, 7, 8, 9] (see also [10, 5]). In particular, he proved that \mathcal{I}_g and $\mathcal{I}_{g,1}$ are finitely generated when $g \geq 3$ and computed their first integral homology. As a consequence of his result, $\mathcal{I}_{g,r}^n$ is finitely generated for all $g \geq 3$ and $n, r \geq 0$ (see [3, 5] for the explicit computation of the first integral homology of $\mathcal{I}_{g,r}^n$). On the contrary, A. Miller and D. McCullough [12] showed that \mathcal{I}_2 is not finitely generated. G. Mess [11] showed that \mathcal{I}_2 is a free group on infinitely many generators. D. Johnson and J. Millson showed that $H_3(\mathcal{I}_3, \mathbf{Z})$ contains a free abelian group of infinite rank (cf. [11]). It is known that $H_2(\mathcal{I}_g, \mathbf{Q})$ and $H_2(\mathcal{I}_g^1, \mathbf{Q})$ are nontrivial for $g \geq 3$ (see [14, 4] for precise statements). To date, it is not known whether $\mathcal{I}_{g,r}^n$ is finitely presented for $g \geq 3$. For other results concerning of the homology of Torelli groups and Torelli spaces, see [10, 14, 11, 5] and references therein. We have proved:

Theorem 1. *For $g \geq 7$ and $n, r \geq 0$, the rational homology $H_*(\mathcal{I}_{g,r}^n, \mathbf{Q})$ of the Torelli group $\mathcal{I}_{g,r}^n$ is infinite dimensional over \mathbf{Q} .*

Corollary. *For $g \geq 7$ and $n, r \geq 0$, the rational homology $H_*(T_{g,r}^n, \mathbf{Q})$ of the Torelli space $T_{g,r}^n$ is infinite dimensional over \mathbf{Q} .*

In other words, we have $\dim_{\mathbf{Q}} H_i(T_{g,r}^n, \mathbf{Q}) = \dim_{\mathbf{Q}} H_i(T_{g,r}^n, \mathbf{Q}) = \infty$ for some $i > 1$, since $T_{g,r}^n$ is a finite dimensional complex manifold. D. Johnson [10] asked whether the Torelli space T_g^1 is homotopy

The author is supported by Grand-in-Aid for Encouragement of Young Scientists (No. 09740072), the Ministry of Education, Science, Sports and Culture of Japan.

equivalent to a finite complex. Corollary answers to his question in the negative.

For $n + r \leq 1$, let $\mathcal{K}_{g,r}^n$ be the subgroup of $\mathcal{I}_{g,r}^n$ generated by all the Dehn twists along separating simple closed curves. The groups $\mathcal{K}_{g,r}^n$ are characterized as the kernel of Johnson homomorphisms when $g \geq 3$ [8] and are related to the Casson invariant of homology 3-spheres and secondary characteristic classes of surface bundles through the work of S. Morita [13, 15, 16]. For $g = 2$, \mathcal{K}_2 is equal to \mathcal{I}_2 so that it is a free group on infinitely many generators. In contrast, the group \mathcal{K}_g is not free for $g \geq 3$ and almost nothing is known about the structure of these groups apart from the results of S. Morita [15]: He proved that $H^1(\mathcal{K}_{g,1}, \mathbf{Z})^{\Gamma_{g,1}} \cong \mathbf{Z} \oplus \mathbf{Z}$ and $H^1(\mathcal{K}_g, \mathbf{Z})^{\Gamma_g} \cong \mathbf{Z}$. We have proved:

Theorem 2. *For $g \geq 2$, the rational homology $H_*(\mathcal{K}_g, \mathbf{Q})$, $H_*(\mathcal{K}_g^1, \mathbf{Q})$, and $H_*(\mathcal{K}_{g,1}, \mathbf{Q})$ are infinite dimensional over \mathbf{Q} .*

Acknowledgement. The author thanks to Prof. Shigeyuki Morita for his helpful comments.

References

- [1] T. Akita: Homological infiniteness of Torelli groups (1998) (preprint).
- [2] S. Garoufalidis and J. Levine: Finite type 3-manifold invariants and the structure of the Torelli group. I. *Invent. Math.*, **131**, 541–594 (1998).
- [3] R. Hain: Torelli groups and geometry of moduli spaces of curves. In *Current Topics in Complex Algebraic Geometry* (eds. C. H. Clemens and J. Kollar). MSRI Publications, **28**, Cambridge University Press, Cambridge, 97–143 (1995).
- [4] R. Hain: Infinitesimal presentations of the Torelli groups. *J. Amer. Math. Soc.*, **10**, 597–651 (1997).
- [5] R. Hain and E. Looijenga: Mapping class groups and moduli spaces of curves. *Algebraic geometry—Santa Cruz 1995. Proc. Symp. Pure Math.*, Amer. Math. Soc., Providence, **62**, 97–142 (1997).
- [6] D. Johnson: An abelian quotient of the mapping class group \mathcal{I}_g . *Math. Ann.*, **249**, 225–242 (1980).
- [7] D. Johnson: The structure of the Torelli group I: A finite set of generators for \mathcal{I} . *Ann. of Math.*, **118**, 423–442 (1983).
- [8] D. Johnson: The structure of the Torelli group—II: A characterization of the group generated by twists on bounding curves. *Topology*, **24**, 113–126 (1985).
- [9] D. Johnson: The structure of the Torelli group—III: The abelianization of \mathcal{I} . *Topology*, **24**, 127–144 (1985).
- [10] D. Johnson: A survey of the Torelli group. *Low Dimensional Topology. Contemp. Math.*, **20**, Amer. Math. Soc., Providence, 165–179 (1983).
- [11] G. Mess: The Torelli groups for genus 2 and 3 surfaces. *Topology*, **31**, 775–790 (1992).
- [12] A. Miller and D. McCullough: The genus 2 Torelli group is not finitely generated. *Topology Appl.*, **22**, 43–49 (1986).
- [13] S. Morita: Casson’s invariant for homology 3-spheres and characteristic classes of surface bundles. I. *Topology*, **28**, 305–323 (1989).
- [14] S. Morita: On the structure and the homology of the Torelli group. *Proc. Japan Acad.*, **65A**, 147–150 (1989).
- [15] S. Morita: On the structure of the Torelli group and the Casson invariant. *Topology*, **30**, 603–621 (1991).
- [16] S. Morita: Casson invariant, signature defect of framed manifolds and the secondary characteristic classes of surface bundles. *J. Differential Geom.*, **47**, 560–599 (1997).