

Non-Commutative Ternary Nambu-Poisson Algebras and Ternary Hom-Nambu-Poisson Algebras

Makhlouf A^{1*} and Amri A²

¹University of Haute Alsace, Laboratory of Mathematics, IT and Applications, 4 Rue des Freres Lumiere, France

²Badji Mokhtar University Annaba, BP 12 Department of Maths, Science Faculty, Algeria

Abstract

The main purpose of this paper is to study non-commutative ternary Nambu-Poisson algebras and their Hom-type version. We provide construction results dealing with tensor product and direct sums of two (non-commutative) ternary (Hom-) Nambu-Poisson algebras. Moreover, we explore twisting principle of (non-commutative) ternary Nambu-Poisson algebras along with algebra morphism that lead to construct (non-commutative) ternary Hom-Nambu-Poisson algebras. Furthermore, we provide examples and a 3-dimensional classification of non-commutative ternary Nambu-Poisson algebras.

Keywords: Hom-nambu poisson algebra; Ternary nambu poisson; Non-commutative ternary; n -ary

Introduction

In the 70's, Nambu proposed a generalized Hamiltonian system based on a ternary product, the Nambu-Poisson bracket, which allows to use more than one hamiltonian [1]. More recent motivation for ternary brackets appeared in string theory and M-branes, ternary Lie type structure was closely linked to the super-symmetry and gauge symmetry transformations of the world-volume theory of multiple coincident M2-branes and was applied to the study of Bagger-Lambert theory. Moreover ternary operations appeared in the study of some quarks models. In 1996, quantizations of Nambu-Poisson brackets were investigated [2], it was presented in a novel approach of Zariski, and this quantization is based on the factorization on \mathbb{R} of polynomials of several variables.

The algebraic formulation of Nambu mechanics was discussed [3] and Nambu algebras was studied [4] as a natural generalization of a Lie algebra for higher-order algebraic operations. By definition, Nambu algebra of order n over a field \mathbb{K} of characteristic zero consists of a vector space V over \mathbb{K} together with a \mathbb{K} -multilinear skew-symmetric operation $[\cdot, \dots, \cdot]: \Lambda^n V \rightarrow V$, called the Nambu bracket that satisfies the following generalization of the Jacobi identity. Namely, for any $x_1, \dots, x_{n-1} \in V$ define an adjoint action $ad(x_1, \dots, x_{n-1}): V \rightarrow V$ by $ad(x_1, \dots, x_{n-1})x_n = [x_1, \dots, x_{n-1}, x_n]$, $x_n \in V$. Then the fundamental identity is a condition saying that the adjoint action is a derivation with respect to the Nambu bracket, i.e. for all $x_1, \dots, x_{n-1}, y_1, \dots, y_n \in V$

$$ad(x_1, \dots, x_{n-1})[y_1, \dots, y_n] = \sum_{k=1}^n [y_1, \dots, ad(x_1, \dots, x_{n-1})y_k, \dots, y_n]. \quad (0.1)$$

When $n=2$, the fundamental identity becomes the Jacobi identity and we get a definition of a Lie algebra.

Different aspects of Nambu mechanics, including quantization, deformation and various algebraic constructions for Nambu algebras have recently been studied. Moreover a twisted generalization, called Hom-Nambu algebras, was introduced [5]. This kind of algebras called Hom-algebras appeared as deformation of algebras of vector fields using σ -derivations. The first examples concerned q -deformations of Witt and Virasoro algebras. Then Hartwig, Larsson and Silvestrov introduced a general framework and studied Hom-Lie algebras [6], in which Jacobi identity is twisted by a homomorphism. The corresponding associative algebras, called Hom-associative algebras

were introduced [7]. Non-commutative Hom-Poisson algebras were discussed [8]. Likewise, n -ary algebras of Hom-type were introduced [5,9-13].

We aim in this paper to explore and study non-commutative ternary Nambu-Poisson algebras and their Hom-type version. The paper includes five Sections. In the first one, we summarize basic definitions of (non-commutative) ternary Nambu-Poisson algebras and discuss examples. In the second Section, we recall some basics about Hom-algebra structures and introduce the notion of (non-commutative) ternary Hom-Nambu-Poisson algebra. Section 3 is dedicated to construction of (non-commutative) ternary Hom-Nambu-Poisson algebras using direct sums and tensor products. In Section 4, we extend twisting principle to ternary Hom-Nambu-Poisson algebras. It is used to build new structures with a given ternary (Hom-) Nambu-Poisson algebra and algebra morphism. This process is used to construct ternary Hom-Nambu-Poisson algebras corresponding to the ternary algebra of polynomials where the bracket is defined by the Jacobian. We provide in the last section a classification of 3-dimensional ternary Nambu-Poisson algebras and then compute corresponding Hom-Nambu-Poisson algebras using twisting principle. Notice that a complete classification of 3-dimensional Hom-Nambu-Poisson algebras is difficult to obtain since so far the classification of 3-dimensional Hom-Nambu-Lie algebras is not known.

Ternary (Non-Commutative) Nambu-Poisson Algebra

In the section we review some basic definitions and fix notations. In the sequel, A denotes a vector space over \mathbb{K} , where \mathbb{K} is an algebraically closed field of characteristic zero. Let $\mu: A \times A \rightarrow A$ be a bilinear map, we denote by $\mu^{op}: A \times A \rightarrow A$ the opposite map, i.e., $\mu^{op} = \mu \circ \tau$ where $\tau: A \times A \rightarrow A \times A$ interchanges the two variables.

***Corresponding author:** Makhlouf A, University of Haute Alsace, Laboratory of Mathematics, IT and Applications, 4 Rue des Freres Lumiere, 68093 Mulhouse Cedex, France, Tel: +33 3 89 33 65 00; E-mail: abdenacer.makhlouf@uha.fr

Received October 17, 2014; **Accepted** June 30, 2015; **Published** July 07, 2015

Citation: Makhlouf A, Amri A (2015) Non-Commutative Ternary Nambu-Poisson Algebras and Ternary Hom-Nambu-Poisson Algebras. J Generalized Lie Theory Appl 9: 221. doi:10.4172/1736-4337.1000221

Copyright: © 2015 Makhlouf A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

A ternary algebra is given by a pair (A, m) , where m is a ternary operation on A , that is a trilinear map $m : A \times A \times A \rightarrow A$, which is denoted sometimes by brackets.

Definition 1.1. A ternary Nambu algebra is a ternary algebra $(A, \{, , \})$ satisfying the fundamental identity defined as

$$\{x_1, x_2, \{x_3, x_4, x_5\}\} = \{\{x_1, x_2, x_3\}, x_4, x_5\} + \{x_3, \{x_1, x_2, x_4\}, x_5\} + \{x_3, x_4, \{x_1, x_2, x_5\}\} \quad (1.1)$$

for all $x_1, x_2, x_3, x_4, x_5 \in A$.

This identity is sometimes called Filippov identity or Nambu identity, and it is equivalent to the identity (0.1) with $n=3$.

A ternary Nambu-Lie algebra or 3-Lie algebra is a ternary Nambu algebra for which the bracket is skew-symmetric, that is for all $\sigma \in S_3$, where S_3 is the permutation group,

$$[x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}] = \text{Sgn}(\sigma)[x_1, x_2, x_3].$$

Let A and A' be two ternary Nambu algebras (resp. Nambu-Lie algebras). A linear map $f : A \rightarrow A'$ is a morphism of a ternary Nambu algebras (resp. ternary Nambu-Lie algebras) if it satisfies $f(\{x, y, z\}_A) = \{f(x), f(y), f(z)\}_{A'}$.

Example 1.2. The polynomials of variables x_1, x_2, x_3 with the ternary operation defined by the Jacobian function:

$$\{f_1, f_2, f_3\} = \begin{vmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} \\ \frac{\partial f_3}{\partial x_1} & \frac{\partial f_3}{\partial x_2} & \frac{\partial f_3}{\partial x_3} \end{vmatrix}, \quad (1.2)$$

is a ternary Nambu-Lie algebra.

Example 1.3. Let $V = \mathbb{R}^4$ be the 4-dimensional oriented Euclidian space over \mathbb{R} . The bracket of 3 vectors $\vec{x}, \vec{y}, \vec{z}$ is given by

$$[x, y, z] = \vec{x} \times \vec{y} \times \vec{z} = \begin{vmatrix} x_1 & y_1 & z_1 & e_1 \\ x_2 & y_2 & z_2 & e_2 \\ x_3 & y_3 & z_3 & e_3 \\ x_4 & y_4 & z_4 & e_4 \end{vmatrix},$$

Where $\{e_1, e_2, e_3, e_4\}$ is a basis of \mathbb{R}^4 and $\vec{x} = \sum_{i=1}^3 x_i \bar{e}_i$, $\vec{y} = \sum_{i=1}^3 y_i \bar{e}_i$ and $\vec{z} = \sum_{i=1}^3 z_i \bar{e}_i$. Then $(V, [, , ,])$ is a ternary Nambu - lie algebra.

Now, we introduce the notion of (non-commutative) ternary Nambu-Poisson algebra.

Definition 1.4. A non-commutative ternary Nambu-Poisson algebra is a triple $(A, \mu, \{, , \})$ consisting of a \mathbb{K} -vector space A , a bilinear map $\mu : A \times A \rightarrow A$ and a trilinear map $\{, , \} : A \otimes A \otimes A \rightarrow A$ such that

- (1) (A, μ) is a binary associative algebra,
- (2) $(A, \{, , \})$ is a ternary Nambu-Lie algebra,
- (3) the following Leibniz rule

$$\{x_1, x_2, \mu(x_3, x_4)\} = \mu(x_3, \{x_1, x_2, x_4\}) + \mu(\{x_1, x_2, x_3\}, x_4)$$

holds for all $x_1, x_2, x_3 \in A$.

A ternary Nambu-Poisson algebra is a non-commutative ternary

Nambu-Poisson algebra $(A, \mu, \{, , \})$ for which μ is commutative, then μ is commutative unless otherwise stated.

In a (non-commutative) ternary Nambu-Poisson algebra, the ternary bracket $\{, , \}$ is called Nambu-Poisson bracket.

Similarly, a non-commutative n -ary Nambu-Poisson algebra is a triple $(A, \mu, \{, \dots, \})$

where $(A, \{, \dots, \})$ defines an n -Lie algebra satisfying similar Leibniz rule with respect to μ .

A morphism of (non-commutative) ternary Nambu-Poisson algebras is a linear map that is a morphism of the underlying ternary Nambu-Lie algebras and associative algebras.

Example 1.5. Let $C^\infty(\mathbb{R}^3)$ be the algebra of C^∞ functions on \mathbb{R}^3 and x_1, x_2, x_3 the coordinates on \mathbb{R}^3 . We define the ternary brackets as in (1.2), then $(C^\infty(\mathbb{R}^3), \{, , \})$ is a ternary Nambu-Lie algebra. In addition the bracket satisfies the Leibniz rule: $f g, \{g, f_2, f_3\} = f \{g, f_2, f_3\} + \{f, f_2, f_3\} g$ where $f, g, f_2, f_3 \in C^\infty(\mathbb{R}^3)$ and the multiplication being the point wise multiplication that is $f g(x) = f(x) g(x)$. Therefore, the algebra is a ternary Nambu-Poisson algebra.

This algebra was considered already in 1973 by Nambu [9] as a possibility of extending the Poisson bracket of standard hamiltonian mechanics to bracket of three functions defined by the Jacobian. Clearly, the Nambu bracket may be generalized further to a Nambu-Poisson allowing for an arbitrary number of entries.

In particular, the algebra of polynomials of variables x_1, x_2, x_3 with the ternary operation defined by the Jacobian function in (1.2), is a ternary Nambu-Poisson algebra.

Remark 1.6. The n -dimensional ternary Nambu-Lie algebra of Example 1.3 does not carry a non-commutative Nambu-Poisson algebra structure except that one given by a trivial multiplication.

Hom-type (Non-Commutative) Ternary Nambu-Poisson Algebras

In this section, we present various Hom-algebra structures. The main feature of Hom-algebra structures is that usual identities are deformed by an endomorphism and when the structure map is the identity, we recover the usual algebra structure.

A Hom-algebra (resp. ternary Hom-algebra) is a triple (A, ν, α) consisting of a \mathbb{K} -vector space A , a bilinear map $\nu : A \times A \rightarrow A$ (resp. a trilinear map $\nu : A \times A \times A \rightarrow A$) and a linear map $\alpha : A \rightarrow A$. A Hom-algebra (A, μ, α) is said to be multiplicative if $\alpha \circ \mu = \mu \circ \alpha^{\otimes 2}$ and it is called commutative if $\mu = \mu^{op}$. A ternary Hom-algebra (A, m, α) is said to be multiplicative if $\alpha \circ m = m \circ \alpha^{\otimes 3}$. Classical algebras (resp. ternary algebras) are regarded as Hom-algebras (resp. ternary Hom-algebras) with identity twisting map. We will often use the abbreviation xy for $\mu(x, y)$ when there is no ambiguity. For a linear map $\alpha : A \rightarrow A$, denote by α^n the n -fold composition of n -copies of α , with $\alpha^0 \equiv Id$.

Definition 2.1. A Hom-algebra (A, μ, α) is a Hom-associative algebra if it satisfies the Hom-associativity condition, that is

$$\mu(\alpha(x), \mu(y, z)) = \mu(\mu(x, y), \alpha(z)) \text{ for all } x, y, z \in A.$$

Remark 2.2. When α is the identity map, we recover the classical associativity condition, then usual associative algebras.

Definition 2.3. A ternary Hom-Nambu algebra is a triple $(A, \{, , \}, \tilde{\alpha})$ consisting of a \mathbb{K} -vector space A , a ternary map $\{, , \}$,

$\cdot\} : A \times A \times A \rightarrow A$ and a pair of $\tilde{\alpha} = (\alpha_1, \alpha_2)$ where $\alpha_1, \alpha_2 : A \rightarrow A$, satisfying [5]

$$\begin{aligned} \{\alpha_1(x_1), \alpha_2(x_2), \{x_3, x_4, x_5\}\} &= \{\{x_1, x_2, x_3\}, \alpha_1(x_4), \alpha_2(x_5)\} + \\ \{\alpha_1(x_3), \{x_1, x_2, x_4\}, \alpha_2(x_5)\} &+ \{\alpha_1(x_3), \alpha_2(x_4), \{x_1, x_2, x_5\}\}. \end{aligned} \quad (2.1)$$

We call the above condition the ternary Hom-Nambu identity.

Generally, the n -ary Hom-Nambu algebras are defined by an n -ary bracket and maps $\alpha_1, \dots, \alpha_{n-1}$, satisfying the following Hom-Nambu identity

$$\begin{aligned} \{\alpha_1(x_1), \dots, \alpha_{n-1}(x_{n-1}), \{x_n, \dots, x_{2n-1}\}\} \\ = \sum_{i=1}^{2n-1} \{\alpha_1(x_n), \dots, \alpha_{i-n}(x_{i-1}), \{x_1, \dots, x_{n-1}, x_i\}, \alpha_{i-n+1}(x_{i+1}), \dots, \alpha_{n-1}(x_{2n-1})\} \end{aligned}$$

for all $(x_1, \dots, x_{2n-1}) \in A^{2n-1}$.

Remark 2.4. A Hom-Nambu algebra is a *Hom-Nambu-Lie* algebra if the bracket is skew-symmetric.

We introduce now the definition of non-commutative ternary Hom-Nambu-Poisson algebra in its general form, involving three linear maps. Next, we will discuss the class in which these three maps are equal. This particular case suits to provide a twisting construction.

Definition 2.5. A non-commutative ternary Hom-Nambu-Poisson algebra is a tuple $(A, \mu, \beta, \{., ., .\}, \tilde{\alpha})$ consisting of a vector space A , a ternary operation $\{., ., .\} : A \times A \times A \rightarrow A$, a binary operation $\mu : A \times A \rightarrow A$, a pair of linear maps $\tilde{\alpha} = (\alpha_1, \alpha_2)$

where $\alpha_1, \alpha_2 : A \rightarrow A$, and a linear map $\beta : A \rightarrow A$ such that:

$$(A, \mu, \beta) \text{ is a binary Hom-associative algebra,} \quad (1)$$

$$(A, \{., ., .\}, \tilde{\alpha}) \text{ is a ternary Hom-Nambu-Lie algebra,}$$

$$\{\mu(x_1, x_2), \alpha_1(x_3), \alpha_2(x_4)\} = \mu(\beta(x_1), \{x_2, x_3, x_4\}) + \mu(\{x_1, x_3, x_4\}, \beta(x_2)).$$

The third condition is called Hom-Leibniz identity.

When, all the linear maps are equal $\alpha = \alpha_1 = \alpha_2 = \beta$, we refer to the ternary

Hom-Nambu-Poisson algebra by a quadruple $(A, \mu, \{., ., .\}, \alpha)$.

Remark 2.6. Notice that μ is not assumed to be commutative. When μ is a commutative multiplication, then $(A, \mu, \beta, \{., ., .\}, \tilde{\alpha})$ is said to be a ternary Hom-Nambu-Poisson algebra.

We recover the classical (non-commutative) ternary Nambu-Poisson algebra when $\alpha_1 = \alpha_2 = \beta = Id$.

Similarly, a non-commutative n -ary Hom-Nambu-Poisson algebra is a tuple $(A, \mu, \beta, \{., \dots, .\}, \tilde{\alpha})$ where $(A, \{., \dots, .\}, \tilde{\alpha})$ with linear maps $\tilde{\alpha} = (\alpha_1, \dots, \alpha_{n-1})$ that defines an n -ary Hom-Nambu-Lie algebra satisfying similar Leibniz rule with respect to (A, μ, β) .

In the sequel we will mainly interested in the class of non-commutative ternary Nambu-Poisson algebras where $\alpha = \alpha_1 = \alpha_2 = \beta$.

Definition 2.7. Let $(A, \mu, \{., ., .\}, \alpha)$ be a (non-commutative) ternary Hom-Nambu-Poisson algebra. It is said to be *multiplicative* if

$$\alpha(\{x_1, x_2, x_3\}) = \{\alpha(x_1), \alpha(x_2), \alpha(x_3)\},$$

$$\alpha \circ \mu = \mu \circ \alpha \otimes 2.$$

If in addition α is bijective then it is called *regular*.

Definition 2.8. Let $(A, \mu, \{., ., .\}, \alpha)$ and $(A', \mu', \{., ., .\}', \alpha')$ be two

(non-commutative) ternary Hom-Nambu-Poisson algebras. A linear map $f : A \rightarrow A'$ is a morphism of (non-commutative) ternary Hom-Nambu-Poisson algebras if it satisfies for all $x_1, x_2, x_3 \in A$:

$$f(\{x_1, x_2, x_3\}) = \{f(x_1), f(x_2), f(x_3)\}', \quad (2.2)$$

$$f \circ \mu = \mu' \circ f^{\otimes 2} \quad (2.3)$$

$$f \circ \alpha = \alpha' \circ f. \quad (2.4)$$

It said to be a weak morphism if hold only the two first conditions.

Tensor Product and Direct Sums

In this section we discuss direct sums and define tensor product of ternary (non-commutative) Hom-Nambu-Poisson algebra and a totally Hom-associative sym-metric ternary algebra. In the following, we define a direct sum of two ternary (non-commutative) Hom-Nambu-Poisson algebras.

Theorem 3.1. Let $(A_1, \mu_1, \{., ., .\}_1, \alpha_1)$ and $(A_2, \mu_2, \{., ., .\}_2, \alpha_2)$ be two ternary (non-commutative) Hom-Nambu-Poisson algebras. Let $\mu_{A_1 \oplus A_2}$ be a bilinear map on $A_1 \oplus A_2$ defined for $x_1, y_1, z_1 \in A_1$ and $x_2, y_2, z_2 \in A_2$ by $\mu(x_1 + x_2, y_1 + y_2, z_1 + z_2) = \mu_1(x_1, y_1, z_1) + \mu_2(x_2, y_2, z_2)$, $\{., ., .\}_{A_1 \oplus A_2}$ a trilinear map defined by $\{x_1 + x_2, y_1 + y_2, z_1 + z_2\}_{A_1 \oplus A_2} = \{x_1, y_1, z_1\}_1 + \{x_2, y_2, z_2\}_2$ and $\alpha_{A_1 \oplus A_2}$ a linear map defined by $\alpha_{A_1 \oplus A_2}(x_1 + x_2) = \alpha_1(x_1) + \alpha_2(x_2)$.

Then $(A_1 \oplus A_2, \mu_{A_1 \oplus A_2}, \{., ., .\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2})$ is a ternary (non-commutative) Hom-Nambu-Poisson algebra.

Proof. The commutativity of $\mu_{A_1 \oplus A_2}$ is obvious since μ_1 and μ_2 are commutative. The skew-symmetry of the bracket follows from the skew-symmetry of $\{., ., .\}_1$ and $\{., ., .\}_2$. So it remains to check the Hom-associativity, the Hom-Nambu and the Hom-Leibniz identities. For Hom-associativity identity, we have

$$\begin{aligned} \mu_{A_1 \oplus A_2}(\mu_{A_1 \oplus A_2}(x_1 + x'_1, x_2 + x'_2), \alpha_{A_1 \oplus A_2}(x_3 + x'_3)) \\ = \mu_{A_1 \oplus A_2}(\mu_1(x_1, x_2) + \mu_2(x'_1, x'_2), \alpha_1(x_3) + \alpha_2(x'_3)) \\ = \mu_1(\mu_1(x_1, x_2), \alpha_1(x_3)) + \mu_2(\mu_2(x'_1, x'_2), \alpha_2(x'_3)) \\ = \mu_1(\alpha_1(x_1), \mu_1(x_2, x_3)) + \mu_2(\alpha_2(x'_1), \mu_2(x'_2, x'_3)) \\ = \mu_{A_1 \oplus A_2}(\alpha_1(x_1) + \alpha_2(x'_1), \mu_1(x_2, x_3) + \mu_2(x'_2, x'_3)) \\ = \mu_{A_1 \oplus A_2}(\alpha_{A_1 \oplus A_2}(x_1, x'_1), \mu_{A_1 \oplus A_2}(x_2 + x'_2, x_3 + x'_3)) \end{aligned}$$

Now we prove the Hom-Nambu identity

$$\begin{aligned} \{\alpha_{A_1 \oplus A_2}(x_1 + x'_1), \alpha_{A_1 \oplus A_2}(x_2 + x'_2), \{x_3 + x'_3, x_4 + x'_4, x_5 + x'_5\}_{A_1 \oplus A_2}\}_{A_1 \oplus A_2} \\ = \{\alpha_1(x_1) + \alpha_2(x'_1), \alpha_1(x_2) + \alpha_2(x'_2), \{x_3, x_4, x_5\}_1 + \{x'_3, x'_4, x'_5\}_2\}_{A_1 \oplus A_2} \\ = \{\alpha_1(x_1), \alpha_1(x_2), \{x_3, x_4, x_5\}_1\}_1 + \{\alpha_2(x'_1), \alpha_2(x'_2), \{x'_3, x'_4, x'_5\}_2\}_2 \\ = \{\{x_1, x_2, x_3\}_1, \alpha_1(x_4), \alpha_1(x_5)\}_1 + \{\alpha_1(x_3), \{x_1, x_2, x_4\}_1, \alpha_1(x_5)\}_1 \\ + \{\alpha_1(x_3), \alpha_1(x_4), \{x_1, x_2, x_5\}_1\}_1 + \{\{x'_1, x'_2, x'_3\}_2, \alpha_2(x'_4), \alpha_2(x'_5)\}_2 \\ + \{\alpha_2(x'_3), \{x'_1, x'_2, x'_4\}_2, \alpha_2(x'_5)\}_2 + \{\alpha_2(x'_3), \alpha_2(x'_4), \{x'_1, x'_2, x'_5\}_2\}_2 \\ = \{\{x_1, x_2, x_3\}_1 + \{x'_1, x'_2, x'_3\}_2, \alpha_1(x_4) + \alpha_2(x'_4), \alpha_1(x_5) + \alpha_2(x'_5)\}_{A_1 \oplus A_2} \\ + \{\alpha_1(x_3) + \alpha_2(x'_3), \{x_1, x_2, x_4\}_1 + \{x'_1, x'_2, x'_4\}_2, \alpha_1(x_5) + \alpha_2(x'_5)\}_{A_1 \oplus A_2} \\ + \{\alpha_1(x_3) + \alpha_2(x'_3), \alpha_1(x_4) + \alpha_2(x'_4), \{x_1, x_2, x_5\}_1 + \{x'_1, x'_2, x'_5\}_2\}_{A_1 \oplus A_2} \\ = \{\{x_1 + x'_1, x_2 + x'_2, x_3 + x'_3\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2}(x_4 + x'_4), \alpha_{A_1 \oplus A_2}(x_5 + x'_5)\}_{A_1 \oplus A_2} \\ + \{\alpha_{A_1 \oplus A_2}(x_3 + x'_3), \{x_1 + x'_1, x_2 + x'_2, x_4 + x'_4\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2}(x_5 + x'_5)\}_{A_1 \oplus A_2} \\ + \{\alpha_{A_1 \oplus A_2}(x_3 + x'_3), \alpha_{A_1 \oplus A_2}(x_4 + x'_4), \{x_1 + x'_1, x_2 + x'_2, x_5 + x'_5\}_{A_1 \oplus A_2}\}_{A_1 \oplus A_2} \end{aligned}$$

Finally, for Hom-Leibniz identity we have

$$\begin{aligned} & \{\mu_{A_1 \oplus A_2}(x_1 + x'_1, x_2 + x'_2), \alpha_{A_1 \oplus A_2}(x_3, x'_3), \alpha_{A_1 \oplus A_2}(x_4, x'_4)\}_{A_1 \oplus A_2} \\ &= \{\mu_1(x_1, x_2) + \mu_2(x'_1, x'_2), \alpha_1(x_3) + \alpha_2(x'_3), \alpha_1(x_4) + \alpha_2(x'_4)\}_{A_1 \oplus A_2} \\ &= \{\mu_1(x_1, x_2), \alpha_1(x_3), \alpha_1(x_4)\}_1 + \{\mu_2(x'_1, x'_2), \alpha_2(x'_3), \alpha_2(x'_4)\}_2 \\ &= \mu_1(\alpha_1(x_1), \{x_2, x_3, x_4\}_1) + \mu_1(\{x'_1, x'_3, x'_4\}_1, \alpha_1(x_2)) \\ &+ \mu_2(\alpha_2(x'_1), \{x'_2, x'_3, x'_4\}_2) + \mu_2(\{x'_1, x'_3, x'_4\}_2, \alpha_2(x'_2)) \\ &= \mu_{A_1 \oplus A_2}(\alpha_{A_1 \oplus A_2}(x_1, x'_1), \{x_2 + x'_2, x_3 + x'_3, x_4 + x'_4\}_{A_1 \oplus A_2}) \\ &+ \mu_{A_1 \oplus A_2}(\{x_1 + x'_1, x_3 + x'_3, x_4 + x'_4\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2}(x_2, x'_2)) \end{aligned}$$

This ends the proof.

Proposition 3.2. Let $(A_1, \mu_1, \{., ., .\}_1, \alpha_1)$ and $(A_2, \mu_2, \{., ., .\}_2, \alpha_2)$ be two ternary (non-commutative) Hom-Nambu-Poisson algebras. A linear map $\phi: A_1 \rightarrow A_2$ is a morphism of ternary (non-commutative) Hom-Nambu-Poisson algebras if and only if $\Gamma_\phi \subseteq A_1 \oplus A_2$ is a Hom-Nambu-Poisson subalgebra of

$$(A_1 \oplus A_2, \mu_{A_1 \oplus A_2}, \{., ., .\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2})$$

Where $\Gamma_\phi = \{(x, \phi(x)) : x \in A_1\} \subset A_1 \oplus A_2$.

Proof. Let $\phi: (A_1, \mu_1, \{., ., .\}_1, \alpha_1) \rightarrow (A_2, \mu_2, \{., ., .\}_2, \alpha_2)$ be a morphism of ternary

Hom-Nambu-Poisson algebras.

We have

$$\begin{aligned} \{x_1 + \phi(x_1), x_2 + \phi(x_2), x_3 + \phi(x_3)\}_{A_1 \oplus A_2} &= \{x_1, x_2, x_3\}_1 + \{\phi(x_1), \phi(x_2), \phi(x_3)\}_2 \\ &= \{x_1, x_2, x_3\}_1 + \phi\{x_1, x_2, x_3\}_1 \end{aligned}$$

Then Γ_ϕ is closed under the bracket $\{., ., .\}_{A_1 \oplus A_2}$.

We have also $(\alpha_1 + \alpha_2)(x_1 + \phi(x_1)) = \alpha_1(x_1) + \alpha_2 \circ \phi(x_1) = \alpha_1(x_1) + \phi \circ \alpha_1(x_1)$,

which implies that $(\alpha_1 + \alpha_2)\Gamma_\phi \subseteq \Gamma_\phi$.

Moreover Γ_ϕ is closed under the multiplication indeed

$$\begin{aligned} \mu_{A_1 \oplus A_2}(x_1 + \phi(x_1), x_2 + \phi(x_2)) &= \mu_1(x_1, x_2) + \mu_2(\phi(x_1), \phi(x_2)) \\ &= \mu_1(x_1, x_2) + \phi \circ \mu_1(x_1, x_2) \subseteq \Gamma_\phi \end{aligned}$$

Conversely, if the graph $\Gamma_\phi \subseteq A_1 \oplus A_2$ is a Hom-subalgebra of

$$(A_1 \oplus A_2, \mu_{A_1 \oplus A_2}, \{., ., .\}_{A_1 \oplus A_2}, \alpha_{A_1 \oplus A_2}),$$

Then we have

$$\{\phi(x_1), \phi(x_2), \phi(x_3)\}_2 = \phi\{x_1, x_2, x_3\}_1,$$

and

$$\alpha_1 + \alpha_2(x + \phi(x)) = \alpha_1(x) + \alpha_2 \circ \phi(x) \in \Gamma_\phi$$

$$= \alpha_1(x) + \phi \circ \alpha_1(x).$$

Finally

$$\begin{aligned} \mu_{A_1 \oplus A_2}(x_1 + \phi(x_1), x_2 + \phi(x_2)) &= \mu_1(x_1, x_2) + \mu_2(\phi(x_1), \phi(x_2)) \\ &= \mu_1(x_1, x_2) + \phi \circ \mu_1(x_1, x_2) \subseteq \Gamma_\phi. \end{aligned}$$

Therefore ϕ is a morphism of ternary (non-commutative) Hom-Nambu-Poisson algebras.

Now, we define the tensor product of two ternary Hom-algebras. Moreover, we consider a tensor product of a ternary Hom-Nambu-Poisson algebra and a totally Hom-associative symmetric ternary algebra.

Let $A_1 = (A, m, \alpha)$, where $\alpha = (\alpha_i)_{i=1,2}$ and $A_2 = (A', m', \alpha')$ where $\alpha' = (\alpha'_i)_{i=1,2}$ be two ternary (non-commutative) Hom-algebras of a given he tensor product $A_1 \otimes A_2$ is a ternary Hom-algebra defined by the triple $(A \otimes A', m \otimes m', \alpha \otimes \alpha')$ where $\alpha \otimes \alpha' = (\alpha_i \otimes \alpha'_i)_{i=1,2}$ with

$$m \otimes m'(x_1 \otimes x'_1, x_2 \otimes x'_2, x_3 \otimes x'_3) = m(x_1, x_2, x_3) \otimes m'(x'_1, x'_2, x'_3), \quad (3.1)$$

$$\alpha_i \otimes \alpha'_i(x_1 \otimes x'_1) = \alpha_i(x_1) \otimes \alpha'_i(x'_1), \quad (3.2)$$

Where $x_1, x_2, x_3 \in A$ and $x'_1, x'_2, x'_3 \in A_2$

Recall that (A, m, α) is a totally Hom-associative ternary algebra if

$$m(\alpha_1(x_1), \alpha_2(x_2), m(x_3, x_4, x_5)) = m(\alpha_1(x_1), m(x_2, x_3, x_4), \alpha_2(x_5))$$

$$= m(m(x_1, x_2, x_3), \alpha_1(x_4), \alpha_2(x_5)).$$

for all $x_1 \dots, x_5 \in A$, and the ternary multiplication m is symmetric if

$$m(x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}) = m(x_1, x_2, x_3). \quad (3.3)$$

for all $\sigma \in S_3, x_1, x_2, x_3 \in A$

Lemma 3.3. Let $A_1 = (A, m, \alpha)$ and $A_2 = (A', m', \alpha')$ be two ternary Hom- algebras of given type (Hom-Nambu, totally Hom-associative). If m is symmetric and m' is skew-symmetric then $m \otimes m'$ is skew symmetric.

Proof. Straight forward

Theorem 3.4. Let $(A, \mu, \beta, \{., ., .\}, (\alpha_1, \alpha_2))$ be a ternary (non-commutative) Hom- Nambu-Poisson algebra, $(B, \tau, (\alpha'_1, \alpha'_2))$ be a totally Hom-associative symmetric ternary algebra, and (B, μ', β') be a Hom-associative algebra, then

$$(A \otimes B, \mu \otimes \mu', \beta \otimes \beta', \{., ., .\}_{A \otimes B}, (\alpha_1 \otimes \alpha'_1, \alpha_2 \otimes \alpha'_2))$$

a (non-commutative) ternary Hom-Nambu-Poisson algebra if and only if

$$\tau(\mu'(b_1, b_2), b_3, b_4) = \mu'(b_1, \tau(b_2, b_3, b_4)) = \mu'(\tau(b_1, b_3, b_4), b_2). \quad (3.4)$$

Proof. Since μ and μ' are both Hom-associative multiplication whence a tensor product $\mu \otimes \mu'$ is Hom-associative. Also the commutativity of $\mu \otimes \mu'$, the skew- symmetry of $\{., ., .\}$ and the symmetry of τ simply the skew-symmetry of $\{., ., .\}_{A \otimes B}$.

Therefore, it remains to check Hom-Nambu identity and Hom-Leibniz identity

We have

$$\begin{aligned} LHS &= \{\alpha_1 \otimes \alpha'_1(a_1 \otimes b_1), \alpha_2 \otimes \alpha'_2(a_2 \otimes b_2), \{a_3 \otimes b_3, a_4 \otimes b_4, a_5 \otimes b_5\}_{A \otimes B}\}_{A \otimes B} \\ &= \{\alpha_1(a_1) \otimes \alpha'_1(b_1), \alpha_2(a_2) \otimes \alpha'_2(b_2), \{a_3, a_4, a_5\}_A \otimes \tau(b_3, b_4, b_5)\}_{A \otimes B} \\ &= \underbrace{\{\alpha_1(a_1), \alpha_2(a_2), \{a_3, a_4, a_5\}_A\}}_a \otimes \underbrace{\tau(\alpha'_1(b_1), \alpha'_2(b_2), \tau(b_3, b_4, b_5))}_b, \end{aligned}$$

and

$$RHS = \{\alpha_1 \otimes \alpha'_1(a_1 \otimes b_1, a_2 \otimes b_2, a_3 \otimes b_3)\}_{A \otimes B}, \alpha_1 \otimes \alpha'_1(a_4 \otimes b_4), \alpha_2 \otimes \alpha'_2(a_5 \otimes b_5)\}_{A \otimes B}$$

$$+ \{\alpha_1 \otimes \alpha'_1(a_3 \otimes b_3), \{a_1 \otimes b_1, a_2 \otimes b_2, a_4 \otimes b_4\}_{A \otimes B}, \alpha_2 \otimes \alpha'_2(a_5 \otimes b_5)\}_{A \otimes B}$$

$$+ \{\alpha_1 \otimes \alpha'_1(a_3 \otimes b_3), \alpha_2 \otimes \alpha'_2(a_4 \otimes b_4), \{a_1 \otimes b_1, a_2 \otimes b_2, a_5 \otimes b_5\}_{A \otimes B}\}_{A \otimes B}$$

$$= \{\{a_1, a_2, a_3\}_A \otimes \tau(b_1, b_2, b_3), \alpha_1(a_4) \otimes \alpha'_1(b_4), \alpha_2(a_5) \otimes \alpha'_2(b_5)\}_{A \otimes B}$$

$$+ \{\alpha_1(a_3) \otimes \alpha'_1(b_3), \{a_1, a_2, a_4\}_A \otimes \tau(b_1, b_2, b_4), \alpha_2(a_5) \otimes \alpha'_2(b_5)\}_{A \otimes B}$$

$$+ \{\alpha_1(a_3) \otimes \alpha'_1(b_3), \alpha_2(a_4) \otimes \alpha'_2(b_4), \{a_1, a_2, a_5\}_A \otimes \tau(b_1, b_2, b_5)\}_{A \otimes B}$$

$$= \underbrace{\{\{a_1, a_2, a_3\}_A, \alpha_1(a_4), \alpha_2(a_5)\}}_c \otimes \underbrace{\tau(\tau(b_1, b_2, b_3), \alpha'_1(b_4), \alpha'_2(b_5))}_d$$

$$+ \underbrace{\{\alpha_1(a_3), \{a_1, a_2, a_4\}_A, \alpha_2(a_5)\}}_e \otimes \underbrace{\tau(\alpha'_1(b_3), \tau(b_1, b_2, b_4), \alpha'_2(b_5))}_f$$

$$+ \underbrace{\{\alpha_1(a_3), \alpha_2(a_4), \{a_1, a_2, a_5\}_A\}}_g \otimes \underbrace{\tau(\alpha'_1(b_3), \alpha'_2(b_4), \tau(b_1, b_2, b_5))}_h$$

Using ternary Nambu identity of $\{., ., .\}$ we have $a=c + e + g$, and $b=d=f=h$ using the symmetry of τ and Hom-associativity of μ' , then the left hand side is equal to the right hand side from where the ternary Hom- Nambu identity of bracket $\{., ., .\}_{A \otimes B}$ is verified

For the Hom-Leibniz identity, we have

$$\begin{aligned} LHS &= \{\mu \otimes \mu'(a_1 \otimes b_1, a_2 \otimes b_2), \alpha_1 \otimes \alpha'_1(a_3 \otimes b_3), \alpha_2 \otimes \alpha'_2(a_4 \otimes b_4)\}_{A \otimes B} \\ &= \{\mu(a_1, b_1) \otimes \mu'(a_2, b_2), \alpha_1(a_3) \otimes \alpha'_1(b_3), \alpha_2(a_4) \otimes \alpha'_2(b_4)\}_{A \otimes B} \\ &= \underbrace{\{\mu(a_1, b_1), \alpha_1(a_3), \alpha_2(a_4)\}_A}_{a'} \otimes \underbrace{\{\mu'(a_2, b_2), \alpha'_1(b_3), \alpha'_2(b_4)\}_B}_{b'} \end{aligned}$$

And

$$\begin{aligned} RHS &= \mu \otimes \mu'(\beta \otimes \beta'(a_1 \otimes b_1), \{a_2 \otimes b_2, a_3 \otimes b_3, a_4 \otimes b_4\}_{A \otimes B}) \\ &+ \mu \otimes \mu'(\{a_1 \otimes b_1, a_3 \otimes b_3, a_4 \otimes b_4\}_{A \otimes B}, \beta \otimes \beta'(a_2 \otimes b_2)) \\ &= \mu \otimes \mu'(\beta(a_1) \otimes \beta'(b_1), \{a_2, a_3, a_4\} \otimes \tau(b_2, b_3, b_4)) \\ &+ \mu \otimes \mu'(\{a_1, a_3, a_4\} \otimes \tau(b_1, b_3, b_4), \beta(a_2) \otimes \beta'(b_2)) \\ &= \underbrace{\mu(\beta(a_1), \{a_2, a_3, a_4\})}_{c'} \otimes \underbrace{\mu'(\beta'(b_1), \tau(b_2, b_3, b_4))}_{d'} \\ &+ \underbrace{\mu(\{a_1, a_3, a_4\}, \beta(a_2))}_{e'} \otimes \underbrace{\mu'(\tau(b_1, b_3, b_4), \beta'(b_2))}_{f'} \end{aligned}$$

With Hom-Leibniz identity we have $a' = c' + e'$, and using condition (3.4) we have $b' = d' = f'$, therefore the left hand side is equal to the right hand side and the Hom-Leibniz identity is proved. Then

$$(A \otimes B, \mu \otimes \mu', \beta \otimes \beta', \{., ., .\}_{A \otimes B}, (\alpha_1 \otimes \alpha'_1, \alpha_2 \otimes \alpha'_2))$$

is a (non-commutative) ternary Hom-Nambu-Poisson algebra.

Construction of Ternary Hom-Nambu-Poisson Algebras

In this section, we provide constructions of ternary Hom-Nambu-Poisson algebras using twisting principle.

Theorem 4.1. Let $(A, \mu, \{., ., .\}, \alpha)$ be a (non-commutative) ternary Hom-Nambu-Poisson algebra and $\beta : A \rightarrow A$ be a weak Hom-Nambu-Poisson morphism, then $A_\beta = (A, \{., ., .\}_\beta = \beta \circ \{., ., .\}, \mu\beta = \beta \circ \mu, \beta\alpha)$ is also a ternary (non-commutative) Hom-Nambu-Poisson algebra. Moreover, if A is multiplicative and β is an algebra morphism, then A_β is a multiplicative (non-commutative) Hom-Nambu-Poisson algebra.

Proof. If μ is commutative, then clearly so is μ_β . The rest of the proof applies whether μ is commutative or not. The skew-symmetry follows from the skew-symmetry of the bracket $\{., ., .\}$. It remains to prove Hom-associativity condition, Hom-Nambu-identity and Hom-Leibniz identity. Indeed

$$\begin{aligned} \mu_\beta(\mu_\beta(x, y), \beta\alpha(z)) &= \mu_\beta(\beta(\mu(x, y), \beta\alpha(z))) = \beta^2(\mu(\mu(x, y), \alpha(z))) \\ &= \beta^2(\mu(\alpha(x), \mu(y, z))) = \mu_\beta(\beta\alpha(x), \mu_\beta(y, z)). \end{aligned}$$

We check the Hom-Nambu identity

$$\begin{aligned} \{\beta\alpha(x_1), \beta\alpha(x_2), \{x_3, x_4, x_5\}_\beta\}_\beta &= \beta^2\{\alpha(x_1), \alpha(x_2), \{x_3, x_4, x_5\}\} \\ &= \beta^2(\{\{x_1, x_2, x_3\}, \alpha(x_4), \alpha(x_5)\} + \{\alpha(x_3), \{x_1, x_2, x_4\}, \alpha(x_5)\} \\ &+ \{\alpha(x_3), \alpha(x_4), \{x_1, x_2, x_5\}\}) \\ &= \{\{x_1, x_2, x_3\}_\beta, \beta\alpha(x_4), \beta\alpha(x_5)\}_\beta + \{\beta\alpha(x_3), \{x_1, x_2, x_4\}_\beta, \beta\alpha(x_5)\}_\beta \\ &+ \{\beta\alpha(x_3), \beta\alpha(x_4), \{x_1, x_2, x_5\}_\beta\}_\beta. \end{aligned}$$

Then it remains to show Hom-Leibniz identity

$$\{\mu_\beta(x_1, x_2), \beta\alpha(x_3), \beta\alpha(x_4)\}_\beta = \beta^2(\{\mu(x_1, x_2), \alpha(x_3), \alpha(x_4)\})$$

$$\begin{aligned} &= \beta^2(\mu(\alpha(x_1), \{x_2, x_3, x_4\}) + \mu(\{x_1, x_3, x_4\}, \alpha(x_2))) \\ &= \mu_\beta(\beta\alpha(x_1), \{x_2, x_3, x_4\}_\beta) + \mu_\beta(\{x_1, x_3, x_4\}_\beta, \beta\alpha(x_2)) \end{aligned}$$

Therefore $A_\beta = (A, \{., ., .\}_\beta, \mu_\beta, \beta\alpha)$ is a ternary (non-commutative) Hom-Nambu-Poisson algebra. For the multiplicativity assertion, suppose that A is multiplicative and β is an algebra morphism. We have

$$(\beta\alpha) \circ (\mu_\beta) = \beta\alpha \circ \beta \circ \mu = \mu_\beta \circ \alpha^{\otimes 2} \beta^{\otimes 2} = \mu_\beta \circ (\beta\alpha)^{\otimes 2}$$

And

$$\beta\alpha \circ \{., ., .\}_\beta = \beta\alpha \circ \beta \circ \{., ., .\} = \{., ., .\} \circ (\beta\alpha)^{\otimes 3}$$

Then A_β is multiplicative.

Corollary 4.2. Let $(A, \mu, \{., ., .\}, \alpha)$ be a multiplicative ternary (non-commutative) Hom-Nambu-Poisson algebra. Then

$$A^n = (A, \mu^{(n)} = \alpha^n \circ \mu, \{., ., .\}^{(n)} = \alpha^{(n)} \circ \{., ., .\}, \alpha^{n+1})$$

is a multiplicative (non-commutative) ternary Hom-Nambu-Poisson algebra for each integer $n \geq 0$.

Proof. The multiplicativity of A implies that $\alpha^n : A \rightarrow A$ is a Nambu-Poisson algebra morphism. By Theorem 4.1 $A_{\alpha^n} = A^n$ is a multiplicative ternary (non-commutative) Hom-Nambu-Poisson algebra.

Corollary 4.3. Let $(A, \mu, \{., ., .\})$ be a ternary (non-commutative) Nambu-Poisson algebra and $\beta : A \rightarrow A$ be a Nambu-Poisson algebra morphism. Then

$$A_\beta = (A, \mu_\beta = \beta \circ \mu, \{., ., .\}_\beta = \beta \circ \{., ., .\}, \beta)$$

is a multiplicative (non-commutative) ternary Hom-Nambu-Poisson algebra.

Remark 4.4. Let $(A, \mu, \{., ., .\}, \alpha)$ and $(A', \mu', \{., ., .\}', \alpha')$ be two (non-commutative) ternary Nambu-Poisson algebras and $\beta : A \rightarrow A, \beta' : A' \rightarrow A'$ be ternary Nambu-Poisson algebra endomorphisms. If $\varphi : A \rightarrow A'$ is a ternary Nambu-Poisson algebra morphism that satisfies $\varphi\beta = \beta'\varphi$ then

$$\varphi : (A, \mu_\beta, \{., ., .\}_\beta, \beta\alpha) \rightarrow (A', \mu'_{\beta'}, \{., ., .\}'_{\beta'}, \beta'\alpha')$$

is a (non-commutative) ternary hom-nambu poisson algebra morphism.

Indeed, we have

$$\begin{aligned} \varphi \circ \{., ., .\}_\beta &= \varphi \circ \beta \circ \{., ., .\} = \beta' \circ \varphi \circ \{., ., .\} = \beta' \circ \{., ., .\}' \circ \varphi^3 = \{., ., .\}'_{\beta'} \circ \varphi^3 \\ \varphi \circ \mu_\beta &= \varphi \circ \beta \circ \mu = \beta' \circ \varphi \circ \mu = \beta' \circ \mu' \circ \varphi^2 = \mu'_{\beta'} \circ \varphi^2 \end{aligned}$$

In the sequel, we will construct Hom-type version of the ternary Nambu-Poisson algebra of polynomials of three variables $(\mathbb{R}[x, y, z], \cdot, \{., ., .\})$, defined in Example 1.5. The Poisson bracket of three polynomials is defined in (1.2).

The twisted version is given by a structure of ternary Hom-Nambu-Poisson algebra where $(\mathbb{R}[x, y, z], \cdot, \alpha \circ \{., ., .\}_\alpha = \alpha \circ \{., ., .\}, \alpha)$ where $\alpha : \mathbb{R}[x, y, z] \rightarrow \mathbb{R}[x, y, z]$ is an algebra morphism satisfying for all $f, g \in \mathbb{R}[x, y, z]$

$$\alpha(f \cdot g) = \alpha(f) \cdot \alpha(g)$$

$$\alpha\{f, g, h\} = \{\alpha(f), \alpha(g), \alpha(h)\}.$$

Theorem 4.5. A morphism $\alpha : \mathbb{R}[x, y, z] \rightarrow \mathbb{R}[x, y, z]$ which gives a structure of ternary Hom-Nambu-Poisson algebra $(\mathbb{R}[x, y, z], \cdot, \alpha^\circ, \{\dots\}_\alpha = \alpha^\circ\{\dots\}, \alpha)$ satisfies the following equation:

$$1 - \begin{vmatrix} \frac{\partial \alpha(x)}{\partial x} & \frac{\partial \alpha(x)}{\partial y} & \frac{\partial \alpha(x)}{\partial z} \\ \frac{\partial \alpha(y)}{\partial x} & \frac{\partial \alpha(y)}{\partial y} & \frac{\partial \alpha(y)}{\partial z} \\ \frac{\partial \alpha(z)}{\partial x} & \frac{\partial \alpha(z)}{\partial y} & \frac{\partial \alpha(z)}{\partial z} \end{vmatrix} = 0 \quad (4.1)$$

Proof. Let α be a Nambu-Poisson algebra morphism, then it satisfies for all $f, g, h \in \mathbb{R}[x, y, z]$

$$\begin{aligned} \alpha(f \cdot g) &= \alpha(f) \cdot \alpha(g), \\ \alpha\{f, g, h\} &= \{\alpha(f), \alpha(g), \alpha(h)\}. \end{aligned}$$

The first equality shows that it is sufficient to just set α on x, y and z . For the second equality, we suppose by linearity that

$$\begin{aligned} f(x, y, z) &= x^i y^j z^k, \\ g(x, y, z) &= x^l y^m z^p, \\ h(x, y, z) &= x^q y^r z^s. \end{aligned}$$

Then we can write the second equation as follows

$$\alpha \begin{vmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} & \frac{\partial f}{\partial z} \\ \frac{\partial g}{\partial x} & \frac{\partial g}{\partial y} & \frac{\partial g}{\partial z} \\ \frac{\partial h}{\partial x} & \frac{\partial h}{\partial y} & \frac{\partial h}{\partial z} \end{vmatrix} = \begin{vmatrix} \frac{\partial \alpha(f)}{\partial x} & \frac{\partial \alpha(f)}{\partial y} & \frac{\partial \alpha(f)}{\partial z} \\ \frac{\partial \alpha(g)}{\partial x} & \frac{\partial \alpha(g)}{\partial y} & \frac{\partial \alpha(g)}{\partial z} \\ \frac{\partial \alpha(h)}{\partial x} & \frac{\partial \alpha(h)}{\partial y} & \frac{\partial \alpha(h)}{\partial z} \end{vmatrix},$$

which can be simplified to

$$1 = \begin{vmatrix} \frac{\partial \alpha(x)}{\partial x} & \frac{\partial \alpha(x)}{\partial y} & \frac{\partial \alpha(x)}{\partial z} \\ \frac{\partial \alpha(y)}{\partial x} & \frac{\partial \alpha(y)}{\partial y} & \frac{\partial \alpha(y)}{\partial z} \\ \frac{\partial \alpha(z)}{\partial x} & \frac{\partial \alpha(z)}{\partial y} & \frac{\partial \alpha(z)}{\partial z} \end{vmatrix}. \quad (4.2)$$

Example 4.6. We set polynomials:

$$\begin{aligned} \alpha(x) &= P_1(x, y, z) = \sum_{0 \leq i, j, k \leq d} a_{ijk} x^i y^j z^k, \\ \alpha(y) &= P_2(x, y, z) = \sum_{0 \leq i, j, k \leq d} b_{ijk} x^i y^j z^k, \\ \alpha(z) &= P_3(x, y, z) = \sum_{0 \leq i, j, k \leq d} c_{ijk} x^i y^j z^k, \end{aligned}$$

Where $P_1, P_2, P_3 \in \mathbb{R}[x, y, z]$, and d the largest degree for each variable. We assume that $a_0 = b_0 = c_0 = 0$

Case of polynomials of degree one. We take

$$\begin{aligned} P_1(x, y, z) &= a_1 x + a_2 y + a_3 z, \\ P_2(x, y, z) &= b_1 x + b_2 y + b_3 z, \\ P_3(x, y, z) &= c_1 x + c_2 y + c_3 z \end{aligned}$$

Equation (2.5) becomes

$$1 - \begin{vmatrix} \frac{\partial P_1(x, y, z)}{\partial x} & \frac{\partial P_1(x, y, z)}{\partial y} & \frac{\partial P_1(x, y, z)}{\partial z} \\ \frac{\partial P_2(x, y, z)}{\partial x} & \frac{\partial P_2(x, y, z)}{\partial y} & \frac{\partial P_2(x, y, z)}{\partial z} \\ \frac{\partial P_3(x, y, z)}{\partial x} & \frac{\partial P_3(x, y, z)}{\partial y} & \frac{\partial P_3(x, y, z)}{\partial z} \end{vmatrix} = 0, \quad (4.3)$$

Whence

$$1 - \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = 0. \quad (4.4)$$

The polynomials P_1, P_2 and P_3 are of one of this form

$$P_1(x, y, z) = xa_1 + ya_2 + za_3, P_2(x, y, z) = b_2 y - \frac{z}{a_1 c_2}, P_3(x, y, z) = c_2 y. \quad (1)$$

$$P_1(x, y, z) = a_1 x + a_2 y + a_3 z, P_2(x, y, z) = \frac{1 + a_1 b_2 c_2}{a_1 c_3} y + b_3 z, P_3(x, y, z) = c_2 y + c_3 z \quad (2)$$

$$P_1(x, y, z) = a_1 x + a_2 y + a_3 z, P_2(x, y, z) = b_1 x + \frac{1}{a_2 c_1} z, P_3(x, y, z) = c_1 x \quad (3)$$

$$P_1(x, y, z) = a_1 x + a_2 y + a_3 z, P_2(x, y, z) = \frac{-1 + a_2 b_3 c_1}{a_2 c_3} x + b_3 z, P_3(x, y, z) = c_1 x + c_3 z \quad (4)$$

$$P_1(x, y, z) = \frac{a_2 b_1 c_3 + b_2}{c_3 x} + a_2 y + a_3 z, P_2(x, y, z) = b_1 x + b_2 y + b_3 z, P_3(x, y, z) = c_3 z \quad (5)$$

$$P_1(x, y, z) = \frac{1}{b_2 c_3} x + a_2 y + a_3 z, P_2(x, y, z) = b_2 y + b_3 z, P_3(x, y, z) = c_3 z \quad (6)$$

$$P_1(x, y, z) = a_1 x + \frac{1}{b_1 c_3} y + a_3 z, P_2(x, y, z) = b_1 x + b_3 z, P_3(x, y, z) = c_3 z \quad (7)$$

$$P_1(x, y, z) = a_1 x + a_2 y + \frac{1}{b_1 c_2} z, P_2(x, y, z) = b_1 x, P_3(x, y, z) = c_1 x + c_2 y \quad (8)$$

$$P_1(x, y, z) = a_1 x + \frac{-1}{b_1 c_3 + a_3 c_2 c_3} y + a_3 z, P_2(x, y, z) = b_1 x, P_3(x, y, z) = c_1 x + c_2 y + c_3 z. \quad (9)$$

$$P_1(x, y, z) = \frac{a_2 b_1}{b_2} + \frac{1}{b_2 c_3 - b_2 c_2} x + a_2 y + a_3 z, P_2(x, y, z) = b_1 x + b_2 y + b_3 z, P_3(x, y, z) = \frac{b_2 c_2}{b_2} x + c_2 y + c_3 z \quad (10)$$

$$P_1(x, y, z) = \frac{-c_3 + a_2 c_1 c_2}{b_2 c_2^2} x + a_2 y + a_3 z, P_2(x, y, z) = b_3 z, P_3(x, y, z) = c_1 x + c_2 y + c_3 z \quad (11)$$

$$P_1(x, y, z) = a_1 x + a_2 y + \frac{1}{b_1 c_2 - b_2 c_1} z, P_2(x, y, z) = b_1 x + b_2 y, P_3(x, y, z) = c_1 x + c_2 y \quad (12)$$

$$P_1(x, y, z) = \frac{1 + a_1 b_1 c_1 - a_1 b_1 c_2 - a_2 b_2 c_1 + a_2 b_2 c_2}{b_2 c_1 - b_2 c_2} x + a_2 y + a_3 z, P_2(x, y, z) = b_1 x + b_2 y + b_3 z, P_3(x, y, z) = c_1 x + c_2 y + c_3 z \quad (13)$$

$$P_1(x, y, z) = a_1 x + \frac{b_2}{b_3} (a_1 - \frac{1}{b_2 c_2 - b_2 c_1}) y + a_3 z, P_2(x, y, z) = b_1 x + b_2 y + b_3 z, P_3(x, y, z) = c_1 x + c_2 y + \frac{b_2 c_2}{b_2} z = c_1 x + c_2 y + \frac{b_2 c_2}{b_2} z \quad (14)$$

Particular case of polynomials of degree two. We take one of the polynomials of degree two

$$\begin{aligned} P_1(x, y, z) &= a_1 x + a_2 y + a_3 z \\ P_2(x, y, z) &= b_1 x + b_2 y + b_3 z \\ P_3(x, y, z) &= c_1 x + c_2 y + c_3 z + c_4 x^2 \end{aligned}$$

The polynomials P_1, P_2 and P_3 are of one of this form

$$P_1(x, y, z) = \frac{a_2 b_1}{b_2} + \frac{1}{b_2 c_3 - b_2 c_2} x + a_2 y + \frac{a_2 b_1}{b_2} z, P_2(x, y, z) = b_1 x + b_2 y + b_3 z, P_3(x, y, z) = c_1 x^2 + c_2 y + c_3 z. \quad (1)$$

$$P_1(x, y, z) = a_2x + \frac{a_1b_2}{b_3}y + a_3z, P_2(x, y, z) = b_2y + b_3z, P_3(x, y, z) = c_4x^2 + c_1x + c_2y + \frac{1+b_3c_2}{b_2}z \quad (2)$$

$$P_1(x, y, z) = a_2x + a_2y + a_3z, P_2(x, y, z) = b_2y, P_3(x, y, z) = c_4x^2 + c_1x + c_2y + \frac{1}{a_1b_2}z. \quad (3)$$

$$P_1(x, y, z) = (\frac{a_2b_1}{b_3} - \frac{1}{c_2b_3})x + a_3z, P_2(x, y, z) = b_1x + b_2z, P_3(x, y, z) = c_4x^2 + c_1x + c_2y + c_3z \quad (4)$$

$$P_1(x, y, z) = -\frac{1}{b_3c_2}x + a_3z, P_2(x, y, z) = b_2z, P_3(x, y, z) = c_4x^2 + c_1x + c_2y + c_3z. \quad (5)$$

$$P_1(x, y, z) = a_1x - \frac{1}{b_1c_3}y + a_3z, P_2(x, y, z) = b_1x, P_3(x, y, z) = c_4x^2 + c_1x + c_3z \quad (6)$$

$$P_1(x, y, z) = a_1x + \frac{-1}{b_1c_3} + \frac{a_3c_2}{c_3}y + a_3z, P_2(x, y, z) = b_1x, P_3(x, y, z) = c_4x^2 + c_1x + c_2y + c_3z \quad (7)$$

$$P_1(x, y, z) = a_1x + a_2y + \frac{1}{b_1c_2}z, P_2(x, y, z) = b_1x, P_3(x, y, z) = c_4x^2 + c_1x + c_2y \quad (8)$$

$$P_1(x, y, z) = a_1x + a_2y + a_3z, P_2(x, y, z) = \frac{(1+a_2b_3c_1)}{a_2c_3}x + b_3z, P_3(x, y, z) = c_1x + c_3z \quad (9)$$

Classification

In this section, we provide the classification of 3-dimensional ternary non-commutative Nambu-Poisson algebras. By straightforward calculations and using a computer algebra system we obtain the following result.

Theorem 5.1. Every 3-dimensional ternary Nambu-Lie algebra is isomorphic to the ternary algebra defined with respect to basis $\{e_1, e_2, e_3\}$, by the skew-symmetric bracket defined as

$$\{e_1, e_2, e_3\} = e_1$$

Moreover it defines a 3-dimensional ternary non-commutative Nambu-Poisson algebra $(A, \{., ., .\}, \mu)$ if and only if μ is one of the following non-commutative associative algebra defined as

$$\begin{aligned} \mu_1(e_2, e_1) &= ae_1\mu_1(e_2, e_2) = ae_2\mu_1(e_2, e_3) = ae_3 \\ \mu_1(e_3, e_1) &= be_1\mu_1(e_3, e_2) = be_2\mu_1(e_3, e_3) = be_3, \end{aligned} \quad (1)$$

where a, b are parameters.

The opposite algebra of (1) (2)

The multiplication which are not mentioned are equal to zero.

The first statement of the Theorem is due to Filippov [4,13]. The two families are naturally not isomorphic.

Remark 5.2. The 3-dimensional ternary Nambu-Lie algebra is endowed with a commutative Nambu-Poisson algebra structure only when the multiplication is trivial.

Using the twisting principle described in Theorem 4.1, we obtain the following 3-dimensional non-commutative ternary Hom-Nambu-Poisson algebras.

Proposition 5.3. Any 3-dimensional ternary non-commutative Hom-Nambu-Poisson algebra $(A, \{., ., .\}_\alpha, \mu_\alpha, \alpha)$ obtained by a twisting defined with respect to the basis $\{e_1, e_2, e_3\} = e_1$ by the ternary bracket $\{e_1, e_2, e_3\}_\alpha = ce_1$, where c is a parameter, is one of the following binary Hom-associative algebras defined by μ_α and a corresponding structure map

$$\begin{aligned} \mu_{\alpha_1}(e_2, e_1) &= ace_1, & \mu_{\alpha_1}(e_3, e_1) &= bce_1, \\ \mu_{\alpha_1}(e_2, e_2) &= a(de_1 + e_2), & \mu_{\alpha_1}(e_3, e_2) &= b(de_1 + e_2), \\ \mu_{\alpha_1}(e_2, e_3) &= a(he_1 + ge_2 + e_3), & \mu_{\alpha_1}(e_3, e_3) &= b(he_1 + ge_2 + e_3), \end{aligned} \quad (1)$$

With

$$\alpha_1(e_1) = ce_1, \alpha_1(e_2) = de_1 + e_2, \alpha_1(e_3) = he_1 + ge_2 + e_3.$$

$$\begin{aligned} \mu_{\alpha_2}(e_1, e_2) &= ace_1, & \mu_{\alpha_2}(e_3, e_1) &= bce_1, \\ \mu_{\alpha_2}(e_2, e_2) &= a(de_1 + e_2 + le_3), & \mu_{\alpha_2}(e_3, e_2) &= b(de_1 + e_2 + le_3), \\ \mu_{\alpha_2}(e_2, e_3) &= a(he_1 + e_3), & \mu_{\alpha_2}(e_3, e_3) &= b(he_1 + e_3), \end{aligned} \quad (2)$$

With

$$\alpha_2(e_1) = ce_1, \alpha_2(e_2) = de_1 + e_2 + le_3, \alpha_2(e_3) = he_1 + e_3e_3$$

$$\begin{aligned} \mu_{\alpha_3}(e_2, e_1) &= ace_1, & \mu_{\alpha_3}(e_3, e_1) &= bce_1, \\ \mu_{\alpha_3}(e_2, e_2) &= a(de_1 + fe_2 + \frac{a}{b}(1-f)e_3), & \mu_{\alpha_3}(e_3, e_2) &= bde_1 + bfe_2 + a(1-f)e_3, \\ \mu_{\alpha_3}(e_2, e_3) &= ahe_1 + b(f-1)e_2 + \frac{a(b-ga)}{b}e_3, & \mu_{\alpha_3}(e_3, e_3) &= bhe_1 + \frac{b^2(f-1)}{a}e_2 + (b-ga)e_3, \end{aligned} \quad (3)$$

With

$$\alpha_3(e_1) = ce_1, \alpha_3(e_2) = de_1 + fe_2 + \frac{a}{b}(1-f)e_3, \alpha_3(e_3) = he_1 + \frac{b}{a}(f-1)e_2 + \frac{(b-ga)}{b}e_3$$

$$\begin{aligned} \mu_{\alpha_4}(e_1, e_2) &= ace_1, & \mu_{\alpha_4}(e_2, e_3) &= b(de_1 + e_2), \\ \mu_{\alpha_4}(e_1, e_3) &= bce_1, & \mu_{\alpha_4}(e_3, e_2) &= a(he_1 + ge_2 + e_3), \\ \mu_{\alpha_4}(e_2, e_2) &= a(de_1 + e_2), & \mu_{\alpha_4}(e_3, e_3) &= b(he_1 + ge_2 + e_3), \end{aligned} \quad (4)$$

With

$$\alpha_4(e_1) = ce_1, \alpha_4(e_2) = de_1 + e_2, \alpha_4(e_3) = he_1 + ge_2 + e_3.$$

$$\begin{aligned} \mu_{\alpha_5}(e_1, e_2) &= ace_1, & \mu_{\alpha_5}(e_2, e_3) &= b(de_1 + e_2 + le_3), \\ \mu_{\alpha_5}(e_1, e_3) &= bce_1, & \mu_{\alpha_5}(e_3, e_2) &= a(he_1 + e_3), \\ \mu_{\alpha_5}(e_2, e_2) &= a(de_1 + e_2 + le_3), & \mu_{\alpha_5}(e_3, e_3) &= b(he_1 + e_3) \end{aligned} \quad (5)$$

With

$$\alpha_5(e_1) = ce_1, \alpha_5(e_2) = de_1 + e_2 + le_3, \alpha_5(e_3) = he_1 + e_3$$

$$\begin{aligned} \mu_{\alpha_6}(e_1, e_2) &= ace_1, & \mu_{\alpha_6}(e_2, e_3) &= bde_1 + bfe_2 + a(1-f)e_3, \\ \mu_{\alpha_6}(e_1, e_3) &= bce_1, & \mu_{\alpha_6}(e_3, e_2) &= ahe_1 - b(f-1)e_2 + \frac{a(b-ag)}{b}e_3, \\ \mu_{\alpha_6}(e_2, e_2) &= a(de_1 + fe_2 + \frac{a}{b}(1-f)e_3), & \mu_{\alpha_6}(e_3, e_3) &= bhe_1 - \frac{b^2(f-1)}{a}e_2 + (b-ag)e_3, \end{aligned} \quad (6)$$

With

$$\alpha_6(e_1) = ce_1, \alpha_6(e_2) = de_1 + fe_2 + \frac{a}{b}(1-f)e_3, \alpha_6(e_3) = he_1 + \frac{-b}{a}(f-1)e_2 + \frac{b-ag}{b}e_3$$

$$\begin{aligned} \mu_{\alpha_7}(e_1, e_3) &= ace_1, \\ \mu_{\alpha_7}(e_2, e_3) &= a(de_1 + fe_2 + le_3), \\ \mu_{\alpha_7}(e_3, e_3) &= a(he_1 + ge_2 + \frac{1+g+l}{f}e_3) \end{aligned} \quad (7)$$

With

$$\alpha_7(e_1) = ce_1, \alpha_7(e_2) = de_1 + fe_2 + le_3, \alpha_7(e_3) = he_1 + ge_2 + \frac{1+g+l}{f}e_3$$

$$\begin{aligned} \mu_{\alpha_8}(e_1, e_3) &= ace_1, \\ \mu_{\alpha_8}(e_2, e_3) &= a(de_1 + e_2), \\ \mu_{\alpha_8}(e_3, e_3) &= a(he_1 + ge_2 + e_3) \end{aligned} \quad (8)$$

With

$$\alpha_8(e_1) = ce_1, \alpha_8(e_2) = de_1 + e_2, \alpha_8(e_3) = he_1 + ge_2 + e_3.$$

$$\begin{aligned} \mu_{\alpha_9}(e_1, e_3) &= ace_1, \\ \mu_{\alpha_9}(e_2, e_3) &= a(de_1 - \frac{1}{g}e_3), \\ \mu_{\alpha_9}(e_3, e_3) &= a(he_1 + ge_2 + re_3) \end{aligned} \quad (9)$$

With

$$\alpha_9(e_1) = ce_1, \alpha_9(e_2) = de_1 - \frac{1}{g}e_3, \alpha_9(e_3) = he_1 + ge_2 + re_3$$

References

1. Nambu Y (1973) Generalized hamiltonian mechanics. Phys Rev D 7: 2405-2412.
2. Dito G, Flato M, Sternheimer D, Takhtajan L (1997) Deformation quantization and Nambu mechanics. Commun Math Phys 183: 1-22.
3. Takhtajan L (1994) On foundation of the generalized Nambu mechanics. Comm Math Phys 160: 295-315.
4. Filippov VT (1985) n-Lie algebras. Siberian Math J 26: 879-891.
5. Ataguema H, Makhlof A, Silvestrov S (2009) Generalization of n-ary Nambu algebras and beyond. J Math Phys.
6. Hartwig JT, Silvestrov DLS (2003) Deformations of lie algebras using σ -derivations. Journal of Algebra 295: 314-361.
7. Makhlof A, Silvestrov SD (2008) Hom-algebra structures. Journal of Generalized Lie Theory and Applications 2: 51-64.
8. Yau D (2010) Non-commutative Hom-poisson algebras.
9. Ammar F, Makhlof A Silvestrov S (2010) Ternary q-virasoro-witt Hom-Nambu-Lie algebras. Journal of Physics A: Mathenmetical and Theoretical.
10. Ammar A, Mabrouk S, Makhlof A (2011) Representations and cohomology of n-ary multiplicative Hom-Nambu-Lie algebras. J Geom Physics 61: 1898-1913.
11. Arnlind J, Makhlof A, Silvestrov S (2010) Ternary Hom-Nambu-Lie algebras induced by Hom-Lie algebras, J Math Phys.
12. Yau D (2012) On n-ary Hom-Nambu and Hom-Nambu-Lie algebras. J Geom Phys 62: 506-522.
13. Yau D (2010) A Hom-associative analogue of n-ary Hom-Nambu algebras.

Citation: Makhlof A, Amri A (2015) Non-Commutative Ternary Nambu-Poisson Algebras and Ternary Hom-Nambu-Poisson Algebras. J Generalized Lie Theory Appl 9: 221. doi:[10.4172/1736-4337.1000221](https://doi.org/10.4172/1736-4337.1000221)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>

