

## Some Aspects of Semi-Abelian Homology and Protoadditive Functors

Everaert T and Gran M\*

Institut de Recherche en Mathématique et Physique, Université catholique de Louvain, Chemin du Cyclotron 2, Louvain-la-Neuve, 1348, Belgium

### Abstract

In this note some recent developments in the study of homology in semi-abelian categories are briefly presented. In particular the role of protoadditive functors in the study of Hopf formulae for homology is explained.

**Keywords:** Semiabelian homology; Centralisation functor; Protoadditive functors

### Introduction

The discovery of higher Hopf formulae for the homology of a group, due to Ronald Brown and Graham Ellis [1], has naturally led to some new perspectives in non-abelian homological algebra. An important advance in this area was made by George Janelidze, who found a connection between group homology and categorical Galois theory [2-5], the latter being a wide extension of Alexander Grothendieck's theory [6]. Among other things, this cleared the path for the discovery of higher Hopf formulae for the homology of general algebraic structures [7-14]. Here, a crucial role is played by the so-called *higher order central extensions*, which are the covering morphisms with respect to certain Galois structures induced by a reflection

$$\mathbb{B} \begin{array}{c} \xleftarrow{I} \\ \perp \\ \xrightarrow{U} \end{array} \mathbb{A}, \quad (1)$$

whose left adjoint  $I: \mathbb{A} \rightarrow \mathbb{B}$  is sometimes called the “coefficient functor”. Here,  $\mathbb{A}$  could, for instance, be the variety of groups,  $\mathbb{B}$  its subvariety of abelian groups, and  $I$  the abelianisation functor. In this case, the induced higher order central extensions are related to the Brown-Ellis Hopf-formulae, as explained below. More generally, higher order central extensions can be defined for any semi-abelian category  $\mathbb{A}$  ([15] e.g., the varieties of groups, rings, Lie algebras, (pre)crossed modules, compact groups, or any abelian category) and any Birkhoff subcategory (i.e., a reflective subcategory closed under subobjects and regular quotients)  $\mathbb{B}$  of  $\mathbb{A}$ . When, moreover,  $\mathbb{A}$  has enough projectives, one obtains higher Hopf formulae for the homology induced by the reflector (or, coefficient functor)  $I: \mathbb{A} \rightarrow \mathbb{B}$ .

In order to explicitly determine of the Hopf-Brown-Ellis formulae for homology in specific algebraic contexts, it is crucial to find suitable descriptions of the higher central extensions, as for instance in terms of algebraic conditions using “generalised commutators”. In general, this is a non-trivial problem, about which we are going to say more in what follows.

Note that, in fact, different approaches to obtaining higher Hopf formulae exist, which can be used in different categorical contexts, based on the comonadic homology theory of Barr and Beck [7,8,10,16], on the abstract notion of Galois group [13,17,18], or on the theory of satellites [19]. These methods essentially coincide in the situation described above, namely for  $\mathbb{A}$  a semi-abelian category with enough projectives and  $\mathbb{B}$  a Birkhoff subcategory of  $\mathbb{A}$ .

Assume that we are in this situation. In this case, the reflector  $I: \mathbb{A} \rightarrow \mathbb{B}$  induces a first “centralisation functor”  $I_1$  from the category  $\text{Ext}(\mathbb{A})$  of extensions (i.e., regular epimorphisms) in  $\mathbb{A}$  to the full subcategory

$\text{CExt}_{\mathbb{B}}(\mathbb{A})$  of extensions that are central with respect to  $\mathbb{B}$ :

$$\text{CExt}_{\mathbb{B}}(\mathbb{A}) \begin{array}{c} \xleftarrow{I_1} \\ \perp \\ \xrightarrow{U_1} \end{array} \text{Ext}(\mathbb{A}). \quad (2)$$

This functor  $I_1$  is the left adjoint of the inclusion functor  $U_1$ . The notion of centrality comes from categorical Galois theory and depends on the choice of the Birkhoff subcategory  $\mathbb{B}$ . It is defined in purely categorical terms [10,20].

The centralisation  $I_1(f)$  of an extension  $f$  is given by a quotient

$$\begin{array}{ccc} A & \xrightarrow{\eta_f} & A/[Ker(f), A]_{\mathbb{B}} \\ & \searrow f & \swarrow I_1(f) \\ & & B, \end{array}$$

where  $[Ker(f), A]_{\mathbb{B}}$  may be thought of as a commutator of  $Ker(f)$  and  $A$ , defined relatively to  $\mathbb{B}$ . For instance, in the classical case of the reflection

$$\text{Ab} \begin{array}{c} \xleftarrow{ab} \\ \perp \\ \xrightarrow{U} \end{array} \text{Gp}$$

this relative commutator is simply the group-theoretical commutator of normal subgroups:

$$[Ker(f), A]_{\text{Ab}} = [Ker(f), A].$$

Hence, in this case  $I_1$  is the usual centralisation functor from the category of group extensions to its full subcategory of central extensions in the classical sense.

Remark that the commutator  $[Ker(f), A]$  appears as the denominator in the Hopf formula for the second integral homology group: given a free presentation

$$0 \longrightarrow K \longrightarrow F \longrightarrow G \longrightarrow 0,$$

of a group  $G$ , Hopf's formula tells us that the second homology group is given by the quotient

\*Corresponding author: Gran M, Institut de Recherche en Mathématique et Physique, Université catholique de Louvain, Chemin du Cyclotron 2, Louvain-la-Neuve, 1348, Belgium, Tel: +3210472111; E-mail: [marino.gran@uclouvain.be](mailto:marino.gran@uclouvain.be)

Received October 11, 2015; Accepted November 20, 2015; Published November 27, 2015

Citation: Everaert T, Gran M (2015) Some Aspects of Semi-Abelian Homology and Protoadditive Functors. J Generalized Lie Theory Appl 9: 238. doi:10.4172/1736-4337.1000238

Copyright: © 2015 Everaert T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

$$H_2(G, \mathbb{Z}) = \frac{K \cap [F, F]}{[K, F]}$$

This is not a coincidence: a similar phenomenon occurs for the higher-order homology groups, where the subgroups appearing as denominator of the Brown-Ellis Hopf formulae are exactly what is required to transform higher-order extensions into a higher-order central extensions, universally.

To illustrate this idea, consider the case of the *third* homology group. For this, let

$$\begin{array}{ccc} F & \longrightarrow & F/K_2 \\ \downarrow & & \downarrow \\ F/K_1 & \longrightarrow & G \cong F/K_1 \cdot K_2 \end{array} \quad (3)$$

be a double presentation of a group  $G$  so that  $F, F/K_1$  and  $F/K_2$  are free groups and the square is a *double extension*: a pushout of surjective homomorphisms. As shown by Brown and Ellis, the third integral homology group  $H_3(G, \mathbb{Z}) (=H_3(G, \text{Ab}))$  of  $G$  is given by

$$H_3(G, \mathbb{Z}) = \frac{[F, F] \cap K_1 \cap K_2}{[K_1 \cap K_2, F] \cdot [K_1, K_2]} \quad (4)$$

Once more, the denominator gives precisely the normal subgroup by which one has to “quotient out” the group  $F$  in order to make this double extension a double central extension, universally:

$$\begin{array}{ccc} F & & \\ \eta_f^2 \searrow & & \\ & \frac{F}{[K_1 \cap K_2, F] \cdot [K_1, K_2]} & \longrightarrow F/K_2 \\ & \downarrow & \downarrow \\ & F/K_1 & \longrightarrow G \end{array}$$

Once again, the notion of centrality comes from categorical Galois theory, and this time depends on the induced reflection (2).

Now, the formula (4) is a special case of the general Hopf formula for the third homology corresponding to a reflection (1), with  $\mathbb{A}$  an arbitrary semi-abelian category with enough projectives, and  $\mathbb{B}$  any Birkhoff subcategory of  $\mathbb{A}$  [10]: starting from a double presentation  $f$  of the form (3), with  $F, F/K_1$  and  $F/K_2$  regular projective objects of  $\mathbb{A}$ , the third homology object is given by a quotient

$$H_3(G, \mathbb{B}) = \frac{[F, F]_{\mathbb{B}} \cap K_1 \cap K_2}{L_2[f]_{\mathbb{B}}} \quad (5)$$

As in the case of groups, also in a general semi-abelian category  $\mathbb{A}$  the denominator  $L_2[f]_{\mathbb{B}}$  of this generalised Hopf formula relative to  $\mathbb{B}$  is the normal subobject of  $F$  that has to be “quotiented out” in order to universally turn the double extension (3) into a double central extension. Hence, in particular, for  $\mathbb{A} = \text{Gp}$  and  $\mathbb{B} = \text{Ab}$ , we have the equality

$$L_2[f]_{\text{Ab}} = [K_1 \cap K_2, F] \cdot [K_1, K_2],$$

and the formula (4) appears as a special case of (5). In general, for a given Birkhoff reflection (1) in a semi-abelian category,  $L_2[f]_{\mathbb{B}}$  may be difficult to compute.

Similar formulae exist for the higher homology objects, again valid in any semi-abelian category  $\mathbb{A}$  with enough projectives and for any Birkhoff subcategory  $\mathbb{B}$  of  $\mathbb{A}$ . This yields, at least in principle, a description of all homology objects  $H_n(\mathbb{A}, \mathbb{B}) (n \geq 2)$ . In practice, a suitable characterisation of the higher-order central extensions is required.

In some cases it has been possible to compute these formulae explicitly. For example, this has been done in [10] for  $\mathbb{A}$  the variety of precrossed modules and  $\mathbb{B}$  its subvariety of crossed modules, or for  $\mathbb{A}$  the variety of groups and  $\mathbb{B}$  its subvariety of nilpotent groups of a fixed class  $k \geq 1$  [21], or the variety of solvable groups of a fixed class  $k \geq 1$ . Similar results have been obtained in the categories of Leibniz and of Lie  $n$ -algebras in [22].

However, in general, computing the Hopf formulae explicitly is a non-trivial task. One possible strategy is to look for suitable conditions on the coefficient functor  $I: \mathbb{A} \rightarrow \mathbb{B}$  that facilitate such computations. In [12,14] we have shown that a natural such condition is the requirement that the reflector  $I$  is a *protoadditive functor*. This notion extends the one of additive functor to the non-additive context of pointed protomodular categories [23]: when  $\mathbb{A}$  and  $\mathbb{B}$  are pointed protomodular categories (for instance,  $\mathbb{A}$  and  $\mathbb{B}$  could be semi-abelian), a functor  $I: \mathbb{A} \rightarrow \mathbb{B}$  is *protoadditive* if, for any split short exact sequence

$$0 \longrightarrow K \xrightarrow{k} A \xleftarrow{s} B \longrightarrow 0$$

in  $\mathbb{A}$  (i.e.,  $f \circ s = 1_B$  and  $k = \text{Ker}(f)$ ), its image

$$0 \longrightarrow I(K) \xrightarrow{I(k)} I(A) \xleftarrow{I(s)} I(B) \longrightarrow 0$$

under  $I$  is a split short exact sequence in  $\mathbb{B}$ . Whenever the coefficient functor  $I: \mathbb{A} \rightarrow \mathbb{B}$  is protoadditive, explicit Hopf formulae can be established in different algebraic and topological contexts. In particular, the protoadditivity condition is fundamental to explore some new Galois theories induced by torsion theories [24-26].

We refer the interested reader to the articles [12,14] for a thorough study of the theory of protoadditive functors and their use in semi-abelian homological algebra. In the first article we study the homology of  $n$ -fold internal groupoids in a semi-abelian category: these results apply in particular to the so-called *cat<sup>n</sup>*-groups in the sense of Loday [27]. The crucial point there is that the connected components functor  $\pi_0: \text{Grpd}(\mathbb{A}) \rightarrow \mathbb{A}$  is protoadditive whenever  $\mathbb{A}$  is semi-abelian. In [14] the general theory of protoadditive functors is investigated, as well as the so-called *derived torsion theories* of a torsion theory having the reflector to its torsion-free subcategory *protoadditive*. These derived torsion theories induce a chain of Galois structures in the categories of higher order extensions. The results concerning the homology objects can be applied in particular to some new torsion theories in the category of compact groups and, more generally, in any category of compact semi-abelian algebras introduced in [28]. Further developments in this direction, including some new results in the categories of commutative rings and of topological groups, for instance, can be found in [13,18].

#### Acknowledgement

This article is partly based on the text of a talk given by the second author on this joint work at the *Séminaire Itinérant des Catégories* that took place at the Université Paris Diderot on October 25, 2009.

#### References

1. Brown R, Ellis G (1988) Hopf formulae for the higher homology of a group. Bull Lond Math Soc 20: 124-128.

2. Janelidze G (1990) Pure Galois theory in categories. *J Algebra* 132: 270-286.
3. Janelidze G (1991) What is a double central extension? (The question was asked by Ronald Brown). *Cah. Topologie Géom. Différ Catég* 32: 191-201.
4. Janelidze G (1995) Higher dimensional central extensions: A categorical approach to homology theory of groups. Lecture at the International Category Theory Meeting CT95.
5. Borceux F, Janelidze G (2001) Galois theories. Cambridge University Press, Cambridge Studies in Adv Mathematics 72.
6. Grothendieck A (1971) Stalls coverings and fundamental group. SGA1 exposed V player. Notes in Math., Springer 224.
7. Everaert T, Van der Linden T (2004) Baer invariants in semi-abelian categories I: General Theory. *Theory Appl Categ* 12: 1-33.
8. Everaert T, Van der Linden T (2004) Baer invariants in semi-abelian categories II: Homology. *Theory Appl Categ* 12: 195-224.
9. Everaert T, Gran M (2007) On low-dimensional homology in categories, Homology. Homotopy and Applications 9: 275-293.
10. Everaert T, Gran M, Van der Linden T (2008) Higher Hopf Formulae for Homology via Galois Theory. *Adv Math* 217: 2231-2267.
11. Everaert T (2010) Higher central extensions and Hopf formulae. *J Algebra* 324: 1771-1789.
12. Everaert T, Gran M (2010) Homology of  $n$ -fold groupoids. *Theory Appl Categ* 23: 22-41.
13. Duckerts M, Everaert T, Gran M (2012) A description of the fundamental group in terms of commutators and closure operators. *J Pure Appl Algebra* 216: 1837-1851.
14. Everaert T, Gran M (2015) Protoadditive functors, derived torsion theories and homology. *J Pure Appl Algebra* 219: 3629-3676.
15. Janelidze G, Márki L, Tholen W (2002) Semi-abelian categories. *J Pure Appl Algebra* 168: 367-386.
16. Barr M, Beck J (1969) Homology and standard constructions. Seminar on Triples and Categorical Homology Theory, Lectures Notes in Math 80: 245-335.
17. Janelidze G (2008) Galois groups, abstract commutators and Hopf formula. *Appl Categ Struct* 16: 653-668.
18. Duckerts-Antoine M (2015) Fundamental group functors in descent-exact homological categories. Preprint 15-38.
19. Goedecke J, Van der Linden T (2009) On satellites in semi-abelian categories: Homology without projectives. *Math. Proc. Cambridge Philos Soc* 147: 629-657.
20. Janelidze G, Kelly GM (1994) Galois theory and a general notion of central extension. *J Pure Appl Algebra* 97: 135-161.
21. Donadze G, Inassaridze N, Porter T (2005)  $n$ -fold Cech derived functors and generalised Hopf type formulas. *K-Theory* 35: 341-373.
22. Casas JM, Khmaladze E, Ladra M, Van Der Linden T (2011) Homology and central extensions of Leibniz and Lie  $n$ -algebras, Homology. Homotopy and Applications 13: 59-74.
23. Bourn D (1991) Normalization equivalence, kernel equivalence and affine categories. *Lecture Notes Math* 1488: 43-62.
24. Bourn D, Gran M (2006) Torsion theories in homological categories. *J Algebra* 305: 18-47.
25. Gran M, Janelidze G (2009) Normal coverings and monadic extensions of Galois structures associated with torsion theories. *Cah. Topologie Géom Différ Catég* 3: 171-188.
26. Gran M, Rossi V (2007) Torsion theories and Galois coverings of topological groups. *J Pure Appl Algebra* 208: 135-151.
27. Loday JL (1982) Spaces with finitely many non-trivial homotopy groups. *J Pure Appl Algebra* 24: 179-202.
28. Borceux F, Clementino MM (2005) Topological semi-abelian algebras. *Adv in Math* 190: 425-453.

**Citation:** Everaert T, Gran M (2015) Some Aspects of Semi-Abelian Homology and Protoadditive Functors. *J Generalized Lie Theory Appl* 9: 238.  
doi:10.4172/1736-4337.1000238

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>