

On algebraic curves for commuting elements in q -Heisenberg algebras

Johan RICHTER and Sergei SILVESTROV

Centre for Mathematical Sciences, Lund University, P.O. Box 118,
SE-221 00 Lund, Sweden

E-mails: johanr@maths.lth.se, ssilvest@maths.lth.se

Abstract

In the present article we continue investigating the algebraic dependence of commuting elements in q -deformed Heisenberg algebras. We provide a simple proof that the 0-chain subalgebra is a maximal commutative subalgebra when q is of free type and that it coincides with the centralizer (commutant) of any one of its elements different from the scalar multiples of the unity. We review the Burchnell-Chaundy-type construction for proving algebraic dependence and obtaining corresponding algebraic curves for commuting elements in the q -deformed Heisenberg algebra by computing a certain determinant with entries depending on two commuting variables and one of the generators. The coefficients in front of the powers of the generator in the expansion of the determinant are polynomials in the two variables defining some algebraic curves and annihilating the two commuting elements. We show that for the elements from the 0-chain subalgebra exactly one algebraic curve arises in the expansion of the determinant. Finally, we present several examples of computation of such algebraic curves and also make some observations on the properties of these curves.

2000 MSC: 16S99, 81S05, 39A13

1 Introduction

In 1994, one of the authors of the present paper, S. Silvestrov, based on consideration of the previous literature and a series of trial computations, made the following three-part conjecture.

- The first part of the conjecture stated that the Burchnell-Chaundy-type result on algebraic dependence of commuting elements can be proved in greater generality, that is, for much more general classes of noncommutative algebras and rings than the Heisenberg algebra and related algebras of differential operators treated by Burchnell and Chaundy and in subsequent literature [1, 2, 3, 7, 8, 11].
- The second part stated that the Burchnell-Chaundy eliminant construction of annihilating algebraic curves formulated in determinant (resultant) form works after some appropriate modifications for most or possibly all classes of algebras where the Burchnell-Chaundy-type result on algebraic dependence of commuting elements can be proved.
- Finally, the third part of the conjecture stated that the proof of the vanishing of the corresponding determinant algebraic curves on the commuting elements can be performed in a purely algebraic way for all classes of algebras or rings where this fact is true, that is, using only the internal structure and calculations with the elements in the corresponding algebras or rings and the algebraic combinatorial expansion formulas for the corresponding determinants without any need of passing to operator representations and use of analytic methods as in the Burchnell-Chaundy-type proofs.

This third part of the conjecture remains widely open with no general such proofs available for any classes of algebras and rings, even in the case of the usual Heisenberg algebra and differential operators, and with only a series of examples calculated for the Heisenberg algebra, q -Heisenberg algebra, and some more general algebras, all supporting the conjecture. In the first and second parts of the conjecture progress has been made. In [4], the key Burchnell-Chaundy-type theorem on algebraic dependence of commuting elements in q -deformed Heisenberg algebras (and thus as a corollary for q -difference operators as operators representing q -deformed Heisenberg algebras) was obtained. The result and the methods have been extended to more general algebras and rings generalizing q -deformed Heisenberg algebras (generalized Weyl structures and graded rings) in [5]. The proof in [4] is totally different from the Burchnell-Chaundy-type proof. It is an existence argument based only on the intrinsic properties of the elements and internal structure of q -deformed Heisenberg algebras, thus supporting the first part of the conjecture. It can be used successfully for an algorithmic implementation for computing the corresponding algebraic curves for given commuting elements. However, it does not give any specific information on the structure or properties of such algebraic curves or any general formulae. It is thus important to have a way of describing such algebraic curves by some explicit formulae, as, for example, those obtained using the Burchnell-Chaundy eliminant construction for the $q = 1$ case, that is, for the classical Heisenberg algebra. In [10], a step in that direction was taken by offering a number of examples, all supporting the claim that the eliminant determinant method should work in the general case. However, no general proof for this was provided. The complete proof following the Burchnell-Chaundy approach in the case of q not a root of unity has been recently obtained [6], by showing that the determinant eliminant construction, properly adjusted for the q -deformed Heisenberg algebras, gives annihilating curves for commuting elements in the q -deformed Heisenberg algebra when q is not a root of unity, thus confirming the second part of the conjecture for these algebras. That proof was obtained by adapting the Burchnell-Chaundy eliminant determinant method of the case $q = 1$ of differential operators to the q -deformed case, after passing to a specific faithful representation of the q -deformed Heisenberg algebra on Laurent series and then performing a detailed analysis of the kernels of arbitrary operators in the image of this representation. While exploring the determinant eliminant construction of the annihilating curves, we also obtained some further information on such curves and some other results on dimensions and bases in the eigenspaces of the q -difference operators in the image of the chosen representation of the q -deformed Heisenberg algebra. Recently, a further extension of Burchnell-Chaundy eliminant determinant method to the context of σ -derivations and Ore extension rings has been considered in [9]. In the case of q being a root of unity the algebraic dependence of commuting elements holds only over the center of the q -deformed Heisenberg algebra [4], and it is still unknown how to modify the eliminant determinant construction to yield annihilating curves for this case.

In the present article we continue investigation of the algebraic dependence of commuting elements in q -deformed Heisenberg algebras within the context of [4, 6, 10]. In Section 2, following [4], we recall some preliminaries on q -deformed Heisenberg algebra, including degree function, decomposition into the direct sum of the “chain” subspaces indexed by the integers and corresponding to this decomposition the upper and lower chain functions. In Section 3, we consider in more detail the 0-chain subspace (indexed by zero). This subspace is a commutative subalgebra in the q -deformed Heisenberg algebra playing a pivotal role for the structure of this algebra [4]. We provide a simple proof that this subalgebra is a maximal commutative subalgebra when q is of free type, and that it coincides with the centralizer (commutant) of any one of its elements different from the scalar multiples of the

unity. In Section 4, we review the Burchall-Chaundy-type construction for proving algebraic dependence and obtaining corresponding algebraic curves for commuting elements in the q -deformed Heisenberg algebra following [6] but putting it into general context of the elements of the q -deformed Heisenberg algebra rather than operators of a specific representation. The construction is based on computing a certain determinant of a matrix with entries depending on two commuting variables and containing one of the generators of the q -deformed Heisenberg algebra. This matrix is constructed from commuting elements. The coefficients in front of the powers of the generator in the expansion of the determinant are polynomials in the two variables defining some algebraic curves. The commuting elements satisfy the equations of these algebraic curves [6]. In Section 5, we show that for the elements from the 0-chain subalgebra exactly one algebraic curve arises via this construction in the expansion of the determinant and then present several examples of computations of such algebraic curves and also make some observations on the properties of these curves based on these examples and further computer experiments.

2 Preliminaries

Let K be a field of characteristic 0, and q a nonzero element of K . We say that q is of free type if it is 1 or not a root of unity. If q is a root of unity, we say it is of torsion type. We define the q -deformed Heisenberg algebra over K as

$$H(q) = K\langle A, B \rangle / (AB - qBA - I)$$

The identity element will be denoted by I . For $q = 1$ we recover the classical Heisenberg algebra (called also Weyl algebra). One can define degree functions \deg_A and \deg_B with respect to A and B on $H(q)$ just as on the commutative algebra of polynomials. One computes these functions by inspection just as one would in a commutative algebra. That the functions are well defined and does not depend on how the elements are written is proved in [4, Chapter 4]. We also define the total degree function $\deg(\alpha) = \deg_A(\alpha) + \deg_B(\alpha)$. In [4, Chapter 4] the following theorem is proved.

Theorem 2.1. *Let $\alpha, \beta \in H(q)$ for some $q \neq 0$ and let $V \in \{A, B\}$. Then*

$$\deg_V(\alpha\beta) = \deg_V(\alpha) + \deg_V(\beta)$$

We define the sets R_n for all integers n by

$$R_n = \left\{ \sum_{\substack{j \geq 0 \\ j+n \geq 0}} a_j B^{j+n} A^j \mid a_j \in K, a_j \neq 0 \text{ for at most finitely many } j \right\}$$

If the element $\alpha \in H(q)$ belongs to some R_n , we say that it is homogeneous. We also define a function

$$\chi : \{ \alpha \in H(q) \mid \alpha \text{ is nonzero and homogeneous} \} \longrightarrow \mathbb{Z}$$

by defining $\chi(\alpha)$ to be the unique integer such that $\alpha \in R_{\chi(\alpha)}$. This function is called the chain function.

All R_i are vector spaces over K . Further $H(q)$ is the direct sum of all the R_i . We can use this to define a projection operation. Let α be an element of $H(q)$. We can write $\alpha = \sum_j \alpha_j$,

where $\alpha_j \in R_j$. This decomposition is unique. We then define the projection of α on R_n by $\alpha \sqcap R_n = \alpha_n$. The notation is intended to recall the notation for intersection. At this point we define two new functions. They are defined for all nonzero elements of $H(q)$.

$$\bar{\chi}(\alpha) = \max \{n \in \mathbb{Z} \mid \alpha \sqcap R_n \neq 0\}, \quad \underline{\chi}(\alpha) = \min \{n \in \mathbb{Z} \mid \alpha \sqcap R_n \neq 0\}$$

These functions are known as the upper and lower chain functions, respectively.

3 R_0 is maximal commutative

We begin by noting that all elements of R_0 commute with each other [4]. Furthermore, the products of two elements $\alpha, \beta \in R_0$ are in R_0 . So R_0 is a commutative subalgebra. We want to show that it is in fact a maximal commutative subalgebra.

For an element $\alpha \in H(q)$ we define $Cen(\alpha) = \{\beta \in H(q) \mid [\alpha, \beta] = 0\}$.

In [4, Chapter 6] the following theorem is proved (as a part of Theorem 6.10).

Theorem 3.1. *Let q be of free type. Let α, β be two commuting elements in $H(q)$. Then the following is true:*

- If $\bar{\chi}(\alpha) = 0$, then $\bar{\chi}(\beta) = 0$ or $\alpha \sqcap R_0 = cI$ for some $c \in K$.
- If $\underline{\chi}(\alpha) = 0$, then $\underline{\chi}(\beta) = 0$ or $\alpha \sqcap R_0 = cI$ for some $c \in K$.

We now describe the centralizer of an element in R_0 .

Theorem 3.2. *Let q be of free type and $\alpha \in R_0 \subset H(q)$. Assume further that $\alpha \neq cI$ for all $c \in K$. Then $Cen(\alpha) = R_0$.*

Proof. As we noted above $R_0 \subseteq Cen(\alpha)$. It remains to show the other inclusion. Let β be an arbitrary nonzero element of $Cen(\alpha)$. By Theorem 3.1 we must have $\bar{\chi}(\beta) = 0$, since $\alpha \sqcap R_0 \neq cI$. Similarly we must have that $\underline{\chi}(\beta) = 0$. So in the direct sum decomposition only elements in R_0 occurs. Thus $\beta \in R_0$. □

Corollary 3.3. *R_0 is maximal commutative.*

Proof. Let β be an element that commutes with everything in R_0 . Then in particular it must commute with BA . But $Cen(BA) = R_0$ by the preceding theorem. Thus $\beta \in R_0$. □

4 Annihilating polynomials

As mentioned in the introduction any two commuting elements in $H(q)$ must be algebraically dependent when q is of free type. More formally, we have the following.

Theorem 4.1. *Let $q \in K$ be of free type. If $\alpha, \beta \in H(q)$ commute, then there exists a nonzero $P \in K[x, y]$ such that $P(\alpha, \beta) = 0$.*

We now describe an explicit construction of this polynomial. We let s and t be variables that take values in the base field K . We write the commuting elements α, β as

$$\alpha = \sum_{i=0}^n p_i(B)A^i, \quad \beta = \sum_{i=0}^m r_i(B)A^i$$

where the p_i and r_i are polynomials. We will form an $n + m$ determinant that will give us the annihilating polynomial.

Consider the expressions obtained by reordering all A to the right of B in

$$A^k(\alpha - sI) = \sum_{i=0}^{n+k} \theta_{i,k} A^i, \quad k = 0, 1, \dots, m - 1$$

$$A^k(\beta - tI) = \sum_{i=0}^{m+k} \omega_{i,k} A^i, \quad k = 0, 1, \dots, n - 1$$

where $\theta_{i,k}$ and $\omega_{i,k}$ are functions of B, s, t arising after reordering. The coefficients of the powers of A will be the elements in the determinant that we compute. $\theta_{i,k}$ will be placed as the element in row $k + 1$ and column i . $\omega_{i,k}$ will be placed in row $k + m + 1$ and column i . The determinant will thus be a polynomial in s, t and B . This polynomial, which we will call the eliminant of α and β , can be written as $\sum_i \delta_i(s, t) B^i$. Every such δ_i will satisfy $\delta_i(\alpha, \beta) = 0$ and at least one of them will not be identically zero.

A more precise formulation with additional information about the construction can be found in the following.

Theorem 4.2. *Let*

$$\alpha = \sum_{j=0}^m p_j(B) A^j, \quad \beta = \sum_{j=0}^n r_j(B) A^j$$

be two commuting elements, the p_j and r_j being polynomials, and denote their eliminant by $\Delta_{\alpha, \beta}(B, s, t)$. Then $\Delta_{\alpha, \beta} \neq 0$. Furthermore $\Delta_{\alpha, \beta}$ has degree n seen as polynomial in s . If $r_n(B) = \sum_i a_i B^i$ ($a_i \in K$), then $\Delta_{\alpha, \beta}$ has leading coefficient $(-1)^n \prod_{k=0}^{m-1} (\sum_i a_i q^{ki} B^i)$, once again seen as a polynomial in s . Symmetrically, $\Delta_{\alpha, \beta}$ will have degree m seen as a polynomial in t . The coefficient of t^m will be $(-1)^m \prod_{k=0}^{n-1} (\sum_i b_i q^{ki} B^i)$ if $p_m(B) = \sum_i b_i B^i$.

Let $g = n \max_j \deg(p_j) + m \max_j \deg(r_j)$. We can write

$$\Delta_{\alpha, \beta}(B, s, t) = \sum_{i=0}^g \delta_i(s, t) B^i$$

Then at least one $\delta_i(s, t) \neq 0$ and $\delta_i(\alpha, \beta) = 0$ for all i .

5 The eliminant when the elements belong to R_0

In the general case the theorem does not rule out that one can get several nonzero δ_i in the expansion of the eliminant, $\Delta_{\alpha, \beta}$. This does not, however, occur when α and β belong to R_0 .

Theorem 5.1. *Let $\alpha = \sum_{k=0}^n p_k B^k A^k$ and $\beta = \sum_{l=0}^m r_l B^l A^l$. Then, with the same notation as before, there will be only one nonzero δ_i when the eliminant is computed and this i will equal nm .*

Proof. We begin by noting that $A^n(\alpha - sI)$ will be of the form $\sum_{i=n}^{n+k} a_i B^{i-k} A^i$, where the a_i belong to K .

We use this result to describe the structure of the eliminant. Denote the element in row u and column v by $e_{u,v}$. Then we will have $e_{u,v} = \phi_{i,j}(s, t) B^{v-u}$ if $u \leq m$ (that is in the first m rows) and $e_{u,v} = \phi_{i,j} B^{v-u+m}$ otherwise (in the last n rows), where the $\phi_{i,j}(s, t)$ are polynomials over K . Many of them will of course be zero, in particular those where B would otherwise occur with a negative exponent.

We know, from ordinary linear algebra, that

$$\Delta_{\alpha,\beta}(B, s, t) = \sum_{\sigma} \text{sign}(\sigma) \prod_{g=1}^{m+n} e_{g,\sigma(g)}$$

where σ denotes a permutation. But looking at an arbitrary term of the sum we find that it can be written as

$$\begin{aligned} & \text{sign}(\sigma) \prod_{g=1}^{g=m} (\phi_{g,\sigma(g)}(s, t) B^{\sigma(g)-g}) * \prod_{g=m+1}^{m+n} \phi_{g,\sigma(g)}(s, t) B^{\sigma(g)-g+m} \\ & = \Phi(s, t) * B^{\sum_{g=1}^{m+n} (\sigma(g)) - \sum_{g=1}^{n+m} (g) + mn} \end{aligned}$$

for a polynomial $\Phi(s, t)$. But the two sums in the exponent cancel, since they have the same terms in different order, and we conclude that we get the exponent mn . Since we picked an arbitrary term, we are done. □

5.1 Examples

We will include some examples here to give a feeling for the construction of the eliminant and our result. Let $\alpha = BA$ and $\beta = B^2A^2$. Then

$$\Delta_{\alpha,\beta}(B, s, t) = \begin{vmatrix} -s & B & 0 \\ 0 & 1-s & qB \\ -t & 0 & B^2 \end{vmatrix}$$

On computing the determinant we find that the annihilating polynomial is $s^2 - s - tq$. This is only a slight modification of the classical case when $q = 1$. (We note that it makes no difference whether we set $q = 1$ at the beginning of the calculation or the end.)

For our next example let α be as before and let $\beta = B^3A^3$. Then we find that

$$\Delta_{\alpha,\beta}(B, s, t) = \begin{vmatrix} -s & B & 0 & 0 \\ 0 & 1-s & qB & 0 \\ 0 & 0 & 1+q-s & q^2B \\ -t & 0 & 0 & B^3 \end{vmatrix}$$

We get the annihilating polynomial

$$s^3 - (2+q)s^2 + (1+q)s - q^3t$$

Once again no essential simplification occurs if we let q approach 1.

Now set $\beta = B^2A^2$ and $\alpha = B^3A^3$. The determinant becomes

$$\Delta_{\alpha,\beta}(B, s, t) = \begin{vmatrix} -s & 0 & 0 & B^3 & 0 \\ 0 & -s & 0 & (1+q+q^2)B^2 & q^3B^3 \\ -t & 0 & B^2 & 0 & 0 \\ 0 & -t & (1+q)B & q^2B^2 & 0 \\ 0 & 0 & 1+q-t & q(1+2q+q^2)B & q^4B^2 \end{vmatrix}$$

and we get the annihilating polynomial

$$q^3s^2 + (q + 2q^2)st + (1 + q)t^2 - t^3$$

In the classical case this polynomial becomes

$$s^2 + 3st + 2t^2 - t^3$$

As a final example we can take $\alpha = B^2A^2$ and $\beta = B^4A^4$. The eliminant is

$$\Delta_{\alpha,\beta}(B, s, t) = \begin{vmatrix} -s & 0 & B^2 & 0 & 0 & 0 \\ 0 & -s & (1+q)B & q^2B^2 & 0 & 0 \\ 0 & 0 & 1+q-s & q(1+2q+q^2)B & q^4B^2 & 0 \\ 0 & 0 & 0 & 1+2q+2q^2+q^3-s & q^2(1+2q+2q^2+q^3)B & q^6B^2 \\ -t & 0 & 0 & 0 & B^4 & 0 \\ 0 & -t & 0 & 0 & (1+q+q^2+q^3)B^3 & q^4B^4 \end{vmatrix}$$

We then get the annihilating polynomial

$$q^8t^2 - 2q^6ts - 3q^5ts - 2q^4ts^2 - 2q^4ts - q^3ts + q^4s^2 - q^3s^3 + 3q^3s^2 - 2q^2s^3 + 4q^2s^2 - 3qs^3 + 3qs^2 + s^2 + s^4 - 2s^3$$

The limit when q goes towards 1 is

$$t^2 - 8ts - 2s^2t + 12s^2 - 8s^3 + s^4$$

This is a simpler expression but only because the coefficients are simpler. No coefficient has become zero.

This illustrates that the complexity of the resulting polynomial grows pretty fast.

Computer experiments indicate that Theorem 5.1 can be generalized substantially. We would be also interested to know whether the annihilating polynomials always have genus 0, a conjecture we have been unable to find any counterexamples to.

Acknowledgements

This work was supported by the Swedish Research Council, the Swedish Foundation for International Cooperation in Research and Higher Education (STINT), the Crafoord Foundation, the Royal Physiographic Society in Lund, and the Royal Swedish Academy of Sciences.

References

- [1] J. L. Burchnell and T. W. Chaundy. Commutative ordinary differential operators. *Proc. Lond. Math. Soc. (2)*, **21** (1922), 420–440.
- [2] J. L. Burchnell and T. W. Chaundy. Commutative ordinary differential operators. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, **118** (1928), 557–583.
- [3] J. L. Burchnell and T. W. Chaundy. Commutative ordinary differential operators. II. The Identity $P^n = Q^m$. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, **134** (1932), 471–485.
- [4] L. Hellström and S. Silvestrov. *Commuting Elements in q -Deformed Heisenberg Algebras*. World Scientific Publishing, River Edge, NJ, 2000.

- [5] L. Hellström and S. Silvestrov. Ergodipotent maps and commutativity of elements in non-commutative rings and algebras with twisted intertwining. *J. Algebra*, **314** (2007), 17–41.
- [6] M. de Jeu, C. Svensson, and S. Silvestrov. Algebraic curves for commuting elements in the q -deformed Heisenberg algebra. *J. Algebra*, **321** (2009), 1239–1255.
- [7] I. Krichever. Integration of nonlinear equations by the methods of algebraic geometry (Russian). *Funktsional. Anal. i Priložen*, **11** (1977), 15–31.
- [8] I. Krichever. Methods of algebraic geometry in the theory of nonlinear equations (Russian). *Uspekhi Mat. Nauk*, **32** (1977), 183–208.
- [9] D. Larsson. Burchnell-Chaundy theory, Ore extensions and σ -differential operators. U.U.D.M. Report 2008:45, Department of Mathematics, Uppsala University, 2008.
- [10] D. Larsson and S. D. Silvestrov. Burchnell-Chaundy theory for q -difference operators and q -deformed Heisenberg algebras. *J. Nonlinear Math. Phys.*, **10** (2003), Suppl. 2, 95–106.
- [11] D. Mumford. An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg deVries equation and related nonlinear equation. Proceedings of the International Symposium on Algebraic Geometry (Kyoto University, Kyoto, 1977), 1978, 115–153.

Received May 21, 2009

Revised September 20, 2009