

MINIMAL IMBEDDINGS OF R -SPACES

MASARU TAKEUCHI & SHOSHICHI KOBAYASHI

1. Introduction

Let G be a connected real semi-simple Lie group without center and U a parabolic subgroup of G . The quotient space G/U is called an R -space. A maximal compact subgroup K of G is transitive on G/U so that an R -space is necessarily compact. Let $\mathfrak{G} = \mathfrak{K} + \mathfrak{P}$ be a Cartan decomposition of the Lie algebra \mathfrak{G} of G with respect to the Lie algebra \mathfrak{K} of K . The main purpose of this paper is to construct a natural imbedding φ of an R -space G/U into \mathfrak{P} with the following properties:

- (1) φ is K -equivariant;
- (2) φ has minimum total curvature;
- (3) If G is simple and $K/K \cap U$ is a symmetric space, then φ is isometric and $\varphi(G/U)$ is a minimal submanifold of a hypersphere in \mathfrak{P} in the sense that its mean curvature normal is zero.

In general, an n -dimensional submanifold M of the hypersphere $S^N(r)$ of radius r about the origin in the Euclidean space \mathbf{R}^{N+1} is a minimal submanifold if and only if

$$\Delta y^i = -\frac{n}{r^2} y^i \quad \text{on } M \text{ for } i = 1, \dots, N+1,$$

where (y^1, \dots, y^{N+1}) is a coordinate system for \mathbf{R}^{N+1} and Δ is the Laplacian of M . For many symmetric R -spaces we verify that the Laplacian Δ for functions has no eigen-value between 0 and $-n/r^2$. We do not know whether this is true or not in general for all symmetric R -spaces.

Previously, it was known that φ has minimum total curvature if G/U is a Kählerian C -space (Kobayashi [6]) or if G/U is a symmetric space of rank 1 (Tai [15]). For a symmetric R -space G/U , the imbedding φ has been considered by Nagano [13], and has also been conjectured to have minimum total curvature (Kobayashi [7]). The class of symmetric R -spaces includes

- (i) all hermitian symmetric spaces of compact type;
- (ii) Grassmann manifolds $O(p+q)/O(p) \times O(q)$, $Sp(p+q)/Sp(p) \times Sp(q)$;

Received November 9, 1967 and, in revised form, February 5, 1968. The work of the first author was partially supported by NSF Grant GP-6974, and that of the second author by NSF Grant GP-5798.

- (iii) the classical groups $SO(m)$, $U(m)$, $Sp(m)$;
- (iv) $U(2m)/Sp(m)$, $U(m)/O(m)$;
- (v) $(SO(p+1) \times SO(q+1))/S(O(p) \times O(q))$, where $S(O(p) \times O(q))$ is the subgroup of $SO(p+1) \times SO(q+1)$ consisting of matrices of the form

$$\begin{pmatrix} \varepsilon & 0 \\ 0 & A \\ & \varepsilon & 0 \\ & 0 & B \end{pmatrix}, \quad \varepsilon = \pm 1, \quad A \in O(p), \quad B \in O(q);$$

(This R -space is covered twice by $S^p \times S^q$.)

- (vi) the Cayley projective plane and three exceptional spaces.

An explicit formula for the imbedding φ of a symmetric R -space of classical type in \mathfrak{B} in terms of matrices can be found in Kobayashi [7].

In §3 we recall briefly the concept of minimum imbedding without mentioning that of total curvature. For the latter we refer the reader to Chern and Lashof [1], [2], Kuiper [9], [10] and references therein.

The result of this paper on the total curvature of φ relies heavily on the cellular decomposition of an R -space obtained by Takeuchi [16].

Our result on minimal submanifolds of a hypersphere is somewhat related to those of Takahashi [7] and Hsiang [4], and Proposition 5.1 on minimal submanifolds appears in Takahashi [17].

2. Parabolic subgroups and R -spaces

Let G be a connected real semi-simple Lie group without center, and \mathfrak{G} its Lie algebra. Let \mathfrak{G}_C be the complexification of \mathfrak{G} , and G_C the connected complex semi-simple Lie group without center generated by the Lie algebra \mathfrak{G}_C . Then we may consider G as a subgroup of G_C . The complex conjugation σ of \mathfrak{G}_C with respect to \mathfrak{G} generates an automorphism σ of G_C which leaves G elementwise fixed.

A subgroup of G_C is called a *parabolic subgroup* of G_C if it contains a maximal solvable subgroup of G_C ; it is always connected. A subgroup of G is called a *parabolic subgroup* of G if it is the intersection of G and a σ -invariant parabolic subgroup of G_C . A parabolic subgroup of G may not be connected, but it is still uniquely determined by its Lie algebra alone. A subalgebra of \mathfrak{G} is called a *parabolic subalgebra* if it is the Lie algebra of a parabolic subgroup of G . If Z is an element of \mathfrak{G} such that $ad Z$ is a semi-simple endomorphism of \mathfrak{G} whose eigen-values are all real, then the direct sum \mathfrak{U} of all eigen-spaces corresponding to the non-negative eigen-values of $ad Z$ is a parabolic subalgebra of \mathfrak{G} . Conversely, every parabolic subalgebra of \mathfrak{G} can be obtained in this fashion (cf. Matsumoto [11]).

An R -space is, by definition, a quotient space $M = G/U$, where G is a connected real semi-simple Lie group without center and U is a parabolic subgroup of G . Given an R -space $M = G/U$, we choose once and for all an

element $Z \in \mathfrak{G}$ which determines the parabolic subalgebra \mathfrak{U} , the Lie algebra of U , in the manner described above. (Such an element Z is not unique.) We choose also a maximal compact subgroup K of G such that Z is perpendicular to the Lie algebra \mathfrak{K} of K with respect to the Killing form (\cdot, \cdot) of \mathfrak{G} . In the Cartan decomposition $\mathfrak{G} = \mathfrak{K} + \mathfrak{P}$, Z is then contained in \mathfrak{P} . We choose a maximal abelian subalgebra \mathfrak{A} of \mathfrak{P} , which contains Z , and introduce a linear order in the dual space of \mathfrak{A} in such a way that $\gamma(Z) \geq 0$ for all positive roots γ of \mathfrak{G} with respect to \mathfrak{A} . Let \mathfrak{N} be the direct sum of the root spaces corresponding to the positive roots. Then \mathfrak{N} is a nilpotent subalgebra of \mathfrak{G} . Let N be the connected subgroup of G generated by \mathfrak{N} , and set

$$K_0 = \{k \in K; (Ad k)Z = Z\}.$$

Then we have (Takeuchi [16])

Proposition 2.1. (i) $KU = G$ and $K \cap U = K_0$ so that $M = K/K_0$; (ii) If we denote by $N_K(\mathfrak{A})$ (resp. $N_{K_0}(\mathfrak{A})$) the normalizer of \mathfrak{A} in K (resp. in K_0), then $N_K(\mathfrak{A})/N_{K_0}(\mathfrak{A})$ is finite. If $k_1, \dots, k_b \in N_K(\mathfrak{A})$ are complete representatives of $N_K(\mathfrak{A})/N_{K_0}(\mathfrak{A})$ and if o denotes the origin of G/U , then the orbits $Nk_1o, \dots, Nk_b o$ of N through $k_1o, \dots, k_b o$ give a cellular decomposition of M , and these cells are all cycles mod 2.

As a consequence, we have $\sum_i \dim H_i(M, \mathbb{Z}_2) = b$. From (i) we see that the mapping $\varphi: M = K/K_0 \rightarrow \mathfrak{P}$ defined by

$$\varphi(kK_0) = (Ad k)Z, \quad kK_0 \in K/K_0$$

is a K -equivariant imbedding of M into \mathfrak{P} . The purpose of this paper is to study geometric properties of this imbedding φ .

Proposition 2.2. Let X be a regular element of \mathfrak{P} . Then the number of zero points of the vector field on M generated by X coincides with the number b of the elements in $N_K(\mathfrak{A})/N_{K_0}(\mathfrak{A})$.

Proof. We first prove

Lemma. If we set $\mathfrak{P}_0 = \{X \in \mathfrak{P}; [Z, X] = 0\}$, then $\mathfrak{U} \cap \mathfrak{P} = \mathfrak{P}_0$.

Proof of Lemma. From the definitions of \mathfrak{U} and \mathfrak{P}_0 we have clearly $\mathfrak{P}_0 \subset \mathfrak{U} \cap \mathfrak{P}$. Let $X \in \mathfrak{U} \cap \mathfrak{P}$ and write

$$X = X_0 + X_+,$$

where $[Z, X_0] = 0$ and X_+ is in the direct sum of the eigen-spaces corresponding to the positive eigen-values of $ad Z$. We wish to show $X_+ = 0$. Let τ be the involutive automorphism of \mathfrak{G} such that $\tau|_{\mathfrak{K}} = \text{identity}$ and $\tau|_{\mathfrak{P}} = -\text{identity}$. Then $\tau Z = -Z$ and hence $\tau \circ (ad Z) = -(ad Z) \circ \tau$. It follows that $[Z, \tau X_+] = 0$ and that τX_+ is in the direct sum of the eigen-spaces corresponding to the negative eigen-values of $ad Z$. On the other hand, since X

is in \mathfrak{B} , we have $\tau X = -X$ and $\tau X \in \mathfrak{U} \cap \mathfrak{B}$. Since $\tau X = \tau X_0 + \tau X_+$ is in \mathfrak{U} , it follows that $X_+ = 0$. This completes the proof of the lemma.

Let X be a regular element of \mathfrak{B} . For each $k \in K$, X and $(Ad k)X$ generate vector fields on M with the same number of zero points on M . Since $(Ad k)X \in \mathfrak{A}$ for a suitable k , we may assume that X is a regular element of \mathfrak{A} . It suffices therefore to prove that, for a regular element X of \mathfrak{A} , the zero points of the vector field generated by X coincide with the orbit $N_K(\mathfrak{A})o$ of $N_K(\mathfrak{A})$ through the origin o of $M = K/K_0$. Let ko ($k \in K$) be a zero point of the vector field generated by X . Then $X \in (Ad k)\mathfrak{U}$ and hence $(Ad k^{-1})X \in \mathfrak{U}$. Since $(Ad k^{-1})X \in \mathfrak{B}$, the lemma above implies $(Ad k^{-1})X \in \mathfrak{B}_0$. If we set $\mathfrak{G}_0 = \{Y \in \mathfrak{G}; [Z, Y] = 0\}$, then \mathfrak{G}_0 is a reductive Lie algebra, and $\mathfrak{G}_0 = \mathfrak{K}_0 + \mathfrak{B}_0$ is a Cartan decomposition of \mathfrak{G}_0 . Since \mathfrak{A} is a maximal abelian subalgebra of \mathfrak{B}_0 , there exists an element $k_0 \in K_0$ such that $(Ad k_0^{-1})(Ad k^{-1})X \in \mathfrak{A}$. If we set $k' = kk_0$, then $(Ad k'^{-1})X \in \mathfrak{A}$. Since X is a regular element of \mathfrak{A} , k' lies in $N_K(\mathfrak{A})$. On the other hand, $k'o = kk_0o = ko$. It is easy to see the converse that $N_K(\mathfrak{A})o$ is contained in the set of zero points of the vector field generated by X .

3. Minimum imbeddings

Let M be a compact manifold, and \mathcal{F} the set of C^∞ functions f on M whose critical points are all isolated and non-degenerate. For each $f \in \mathcal{F}$, we denote by $\beta(f)$ the number of the critical points of f on M . Set

$$\beta = \inf_{f \in \mathcal{F}} \beta(f).$$

Then β depends only on the differentiable structure of M , and the theory of Morse tells us that, for any coefficient field F , the following inequality holds:

$$\beta \geq \sum_i \dim H_i(M, F).$$

Let φ be an imbedding of M into a real vector space V . Then for almost¹ all linear functional u on V , the function $u \circ \varphi$ belongs to the family \mathcal{F} . We say that the imbedding $\varphi: M \rightarrow V$ is *minimum* if $\beta = \beta(u \circ \varphi)$ for almost all linear functionals u on V such that $u \circ \varphi$ belongs to the family \mathcal{F} . Since $\beta(u \circ \varphi) \geq \beta \geq \sum_i \dim H_i(M, F)$ always, φ is minimum if $\beta(u \circ \varphi) = \sum_i \dim H_i(M, F)$ for some coefficient field F and almost all linear functionals u such that $(u \circ \varphi) \in \mathcal{F}$.

We shall prove the following theorem:

Theorem 3.1. *Let $M = G/U$ be an R-space, and $\varphi: M \rightarrow \mathfrak{B}$ the imbedding defined in § 2. Then φ is minimum, and*

¹ in the sense of measure.

$$\beta = \sum_i H_i(M, Z_2).$$

We shall first outline the proof. Let X be any element of \mathfrak{P} , and u_X the linear functional on \mathfrak{P} which corresponds to X under the duality defined by the Killing form $(,)$ of \mathfrak{G} . We define a suitable Riemannian metric \ll, \gg and show that the 1-form $d(u_X \circ \varphi)$ corresponds to the vector field generated by X by the duality defined by \ll, \gg . Then the critical points of $u_X \circ \varphi$ coincide with the zero points of the vector field generated by X . Since the singular elements of \mathfrak{P} form a set of measure zero, the theorem will then follow immediately from Propositions 2.1 and 2.2. We now give the details of the proof.

Let \mathfrak{R}_0 be the Lie algebra of K_0 . The Killing form $(,)$ of \mathfrak{G} is negative definite on \mathfrak{R} . Let \mathfrak{M} be the orthogonal complement of \mathfrak{R}_0 in \mathfrak{R} with respect to the Killing form $(,)$. Then \mathfrak{M} is invariant by $Ad K_0$. As in the proof of Lemma for Proposition 2.2, let τ be the involutive automorphism of \mathfrak{G} defined by $\tau|_{\mathfrak{R}} = \text{identity}$ and $\tau|_{\mathfrak{P}} = -\text{identity}$. Since $\tau \circ (ad Z) = -(ad Z) \circ \tau$ as we have shown earlier in the proof of Proposition 2.2, we have $\tau \circ (ad Z)^2 = (ad Z)^2 \circ \tau$. Hence $(ad Z)^2$ leaves \mathfrak{R} and \mathfrak{P} invariant. Since $ad Z$ leaves the Killing form $(,)$ invariant, $(ad Z)^2$ is a symmetric endomorphism of \mathfrak{G} with respect to $(,)$. If we denote by \mathfrak{P}_+ the direct sum of the eigen-spaces corresponding to the positive eigen-values of $(ad Z)^2|_{\mathfrak{P}}$, then $\mathfrak{P} = \mathfrak{P}_0 + \mathfrak{P}_+$, and \mathfrak{P}_0 and \mathfrak{P}_+ are mutually orthogonal with respect to the Killing form $(,)$. Since $(ad Z)^2$ maps \mathfrak{R}_0 into 0, $(ad Z)^2$ leaves \mathfrak{M} invariant. Let $\gamma_1, \dots, \gamma_n$ be the set of roots γ (multiplicity counted) of \mathfrak{G} with respect to \mathfrak{A} such that $\gamma(Z) > 0$. Then we know (Takeuchi [16]) that there exist a basis S_1, \dots, S_n for \mathfrak{M} and a basis T_1, \dots, T_n for \mathfrak{P}_+ such that

$$\begin{aligned} &-(S_i, S_j) = \delta_{ij}, \quad (T_i, T_j) = \delta_{ij} \quad \text{for } 1 \leq i, j \leq n; \\ (*) \quad &[H, S_i] = \gamma_i(H)T_i, \quad [H, T_i] = \gamma_i(H)S_i \quad \text{for } H \in \mathfrak{A} \text{ and } 1 \leq i \leq n; \\ &S_i + T_i \in \mathfrak{U} \quad \text{for } 1 \leq i \leq n. \end{aligned}$$

By setting $H = Z$ in $(*)$, we see that $[Z, \mathfrak{M}] = \mathfrak{P}_+$ and $[Z, \mathfrak{P}_+] = \mathfrak{M}$ and that $(ad Z)^2|_{\mathfrak{M}}$ is a positive definite symmetric endomorphism of \mathfrak{M} with respect to $-(,)$. Let ζ be a positive definite symmetric endomorphism of \mathfrak{M} with respect to $-(,)$ such that $\zeta^2 = (ad Z)^2|_{\mathfrak{M}}$. Then $\zeta S_i = \gamma_i(Z)S_i$ for $1 \leq i \leq n$. Since $(Ad k)Z = Z$ for $k \in K_0$, we have $(Ad k)\zeta X = \zeta(Ad k)X$ for $X \in \mathfrak{M}$ and $k \in K_0$.

Lemma 1. $X + \zeta^{-1}[Z, X] \in \mathfrak{U}$ for $X \in \mathfrak{P}_+$.

Proof of Lemma 1. It suffices to verify for $X = T_i$ ($1 \leq i \leq n$). From $(*)$ we obtain

$$T_i + \zeta^{-1}[Z, T_i] = T_i + \zeta^{-1}\gamma_i(Z)S_i = T_i + \zeta^{-1}\zeta S_i = T_i + S_i \in \mathfrak{U},$$

which proves Lemma 1.

We shall now construct K -invariant Riemannian metric $\langle\langle \cdot, \cdot \rangle\rangle$ on $M = K/K_0$. Let $T_o(M)$ be the tangent space of $M = K/K_0$ at the origin o . Under the natural identification of \mathfrak{M} with $T_o(M)$, the adjoint action of K_0 on \mathfrak{M} corresponds to the linear isotropy representation of K_0 on $T_o(M)$. We set

$$\langle\langle X, Y \rangle\rangle = -(\zeta X, Y) \quad \text{for } X, Y \in \mathfrak{M}.$$

Since (\cdot, \cdot) is negative definite on \mathfrak{R} and ζ commutes with $Ad k$ on \mathfrak{M} for every $k \in K_0$, it follows that $\langle\langle \cdot, \cdot \rangle\rangle$ is a K_0 -invariant positive definite symmetric bilinear form on \mathfrak{M} . Hence $\langle\langle \cdot, \cdot \rangle\rangle$ can be extended uniquely to a K -invariant Riemannian metric $\langle\langle \cdot, \cdot \rangle\rangle$ on $M = K/K_0$.

Let $X \in \mathfrak{P}$ and let u_X denote the linear functional on \mathfrak{P} defined by $u_X(Y) = (Y, X)$ for $Y \in \mathfrak{P}$. Let φ be the imbedding of M into \mathfrak{P} defined in §2, and set $f_X = u_X \circ \varphi$. In other words, f_X is defined by

$$f_X(ko) = ((Ad k)Z, X) \quad \text{for } k \in K.$$

Lemma 2. *For every $X \in \mathfrak{P}$, df_X is the 1-form (i.e., the covariant vector) corresponding to the vector field (i.e., the contravariant vector) generated by X under the duality defined by the Riemannian metric $\langle\langle \cdot, \cdot \rangle\rangle$.*

Proof of Lemma 2. We denote by the same letter X the vector field on M generated by X . The value of X at a point ko of M will be denoted by Xko . Similarly, for $Y \in \mathfrak{M}$, kYo denotes the vector at ko obtained from the vector $Yo \in T_o(M)$ by a transformation $k \in K$. Then Lemma 2 may be stated as follows:

$$\langle (df_X)_{ko}, kYo \rangle = \langle\langle Xko, kYo \rangle\rangle \quad \text{for } Y \in \mathfrak{M} \quad \text{and } k \in K.$$

We calculate the left hand side first.

$$\begin{aligned} \langle (df_X)_{ko}, kYo \rangle &= \frac{d}{dt} f_X((k \cdot \exp tY)o)|_0 = \frac{d}{dt} ((Ad k \cdot \exp tY)Z, X)|_0 \\ &= \frac{d}{dt} ((Ad \exp tY)Z, (Ad k^{-1})X)|_0 = ([Y, Z], (Ad k^{-1})X) \\ &= (Y, [Z, (Ad k^{-1})X]). \end{aligned}$$

We decompose $(Ad k^{-1})X \in \mathfrak{P}$ as follows: $(Ad k^{-1})X = X_0 + X_+$, where $X_0 \in \mathfrak{P}_0$ and $X_+ \in \mathfrak{P}_+$. Then we have

$$\langle (df_X)_{ko}, kYo \rangle = (Y, [Z, X_+]).$$

We now calculate the right hand side.

$$\langle\langle Xko, kYo \rangle\rangle = \langle\langle (Ad k^{-1})Xo, Yo \rangle\rangle.$$

Since we have $((Ad k^{-1})X)o = (-\zeta^{-1}[Z, X_+])o$ by Lemma 1, we obtain

$$\langle\langle Xko, kYo \rangle\rangle = -\langle\langle \zeta^{-1}[Z, X_+], Y \rangle\rangle = ([Z, X_+], Y).$$

This completes the proof of Lemma 2.

Theorem 3.1 now follows from Propositions 2.1 and 2.2 and from Lemma 2 just proved.

Remark 1. Given an R -space $M = G/U$ we may assume without loss of generality that G acts effectively on M , i.e., U contains no nontrivial normal subgroup of G . Then the minimum imbedding $\varphi: M \rightarrow \mathfrak{P}$ is substantial in the sense that $\varphi(M)$ is not contained in any (affine) hyperplane of \mathfrak{P} ; otherwise there would exist a nonzero linear functional u_X of \mathfrak{P} such that the function $f_X = u_X \circ \varphi$ is constant on M . But Lemma 2 says that if $df_X = 0$ on M , then the vector field on M generated by X also vanishes identically on M . Hence, $X = 0$.

Remark 2. Since $\beta \geq \sum \dim H_i(M, Z_p)$ by Morse theory, we may conclude that, for any R -space $M = G/U$, the inequality

$$\sum \dim H_i(M, Z_2) \geq \sum \dim H_i(M, Z_p)$$

holds for all prime numbers p .

4. Symmetric R-spaces and minimal submanifolds of spheres

Let G be a connected real semi-simple Lie group without center, and Z an element of \mathfrak{G} such that $ad Z$ is a semi-simple endomorphism of \mathfrak{G} with eigenvalues $-1, 0$ and 1 . Let $\mathfrak{G} = \mathfrak{G}_{-1} + \mathfrak{G}_0 + \mathfrak{G}_1$ be the corresponding eigenspace decomposition, and U the parabolic subgroup of G with Lie algebra $\mathfrak{u} = \mathfrak{G}_0 + \mathfrak{G}_1$. Taking a Cartan decomposition $\mathfrak{G} = \mathfrak{K} + \mathfrak{P}$ such that $Z \in \mathfrak{P}$, let K be the maximal compact subgroup of G generated by \mathfrak{K} . Let $K_0 = \{k \in K; (ad k)Z = Z\}$ and $\mathfrak{K} = \mathfrak{K}_0 + \mathfrak{M}$ as in §§ 2 and 3. Let \mathfrak{G}_c be the complexification of \mathfrak{G} and G_c the complex semi-simple Lie group without center generated by \mathfrak{G}_c . Let θ denote the restriction to K of the inner automorphism of \mathfrak{G}_c defined by $\exp(\pi i Z) \in G_c$. If we set $K_\theta = \{k \in K; \theta k = k\}$, then K_0 lies between K_θ and the identity component of K_θ . It follows that $M = K/K_0$ is a symmetric space defined by the involutive automorphism θ of K . (By results of Nagano [13] (cf. also Kobayashi-Nagano [8] and Takeuchi [16]), the converse is also true; namely, if $M = G/U$ is an R -space such that $M = K/K_0$ is symmetric, then U is determined by an element $Z \in \mathfrak{G}$ such that $ad Z$ has eigenvalues $-1, 0, 1$.) Throughout this section we shall consider a symmetric R -space $M = G/U = K/K_0$, where U is determined by such a $Z \in \mathfrak{G}$. The main purpose of this section is to prove that, with respect to the imbedding $\varphi: M \rightarrow \mathfrak{P}$ defined in § 2, $\varphi(M)$ is a minimal submanifold of the sphere of radius $\sqrt{2n}$ in \mathfrak{P} , where $n = \dim M$.

With our notations in § 3, we have $\gamma_i(Z) = 1$ for $1 \leq i \leq n$ and $\zeta(X) = X$ for all $X \in \mathfrak{M}$. The Riemannian metric $\langle \cdot, \cdot \rangle$ on M is defined by $\langle X, Y \rangle = -(X, Y)$ for $X, Y \in \mathfrak{M} = T_o(M)$. From the formulas (*) in § 3 it follows that the imbedding $\varphi: M \rightarrow \mathfrak{P}$ is isometric with respect to the Riemannian metric $\langle \cdot, \cdot \rangle$ and the restriction of the Killing form (\cdot, \cdot) of \mathfrak{G} to \mathfrak{P} .

From the definition of the imbedding $\varphi: M \rightarrow \mathfrak{P}$ it is clear that its image $\varphi(M)$ lies on the sphere of radius $(Z, Z)^{\frac{1}{2}}$ with center at the origin of \mathfrak{P} .

Proposition 4.1. *For a symmetric R-space $M = G/U$, we have $(Z, Z) = 2n$, where $n = \dim M$.*

$$\text{Proof. } (Z, Z) = \text{Tr}(ad Z)^2 = \sum_{i=1}^n \gamma_i(Z)^2 + \sum_{i=1}^n (-\gamma_i(Z))^2 = 2n.$$

Theorem 4.2. *Let $M = G/U = K/K_0$ be a symmetric R-space with G simple. Then $\varphi(M)$ is a minimal submanifold of the sphere of radius $\sqrt{2n}$ about the origin in \mathfrak{P} , where $n = \dim M$.*

Proof. We identify $\varphi(M)$ with M . Let S denote the sphere of radius $\sqrt{2n}$ about the origin in \mathfrak{P} , and α be the second fundamental form of M in S ; at each point $x \in M$, it defines a symmetric bilinear mapping $T_x(M) \times T_x(M) \rightarrow T_x^\perp$, where T_x^\perp denotes the normal space to M in S at x . Choosing an orthonormal basis e_1, \dots, e_n for $T_x(M)$, we define the mean curvature normal ξ_x by

$$\xi_x = \sum_{i=1}^n \alpha(e_i, e_i).$$

Then ξ_x is independent of the choice of e_1, \dots, e_n . The submanifold M is minimal if and only if $\xi_x = 0$ at every point x of M . In the present case, since the imbedding φ is K -equivariant, the field ξ of mean curvature normals is invariant by the adjoint action of K in \mathfrak{P} . It suffices therefore to prove that ξ vanishes at the origin o of M . The tangent space $T_o(M)$ is parallel to $[Z, \mathfrak{M}] = \mathfrak{P}_+$ in \mathfrak{P} (cf. formulas (*) in § 3). Since Z is normal to the sphere S at o , ξ_o is perpendicular to Z as well as to \mathfrak{P}_+ . Hence ξ_o can be identified with an element of \mathfrak{P}_0 which is perpendicular to Z and is invariant by the adjoint action of K_0 in \mathfrak{P}_0 . The proof of the theorem is now reduced to that of the following lemma.

Lemma. *Let $M = G/U$ be a symmetric R-space with G simple. Then the space $\{X \in \mathfrak{P}_0; (Ad k)X = X \text{ for all } k \in K_0\}$ is spanned by Z .*

Proof of Lemma. Consider first the case where the complexification $\mathfrak{G}_\mathbb{C}$ of \mathfrak{G} is not simple. In this case, \mathfrak{R} is compact and simple, and \mathfrak{G} admits a complex structure J such that $\mathfrak{P} = J\mathfrak{R}$ and $\mathfrak{P}_0 = J\mathfrak{R}_0$. Moreover, \mathfrak{R}_0 has center of dimension 1 (cf. Helgason [3]). Our lemma is clearly true in this case.

Consider now the case where $\mathfrak{G}_\mathbb{C}$ is simple. In this case, the center of \mathfrak{G}_0 is spanned by Z (cf. Kobayashi-Nagano [8] and Takeuchi [16]). Let $\mathfrak{G}'_0 = [\mathfrak{G}_0, \mathfrak{G}_0]$ and $\mathfrak{P}'_0 = \mathfrak{G}'_0 \cap \mathfrak{P}_0$. Then $\mathfrak{G}'_0 = \mathfrak{R}_0 + \mathfrak{P}'_0$ is a Cartan decomposition

of a semi-simple Lie algebra \mathfrak{G}'_0 . It follows that no nonzero element of \mathfrak{P}'_0 is invariant by \mathfrak{R}_0 (cf. Helgason [3]). Since the center of \mathfrak{G}_0 is spanned by Z , we have $\mathfrak{P}_0 = \mathfrak{P}'_0 + \{Z\}_R$.

Remark. The lemma above may be derived also from Frobenius reciprocity and the theorem of E. Cartan to the effect that every complex irreducible representation of K appears with multiplicity at most 1 in the regular representation of K on K/K_0 .

5. Eigen-values of the Laplacian

Let R^{N+1} be a Euclidean space of dimension $N + 1$ with natural coordinate system $y = (y^1, \dots, y^{N+1})$. Let $S^N(r)$ be the sphere of radius r about the origin of R^{N+1} , M an n -dimensional submanifold of $S^N(r)$ with local coordinate system x^1, \dots, x^n , and

$$y = y(x^1, \dots, x^n)$$

the local equation defining M . At each point of M , we choose an orthonormal system of unit vectors $\xi_0, \xi_1, \dots, \xi_{N-n}$ such that ξ_0 is normal to $S^N(r)$ and ξ_1, \dots, ξ_{N-n} are tangent to $S^N(r)$ but normal to M . Then

$$\frac{\partial^2 y}{\partial x^j \partial x^k} = \sum_i \Gamma^i_{jk} \frac{\partial y}{\partial x^i} + \sum_{\lambda=1}^{N-n} b^{\lambda}_{jk} \xi_\lambda + b^0_{jk} \xi_0.$$

If we set $g_{jk} = \left(\frac{\partial y}{\partial x^j}, \frac{\partial y}{\partial x^k} \right)$ and denote by (g^{jk}) the inverse matrix of (g_{jk}) , then the Laplacian of $y = (y^1, \dots, y^{N+1})$ as a system of functions on M is given by

$$\Delta y = \sum_{j,k} g^{jk} \nabla_j \nabla_k y = \sum_{\lambda, j, k} g^{jk} b^{\lambda}_{jk} \xi_\lambda + \sum_{j,k} g^{jk} b^0_{jk} \xi_0,$$

where ∇_j denotes the covariant differentiation with respect to $\partial/\partial x^j$. The first term on the right hand side is nothing but the so-called mean curvature normal on M as a submanifold of $S^N(r)$. Hence, M is a minimal submanifold of $S^N(r)$ if and only if

$$\Delta y = \sum_{j,k} g^{jk} b^0_{jk} \xi_0.$$

To simplify the right hand side, we note that

$$\begin{aligned} (y, y) &= r^2, \quad \left(\frac{\partial y}{\partial x^j}, y \right) = 0, \\ \left(\frac{\partial^2 y}{\partial x^j \partial x^k}, y \right) + \left(\frac{\partial y}{\partial x^j}, \frac{\partial y}{\partial x^k} \right) &= 0. \end{aligned}$$

Since $y = r\xi_0$ on M , the last equality above may be rewritten as follows :

$$rb_{jk}^0 + g_{jk} = 0 .$$

Hence, $\sum_{j,k} g^{jk} b_{jk}^0 \xi_0 = -\frac{n}{r^2} y$. We may now conclude

Proposition 5.1. *A submanifold M of $S^N(r)$ is a minimal submanifold of $S^N(r)$ if and only if*

$$\Delta y = -\frac{n}{r^2} y ,$$

where $n = \dim M$.

From Theorem 4.2 and Proposition 5.1 we obtain

Theorem 5.2. *Let $M = G/U = K/K_0$ be a symmetric R -space with G simple, and $\varphi : M \rightarrow \mathfrak{P}$ the imbedding defined in § 2. For each linear functional u of \mathfrak{P} , we set $f = u \circ \varphi$. Then with respect to the metric \ll, \gg on M , f satisfies $\Delta f = -\frac{1}{2}f$.*

Remark. The fact that $\Delta f = \lambda f$ for some λ (independent of f) may be derived from the theorem of Cartan quoted in the remark at the end of § 4. We can then verify $\lambda = -1/2$ using the special function $f_z = u \circ \varphi$.

We wish to relate this eigen-value $-1/2$ with the scalar curvature of M . We denote by $(,)_{\mathfrak{G}}$ and $(,)_{\mathfrak{R}}$ the Killing forms of \mathfrak{G} and \mathfrak{R} , respectively. The curvature tensor R of the symmetric space $M = K/K_0$ is given by

$$R(V, X)Y = -[[V, X], Y] \quad \text{for } V, X, Y \in \mathfrak{M};$$

its Ricci tensor S is given by

$$\begin{aligned} S(X, Y) &= \text{trace of the map } V \rightarrow R(V, X)Y \\ &= \text{trace of the map } V \rightarrow -[[V, X], Y] . \\ &= -\text{trace} ((ad Y)(ad X))|_{\mathfrak{M}} . \end{aligned}$$

If we construct an orthonormal basis for \mathfrak{R} with respect to $-(,)_{\mathfrak{G}}$ by choosing first an orthonormal basis for \mathfrak{K}_0 and then one for \mathfrak{M} , $ad X$ acting on \mathfrak{R} is given by a matrix of the form

$$\begin{pmatrix} 0 & A(X) \\ -{}^t A(X) & 0 \end{pmatrix} .$$

Hence, $(ad Y)(ad X)$ acting on \mathfrak{R} is given by a matrix of the form

$$\begin{pmatrix} -A(Y){}^t A(X) & 0 \\ 0 & -{}^t A(Y)A(X) \end{pmatrix} .$$

It follows that

$$\begin{aligned} (X, Y)_{\mathfrak{R}} &= \text{trace} (ad Y)(ad X)|_{\mathfrak{R}} = -2(\text{trace } {}^t A(Y)A(X)) \\ &= 2 \text{trace} (ad Y)(ad X)|_{\mathfrak{M}} = -2S(X, Y) . \end{aligned}$$

Proposition 5.3. *The Ricci tensor S of a symmetric space $M = K/K_0$ is given by*

$$S(X, Y) = -\frac{1}{2}(X, Y)_{\mathfrak{R}} \quad \text{for } X, Y \in \mathfrak{M} .$$

If we multiply the metric tensor of M by a positive constant a , then both the scalar curvature c of M and the Laplacian Δ of M are multiplied by $1/a$. It is therefore desirable to express the eigen-values of Δ in terms of c . Now we calculate c for some R -spaces. If there exists a positive number μ such that

$$(X, Y)_{\mathfrak{R}} = \mu \cdot (X, Y)_{\mathfrak{G}} \quad \text{for } X, Y \in \mathfrak{R} ,$$

then the scalar curvature c is given by

$$c = \frac{1}{2}n\mu \quad (n = \dim M) .$$

In fact, for $X, Y \in \mathfrak{M}$, we have

$$S(X, Y) = -\frac{1}{2}(X, Y)_{\mathfrak{R}} = -\frac{\mu}{2}(X, Y)_{\mathfrak{G}} = -\frac{\mu}{2}\langle\langle X, Y \rangle\rangle ,$$

and hence $c = \frac{1}{2}n\mu$. For the following six classes of symmetric spaces, this method enables us to calculate the scalar curvature c . (For calculation of μ , we refer the reader to Iwahori [5].)

(1) Irreducible hermitian symmetric space of compact type:

$$\mu = \frac{1}{2}, \quad c = \frac{n}{4} .$$

(2) Real Grassmann manifold of non-oriented p -planes in R^{p+q} , ($p + q > 2$):

$$\mu = \frac{p + q - 2}{2(p + q)}, \quad c = \frac{pq(p + q - 2)}{4(p + q)} .$$

(3) Quaternionic Grassmann manifold of p -planes in quaternionic vector space of dimension $p + q$:

$$\mu = \frac{p + q + 1}{2(p + q)}, \quad c = \frac{pq(p + q + 1)}{p + q} .$$

(4) Group manifold $SO(m)$, ($m > 2$):

$$\mu = \frac{m-2}{2m-2}, \quad c = \frac{1}{8}m(m-2).$$

(5) Group manifold $Sp(m)$:

$$\mu = \frac{m+1}{2m+1}, \quad c = \frac{1}{2}m(m+1).$$

(6) n -sphere, ($n > 1$):

$$\mu = \frac{n-1}{n}, \quad c = \frac{1}{2}(n-1).$$

By calculating the eigen-values of the Casimir operator, Nagano [12] determined the eigen-values of the Laplacian Δ acting on the space of functions on a compact symmetric space K/K_0 with K simple and K/K_0 simply connected (with respect to the invariant Riemannian metric induced from the Killing form of \mathfrak{K}). From Nagano's table we see that, for (1), (3) and (6), there is no eigen-value of Δ between 0 and $-\frac{1}{2}(= -c/(n\mu))$. Every eigen-value of Δ for functions on the Grassmann manifold of non-oriented p -planes in \mathbb{R}^{p+q} appears as an eigen-value of Δ for functions on the Grassmann manifold of oriented p -planes in \mathbb{R}^{p+q} , but not vice versa. From Nagano's table we see that the Laplacian Δ for functions on the Grassmann manifold of non-oriented p -planes in \mathbb{R}^{p+q} has no eigen-value between 0 and $-\frac{1}{2}\left(= -\frac{2c(p+q)}{pq(p+q-2)}\right)$ at least if $p \geq 3$ and $p+q \geq 17$. But we do not know if this is true for all p and q . By the same method we can verify that the Laplacian acting on the space of functions on the group manifold $SO(m)$ (resp. $Sp(m)$) has no eigen-value between 0 and $-\frac{1}{2}\left(= -\frac{4c}{m(m-2)}\right)$ (resp. 0 and $-\frac{1}{2}\left(= -\frac{c}{m(m+1)}\right)$). For eigen-values of the Laplacian for the spaces (1) and (6), see also Obata [14].

Bibliography

- [1] S. S. Chern & R. K. Lashof, *On the total curvature of immersed manifolds*, Amer. J. Math. **79** (1957) 306-318.
- [2] —, *On the total curvature of immersed manifolds II*, Michigan Math. J. **5** (1958) 5-12.
- [3] S. Helgason, *Differential geometry and symmetric spaces*, Academic Press, New York, 1962.
- [4] W. Y. Hsiang, *On the compact homogeneous minimal submanifolds*, Proc. Nat. Acad. Sci. U. S. A. **56** (1966) 5-6.
- [5] N. Iwahori, *Theory of Lie Groups. II* (in Japanese), Iwanami, 1957.
- [6] S. Kobayashi, *Imbeddings of homogeneous spaces with minimum total curvature*, Tôhoku Math. J. **19** (1967) 63-70.

- [7] —, *Isometric imbeddings of compact symmetric spaces*, Tôhoku Math. J. **20** (1968) 21–25.
- [8] S. Kobayashi & T. Nagano, *On filtered Lie algebras and geometric structures I*, J. Math. & Mech. **13** (1964) 875–908.
- [9] N. H. Kuiper, *Immersions with minimal total absolute curvature*, Colloque de geometrie differentielle globale, Bruxelles, 1958, 75–88.
- [10] —, *Der Satz von Gauss-Bonnet für Abbildungen in E^N und damit verwandte Probleme*, Jahresbericht der DMV **69** (1967) 77–88.
- [11] H. Matsumoto, *Quelques remarques sur les groupes de Lie algébriques réels*, J. Math. Soc. Japan **16** (1964) 419–446.
- [12] T. Nagano, *On the minimum eigenvalues of the Laplacians in Riemannian manifolds*, Sci. Papers College Gen. Educ. Univ. Tokyo **11** (1961) 177–182.
- [13] —, *Transformation groups on compact symmetric spaces*, Trans. Amer. Math. Soc. **118** (1965) 428–453.
- [14] M. Obata, *Riemannian manifolds admitting a solution of a certain system of differential equations*, Proc. United States-Japan Seminar in Differential Geometry, Kyoto, 1965, 101–114.
- [15] S. S. Tai, *On minimum imbeddings of compact symmetric spaces of rank one*, J. Differential Geometry **2** (1968) 55–66.
- [16] M. Takeuchi, *Cell decompositions and Morse equalities on certain symmetric spaces*, J. Fac. Sci. Univ. Tokyo, Sec. I, **12** (1965) 81–192.
- [17] T. Takahashi, *Minimal immersions of Riemannian manifolds*, J. Math. Soc. Japan **18** (1966) 380–385.

UNIVERSITY OF TOKYO
UNIVERSITY OF CALIFORNIA, BERKELEY

