

ERRATUM FOR “THE DEGREE THEOREM IN HIGHER RANK”

CHRIS CONNELL & BENSON FARB

Abstract

The purpose of this erratum is to correct a mistake in the proof of Theorem 4.1 of [CF].

1. Introduction

In this note we fix a mistake in Theorem 4.1 of [CF]. This error was pointed out to us by Inkang Kim and Sungwoon Kim, to whom we are extremely grateful. The error occurs in the final step of the proof of Theorem 4.4 in [CF]: the stated angle inequality should hold not just for a subspace V , but for each individual vector in an orthonormal k -frame. The problem with the proof in [CF] occurs at the very end of §5, at the top of page 52. Lemma 5.4 in [CF] applies to all of V'_k , but one needs to justify that this lemma applies to the subspace W' .

The fix. Throughout the present paper we use the notation and terminology of [CF]. The setup is as follows. Let $X = G/K$ be a symmetric space of noncompact type with no local \mathbb{R}, \mathbb{H}^2 or $\mathrm{SL}_3(\mathbb{R})/\mathrm{SO}(3)$ factors. We also assume (cf. §4.2 of [CF]) that X is irreducible. We fix a point $x \in X$ and a maximal flat \mathcal{F} through $x \in X$. The stabilizer of x in G is (after conjugation) K , and K acts by the derivative action on the tangent space $T_x X$, which we identify as a subspace \mathfrak{p} of the Lie algebra $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ of G , endowed with the standard inner product coming from the Killing form B . We identify \mathcal{F} and \mathcal{F}^\perp with their corresponding tangent spaces in $T_x X$. As in §4.4 of [CF], define the *angle* between two subspaces $V, W \subseteq T_x X$ as

$$\angle(V, W) :=$$

$$\inf\{d_{\mathrm{SO}(T_x X)}(I, P) : P \in \mathrm{SO}(H) \text{ with } PV \subset W \text{ or } PW \subset V\}.$$

The first author was in part supported by the Simons Foundation and the CNRS. The second author was in part supported by the NSF.

Received 12/05/2014.

We do not see how to prove Theorem 4.4 of [CF], called there the “Eigenvalue Matching Theorem”, as stated. We instead prove the following result. Call a set of vectors $\{w_1, \dots, w_k\}$ is a δ -*orthonormal k -frame* if $\langle w_i, w_j \rangle < \delta$ for all $1 \leq i < j \leq k$.

Theorem 1.1 (Weak eigenvalue matching). *For each symmetric space X as above, there are constants C_1 and C so that the following holds. Given any $\epsilon < 1/(\text{rank}(X)+1)^2$, for any orthonormal k -frame v_1, \dots, v_k in $T_x X$ with $k \leq \text{rank}(X)$, whose span V satisfies $\angle(V, \mathcal{F}) \leq \epsilon$ there is a $C_1\epsilon$ -orthonormal $2k$ -frame given by vectors $v'_1, v''_1, \dots, v'_k, v''_k$, such that for $i = 1, \dots, k$:*

$$(1.1) \quad \angle(hv'_i, \mathcal{F}^\perp) \leq C\angle(hv_i, \mathcal{F})$$

and

$$(1.2) \quad \angle(hv''_i, \mathcal{F}^\perp) \leq C\angle(hv_i, \mathcal{F})$$

for every $h \in K$, where hv is the linear (derivative) action of K on $v \in T_x X$.

Fortunately, Theorem 1.1 is enough to deduce the main theorem (Theorem 4.1) of [CF]. We make this deduction in §4 below.

Acknowledgements. We are extremely grateful to Inkang Kim and Sungwoon Kim for finding the mistake in [CF], and for their careful readings, questions and comments on the present paper. We are also grateful to the referee for making a number of corrections and suggestions that improved the paper.

2. Proof of Theorem 1.1 when $\epsilon = 0$

By extending a given orthonormal k -frame to an orthonormal basis, it is enough to prove the theorem for the case $k = n$, where $n = \text{rank}(X)$. So let $\{v_1, \dots, v_n\}$ be an orthonormal n -frame in $T_x X$, and let V denote its span. In this section we prove will prove the following:

Theorem 1.1 holds in the special case when $\epsilon = 0$, that is when $V \subset \mathcal{F}$. Further, in this special case, the theorem holds with only the assumption that $\{v_i\}$ spans all of \mathcal{F} , not necessarily that $\{v_i\}$ forms an orthonormal frame.

We thus assume throughout this section that $V \subset \mathcal{F}$. Let K_i denote the stabilizer of v_i . Recall that the Lie algebra of K_i is $\mathfrak{m} \oplus_j \mathfrak{k}_{\alpha_{i_j}}$, where \mathfrak{m} is the Lie algebra of the stabilizer of \mathcal{F} and the sum is taken over the family of all one-dimensional spaces $\mathfrak{k}_{\alpha_{i_j}} \subset \mathfrak{k}$ such that v_i belongs to the kernel of the (positive) root α_{i_j} . We define for each $1 \leq i \leq n$ the subspace

$$Q_i := (\text{span}\{K_i \cdot \mathcal{F}\})^\perp.$$

For each positive root α , we have $[\mathfrak{k}_\alpha, \mathfrak{a}] \subset \mathfrak{p}_\alpha \oplus \mathfrak{a} \subset \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \oplus \mathfrak{g}_0$. In particular, Q_i is spanned by the set of \mathfrak{p}_α for those positive roots $\alpha \neq 0$ such that $\mathfrak{k}_\alpha \not\subset \mathfrak{k}_{v_i}$.

Lemma 2.1 (Vectors in Q_i satisfy (1.1)). *There exists a constant $C > 0$, depending only on $\dim(X)$, so that for any $w \in Q_i$ and any $h \in K$:*

$$(2.1) \quad \angle(w, h \cdot \mathcal{F}^\perp) \leq C \angle(v_i, h \cdot \mathcal{F})$$

where h acts via the derivative action of K on $v \in T_x X$.

Proof. This exact fact was proven in Lemma 5.3 of [CF]: take $V := \text{span}\{v_i\}$ and $V' := \text{span}\{w\}$ and apply the proof of that lemma verbatim starting with the line “If no such constant” The earlier part of the lemma was meant only to produce such a V' . q.e.d.

Let $K_{\mathcal{F}} < K$ denote the stabilizer of \mathcal{F} , and let $m = \dim K/K_{\mathcal{F}} = \dim K - \dim K_{\mathcal{F}}$. Note that $n + m = \dim(X)$. For each positive root α , let $\mathfrak{p}_\alpha := (\mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}) \cap \mathfrak{p}$. Now $[\mathfrak{k}_\alpha, \mathfrak{a}] \subset \mathfrak{p}_\alpha \oplus \mathfrak{a} \subset \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \oplus \mathfrak{g}_0$. For each i choose $a_i \in \mathfrak{a}$ such that $b_i := [k_i, a_i]$ spans $\mathfrak{p}_{\alpha_i} := (\mathfrak{g}_{\alpha_i} \oplus \mathfrak{g}_{-\alpha_i}) \cap \mathfrak{p}$. Replacing b_i by $b_i/\|b_i\|$, we can assume that each b_i has length 1. Note that $b_i \notin \mathfrak{a}$ since $[\mathfrak{k}, \mathfrak{a}] \cap \mathfrak{a} = 0$.

For two distinct roots α, β with $\alpha + \beta \neq 0$, the Killing form satisfies $B(\mathfrak{g}_\alpha, \mathfrak{g}_\beta) = 0$. It follows that $b_i \in \mathcal{F}^\perp$ and $\{b_i\}$ is orthonormal. In particular, since this set has cardinality $\dim(\mathcal{F}^\perp)$, it forms an orthonormal basis for \mathcal{F}^\perp .

Denote the Lie algebra of K_i by \mathfrak{k}_i . Now

$$Q_i = (\text{span}\{K_i \cdot \mathcal{F}\})^\perp = \text{span}\{b_j : b_j \notin [\mathfrak{k}_i, \mathfrak{a}]\}.$$

Since K_i is a proper subgroup of K , for each j there exists i so that $b_j \notin [\mathfrak{k}_i, \mathfrak{a}]$; in particular $b_j \in Q_i$. Note that this i is not necessarily unique. Thus, to summarize, the basis $\{b_i\}$ is adapted to the Q_i in the sense that each b_j belongs to some Q_i and each Q_i is spanned by the collection of b_j 's that it contains.

Lemma 2.2. *There exists a subset of $\{b_i\}$ consisting of $2n$ distinct elements, two from each $Q_i, 1 \leq i \leq n$.*

We prove Lemma 2.2 below. Assuming this for now, let v_i, v'_i denote the pair of vectors in Q_i guaranteed by the lemma. We claim that $\{v'_1, v''_1, \dots, v'_n, v''_n\}$ satisfy the conclusion of Theorem 1.1, thus proving that theorem in the special case $V \subset \mathcal{F}$, which we have assumed throughout this section. To see this, first note that, by definition, the set $\{b_i\}$ is orthonormal, and its span is also orthogonal to V . Since $v'_i, v''_i \in Q_i$, the inequality (2.1) of Lemma 2.1 gives exactly inequalities (1.1) and (1.2) of the theorem, as desired. The rest of this paper is devoted to proving Lemma 2.2.

2.1. Combinatorial Translation. To prove Lemma 2.2 we first translate it into a problem that is purely combinatorial. To this end, let $A = A_G = (a_{ij})$ be the $n \times m$ matrix whose i, j entry is 1 if b_j belongs to Q_i and 0 otherwise. Lemma 2.2 is then the statement that we can pick two 1 entries from each row of A , so that all of our $2n$ choices are in different columns. More formally:

Key Claim (Lemma 2.2 restated): *For each $1 \leq i \leq n$ there exists $1 \leq j_i, k_i \leq m$ with $j_i \neq k_i$ so that each $a_{ij_i} = a_{ik_i} = 1$ and $\bigcup_{i=1}^n \{j_i, k_i\}$ has cardinality $2n$.*

Given the Key Claim, we set $v'_i := b_{j_i}$ and $v''_i := b_{k_i}$, proving (as explained above) Theorem 1.1. The rest of this paper is devoted to proving the Key Claim.

Note that the Key Claim is true for A if and only if it is true for any matrix obtained from A by permuting its rows or columns, as these operations correspond to just re-ordering the Q_i 's and b_j 's, respectively. We think of the $n \times m$ matrix of being a list u_1, \dots, u_n of n row vectors, each in $\{0, 1\}^m$. Before proving properties of the matrix A , we will need the following lemma from Lie theory.

Lemma 2.3 (Codimension of proper Lie subgroups of K). *Let $G \neq \mathrm{SL}(3, \mathbb{R})$ be a connected, simply-connected, simple Lie group with $n := \mathrm{rank}_{\mathbb{R}}(G) \geq 2$. Let K denote the maximal compact subgroup of G . Let $H < K$ be the stabilizer of a vector in \mathcal{F} , and let $d(H) := \dim K - \dim H$. Then:*

- 1) *If $K = \mathrm{SO}(n+1)$ then either $d(H) \geq 2n - 2$ or $d(H) = n$ and H is locally isomorphic to $\mathrm{SO}(n)$.*
- 2) *If $K = \mathrm{SO}(n) \times \mathrm{SO}(n+r)$ with $r \geq 0$ then $d(H) \geq 2n - 2 + r$.*
- 3) *For all other K we have $d(H) \geq 2n - 1$.*

Lemma 2.3 should not be a surprise since the dimension of a rank n compact Lie group K grows quadratically in n , and typically the rank of a proper Lie subgroup of K has rank $< n$, and $(n+1)^2 - n^2 = 2n + 2$.

Proof. The list of maximal compact subgroups K of all possible real and complex, connected, simply-connected, simple Lie groups, including exceptional groups, is given on pages 684–718 of [Kna]. Since we are bounding codimension from below, we can assume that H is a subgroup of a maximal proper subgroup of K .

For each of the K coming from the classical algebras, the list of possible connected Lie subgroups is given in Tables 5–8 found on pages 1018–1027 of [AFG]. (For a simpler list that is sufficient in our case, the maximal Lie subalgebras are found in Tables 1–4 on pages 987–1010 of that same paper.) The list of the exceptional cases can be found in Table 1.7 on page 37 of [Ant], where one interprets the list respectively

as either the complex or real compact form, ignoring the noncompact real split cases.

The dimension of each of these groups is the sum of the dimensions of its simple factors. The dimensions for these can be computed for example, from Table 1 on page 66 of [Hum]. (Note that for the complex groups, to compute the real dimension one must multiply the number of positive roots by two and add the rank.) The possible dimensions of maximal compact lie subgroups follows by going through each case and plugging in the numbers from these tables.

Now there is one general case where the codimensions of maximal proper subgroups do not agree with the codimension bounds listed in the statement, namely there are maximal subgroups of $K = SO(n) \times SO(n+r)$ which are locally isomorphic to $SO(n-1) \times SO(n+r)$ which only have codimension $n-1$ in K . (This case arises for $G = SO(n, n+r)$.) However, any maximal stabilizer of a vector in \mathcal{F} must be locally isomorphic to a subgroup of $SO(n-1) \times SO(n+r-1)$ since neither factor group stabilizes any vector of \mathcal{F} and maximal proper subgroups of $SO(k)$ for any $k > 1$ are locally isomorphic to subgroups of $SO(k-1)$. Hence the codimension of $H < K$ is $2n-2+r$ in this case. q.e.d.

We remark that the data we need from all these tables is really originally due to Dynkin [Dyn, Dyn2] who computed these in the algebraically closed case. However, it is a tedious exercise to extract all of the real reductive, split and compact form cases that arise. This has been done for us in the more modern references cited above.

With Lemma 2.3 in hand, we are now ready to prove properties of the matrix A . Recall that we have reduced the situation to the case where X is irreducible. Thus we only care about simple Lie group G not locally isomorphic to $SL_3(\mathbb{R})$. Let $|u_i|$ denote the number of 1 entries of u_i .

Lemma 2.4 (Properties of A). *Let $G \neq SL_3(\mathbb{R})$ be a connected, simply-connected, simple Lie group with $n := \text{rank}_{\mathbb{R}}(G) \geq 2$. Let K denote the maximal compact subgroup of G . Let $A = A_G$ be defined as above, with row vectors u_1, \dots, u_n . Then the following hold.*

- 1) *Each column of A has at least one entry equal to 1.*
- 2) *$|u_i| \geq n$ for each i .*
- 3) *If K is not locally isomorphic to $SO(n+1)$, then $|u_i| \geq 2n-2$ for each i .*
- 4) *If $u_i = u_j$ then $|u_i| = |u_j| \geq 2n-1$.*
- 5) *For $i \neq j$, if $|u_i| < 2n-2$ and $|u_j| < 2n-2$, then there is at most one k with $a_{ik} = a_{jk} = 1$.*

Proof. If (1) does not hold, then there is some $b \in \{b_i\}$ that does not lie in Q_i for any i . We can write $b = [k, a]$ where k (resp. a) is a positive root vector in \mathfrak{k} (resp. \mathfrak{a}). Let \mathfrak{k}_i be the Lie algebra of K_i .

Since $b \in \{b_i\}$ but $b \notin Q_i = (\text{span}\{K_i \cdot \mathcal{F}\})^\perp$, the fact that $\{b_i\}$ is an orthonormal basis for \mathcal{F}^\perp implies that $b \in \text{span}\{K_i \cdot \mathcal{F}\}$, so we can write $b \in [\mathfrak{k}_i, \mathfrak{a}]$. Since this is true for each i , it follows that $k \in \cap_i \mathfrak{k}_i$. However, since the entire frame $\{v_i\}$ forms a basis for \mathcal{F} and $\exp k$ stabilizes each vector simultaneously, k belongs to the stabilizer of \mathfrak{a} in \mathfrak{k} . In other words, $k \in \mathfrak{k} \cap \mathfrak{g}_0$, contradicting the fact that k belongs to a positive root space. Thus it must be that each b_j lies in some Q_i , proving (1).

To prove (2), we first note that $X = K \cdot \mathcal{F}$. Thus

$$\dim Q_i = \dim \text{span}(\{K_i \cdot \mathcal{F}\})^\perp = \dim(X) - \dim \text{span}\{K_i \cdot \mathcal{F}\}.$$

Since $K_{\mathcal{F}}$ is an extension of the pointwise stabilizer $K'_{\mathcal{F}}$ of \mathcal{F} by a finite group (the Weyl group of G), and so $\dim K_{\mathcal{F}} = \dim K'_{\mathcal{F}}$, we also have

$$\dim X = \dim \text{span}\{K \cdot \mathcal{F}\} = \dim K + \dim \mathcal{F} - \dim K_{\mathcal{F}}$$

and

$$\dim \text{span}\{K_i \cdot \mathcal{F}\} = \dim K_i + \dim \mathcal{F} - \dim K_{\mathcal{F}}$$

since $K_{\mathcal{F}}$ is contained in K_i . Combining the above three equations gives

$$(2.2) \quad \dim Q_i = \dim K - \dim K_i.$$

Items (2) and (3) now follow by applying Lemma 2.3.

We now prove (4). If $u_i = u_j$ then $K_i = K_j$ and therefore v_i and v_j belong to the same singular subspace $W \subseteq \mathcal{F}$, and neither lies in a more singular subspace. Hence $\dim(W) \geq 2$, and so W contains a 1-dimensional subspace that is K' -invariant for some proper subgroup K' of K properly containing K_i . Lemma 2.3 implies that if K' does not already have codimension at least $2n - 1$, then either K is isomorphic to $SO(n+1)$ and K' is necessarily locally isomorphic to $SO(n)$ or else $K = SO(n) \times SO(n)$ and K' is locally isomorphic to $SO(n-1) \times SO(n-1)$. In the second case any proper Lie subgroup of K' already has codimension $2n - 1$ in K . In the first case, K_i can have dimension no larger than that of $SO(n - 1)$, corresponding to a proper Lie subgroup of K' . Therefore K_i has codimension at least $n - 1$ in K' and codimension at least $2n - 1$ in K , as indicated. This proves (4).

For (5), we first note that, by Proposition 2.20.5 of [Ebe], both K_i and K_j are proper semisimple subgroups of K . From Lemma 2.3, we note that the only case where there can exist u_r with $|u_r| < 2n - 2$ is when K is locally isomorphic to $SO(n + 1)$ and K_r is locally isomorphic to $SO(n)$. We assume this is the case, and hence K_i and some K_j are locally isomorphic to $SO(n)$.

From the discussion above, it remains to show that the dimension of $Q_i \cap Q_j$, or the intersection of $\mathfrak{k} \ominus \mathfrak{k}_i$ and $\mathfrak{k} \ominus \mathfrak{k}_j$, is at most one. We note that the basis of Q_i consists of all of those b_r 's whose corresponding $k_r \in \mathfrak{k}$ is not in \mathfrak{k}_i ; similarly, Q_j consists of those b_r for which $k_r \notin \mathfrak{k}_j$.

Hence $Q_i + Q_j$ is spanned by the set of those b_r with $b_r \notin \mathfrak{k}_i \cap \mathfrak{k}_j$. This corresponds exactly to $(\text{span}\{(K_i \cap K_j) \cdot \mathcal{F}\})^\perp$.

The dimension of $Q_i + Q_j$ is therefore the codimension in K of $K_i \cap K_j$. The subgroups K_i and K_j are distinct by (4). The intersection of two distinct copies of $\mathfrak{so}(n)$ in $\mathfrak{so}(n + 1)$ is isomorphic to a subalgebra of $\mathfrak{so}(n - 1)$. Hence $\dim(Q_i + Q_j) \geq 2n - 1$ and hence $\dim Q_i \cap Q_j = \dim Q_i + \dim Q_j - \dim(Q_i + Q_j) \leq 2n - (2n - 1) = 1$, completing the proof. q.e.d.

2.2. Solving the combinatorial problem. By re-ordering and relabeling the rows, we can and will assume $|u_i| \leq |u_{i+1}|$ for all i . Lemma 2.4(2), Lemma 2.4(3) and Lemma 2.3(1) together imply that for each i either $|u_i| = n$ or else $|u_i| \geq 2n - 2$.

We will now describe an algorithm which takes input a subset of row vectors $\{u_i\}$, and at each stage removes one of the vectors, and changes each vector by removing two of its entries. We still call the remaining vectors by the same names u_i . First consider the case that there exists $p > 0$ so that $|u_i| = n$ for each $1 \leq i \leq p$. Set $t = 1$. For any t let $N(i, t)$ denote the number of 1’s left in the vector u_i at the start of Stage t of the algorithm. So for example $N(i, 1) = n$ for all $1 \leq i \leq p$. Now, starting with $t = 1$, perform “Stage t ” of the following algorithm on the row vectors $\{u_1, \dots, u_p\}$:

- Step 1: Re-order the rows so that $N(i, t) \leq N(i + 1, t)$ for each $1 \leq i \leq p - t$.
- Step 2: Choose two 1 entries of the top row; and let j_t, k_t be the column numbers of these two entries.
- Step 3: Delete the top row and the columns j_t and k_t , still calling the remaining vectors u_j by their original name. Now increase the counter t by 1, and go to Step 1.

At each stage we remove two columns corresponding to the columns of two 1 entries of the top row. By Lemma 2.4(5), this implies that at most one 1 is removed from any of the other rows. We thus have that if the vector u_i remains at Stage t then

$$(2.3) \quad N(i, t) \geq N(i, t - 1) - 1.$$

The algorithm can only fail at Step 2. Let d be the smallest t for which Stage t of the algorithm fails. A has at most n rows, so $d \leq n$. Let u_j denote the top row after performing Step 1 at Stage $t = d$. The assumption of failure is then $N(j, d) \leq 1$. Lemma 2.4(5) implies that at each stage $t = 1, \dots, d - 1$, at most one 1 was removed from u_j . But $N(j, 1) \geq n$, so that

$$1 \geq N(j, d) \geq n - (d - 1) = n - d + 1$$

and so $d \geq n$, so that $d = n$ and $N(j, n) = 1$. In other words, the algorithm will succeed in choosing two 1’s from each row u_i with $|u_i| =$

n , except possibly if $|u_i| = n$ for each $1 \leq i \leq n$, in which case the algorithm can only possibly fail at Stage n , at the final row vector u_j .

If $N(j, n) = 2$ we are done, so assume $N(j, n) \leq 1$. Application of (2.3) gives $N(j, n-1) \leq 2$. By our ordering in Step 1, the top row u_k at Stage $t = n-1$ has $N(k, n-1) \leq 2$. A repeated application of (2.3) gives that $N(i, 2) \leq n-1$ for all $2 \leq i \leq p$. This means that each u_2, \dots, u_p must have a 1 in one of the two columns removed from u_1 during Step 2 of Stage 1. Since $N(1, 1) = n > 2$, there exists an entry of u_1 that does not overlap with any other u_j . Instead of choosing the entry of u_1 that overlaps with u_j , choose this entry to remove. Then we can choose the original entry from u_j , so that we do not fail at the last stage.

We have thus shown that the above algorithm always succeeds: we can choose two 1's from each u_i with $|u_i| = n$, all satisfying the Key Claim. Since we are done if every row is of this form, we can now assume that $|u_n| \geq 2n-2$. Note that it may be that $|u_i| = 2n-2$ for each $1 \leq i \leq n$.

Having performed the algorithm successfully on the (possibly empty) $\{u_i : |u_i| = n\}$, we now continue with the algorithm on the remaining vectors $\{u_i : |u_i| \geq 2n-2\}$, not resetting $t = 1$. Since at most two columns are removed at any stage of the algorithm, the only way for the algorithm to fail with some vector u_j with $|u_j| = 2n-2$ or $|u_j| = 2n-1$ in the top row is at Stage $t = n$. If this happens then of the $2n-2$ columns removed in the first $n-1$ stages, at least $2n-3$ of them must have been in columns in which u_j has a 1.

Lemma 2.4(1) states that each column of A has a 1. The number of columns of A is the dimension of the symmetric space X minus the rank. From Table II on page 354 of [Hel], we see that for any given rank $n \geq 2$ the $\dim(X) - n$ is at least $n(n+1)/2$, equality occurring only for the case when $G = \mathrm{SL}(n+1, \mathbb{R})$. Hence the total number of columns of A is always at least $n(n+1)/2$. Thus, after having removed at most $2n-2$ columns, there must be at least $n(n+1)/2 - (2n-2) = (n^2 - 3n + 4)/2 \geq 2$ (for $n \geq 3$ - this is where we are using the hypothesis $G \not\approx \mathrm{SL}_3(\mathbb{R})$) columns not yet removed, which have an entry with 1. Call two of these columns c_1, c_2 . These 1 entries are entries in row vectors u_p, u_q for some $p, q \neq j$, with $p = q$ possible. At some stage u_p was the top row, and two columns were removed corresponding to two 1 entries of u_p . Put back one of these columns and remove c_1 instead. Do the same thing with u_p replaced by u_q and c_1 replaced by c_2 .

We claim that there is not a failure at stage n , with row vector u_j . If $N(j, n) = 0$ then precisely two 1's from u_j were removed at each stage $1, \dots, n-1$, so that the two columns we just replaced now each give a 1 back to u_j , so that the algorithm doesn't fail at u_j . If $N(j, n) = 1$ then it is still the case that of the $2n-2$ columns removed, at most one such

column of u_j did not have a 1 entry. In particular u_j had a 1 removed from one of the columns c_1 or c_2 . Since we replaced this column, and since $N(j, n) = 1$, the replacement gives two 1 entries for u_j , and again the algorithm does not fail at stage n .

We have thus shown that the modified algorithm given above terminates with the choices proving the Key Claim.

3. Finishing the proof of Theorem 1.1

In this section we complete the proof of prove Theorem 1.1.

Let $k_i \in K$ be a closest element to the identity such that $\widehat{w}_i := k_i^{-1}v_i$ lies in \mathcal{F} . If it happens that there is a more singular vector in \mathcal{F} very nearby to \widehat{w}_i , then it may be that k_i could be large (say on the order of π) as it moves \widehat{w}_i through a large rotation around the singular vector, but keeping it very close to \mathcal{F} . Hence we begin by replacing \widehat{w}_i with the most singular vector w_i in the ball of radius $\varepsilon_o = 1/(\text{rank}(X) + 1)^2$ about \widehat{w}_i and that is closest to \widehat{w}_i . (This vector will be unique as the singular subspaces form linear flags.) By the choice of ε_o , the new w_i will be the most singular in its ε_o -ball.

Set $K_i = \text{Stab}_K(w_i)$ and note that K_i will contain the stabilizer of \widehat{w}_i . Since each element k of a stabilizer subgroup not belonging to K_i stabilizes a vector at least $3\varepsilon_o$ away from \widehat{w}_i , it follows that k moves \widehat{w}_i at least a distance of $\frac{\varepsilon_o}{4\pi}d_K(k, 1)$ away from \widehat{w}_i , provided $d_K(k, 1) < \frac{\pi}{4}$.

We will show that there is a small element of K that moves v_i into $K_i \cdot \mathcal{F}$, as follows. Since the derivative at 0 of the exponential map $\exp : \mathfrak{g} \rightarrow G$ is the identity map, we can transport metric estimates to \mathfrak{g} . Therefore, setting \widehat{a}_i to be the lift to \mathfrak{a} of \widehat{w}_i , there is a c_o depending only on ε_o such that each element $u \in \mathfrak{k}$ orthogonal to \mathfrak{k}_i and with $|u| < 1$ has $||[u, \widehat{a}_i]|| \geq c_o|u|$. In particular, the $\frac{\varepsilon_o}{c_o}$ -neighborhood U of 0 in \mathfrak{k} has the property that $[U + \mathfrak{k}_i, \mathfrak{a}] + \mathfrak{a}$ contains the ε -neighborhood of \widehat{a}_i in \mathfrak{p} . Consequently, descending back to X , there is a constant c_1 depending only on ε_o (or equivalently $\text{rank}(X)$) such that smallest element $k'_i = \exp(u) \in K$ such that $v_i \in k'_i K_i \cdot \mathcal{F}$ has $d_K(k'_i, 1) < c_1\varepsilon$.

We also have $\angle(w_i, v_i) < \varepsilon_o + \varepsilon < 2\varepsilon_o$. Since $\{v_i\}$ is orthonormal, it follows that $\{w_i\}$ is $4\varepsilon_o$ -orthonormal, and in particular it is still a frame.

Since $\{w_i\} \subset \mathcal{F}$, we can apply the special case $\epsilon = 0$ of Theorem 1.1 proved in §2. (Recall that for this case, we did not require the $\{w_i\}$ to be orthonormal.) This produces an orthonormal (since $\epsilon = 0$) $2k$ -frame $\{w'_i, w''_i\}$ satisfying the angle inequalities of Theorem 1.1 with v_i, v'_i, v''_i replaced by w_i, w'_i, w''_i . (Observe that w'_i and w''_i also satisfy the angle inequalities with w_i replaced by \widehat{w}_i as well since w'_i and w''_i are orthogonal to all of $K_i \cdot \mathcal{F}$ and K_i contains the stabilizer of \widehat{w}_i .)

Moreover, as proved in the $\epsilon = 0$ case of Theorem 1.1, $w'_i, w''_i \in (K_i \mathcal{F})^\perp$ for each i . Now let $v'_i = k'_i w'_i$, let $v''_i = k'_i w''_i$ and let $z_i = v'_i - w'_i$.

Since $d_K(k'_i, 1) < c_1\varepsilon$ it follows that $|z_i| < c_1\varepsilon$ and

$$\begin{aligned} |\langle v'_i, v'_j \rangle| &= |\langle w'_i + z_i, w'_j + z_j \rangle| \\ &= |0 + \langle w'_i, z_j \rangle + \langle z_i, w'_j \rangle + \langle z_i, z_j \rangle| \\ &\leq 3c_1\varepsilon \end{aligned}$$

for all $1 \leq i, j \leq k$. The same bound holds for $\langle v'_i, v''_j \rangle$ and $\langle v''_i, v''_j \rangle$ by the same computation. Now set $C_1 := 3c_1$. (Note that v'_i and v''_i are also orthogonal to v_i since $k'_i(K_i\mathcal{F})^\perp = (k''_i K_i\mathcal{F})^\perp$.)

Finally $\angle(hv'_i, \mathcal{F}^\perp) = \angle(hk'_i w'_i, \mathcal{F}^\perp) \leq C\angle(hk'_i \widehat{w}_i, \mathcal{F}) = C\angle(hv_i, \mathcal{F})$, and similarly for v''_i . This completes the proof of Theorem 1.1.

4. Proving Theorem 4.1 of [CF]

In this section we prove the main theorem (Theorem 4.1) of [CF]. The proof as given in §4.5 of [CF] needs to be slightly modified, given that we do not know Theorem 4.4 of [CF] as stated, but only the slightly weakened form, Theorem 1.1 above.

On page 41 of [CF] we choose $\varepsilon = 1/(\text{rank}(X) + 1)$. We now instead choose ε so small that $\varepsilon < 1/(\text{rank}(X) + 1)^2$ and so that for any t , when $\sin(t) < \varepsilon$ then $\sin(t) > t/2$. This new choice of constant of course still depends only on $\text{rank}(X)$, and the only affect of this change will be to change the resulting constants in the proof of the theorem. As in [CF], we let L_1, \dots, L_k be the $k \leq \text{rank}(X)$ eigenvalues of the positive semi-definite quadratic form Q_2 that are strictly less than ε . As stated in [CF], if no such eigenvalues exist then we are done, so we assume $k \geq 1$. Label the L_i so that $0 \leq L_1 \leq \dots \leq L_k$. Denote by v_i the eigenvector associated to L_i .

Plugging the formula $r(v) = \sin^2 \angle(v, \mathcal{F})$, given on page 42 of [CF], into the formula for L_i given on the last line of page 41 of [CF], gives

$$L_i = \int_{\partial_F X} \sin^2 \angle(kv_i, \mathcal{F}) d\sigma_y^s(k).$$

Recall that we are identifying $\partial_F X$ with K/M , whose elements we write as elements of K , remembering that they are really equivalence classes. For each i let

$$A_i := \{k \in \partial_F X : \sin^2(\angle kv_i, \mathcal{F}) \leq \sqrt{L_i}\}$$

and let $B_i := \partial_F X - A_i$.

We claim that for each fixed i , each $k \in A_i$ moves v_i a small angle from \mathcal{F} . To see this, note that for any $k \in A_i$:

$$\sqrt{L_i} \geq \sin^2 \angle(kv_i, \mathcal{F}) > (\angle(kv_i, \mathcal{F})/2)^2$$

so that $\angle(kv_i, \mathcal{F}) \leq 2(L_i)^{1/4}$, as desired. Here we have used our choice of ε to obtain the second inequality.

Now

$$\begin{aligned} L_i &= \int_{\partial_F X} \sin^2 \angle(kv_i, \mathcal{F}) d\sigma_y^s(k) \geq \int_{B_i} \sin^2 \angle(kv_i, \mathcal{F}) d\sigma_y^s(k) \\ &\geq \sqrt{L_i} \cdot \sigma_y^s(B) \end{aligned}$$

so that $\sigma_y^s(B_i) \leq \sqrt{L_i} \leq \sqrt{\epsilon}$ for each i . Since we have chosen $\epsilon < 1/(\text{rank}(X) + 1)^2$, we obtain

$$\sigma_y^s(B_1 \cup \dots \cup B_k) \leq k\sqrt{\epsilon} \leq \text{rank}(X)\sqrt{\epsilon} < \frac{\text{rank}(X)}{\text{rank}(X) + 1} < 1.$$

Since by definition A_i is the complement B_i^c and $\sigma_y^s(\partial_F(X)) = 1$, it follows that $A := A_1 \cap \dots \cap A_k \neq \emptyset$. Any element in A moves all of $V = \text{span}\{v_1, \dots, v_k\}$ to within $2\epsilon^{1/4}$ of \mathcal{F} .

We have just proved that there exists an element $k_0 \in A$ with the property that $\angle(k_0 v_i, \mathcal{F}) \leq 2\epsilon^{1/4}$ for each i . Now apply Theorem 1.1 to $\{k_0 v_i\}$, and note that the vectors $\{v'_i, v''_i\}$ produced satisfy the same inequalities with $k_0 v_i$ replaced by v_i . We now apply these inequalities in the string of inequalities starting on line 2 of Page 43 of [CF], with only one modification, namely, the first line should now read:

$$\det Q_1 \leq C' \prod_{i=1}^k \langle v'_i Q_1, v'_i \rangle \langle v''_i Q_1, v''_i \rangle$$

Where $C' = \frac{1}{(1-C_1\epsilon)^{4k}}$ and C_1 is from Theorem 1.1. This uniform constant will also be carried along in the rest of the inequalities and then absorbed into the final constant C .

References

- [Ant] F. Antoneli, *Subalgebras maximais das Álgebras de Lie semisimples*, Quebra de Qimetria e o Código Genético, Ph.D. Thesis, Instituto de Matemática e Estatística, Universidade de Sao Paulo, 1998.
- [AFG] F. Antoneli, M. Forger & P. Gaviria, *Maximal subgroups of compact Lie groups*, J. Lie Theory, Vol. 22(4) (2012), 949–1024.
- [CF] C. Connell & B. Farb, *The Degree Theorem in Higher Rank*, J. Diff. Geom., Vol. 65 (2003), 19–59.
- [Dyn] E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, (Russian) Mat. Sbornik N.S. Vol. 30(72), (1952). 349–462 (3 plates). MR 0047629.
- [Dyn2] E. B. Dynkin, *On semisimple subalgebras of semisimple Lie algebras*, (Russian) Doklady Akad. Nauk SSSR (N.S.) Vol. 81, (1951). 987–990. MR 0045102.
- [Ebe] P. Eberlein, *Geometry of nonpositively curved manifolds*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1996. vii+449 pp. ISBN: 0-226-18197-9; 0-226-18198-7. MR 1441541.

- [Hel] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Corrected reprint of the 1978 original. Graduate Studies in Mathematics, 34. American Mathematical Society, Providence, RI, 2001. xxvi+641 pp. ISBN: 0-8218-2848-7. MR 1834454.
- [Hum] J. Humphreys, *Linear Algebraic Groups*, Springer GTM, Vol. 21, 1989.
- [Kna] A. W. Knapp, *Lie groups beyond an introduction*, Second edition. Progress in Mathematics, 140. Birkhuser Boston, Inc., Boston, MA, 2002. xviii+812 pp. ISBN: 0-8176-4259-5. MR 1920389.

DEPARTMENT OF MATHEMATICS
INDIANA UNIVERSITY
RAWLES HALL
831 EAST 3RD ST
BLOOMINGTON, IN 47405
E-mail address: connell@indiana.edu

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CHICAGO
5734 S. UNIVERSITY AVE.
CHICAGO, IL 60637
E-mail address: farb@math.uchicago.edu