

ON A QUESTION OF KAPLANSKY II

P.G. WALSH

(Communicated by F. Luca)

*This paper is dedicated to the memory of
Professor Irving Kaplansky*

ABSTRACT. There is a question attributed to Irving Kaplansky concerning the solvability of the quadratic equation $x^2 - py^2 = a$ in the case that the prime $p = a^2 + (2b)^2$. This question was answered in the affirmative by Mollin [1], although according to [3], this result is implicit in the work of Gauss and Legendre. The proof appearing in [1] was later simplified in [4], and it was also shown therein that Kaplansky's question was a special case of a more general result. Using the method of proof in [4], Mollin [2] has recently extended the results of [4], but upon further consideration, it appears that there is a more general phenomenon occurring, and also, that one of the assumptions in the main theorem of [2] is unnecessary. In this paper we prove this generalization, and eliminate one of the assumptions stated in the main result of [2]. The proof is again based on the method described in [4].

1. INTRODUCTION

In an earlier article [4], the author generalized a result of Mollin, and at the same time, simplified the method of proof. Recently, Mollin has used this same elementary approach to further the results of [4]. The purpose of this present paper is to extend the results of [2]. The method remains the same as in [4], with the appropriate modifications described in [2] in order to deal with parity issues.

Theorem 1.1. *Let $d \equiv 1 \pmod{4}$ be a positive integer, and assume that $n = a^2 + db^2$ for positive integers a, b with a odd and $(n, a) = 1$. Assume further that there is a positive integer c , with $(a, c) = 1$ for which the equation*

$$X^2n - Y^2d = c^2$$

is solvable in coprime positive integers X, Y . Then there exists a (possibly trivial) factorization rs of nd , and a divisor f of σc , for which the equation

$$rx^2 - sy^2 = af$$

is solvable in positive integers x, y , where $\sigma = 2$ if n is odd and c is even, and $\sigma = 1$ otherwise.

Received by the editors August 16, 2007 and in revised form, September 11, 2007.
2000 *Mathematics Subject Classification.* Primary: 11D09, 11D85.
Key words and phrases. Pell equation.

For simplicity, Theorem 1 only deals with the case $d \equiv 1 \pmod{4}$. A similar statement holds for the other cases, which we leave as an exercise for the reader.

We note that Theorem 1 not only extends the result of [2], which dealt with the particular case $d = 1$, but moreover removes an unnecessary assumption contained in the statement of the main theorem in [2]. Specifically, it is assumed therein that the quadratic equation $X^2 - nY^2 = -1$ is solvable. As the conclusion of the main theorem in [2] does not place any restriction on the constructed factors r and s of n , there is no need, during the course of the proof, to multiply by a unit of norm -1 . Therefore, we do not need to include an analogous assumption (that the quadratic equation $x^2n - y^2d = 1$ be solvable in positive integers x, y) in the statement of Theorem 1 above.

2. PROOF OF THEOREM 1

Let (T, U) be coprime positive integers which satisfy $T^2n - U^2d = c^2$, and let α be defined $\alpha = T\sqrt{n} + U\sqrt{d}$. Let $\beta = \sqrt{n} + b\sqrt{d}$, and define integers u, v by $u = Tn + Ubd, v = Tb + U$. Then

$$\alpha\beta = (Tn + Ubd) + (Tb + U)\sqrt{nd} = u + v\sqrt{nd}$$

is an element in $\mathbf{Z}[\sqrt{nd}]$ with norm a^2c^2 .

Let $g = (u, v)$, then clearly g divides c^2 , but in fact, g divides c . We provide the details for this assertion, as the reasoning in [2] appears to be flawed. Suppose that p is a prime dividing g , with p^μ properly dividing g ($\mu > 0$), and such that p^μ does not divide c . Note that p divides c because p^μ divides c^2 . It follows that p^μ divides both u and v , hence $p^{2\mu}$ divides $u^2 - v^2nd = a^2c^2$. By assumption, $p^{2\mu}$ does not divide c^2 , and so p must divide a , contradicting the fact that $(a, c) = 1$. We conclude that g divides c , and from the equation $u^2 - v^2nd = a^2c^2$, we deduce that

$$(1) \quad (u/g)^2 - a^2(c/g)^2 = ((u/g) + a(c/g))((u/g) - a(c/g)) = (v/g)^2nd.$$

We now break up the argument into three cases, depending on the relative parities of n and c . We note that n and c cannot both be even, as this would contradict either $(n, a) = 1$ or $(T, U) = 1$.

Case 1: c even, n odd.

In this case, as n, a and d are odd, and $n = a^2 + db^2$, it follows that b is even. Also, the assumption that c is even implies that T and U are odd (as they are coprime), whence it follows that both u and v are odd, which by equation (1) implies that there are integers A, B, r, s , with $v/g = AB$ and $nd = rs$, satisfying

$$(u/g) + a(c/g) = A^2r, \quad (u/g) - a(c/g) = B^2s,$$

from which it follows that

$$A^2r - B^2s = af,$$

with $f = 2(c/g)$.

Case 2: c odd, n even.

In this case, since $d \equiv 1 \pmod{4}$ and $(a, b) = 1$, it follows that $n \equiv 2 \pmod{4}$, and that b is odd. By considering the equation $T^2n - U^2d = c^2$ modulo 4, it is readily verified that both T and U are odd. We conclude that $u = Tn + Ubd$ is odd, and that $v = Tb + U$ is even. Therefore, there are integers A, B, r, s , with $nd = rs$ and $v/g = 2AB$, satisfying

$$(u/g) + a(c/g) = 2A^2r, \quad (u/g) - a(c/g) = 2B^2s,$$

from which it follows that

$$A^2r - B^2s = af,$$

with $f = c/g$.

Case 3: c odd and n odd.

Since $d \equiv 1 \pmod{4}$, it follows that $n \equiv 1 \pmod{4}$, and again by considering the equation $T^2n - U^2d = c^2$ modulo 4 we deduce that T is odd and that U is even. Therefore, in this case we find again that $u = Tn + Ubd$ is odd and that $v = Tb + U$ is even, and the rest of the proof for this case follows as in the previous case.

Acknowledgement The author gratefully acknowledges support for his research from the Natural Sciences and Engineering Research Council of Canada

REFERENCES

- [1] R.A. MOLLIN. *Proof of some conjectures by Kaplansky*, C.R. Math. Rep. Acad. Sci. Canada **23** (2001), 60-64.
- [2] R.A. MOLLIN. *On a generalized Kaplansky conjecture*, Int. J. Contemp. Math. Sciences **2** (2007), 411-416.
- [3] N. Tzanakis [M.R. 1913340 (2003g:11027)], Reviews of the A.M.S., 2003.
- [4] P.G. WALSH, *On a conjecture of Kaplansky*, Amer. Math. Monthly **109** (2002), 60-61.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD ST., OTTAWA, ONTARIO, CANADA, K1N-6N5

E-mail address: gwalsh@mathstat.uottawa.ca