

ON SINGULAR CHAINS AND CYCLES

BY S. LEFSCHETZ

1. *Introduction.* The theory of the topological invariance of the absolute or relative combinatorial characters of a complex, as developed in our Colloquium Lectures on *Topology* (Chapter II), was based, following Alexander and Veblen, upon the concept of singular chain. Our presentation, and indeed any known to us, appears to give rise to many misconceptions which it is proposed to clear up in the present note. Unless otherwise stated the notations are those of *Topology*.

2. *Singular Cells.* Let \mathcal{R} be a topological space and let e_p be a simplicial oriented cell such that there exists a continuous single-valued transformation (=c.s.v.t.) T of the point set e_p into a subset E_p of \mathcal{R} , where $E_p = Te_p$. The symbol (e_p, T, E_p) , associated with the set \bar{E}_p is called a *singular oriented p -cell on \mathcal{R}* . If e_p' is another e_p , there exists a barycentric transformation U of \bar{e}_p' into \bar{e}_p : $U\bar{e}_p' = \bar{e}_p$. If we set $T' = TU$, it is evident that (e_p', T', E_p) defines also a singular oriented p -cell on \mathcal{R} . We shall agree to consider it as identical with the first:

$$(1) \quad (e_p', T', E_p) = (e_p, T, E_p).$$

This has the advantage of freeing the notion of singular cell from a too narrow connection with a specific image e_p .

3. *Singular Chains.* The singular p -chain C_p on \mathcal{R} is now defined as the association of a symbol

$$(2) \quad C_p = \sum t_i(e_p^i, T^i, E_p^i)$$

with coefficients t belonging to one of the three rings (rational numbers, integers, integers mod m) considered in *Topology*, together with the set of all sets \bar{E}_p^i corresponding to t 's $\neq 0$. As a special case the e 's might be cells of a finite complex k such that there exists a c.s.v.t. T of k into a subset of \mathcal{R} . Then the chain symbol may take the form

$$(3) \quad C_p = \sum t_i(e_p^i, T, E_p^i),$$

and C_p may be considered as the image of the subchain

$$(4) \quad c_p = \sum t_i e_p^i$$

of k , but that is not essential. In this instance we might have represented C_p by the symbol $(\sum t_i e_p^i, T, \sum t_i E_p^i)$ analogous to the cell symbol. Observe also that we may find for any chain (2) an equivalent representation (3). For we may take the cells e_p^i to be simplexes in some S_n whose closures do not meet, then define T as coincident with T^i on e_p^i . The closure of the sum of the cells e_p^i will then be k , and (2) will assume the form (3).

If we have several singular chains C_p^i , then $\sum s_i C_p^i$, where the coefficients s_i belong to the same ring as those of the chains, defines a p -chain which is called the linear combination with coefficients s_i of the chains C_p^i . We have thus moduli of singular p -chains wholly analogous to the moduli of subchains of a complex.

4. *Boundary Relations.* Returning to (e_p, T, E_p) , let the boundary relations for e_p be

$$(5) \quad e_p \rightarrow \sum \eta_i e_{p-1}^i = F(e_p).$$

Since T is a transformation of \bar{e}_p into \bar{E}_p , it transforms e_{p-1}^i into a subset E_{p-1}^i of \bar{E}_p and hence $(e_{p-1}^i, T, E_{p-1}^i)$ is a singular $(p-1)$ -cell on \mathcal{R} . The singular $(p-1)$ -chain

$$(6) \quad F(e_p, T, E_p) = \sum \eta_i (e_{p-1}^i, T, E_{p-1}^i)$$

is called the boundary of (e_p, T, E_p) and we write here also

$$(7) \quad (e_p, T, E_p) \rightarrow F(e_p, T, E_p).$$

The boundary of the chain (2) is now by definition

$$(8) \quad F(C_p) = \sum t_i F(e_p^i, T^i, E_p^i),$$

for which we also write

$$(9) \quad C_p \rightarrow F(C_p).$$

Owing to (1) this boundary depends solely on C_p but not on the particular transformations T^i that occur in (8). Let C_p be in the special form (3) with an associated non-singular image (4), and let the boundary relation for c_p be

$$(10) \quad c_p \rightarrow \sum s_i e_{p-1}^i.$$

Then we have well defined singular cells $(e_{p-1}^i, T, E_{p-1}^i)$ and we find that

$$(11) \quad C_p \rightarrow \sum s_i(e_{p-1}^i, T, E_{p-1}^i),$$

which may be described by the statement: the boundary of a singular image of a chain is the singular image of the boundary of the chain.

5. *Degenerate Case.* Let \mathcal{R} undergo a c.s.v.t. U into a new space \mathcal{R}' . Then the singular cell (e_p, T, E_p) will go over into a singular p -cell (e_p, UT, UE_p) and C_p into

$$(12) \quad UC_p = \sum t_i(e_p^i, UT^i, UE_p^i).$$

To different representations of the same singular p -cell on \mathcal{R} there will merely correspond different representations of the same singular p -cell on \mathcal{R}' , and to $F(C_p)$ there will now correspond $F(UC_p)$. In particular also if $C_p \equiv 0$, likewise $UC_p \equiv 0$.

The preceding observations have an immediate application to degenerate cells. Let (e_p, T, E_p) be a singular cell on \mathcal{R} , and let us suppose that there exists a simplex σ_q , $q < p$, and two c.s.v.t.'s T', T'' , such that T' is a simplicial transformation of e_p into σ_q and that $T'' \cdot T' = T$. The cell (e_p, T, E_p) is called a *singular degenerate p -cell* on \mathcal{R} and chains made up exclusively of such cells are called *degenerate chains*. If $\mathcal{R}' = U\mathcal{R}$ as above, the degenerate cells and chains of \mathcal{R} go over into degenerate cells and chains of \mathcal{R}' .

According to *Topology*, Chapter II, No. 2, $F(e_p, T', \sigma_q)$ is a degenerate $(p-1)$ -chain, and hence when (e_p, T, E_p) is degenerate so is its boundary. Hence this holds likewise as regards degenerate p -chains. Let us agree to consider all degenerate chains as identically zero. By the observation just made degenerate chains will then completely disappear from all boundary relations.

6. *Homologies.* From the preceding section it appears clearly that when \mathcal{R} and \mathcal{R}' are homeomorphic, the homeomorphism between them associates respectively to one another their moduli of p -chains, of bounding p -chains and their degenerate p -chains. These are therefore topological and the homology characters derived from the moduli are topological invariants.

Regarding these homologies, we introduce them exactly as for complexes. In particular if A intersects B in a set $A \cdot B$ closed

relatively to B , the neglect of the singular cells (e_p, T, E_p) , such that $E_p \subset A$, leads to the characters of $B \bmod A$.

7. *Invariance of the Combinatorial Homology Characters.* Suppose now that \mathcal{R} itself is a finite simplicial complex K and let ϵ_p^i designate its cells. They can be considered as the singular cells $(\epsilon_p^i, 1, \epsilon_p^i)$ and it is readily seen that the formal singular boundary relations involving these cells alone are the same as the combinatorial relations between the cells ϵ_p^i themselves. Therefore whenever only singular cells of this type are involved, the singular boundary relations (7) for K are reduced to the combinatorial relations.

The invariance of the combinatorial homology characters of K is established by identifying them with the corresponding topological characters. The steps in the proof are as follows.

(a) Let C_p be a singular chain on K which we assume henceforth in the simplified form (11) with T and the non-singular prototype c_p fixed. There exists an $\eta > 0$ depending on K but not on C_p , such that when mesh $C_p < \eta$, the chain can be homotopically deformed into a subchain C_p' of K , the deformation keeping each cell on the closure of the cell of K that carries it. This is the deformation theorem (*Topology*, p. 86). It implies (*loc. cit.*, p. 78) that there are deformation chains, all singular, indicated by \mathcal{D} , such that

$$(13) \quad \begin{aligned} \mathcal{D} C_p &\rightarrow C_p' - C_p - \mathcal{D}F(C_p), \\ C_p' &= \sum s_i E_p^i. \end{aligned}$$

(b) If mesh $C_p > \eta$ the chain has a subdivision chain C_p whose mesh is suitable. Subdivision is defined as in *Topology* (p. 85), by reference to a subdivision of c_p .

(c) Suppose that C_p possesses certain cells (not necessarily p -cells) which belong to K and whose sum is therefore a sub-complex K_q of K . Then the subdivision and deformation in (b) may be so chosen as to leave K_q fixed point for point. The proof indicated in *Topology* (p. 87, Remark I) only shows that C_p may be so modified as to leave the cells of K_q invariant individually but not point for point. The more accurate result, which is of interest for its own sake, is proved as follows. We show by induction as in *Topology* (p. 86) that the deformations there indicated leave K_q invariant point for point provided that

any cell of C_p without vertices on K_q is of diameter $< \eta$, and that a cell having a face in common with K_q has all its points not farther than η from that face. If C_p does not fulfill these conditions we find by reference to c_p that a suitable subdivision of C_p without new vertices on K_q will behave as required. For if c_p is a subchain, say of k , there is a subcomplex k' of k such that $T \cdot k' = K_q$. We can then apply to k a series of subdivisions differing from regular subdivisions only in so far that no new vertices are ever introduced on k' . Given any ζ we can thus obtain a subdivision c_p^* of c_p whose cells fulfill relatively to k' and ζ the two conditions that we wish to impose upon the cells of C_p relatively to K_q and η . Since T is continuous the required result follows for C_p .

Consider now the boundary relations mod L , where L is a subcomplex of K . Let Γ_p be a (singular) cycle mod L . By (a) there is a subdivision Γ_p' of Γ_p homotopically deformable into a subcycle Γ_p^* of K , its points on L remaining on L , and $\Gamma_p' \sim \Gamma_p$ mod L on Γ_p itself (*Topology*, p. 87) and hence a fortiori on K . By (c), if $C_{p+1} \rightarrow \Gamma_p'$ mod L , there is a subdivision C_{p+1}' of C_{p+1} with the same boundary Γ_p' , deformable into a subchain C_{p+1}^* by a homotopy leaving Γ_p' invariant, so that $C_{p+1}' \rightarrow \Gamma_p'$ mod L . Therefore if the initial cycle ~ 0 mod L in the topological sense, the reduced cycle ~ 0 mod L in the combinatorial sense. From this follows immediately as in *Topology* (p. 88), that the topological and combinatorial homology groups of the same types are simply isomorphic and hence have the same numerical invariants. Therefore the combinatorial homology characters are topological invariants.

8. *Remarks.* I. Once the notion of singular cell has become familiar one will naturally abandon the explicit (too explicit) (e, T, E) notation in favor of the simpler E of *Topology*.

II. The following circumstance may arise in connection with our definition of singular cell. Taking for the sake of simplicity $p=2$, let $e_2 = ABC$ be an (oriented) isosceles triangle with $AB = AC$ and let AD be the altitude issued from A . Let U be the symmetry about AD and T a transformation $= 1$ on ADB , $= U$ on ADC . If we set $T' = T \cdot U$, $e_2' = ACB$, we have

$$(e_2, T, E_2) = (e_2', TU, E_2) = (-e_2, TU, E_2) = (-e_2, T, E_2)$$

and therefore

$$2(e_2, T, E_2) = 0.$$

Owing to this, E. Čech, who pointed out this circumstance to us, suggested that in the present and in the similar instance for any p , the singular cell be also considered as degenerate. The more extended meaning to be thus attached to degenerate cells, while justifiable, is not however essential.

PRINCETON UNIVERSITY

VARIABLES CORRELATED IN SEQUENCE*

BY A. T. CRAIG

1. *Introduction.* If each of n variables, x_1, x_2, \dots, x_n , represents a quantitative character of an individual, and if the variables are correlated in sequence, that is, x_1 is correlated with x_2 , x_2 is correlated with x_3 , \dots , and in general x_i is correlated with x_{i+1} , it seems natural to inquire about the correlation between a character, say x_1 , of one individual and a character, say x_3 , of a second individual, with the condition imposed that the two individuals have identical measurements with regard to the character x_2 . It is this problem with which we shall be primarily concerned in the present paper. As we proceed, we shall place appropriate restrictions upon the nature of the correlation which exists between the variables. We shall, however, make no assumptions regarding the correlation between the variables other than that between them in adjacent pairs.

In order to provide a convenient point of departure and to exhibit a set of variables correlated in sequence, we shall first consider a rather elementary problem which arises when measurements are made under a constant law of probability.

2. *The Correlation between Measurements under a Constant Law of Probability.* Let the variable t obey a constant law of probability $f(t) = 1/a$, $0 \leq t \leq a$. Let successive sets of n independent measurements each, say t_1, t_2, \dots, t_n , be made upon t . We may, without loss of generality, suppose the meas-

* Presented to the Society, April 8, 1932, under the title *Some properties of correlated variables.*