

## THE LOCAL RING OF THE GENUS THREE MODULUS SPACE AT KLEIN'S 168 SURFACE

BY H. E. RAUCH<sup>1</sup>

Communicated by E. Calabi, November 28, 1966

1. **Introduction.** In [6], as a synthesis of earlier papers of mine, I give, in the form of a set of prescriptions for local coordinates, a description of  $M^g$ , the space of conformal equivalence classes of compact Riemann surfaces of genus  $g$ , as a complex space. Particular interest attaches to those points (surface classes) of  $M^g$  representing surfaces admitting conformal self-maps (automorphisms) because, outside of certain cases for  $g=1, 2, 3$  (over and above the elliptic and hyperelliptic involutions for  $g=1, 2$ ), these points are singular (non-uniformizable) points in the structure. In particular for  $g \geq 2$ , where one needs  $3g-3$  complex parameters to describe  $M^g$  near a generic point, one needs  $3g-3+\rho$ ,  $\rho > 0$ , near one of the points in question.<sup>2</sup> According to Prescription III ([6, p. 17]) the problem reduces to finding an irreducible basis for the homogeneous nonconstant, polynomial invariants of a finite group of linear transformations in  $3g-3$  variables, namely, the hermitian adjoint of the group induced on the quadratic differentials of a representative surface of the point in question by the conformal automorphism group of that surface.

For a finite nonabelian linear group, while there is an algorithm for computing some basis for the invariants (cf. Prescription III), there is notoriously no known algorithm for computing an *irreducible* basis, i.e., for discarding the superfluous ones. Accordingly I felt it of interest to illustrate the whole phenomenon by a nontrivial example. To anyone who has worked on the subject the one that immediately comes to mind is Klein's surface of genus three admitting as automorphism group a representation of the simple group of order 168 ([1], [3]). This example commends itself in that it is of "maximum complexity" in the sense that it admits its full quota according to Hurwitz [2] of  $84(g-1) = 168$ , ( $g=3$ ) automorphisms (on the subject of such surfaces see the interesting papers [4], [5]). It develops, *vide infra*, that eleven invariants, i.e.,  $11 = 3 \cdot 3 - 3 + 5 = 6 + 5$ , so  $\rho = 5$ , are needed to generate the local ring of  $M^3$  at Klein's surface class.<sup>2</sup>

---

<sup>1</sup> Research partially sponsored by the Air Force Office of Scientific Research, Office of Aerospace Research, U. S. Air Force, under AFOSR Grant No. AF-AFOSR-1077-66.

<sup>2</sup> With  $\rho$  relations on "syzygies," of course.

2. **The quadratic differentials on Klein's surface.** Klein's surface  $S$  is the compact Riemann surface of genus three obtained by identifying the upper half-plane under the principal congruence subgroup of level seven of the inhomogeneous modular group with appropriate conventions at cusps and vertices. The factor group  $G$  acts on  $S$  as its group of conformal automorphisms representing faithfully thereby the unique simple group of order 168.

LEMMA 1 ([3, p. 444]). *There is a basis  $z_1, z_2, z_3$  of the abelian differentials of first kind on  $S$  on which  $G$  induces a representation  $R_3(G)$  generated by*

$$\begin{aligned}
 (1) \quad & T: z'_1 = \epsilon z_1, \quad z'_2 = \epsilon^4 z_2, \quad z'_3 = \epsilon^2 z_3 \\
 & U: z'_1 = az_1 + bz_2 + cz_3 \\
 & \quad z'_2 = bz_1 + cz_2 + az_3 \\
 & \quad z'_3 = cz_1 + az_2 + bz_3,
 \end{aligned}$$

where  $\epsilon = e^{2\pi i/7}$ ,  $\sqrt{-7} = \epsilon + \epsilon^2 + \epsilon^4 - \epsilon^3 - \epsilon^5 - \epsilon^6$ , and  $a = (\epsilon^5 - \epsilon^2)/\sqrt{-7}$ ,  $b = (\epsilon^3 - \epsilon^4)/\sqrt{-7}$ ,  $c = (\epsilon^6 - \epsilon)/\sqrt{-7}$  are all real.  $T$  has order 7, and  $U$  has order 2.

LEMMA 2. *As a basis for the quadratic differentials on  $S$  one can choose*

$$\begin{aligned}
 (2) \quad & \xi_1 = z_1^2, \quad \xi_2 = z_2^2, \quad \xi_3 = z_3^2, \\
 & \xi_4 = \sqrt{2} z_1 z_2, \quad \xi_5 = \sqrt{2} z_2 z_3, \quad \xi_6 = \sqrt{2} z_1 z_3.
 \end{aligned}$$

The representation  $R_6(G)$  induced on (2) by  $R_3(G)$  is unitary.

PROOF.  $S$  is not hyperelliptic ([3, p. 437]), hence Noether's theorem ([6, Lemma 5]) implies that any six distinct quadratic products of the  $z$ 's, in particular (2), span the quadratic differentials. By Lemma 1,  $R_6(G)$  is generated by

$$\begin{aligned}
 (3) \quad & T_2: \xi'_1 = \epsilon^2 \xi_1, \xi'_2 = \epsilon \xi_2, \xi'_3 = \epsilon^4 \xi_3, \xi'_4 = \epsilon^5 \xi_4, \xi'_5 = \epsilon^6 \xi_5, \xi'_6 = \epsilon^3 \xi_6, \\
 & U_2: \xi'_1 = a^2 \xi_1 + b^2 \xi_2 + c^2 \xi_3 + \sqrt{2} ab \xi_4 + \sqrt{2} bc \xi_5 + \sqrt{2} ac \xi_6, \dots, \\
 & \quad \xi'_4 = \sqrt{2} ab \xi_1 + \sqrt{2} bc \xi_2 + \sqrt{2} ac \xi_3 + (ac + b^2) \xi_4 \\
 & \quad + (ab + c^2) \xi_5 + (bc + a^2) \xi_6, \dots,
 \end{aligned}$$

where the dots signify cyclic permutation of  $a, b, c$ .  $T_2$  is clearly unitary.  $U_2$  is real symmetric by inspection and of order 2 by Lemma 1, hence real orthogonal, *a fortiori* unitary.

LEMMA 3. *Define*

$$\begin{aligned}
 (4) \quad & \gamma_k(\xi) = \alpha(\epsilon^{-2k} \xi_1 + \epsilon^{-k} \xi_2 + \epsilon^{-4k} \xi_3) + \beta(\epsilon^{-6k} \xi_4 + \epsilon^{-6k} \xi_5 + \epsilon^{-3k} \xi_6), \\
 & \bar{\gamma}_k(\xi) = \beta(\epsilon^{2k} \xi_1 + \epsilon^k \xi_2 + \epsilon^{4k} \xi_3) + \alpha(\epsilon^{5k} \xi_4 + \epsilon^{6k} \xi_5 + \epsilon^{3k} \xi_6),
 \end{aligned}$$

where  $\alpha^2 = (-1 + \sqrt{-7})/\sqrt{8} = (\epsilon + \epsilon^2 + \epsilon^4)/\sqrt{2}$ ,  $\beta = \bar{\alpha}$ ,  $\beta^2 = (\epsilon^3 + \epsilon^5 + \epsilon^6)/\sqrt{2}$ ,  $\alpha\bar{\alpha} = \beta\bar{\beta} = 1$ , and  $k=0, \dots, 6$ . One has (i)  $R_6(G)$  induces on the  $\gamma_k(\xi)$  and the  $\bar{\gamma}_k$  respectively the degree seven permutation representation  $R_7(G)$  of  $G$  and its inverse  $R_7^{-1}(G)$  and (ii)

$$(5) \quad \Sigma\gamma_k(\xi) = \Sigma\bar{\gamma}_k(\xi) = 0,$$

$$(6) \quad \gamma_{1-k}(\xi) = (1/\sqrt{2})(\bar{\gamma}_k(\xi) + \bar{\gamma}_{k+1}(\xi) + \bar{\gamma}_{k+3}(\xi)),$$

$$\bar{\gamma}_{1-k}(\xi) = (1/\sqrt{2})(\gamma_k(\xi) + \gamma_{k+1}(\xi) + \gamma_{k+3}(\xi)),$$

where  $k$  is computed mod 7, and

$$(7) \quad \xi_1 = (1/7\alpha)\Sigma\epsilon^{2k}\gamma_k(\xi), \dots$$

PROOF. (i) and (ii) follow from the substitution of (2) in the formulae in [1, p. 519] and [1, II, p. 459 and p. 501] and the corresponding facts noted there. However, once aware of (5), (6), (7) one easily verifies them directly. As for (i) one sees immediately that  $T_2$  generates the cyclic permutation (6543210) of the indices of the  $\gamma_k(\xi)$  and the inverse permutation of the  $\bar{\gamma}_k(\xi)$ . Calculation reveals that  $U_2$  induces the (self-inverse) permutation (12)(36) on both  $\gamma_k(\xi)$  and  $\bar{\gamma}_k(\xi)$ . These permutations generate the representations of (i). (7) implies their faithfulness.

### 3. Invariants of $R_6(G)$ and main theorem.

THEOREM A (GORDAN [1]). Let  $x_k, \bar{x}_k, k=0, \dots, 6$  be two sets of variables which satisfy conditions (i) and (ii) of Lemma 3 when formally substituted for  $\gamma_k(\xi), \bar{\gamma}_k(\xi)$  respectively. Then any (nonconstant, homogeneous) polynomial in  $x_0, \dots, x_6$  over the ring of integers which is invariant under  $R_7(G)$  is a polynomial over  $R(\sqrt{2})$  ( $R$  is the rationals) in the power sums  $S_i, \bar{S}_i, i=1, \dots, 7$  of the  $x$ 's and  $\bar{x}$ 's (separately). By assumption  $S_1 = \bar{S}_1 = 0$ . One easily verifies  $S_2 = \bar{S}_2$ . The basis of eleven invariants  $S_2, S_j, \bar{S}_j, j=3, \dots, 7$  is irreducible, i.e., none is a polynomial (over  $\mathbb{C}$ ) in the others.

Gordan also computes explicitly a set of five syzygies but does not prove they are a basis for all syzygies.

Define, for a set of variables  $\lambda_1, \dots, \lambda_6$  and positive integral  $j$

$$(8) \quad S_j(\lambda) = \Sigma(\gamma_k(\lambda))^j, \quad \bar{S}_j = \Sigma(\bar{\gamma}_k(\lambda))^j.$$

THEOREM 1. Let  $t \in T^3$  (Teichmueller space of genus three) lie over  $[S] \in M^3$  (modulus space of genus three), where  $S$  is Klein's surface. One can introduce coordinates in  $T^3$  near  $t$  such that the local ring of  $M^3$  at  $[S]$  is generated by the irreducible basis of eleven distinct homogeneous polynomials obtained by setting  $j=2, \dots, 7$  in (8).

PROOF. Introducing  $\lambda_1, \dots, \lambda_6$  by Prescription III, using (2) as a basis for  $A(S)$  ([6, p. 17—in Proposition 8 the reference should be to Prescription I, not II]), I have to find an irreducible basis for the nonconstant homogeneous polynomials in the  $\lambda$ 's invariant under the hermitian adjoints of the matrices of  $R_6(G)$ . But by Lemma 2 and the group property this will be identical with a basis for the invariants of  $R_6(G)$ . Lemma 3 and Theorem A with

$$(9) \quad x_k = \gamma_k(\lambda), \quad \bar{x}_k = \bar{\gamma}_k(\lambda), \quad k = 0, \dots, 6$$

show that the invariants of Theorem 1 are a basis. The sticky point is that under the specialization (9) they might reduce. However, Gordan in [1, II, p. 461], says that even under the more severe specialization (9) and  $\lambda_1\lambda_2 = 2\lambda_4^2$ , etc., the worst that can happen is  $\beta S_3(\lambda) = \alpha \bar{S}_3(\lambda)$ . A computation shows that this does not happen without the additional specialization.

#### BIBLIOGRAPHY

1. P. Gordan, *Ueber Gleichungen siebenten Grades mit einer Gruppe von 168 Substitutionen*, Math. Ann. 20 (1882), 515–530; II, *ibid.* 25 (1885), 459–521.
2. A. Hurwitz, *Über algebraische Gebilde mit eindeutigen Transformationen in sich*, Mathematische Werke, Bd. I, Birkhäuser, Basel, 1932, pp. 392–430.
3. F. Klein, *Ueber die Transformation siebenter Ordnung der elliptischen Funktionen*, Math. Ann. 14 (1879), 428–471.
4. A. M. Macbeath, *On a theorem of Hurwitz*. Proc. Glasgow Math. Assoc. 5 (1961), 90–96.
5. ———, *On a curve of genus 7*, Proc. London Math. Soc. 15 (1965), 527–542.
6. H. E. Rauch, *A transcendental view of the space of algebraic Riemann surfaces*, Bull. Amer. Math. Soc. 71 (1965), 1–39.

BELFER GRADUATE SCHOOL OF SCIENCE, YESHIVA UNIVERSITY