


TRANSVERSALITY STRUCTURES AND P.L. STRUCTURES ON SPHERICAL FIBRATIONS

BY NORMAN LEVITT AND JOHN W. MORGAN

Communicated by William Browder, May 18, 1972

The first author in [L] introduced the notion of “Poincaré transversality” for an N -dimensional spherical fiber space, $\pi: \xi^N \rightarrow X$. If $T(\xi^N)$ is the Thom space of ξ^N , then we consider $T(\xi^N) = M_\xi \cup c(S(\xi))$ where M_ξ is the mapping cylinder of π and $S(\xi)$ is the total space of ξ^N . A map $f: \Delta^{N+i} \rightarrow T(\xi^N)$ is Poincaré transversal if f is p.l. transversal to $S(\xi) \subset T(\xi)$ with $(f^{-1}(M_\xi), f^{-1}(S(\xi)))$ a codimension 0 submanifold of Δ^{N+i} with the inclusion $f^{-1}(S(\xi)) \subset f^{-1}(M_\xi)$ the spherical fibration induced by f over $f^{-1}(M_\xi)$. This implies $f^{-1}(M_\xi)$ is a Poincaré duality space, (P.D. space), of dimension i with boundary $f^{-1}(M_\xi) \cap \partial\Delta^{N+i}$, and that $(f^{-1}(M_\xi), f^{-1}(S_\xi))$ is its normal tube.



A p.l. manifold M^j mapping by $f: M^j \rightarrow T(\xi^N)$ is Poincaré transversal to ξ^N if and only if $f|(\text{any simplex})$ is. If f is Poincaré transversal then $f^{-1}(M_\xi)$ is a P.D. space with boundary $f^{-1}(M_\xi) \cap \partial M$ and of dimension $j - N$. One of the main results of [L] is to develop a theory to study the problem of when a map $f: M^j \rightarrow T(\xi^N)$ may be shifted to be Poincaré transversal. To do this one introduces the space $W(\xi^N)$, of Poincaré transversal maps of $\Delta^i \rightarrow T(\xi^N)$ for all i . In [L] and [J] it is proved that if F_{ξ^N} denotes the homotopy theoretic fiber of $W(\xi^N) \rightarrow T(\xi^N)$, then $\pi_i(F_{\xi^N}) \cong \pi_{i-N}(G/PL)$ for $i - N \neq 1, 2, \text{ or } 3$. In fact a map of fiber spaces $\xi^N \rightarrow \zeta^N$ induces $F_{\xi^N} \rightarrow F_{\zeta^N}$ and this map is an isomorphism on π_i for $i - N \neq 1, 2, \text{ or } 3$. Also if $f: M^j \rightarrow T(\xi^N)$, then homotopying f until it is Poincaré transversal is equivalent to lifting f up to homotopy to $W(\xi^N)$. In this announcement we shall describe further results in this theory.

AMS 1970 subject classifications. Primary 55F60, 57C50.

Key words and phrases. Transversality, spherical fibrations, p.l. structures.

Copyright © American Mathematical Society 1972

By a theory of transversality, τ , for a spherical fiber space, $\pi: \xi^N \rightarrow X$, we mean an assignment to each $f: \Delta^i \rightarrow T(\xi)$ a deformation of f until it is Poincaré transverse on Δ^i and all its faces. These assignments are required to be compatible with inclusions of faces $\Delta^{i-1} \subset \Delta^i$. Two theories of transversality τ_0 and τ_1 are equivalent (concordant) if there is a theory of transversality for $\xi^N \times I \rightarrow X \times I$ which, when restricted to $X \times i$, is τ_i for $i = 0, 1$. In the language of [L], a theory of transversality is exactly a section of $W(\xi^N) \rightarrow T(\xi^N)$.

A p.l. structure for ξ^N is a p.l. block bundle (see [R-S]), $\pi: E^N \rightarrow X$, and a fiber homotopy equivalence of $S(E^N)$ to ξ^N .

Let $PL(\xi^N)$ denote the concordance classes of p.l. structures on ξ^N . Clearly p.l. transversality in E provides a theory of transversality for ξ^N . This defines a function from $PL(\xi^N)$ to concordance classes of theories of transversalities for ξ^N .

THEOREM A. *Let $\pi: \xi^N \rightarrow X$ be a spherical fiber space with $N \geq 3$ and X four connected, then the function $PL(\xi^N) \rightarrow \{\text{concordance classes of theories of transversalities for } \xi^N\}$ is a bijection. In particular ξ^N has a p.l. structure if and only if it has a theory of transversality.*

This is the bundle analogue of Winkelkemper’s philosophy that “transversality unlocks the secret of a manifold.” The homotopy theoretic formulation of Theorem A states that given X four connected and $\xi^N: X \rightarrow BSG(N)$ then ξ^N lifts to $E: X \rightarrow BSPL(N)$ if and only if $W(\xi^N) \rightarrow T(\xi^N)$ has a section. The latter in turn is equivalent to the existence of a lift in the following diagram:

$$\begin{array}{ccc}
 & & WSG(N) \\
 & \nearrow & \downarrow \\
 T(\xi^N) & \longrightarrow & MSG(N)
 \end{array}$$

The problem of lifting $X \rightarrow BSG(N)$ to $X \rightarrow BSPL(N)$ involves the space $G(N)/PL(N) = G/PL$. The problem of lifting $T(\xi^N) \rightarrow MSG(N)$ to $T(\xi^N) \rightarrow WSG(N)$ involves the space $F_{SG(N)}$. We now consider the relation of these two spaces. The results in [L] show that they have the same homotopy groups shifted by N dimensions (except for low dimensions). To actually compare these two spaces we need to extend G/PL to a connected spectrum. There are several ways of doing this. One is to use Sullivan’s calculation of the homotopy type of G/PL as KO -theory at odd primes and cohomology at 2 (with one low dimensional twist), and then define $G/PL \langle k \rangle$ to be $bo \langle k \rangle$ at odd primes ($bo \langle k \rangle$ is Ω^{4l-k} ($4l$ connected cover of BO)) and cohomology shifted k dimensions (with the twist) at 2; see [S]. Another approach is to define $G/PL \langle k \rangle$ to be Quinn’s

space $L_k(e)$. This is the semisimplicial complex of surgery problems shifted k -dimensions; see [Q]. These approaches lead to the same results.

THEOREM B. *There is a map $\phi: F_{\xi^N} \rightarrow G/PL\langle N \rangle$ which is an isomorphism on π_i for $i \neq N + 1, N + 2,$ and $N + 3$ provided only that $N \geq 3$.*

ϕ is a realization on the space level of the isomorphism on homotopy groups given in [L].

II. Homotopy theoretic reformulations. If $\pi: \xi^N \rightarrow X$ is a spherical fiber space, then we may form a connected spectrum $T(\xi)$. The i th space is $T(\xi^N \oplus \varepsilon^{i-N})$ for $i \geq N$ and the maps are the usual ones $\sum T(\xi^N \oplus \varepsilon^{i-N}) \rightarrow T(\xi^N \oplus \varepsilon^{i-N+1})$. In [L] it is shown that the spaces $\{W(\xi^N \oplus \varepsilon^{i-N})\}$ also form a spectrum, $W(\xi)$, and that there is a map of spectra

$$F_\xi \rightarrow W(\xi) \rightarrow T(\xi).$$

The spectrum fiber (or cofiber) F_ξ is made up of the spaces $F_{\xi^N \oplus \varepsilon^{i-N}}$. Let G/PL be the spectrum whose i th space is $G/PL\langle i \rangle$. Thus we have $\phi: F_\xi \rightarrow G/PL$ with $\phi_\#$ an isomorphism on π_i for $i \geq 4$.

We can give another description of the spectrum F_ξ . Let $\gamma^i \rightarrow BSG(i)$ be the universal i -dimensional spherical fiber space. Then $T(\gamma^i) = MSG$. $p: W(\gamma^i) \rightarrow T(\gamma^i)$ with $W(\gamma^i) = p^{-1}(M_{\gamma^i}) \cup p^{-1}(c(S(\gamma^i)))$. Let $E_i = p^{-1}(c(S(\gamma^i)))$. The $\{E_i\}$ form a spectrum ε .

PROPOSITION. $\varepsilon \cong F_\gamma$.

This follows from

THEOREM C.

$$\begin{array}{ccc} p^{-1}(S(\gamma^i)) & \xrightarrow{p} & S(\gamma^i) \\ \cap & & \cap \\ p^{-1}(M_{\gamma^i}) & \xrightarrow{p} & M_{\gamma^i} \end{array}$$

is a homotopy equivalence of pairs in dimensions less than $2i$.

Thus $\varepsilon \rightarrow W(\gamma) \rightarrow T(\gamma)$ ($= MSG$) is a cofibration of spectra. This proves $\varepsilon \cong F_\gamma$.

III. Idea of proofs of Theorems A and B. The proof of Theorem A reduces inductively to the following. Let $\eta = \xi | X^{(n)}$ be given a p.l. structure, $n \geq 5$.

$$\begin{array}{ccc} E & \longrightarrow & \xi | X^{(n)} \\ \downarrow & & \downarrow \\ X^{(n)} & = & X^{(n)} \end{array}$$

Thus for any $(n + 1)$ cell τ_{n+1} in X we have $\xi | \tau_{n+1}$ with a canonical trivialization and $E | \partial\tau_{n+1} \rightarrow \xi | \partial\tau_{n+1}$ a p.l. structure on the boundary. This is equivalent to a p.l. bundle, $E | \partial\tau_{n+1}$, and a fiber homotopy trivialization, i.e., an element in $\pi_n(G/PL)$. This defines a cocycle σ whose class in $H^{n+1}(X; \pi_n(G/PL))$ is the obstruction to extending the p.l. structure relative to $X^{(n-1)}$ over $X^{(n+1)}$.

The p.l. structure defines a theory of transversality for $\xi | X^{(n)}$. Again considering τ_{n+1} , we have $\xi | \tau_{n+1}$ with a theory of transversality for $\xi | \partial\tau_{n+1}$. $\xi | \tau_{n+1}$ is canonically trivial as a spherical fiber space, and thus there is a map $S^N \times S^N \rightarrow T(\xi^N | \tau_{n+1})$. Using the theory of transversality over the boundary, we get $f^{-1}(M_{\xi|\partial\tau})$, $f^{-1}(S_{\xi|\partial\tau})$ an n -dimensional G -framed P.D. space in $S^N \times S^N$. But G -framed P.D. bordism in dimension n is isomorphic to $\pi_n(G/PL)$ for $n \geq 4$. The isomorphism assigns to $(Y, Z: \nu_Y \rightarrow \mathbf{R}^N)$ the surgery obstruction of the problem $Z^{-1}(0) \rightarrow Y$ where Z is shifted p.l. transverse to 0. Thus we have a cocycle $\sigma' \in C^{N+n+1}(T(\xi^N)) \otimes \pi_n(G/PL)$. Theorem A is proved by showing $\sigma' = \Phi(\sigma)$ where Φ is the Thom isomorphism.

To prove Theorem B, we need only calculate F_{ε^N} since in [L] it is proved that all the fibers F_{ε^N} are the same homotopy type. Using the characteristic homomorphism (with a shift of N dimensions) definition of $G/PL\langle N \rangle$, we need only assign to $M^{N+i} \rightarrow F_{\varepsilon^N}$ a surgery problem between P.D. spaces. To define the homomorphism we take the usual surgery obstruction of the problem. We do not need to know that this is the only obstruction to doing surgery on P.D. spaces, only that it is an obstruction. A map $M^{N+i} \rightarrow F_{\varepsilon^N}$ is equivalent to a transversal map $f: M^{N+i} \rightarrow T(\varepsilon^N)$ together with a homotopy of f to zero in $T(\varepsilon^N)$, $F: M^{N+i} \times I \rightarrow T(\varepsilon^N)$. Shift F slightly until it is p.l. transverse regular to $pt \in T(\varepsilon^N)$. Let V^{i+1} be the pre-image. $f^{-1}(M_{\varepsilon^N})$ is an i -dimensional P.D. space and if we have not shifted F too much then $V^{i+1} \cap M^{N+i} \times 0 \subset f^{-1}(M_{\varepsilon^N})$ is a degree one normal map.

IV. Applications.

COROLLARY 4.1. *A stable spherical fiber space $\xi \rightarrow X$ has a p.l. structure if and only if it has an MSPL orientation lifting its MSG orientation.*

PROOF. A p.l. bundle has a natural MSPL orientation. Conversely, $MSPL \rightarrow MSG$ factors through $W(\gamma)$. Thus lifting the canonical MSG orientation to an MSPL orientation also lifts it to a $W(\gamma)$ orientation.

COROLLARY 4.2. *Let M^n be a 4-connected P.D. space, then M^n is homotopy equivalent to a closed p.l. manifold if and only if M^n satisfies Poincaré duality for the homology and cohomology theories associated*

to MSPL with a fundamental class which reduces to the MSG fundamental class.

PROOF. Use S-duality and 4.1.

COROLLARY 4.3. (THE INTRINSIC BROWDER-NOVIKOV THEOREM). *Let M^n be a 4-connected P.D. space. A necessary and sufficient condition for M^n to be the homotopy type of a closed p.l. manifold is that M^n have two regular neighborhoods $R' \subset R \subset S^{n+K}$, for some large K , with R triangulated so that for every simplex σ^j , $(\sigma^j \cap R', \partial\sigma \cap R')$ is a codimension 0 submanifold which is the tube of a P.D. pair of formal dimension $j - K$.*

Thus M^n is homotopy equivalent to a closed p.l. manifold if and only if M^n may be put in "Poincaré general position" with respect to the simplices of some triangulation of its regular neighborhood. See also [J].

COROLLARY 4.4. *Let $f: N^n \rightarrow M^n$ be a homotopy equivalence of 4-connected manifolds and \mathcal{M}_f be the mapping cylinder of f . Then $(\mathcal{M}_f, N \cup M)$ is a Poincaré pair. If the section $T(v(N \cup M)) \rightarrow W(v(N \cup M))$ corresponding to the natural manifold structure on $N \cup M$ extends to a section $T(v(\mathcal{M}_f)) \rightarrow W(v(\mathcal{M}_f))$ then f is homotopic to a p.l. homeomorphism.*

REFERENCES

- [J] L. Jones, *Patch spaces* (to appear).
- [L] N. Levitt, *Poincaré duality cobordisms*, Ann. of Math. (to appear).
- [S] D. Sullivan, *Geometric topology*, Mimeographed notes, Princeton University, Princeton, N.J., 1967.
- [Q] F. Quinn, Thesis, Princeton University, Princeton, N.J., 1970.
- [R-S] C. P. Rourke and B. Sanderson, *Block bundles*. I, Ann. of Math. (2) **87** (1968), 1-28. MR **37** #2234a.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540

Current address (Norman Levitt): Department of Mathematics, Rutgers University, New Brunswick, New Jersey 08903

Current address (John W. Morgan): Department of Mathematics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139