

INITIAL BOUNDARY VALUE PROBLEM FOR COMPRESSIBLE EULER EQUATIONS WITH RELAXATION

QIANGCHANG JU*

Institute of Applied Physics and Computational Mathematics

P.O. Box 8009-28

Beijing, 100088, P. R. China

YONG LI†

College of Science

Beijing University of Technology

Beijing, 100022, P. R. China

RONGHUA PAN‡

School of Mathematics

Georgia Institute of Technology

Atlanta, GA 30332, USA

(Communicated by Toka Diagana)

Abstract

In this paper, we study the global existence of L^∞ weak entropy solution to the initial boundary value problem for compressible Euler equations with relaxation and the large time asymptotic behavior of the solution. Motivated by the sub-characteristic conditions, we proposed some structural conditions on the relaxation term comparing with the pressure function. These conditions are proved to be sufficient to construct the global L^∞ entropy weak solution and to prove the equilibrium state is the global attractor of all physical weak solutions. Furthermore, the convergence rate is proved to be exponential in time. The proof is based on the entropy dissipation principle.

AMS Subject Classification: 35Q35, 35L04, 35L65.

Keywords: Compressible Euler Equations, Relaxation, Entropy weak solutions, Large time behavior, Compensated compactness.

*E-mail address: qiangchang_ju@yahoo.com

†E-mail address: liy@bjut.edu.cn, Li's research is supported by NSFC grant 10541001.

‡E-mail address: panrh@math.gatech.edu, Pan's research is supported by NSF under grant DMS-0807406.

1 Introduction

The relaxation phenomena occurs in many physical applications, including gas dynamics away from thermo-equilibrium, chromatography, river flow, traffic flow, reacting flow and etc; see for instance [33]. In this paper, we consider the compressible Euler equations with relaxation

$$\begin{cases} \rho_t + (\rho u)_x = 0, \\ (\rho u)_t + (\rho u^2 + P(\rho))_x = \frac{Q(\rho) - \rho u}{\tau}. \end{cases} \quad (1.1)$$

Here ρ, u and P denote the density, velocity and pressure. Assuming the flow is a polytropic perfect gas with the adiabatic exponent γ , then $P(\rho) = P_0 \rho^\gamma$, $\gamma > 1$, where P_0 is a positive constant. τ denotes the relaxation time. Without loss of generality, we take $P_0 = \frac{1}{\gamma}$, $\tau = 1$ throughout this paper. Since the particle velocity is not well-defined at vacuum, the momentum $m = \rho u$ is often used to rewrite (1.1) as follows:

$$\begin{cases} \rho_t + m_x = 0, \\ m_t + \left(\frac{m^2}{\rho} + P(\rho) \right)_x = Q(\rho) - m. \end{cases} \quad (1.2)$$

The system (1.2) is supplemented by the following initial value and boundary conditions:

$$\begin{cases} \rho(x, 0) = \rho_0(x), m(x, 0) = m_0(x), 0 < x < 1, \\ m|_{x=0} = 0, m|_{x=1} = 0, t \geq 0, \\ m_0(0) = m_0(1) = 0, \\ \int_0^1 \rho_0(x) dx = \rho^* > 0. \end{cases} \quad (1.3)$$

The last condition in (1.3) is imposed to avoid the trivial case: $\rho \equiv 0$.

It is clear that the system (1.2) is hyperbolic with characteristic speeds

$$\lambda_1 = \frac{m}{\rho} - \rho^\theta, \lambda_2 = \frac{m}{\rho} + \rho^\theta, \quad (1.4)$$

with $\theta = \frac{\gamma-1}{2}$. The corresponding Riemann invariants are

$$w = \frac{m}{\rho} + \frac{\rho^\theta}{\theta}, z = \frac{m}{\rho} - \frac{\rho^\theta}{\theta}. \quad (1.5)$$

The nonlinearity of the problem, along with the degeneracy at vacuum where $\rho = 0$ and relaxation effect in the source term, gives the system (1.2) rich phenomena to study. The vanishing relaxation limit of (1.1) as $\tau \rightarrow 0$ toward the equilibrium system

$$\rho_t + Q(\rho)_x = 0 \quad (1.6)$$

and the related wave stability problem in large time are two significant problems. In the past two decades, the mathematical theories on hyperbolic conservation laws with relaxation have been developed extensively since the pioneer work of Liu [22]. Both the zero relaxation limit and the nonlinear stability of elementary waves were studied in a great deal,

see [2], [3], [17], [8],[9], [10], [11], [12], [16], [17], [18], [19], [20], [21], [25], [26], [27], [28], [29], [34], [35], [37] and [38] for a limited partial list of references. The purpose of current paper is to study the fundamental problems on the global existence of weak solution and the large time behavior of the the solutions to (1.2)–(1.3).

It has been a general belief that the nonlinearity and hyperbolicity lead to the blow up of smooth solutions in finite time when the dissipation in the source term can not win the competition for some initial data. For the system (1.2), the study of [34] and [19] clearly shows that when initial data is not restricted in certain range, the solution will break down in finite time. Therefore, as in current paper, when the intial data could be large and rough, the weak solution is the choice. We now give the defintion of the weak solution to our problem.

Definition 1.1. For every $T > 0$, we define a weak solution of (1.2)-(1.3) to be a pair of bounded measurable functions $v(x, t) = (\rho(x, t), m(x, t))$ satisfying the following pair of integral identities:

$$\int_0^T \int_0^1 (\rho \phi_t + m \phi_x) dx dt + \int_{t=0} \rho_0 \phi dx = 0, \quad (1.7)$$

$$\begin{aligned} & \int_0^T \int_0^1 \left(m \phi_t + \left(\frac{m^2}{\rho} + P(\rho) \right) \phi_x \right) dx dt \\ & + \int_0^T \int_0^1 (Q(\rho) - m) \phi dx dt + \int_{t=0} m_0 \phi dx = 0, \end{aligned} \quad (1.8)$$

for all $\phi \in C_0^\infty(I_T)$ satisfying $\phi(x, T) = 0$ for $0 \leq x \leq 1$ and $\phi(0, t) = \phi(1, t) = 0$ for $t \geq 0$, where $I_T = (0, 1) \times (0, T)$. Moreover, (ρ, m) satisfy the initial boundary conditions (1.3) in the sense of trace.

In order to identify the physical relevant weak solutions, the entropy admissible condition is often imposed, motivated by the second law of thermodynamics. For system (1.2), the entropy and entropy flux pairs are defined as follows.

Definition 1.2. A pair of functions $\eta(\rho, m)$ and $q(\rho, m)$ is called an entropy-entropy flux pair if it satisfies the following equations

$$\nabla q = \nabla \eta \nabla f,$$

where

$$f = \left(m, \frac{m^2}{\rho} + P(\rho) \right).$$

Among all entropies, the most natural entropy is the mechanical energy

$$\eta_e(\rho, m) = \frac{m^2}{2\rho} + \frac{\rho^\gamma}{\gamma(\gamma-1)},$$

which plays a very important role in estimates for entropy dissipation measures.

Definition 1.3. The weak solution $v(x, t) = (\rho(x, t), m(x, t))$ defined in Definition 1.1 is said to be entropy admissible if for any convex entropy η and associated entropy flux q , the following entropy inequality holds

$$\eta_t + q_x + \eta_m(m - Q(\rho)) \leq 0, \quad (1.9)$$

in the sense of distribution.

Regarding the study of the L^∞ weak entropy solution of hyperbolic system with source term, very few references are available. Concerning to our system (1.2), a closely related system is the compressible Euler equations with linear damping where $Q(\rho) = 0$. For the latter system, the existence of weak entropy solutions and large time behavior has been studied in [4, 13, 14, 32] for Cauchy problem, and in [30] for initial boundary value problem. Motivated by the method in [13, 14, 30], we shall construct the global L^∞ entropy weak solutions to (1.2)-(1.3) by means of Godunov scheme and the compensation compactness frameworks see [5, 6, 23, 24]. Based on the entropy dissipation and energy method, the exponential decay of any L^∞ weak entropy solutions to the equilibrium state is shown under some assumptions on relaxation term $Q(\rho)$.

On the way to our goal, one interesting issue is to solicit the appropriate conditions to ensure the uniform L^∞ estimate for the solutions and the stability of the equilibrium $(\rho^*, 0)$ under any perturbation with finite amplitude. In the traditional setting, the sub-characteristic condition

$$\lambda_1 < Q'(\rho) < \lambda_2 \quad (1.10)$$

is often proposed to ensure at least the linear stability of the equilibrium with small perturbation; see for instance [3] and [33]. However, in the context of stability with large amplitude perturbation, the sub-characteristic condition (1.10) seems not enough to ensure the stability. Indeed, since one expects $Q(\rho)$ and m approach to each other, some deeper relations are waiting for further investigations. We shall propose some in this paper as an attempt in this direction.

For this purpose, we define the following quantity

$$\alpha_0 = \max\left\{\sup_x w_0(x), -\inf_x z_0(x)\right\}, \quad (1.11)$$

which will measure the L^∞ bounds of the solution. Here, w_0 and z_0 are initial Riemann invariants. For the convenience of presentation, we also introduce the following notations

$$\begin{aligned} f_1(\rho, \rho^*) &= P(\rho) - P(\rho^*) - P'(\rho^*)(\rho - \rho^*) \equiv f_2(\rho, \rho^*)(\rho - \rho^*)^2, \\ f_3(\rho, \rho^*) &= [P(\rho) - P(\rho^*)](\rho - \rho^*) \equiv f_4(\rho, \rho^*)(\rho - \rho^*)^2, \\ f_5(\rho, \rho^*) &= \frac{Q(\rho) - Q(\rho^*)}{\rho - \rho^*}. \end{aligned} \quad (1.12)$$

Clearly, the above functions f_i are well defined when $\rho \neq \rho^*$, the difference quotients will be replaced by the corresponding derivatives when $\rho = \rho^*$ for the definition of f_2 , f_4 and f_5 .

In section 3, the following assumption on $Q(\rho)$ plays an important role in the proof of uniform L^∞ bound for solutions to (1.2)-(1.3).

(A1) For $0 \leq \rho \leq (\theta\alpha_0)^{1/\theta}$, $Q(\rho)$ is C^2 and it holds that

$$|Q(\rho)| \leq \rho \left(\alpha_0 - \frac{\rho^\theta}{\theta} \right), \quad (1.13)$$

where α_0 is defined in (1.11).

Remark 1.4. This assumption simply asks the bounds on $Q(\rho)$ near vacuum and the large ρ . One observes that from (1.13) that $Q(0) = 0$ which is necessary to confirm that $(0, 0)$ is a solution to (1.2)-(1.3). Since $\frac{Q(\rho)}{\rho}$ is the difference quotient of $Q(\rho)$ over the interval $[0, \rho]$, the inequality (1.13) is closely related to the sub-characteristic condition (1.10). The existence of such $Q(\rho)$ is obvious. For example, we can take $Q(\rho) = \varepsilon\rho(\alpha_0 - \frac{\rho^\theta}{\theta})$, or $Q(\rho) = \varepsilon\rho(\frac{\rho^\theta}{\theta} - \alpha_0)$, where $0 \leq \varepsilon \leq 1$. It is also worthy to remark that the right hand side of (1.13) with $\theta = 1$ appears in the traffic flow models, see [33].

To investigate the large time behavior of weak entropy solutions to the initial boundary value problem (1.2)-(1.3), we will need the following strong **sub-slope condition** on $Q(\rho)$.

(A2) There are $a_1 > 0$, $a_2 > 0$, $a_3 > 0$ and $1/a_1 + 1/a_2 + 1/a_3 = 1$ such that, for $0 < \rho \leq M$, it holds that

$$f_4(\rho, \rho^*) > \frac{a_1 M}{\rho} [f_5(\rho, \rho^*)]^2, \quad (1.14)$$

and

$$\frac{8C^*}{a_2 a_3} f_4(\rho, \rho^*) \geq [f_5(\rho, \rho^*)]^2, \quad (1.15)$$

where $C^* = \frac{1}{\gamma}(\rho^*)^{\gamma-1}$.

Remark 1.5. This condition states some relation on the slopes of P and Q . We thus call it the **strong sub-slope condition**. The assumption (1.14) implies that

$$\frac{P(\rho) - P(\rho^*)}{\rho - \rho^*} \geq \left(\frac{Q(\rho) - Q(\rho^*)}{\rho - \rho^*} \right)^2,$$

which implies the sub-characteristic condition (1.10) when $\frac{m}{\rho} = 0$. We remark that sub-characteristic condition involves the velocity $\frac{m}{\rho}$, which does not appear in the limiting equation (1.6). However, at the equilibrium where $m = Q$, (A2) is very close to the sub-characteristic condition. Condition (A2) gives a global picture between P and Q without involving the velocity, which will give us some advantage in the proof of large time asymptotic behavior of weak solutions in section 4.

The plan of the rest of this paper is organized as follows. In section 2 we give some elementary notations and basic facts. The main results will also be stated there. In section 3, the global existence of L^∞ weak entropy solutions will be proved. Finally, we will investigate the large time behavior of any L^∞ weak solutions in section 4.

2 Preliminaries and main results

In this section, we will present some preliminaries for the foundation of our studies in next sections.

The homogeneous system corresponding to system of (1.2) reads

$$\begin{cases} \rho_t + m_x = 0, \\ m_t + \left(\frac{m^2}{\rho} + P(\rho) \right)_x = 0. \end{cases} \quad (2.1)$$

For a smooth solution, (2.1) can be rewritten as

$$v_t + \nabla f(v)v_x = 0, \quad (2.2)$$

where $v = (\rho, m)^T$, $f(v) = (m, m^2/\rho + \rho^\gamma/\gamma)^T$, and

$$\nabla f = \begin{pmatrix} 0 & 1 \\ -\frac{m^2}{\rho^2} + \rho^{\gamma-1} & \frac{2m}{\rho} \end{pmatrix}. \quad (2.3)$$

The eigenvalues of (2.3) are

$$\lambda_1 = \frac{m}{\rho} - \rho^\theta, \quad \lambda_2 = \frac{m}{\rho} + \rho^\theta, \quad (2.4)$$

and the Riemann invariants are

$$w = \frac{m}{\rho} + \frac{\rho^\theta}{\theta}, \quad z = \frac{m}{\rho} - \frac{\rho^\theta}{\theta},$$

for $\theta = (\gamma - 1)/2$. We note that (w, z) satisfies

$$\begin{cases} z_t + \lambda_1 z_x = \frac{Q(\rho)}{\rho} - \frac{m}{\rho} \\ w_t + \lambda_2 w_x = \frac{Q(\rho)}{\rho} - \frac{m}{\rho} \end{cases} \quad (2.5)$$

For the Riemann problem

$$\begin{cases} (2.2), & t > 0, \quad x \in \mathbb{R}, \\ (\rho, m)|_{t=0} = \begin{cases} (\rho_l, m_l), & x < 0, \\ (\rho_r, m_r), & x > 0, \end{cases} \end{cases} \quad (2.6)$$

where ρ_l, ρ_r, m_l and m_r are constants satisfying $0 \leq \rho_l, \rho_r, |m_l/\rho_l|, |m_r/\rho_r| < \infty$, there are two distinct types of rarefaction waves and shock waves, called elementary waves, which are labelled 1-rarefaction or 2-rarefaction waves and 1-shock or 2-shock waves, respectively.

Lemma 2.1. *There exists a global weak entropy solution of (2.6) which is piecewise smooth function satisfying*

$$\begin{aligned} w(x, t) &= w\left(\frac{x}{t}\right) \leq \max\{w(\rho_l, m_l), w(\rho_r, m_r)\}, \\ z(x, t) &= z\left(\frac{x}{t}\right) \geq \min\{z(\rho_l, m_l), z(\rho_r, m_r)\}, \\ w(x, t) - z(x, t) &\geq 0. \end{aligned}$$

It follows that the region $\Lambda = \{(\rho, m) : w \leq w_0, z \geq z_0, w - z \geq 0\}$ is an invariant region for the Riemann problem (2.6).

Lemma 2.2. *If $\{(\rho, m) : a \leq x \leq b\} \subset \Lambda$, then*

$$\left(\frac{1}{b-a} \int_a^b \rho \, dx, \quad \frac{1}{b-a} \int_a^b m \, dx \right) \in \Lambda. \quad (2.7)$$

Lemma 2.3. *For the mixed problem*

$$\begin{cases} (2.2), & t > 0, \quad x > 0, \\ (\rho, m)|_{t=0} = (\rho_0, m_0), & x > 0, \\ m|_{x=0} = 0, & t \geq 0, \end{cases} \quad (2.8)$$

where (ρ_0, m_0) are constants, there exists a weak entropy solution in the region $\{(x, t) : x \geq 0, t \geq 0\}$ satisfying the following estimates

$$\begin{aligned} w(x, t) &\leq \max\{w(\rho_0, m_0), -z(\rho_0, m_0)\}, \\ z(x, t) &\geq z(\rho_0, m_0) \quad \text{and} \quad w(x, t) - z(x, t) \geq 0. \end{aligned}$$

Similarly, we can solve the following mixed problem in the region $\{(x, t) : x \leq 1, t \geq 0\}$

$$\begin{cases} (2.2), & t > 0, \quad x < 1, \\ (\rho, m)|_{t=0} = (\rho_0, m_0), & x < 1, \\ m|_{x=1} = 0, & t \geq 0. \end{cases} \quad (2.9)$$

The weak solution for (2.9) satisfies the following estimates

$$\begin{aligned} z(x, t) &\geq \min\{z(\rho_0, m_0), -w(\rho_0, m_0)\}, \\ w(x, t) &\leq w(\rho_0, m_0) \quad \text{and} \quad w(x, t) - z(x, t) \geq 0. \end{aligned}$$

Lemma 2.4. *Suppose that $(\rho(x, t), m(x, t))$ is a solution to (2.6) or (2.8) or (2.9). Then the jump strength of $m(x, t)$ across an elementary wave can be dominated by that of $\rho(x, t)$ across the same elementary wave, i.e.,*

$$\begin{aligned} \text{across a shock wave:} \quad & |m_r - m_l| \leq C|\rho_r - \rho_l|, \\ \text{across a rarefaction wave:} \quad & |m - m_l| \leq C|\rho - \rho_l| \leq C|\rho_r - \rho_l|, \end{aligned}$$

where C depends only on the bounds of ρ and $|m|$.

Lemma 2.5. *For any $\varepsilon > 0$, there exist constants $h > 0$ and $k > 0$ such that the solution of (2.6) in the region $\{(x, t) : |x| < h, 0 \leq t < k\}$ satisfies*

$$\int_{-h}^h |\rho(x, t) - \rho(x, 0)| \, dx \leq Ch\varepsilon, \quad 0 \leq t \leq k, \quad (2.10)$$

where C depends only on the bounds of ρ and $|m|$, and the mesh lengths h and k satisfy $\max_{i=1,2} \sup |\lambda_i(\rho, m)| < \frac{h}{2k}$.

Theorem 2.6. Assume that the initial data (ρ_0, m_0) satisfy the following conditions

$$0 \leq \rho_0(x) \leq M_1, \rho_0 \not\equiv 0, |m_0(x)| \leq M_2 \rho_0(x),$$

for some positive constants $M_i (i = 1, 2)$. And assume $Q(\rho)$ satisfies the assumption (A1). Then, for $\gamma > 1$, there exists a positive constant M , the initial boundary value problem (1.2)-(1.3) has a global weak solution $(\rho(x, t), m(x, t))$ satisfying the following estimates and entropy condition

$$0 \leq \rho(x, t) \leq M, |m(x, t)| \leq M\rho \quad \text{a.e.},$$

$$\int_0^T \int_0^1 (\eta(\rho, m)\psi_t + q(\rho, m)\psi_x) dxdt + \int_0^T \int_0^1 \eta_m(\rho, m)(Q(\rho) - m)\psi dxdt \geq 0,$$

for all weak and convex entropy pairs (η, q) for (1.2)-(1.3) and for all nonnegative smooth functions $\psi \in C_0^1(I_T)$.

Theorem 2.7. Let (ρ, m) be any L^∞ entropy weak solution of the initial boundary problem (1.2)-(1.3), satisfying $\int_0^1 \rho_0(x) dx = \rho^*$, $Q(\rho^*) = 0$ and

$$0 \leq \rho(x, t) \leq M < \infty, |m(x, t)| \leq M_1 \rho(x, t),$$

where M and M_1 are positive constants. And assume $Q(\rho)$ satisfies the assumption A(2). Then, there exist constants $C > 0$ and $\delta > 0$ depending on γ, ρ^*, M and initial data such that

$$\|(\rho - \rho^*, m)(\cdot, t)\|_{L^2([0,1])}^2 \leq Ce^{-\delta t}.$$

3 Global existence of weak entropy solutions

We begin with the construction of approximate solution by modified Godunov Scheme in the spirit of operator splitting. Let us take the space mesh length $h = 1/N$, where N is a positive integer. The time mesh length $k = k(h)$ will be determined later so that the Courant-Friedrich-Lewy condition

$$\max_{i=1,2}(\sup |\lambda_i(v)|) < \frac{h}{2k} \quad (3.1)$$

holds for a given $T > 0$. We partition the interval $[0, 1]$ into cells, with the j -th cell centered at $x_j = jh$, $j = 1, \dots, N-1$, and denote t_i by ik . Set $x_0 = 0$ and $x_N = 1$. Now we consider the solution $\underline{v}_h = (\underline{\rho}_h, \underline{m}_h)^T$ of the Riemann problems (2.6) in the region $R_j^1 \equiv \{(x, t) : x_{j-\frac{1}{2}} \leq x < x_{j+\frac{1}{2}}, 0 \leq t < k\}$:

$$\begin{cases} \frac{\partial}{\partial t} \underline{v}_h + \frac{\partial}{\partial x} f(\underline{v}_h) = 0, \\ \underline{v}_h|_{t=0} = \begin{cases} (\rho_j^0, m_j^0), & x < x_j, \\ (\rho_{j+1}^0, m_{j+1}^0), & x > x_j, \end{cases} \quad j = 1, \dots, N-1, \end{cases} \quad (3.2)$$

where

$$\rho_j^0 = \frac{1}{h} \int_{x_{j-1}}^{x_j} \rho_0(x) dx, \quad m_j^0 = \frac{1}{h} \int_{x_{j-1}}^{x_j} m_0(x) dx, \quad \text{for } j = 1, \dots, N. \quad (3.3)$$

At the same time we also solve the mixed problem (2.8) and (2.9) with (ρ_1^0, m_1^0) and (ρ_N^0, m_N^0) , in regions $\{(x, t) : 0 \leq x < x_{1/2}, 0 \leq t < k\}$ and $\{(x, t) : x_{N-1/2} \leq x < 1, 0 \leq t < k\}$, respectively. Then we set, for $0 \leq x \leq 1, 0 \leq t < k$,

$$\begin{cases} v_h(x, t) = (\rho_h(x, t), m_h(x, t))^T, \\ \rho_h(x, t) = \underline{\rho}_h(x, t), \\ m_h(x, t) = \underline{m}_h(x, t) - \underline{m}_h(x, t)t + Q(\underline{\rho}_h(x, t))t \end{cases} \quad (3.4)$$

and

$$v_j^1 = \frac{1}{h} \int_{x_{j-1}}^{x_j} v_h(x, t_1 - 0) dx, \quad j = 1, \dots, N. \quad (3.5)$$

Next we will define approximate solutions v_h for $t_i \leq t < t_{i+1}$ through using approximate solutions defined in $0 \leq t < t_i$. Suppose that we have defined approximate solutions $v_h(x, t)$ for $0 \leq t < t_i$. we then define approximate solutions for $0 \leq x \leq 1, t_i \leq t < t_{i+1}$ as follows

$$\begin{cases} v_h(x, t) = (\rho_h(x, t), m_h(x, t))^T, \\ \rho_h(x, t) = \underline{\rho}_h(x, t), \\ m_h(x, t) = \underline{m}_h(x, t) - \underline{m}_h(x, t)(t - t_i) + Q(\underline{\rho}_h(x, t))(t - t_i) \end{cases} \quad (3.6)$$

where $\underline{v}_h(x, t) = (\underline{\rho}_h(x, t), \underline{m}_h(x, t))$ are piecewise-smooth functions defined as solutions of Riemann problems in the region $R_j^{i+1} \equiv \{(x, t) : x_{j-\frac{1}{2}} \leq x < x_{j+\frac{1}{2}}, t_i \leq t < t_{i+1}\}$:

$$\begin{cases} (2.2), \\ \underline{v}_h(x, t)|_{t=t_i} = \begin{cases} v_j^i, & x < x_j, \\ v_{j+1}^i, & x > x_j, \end{cases} \quad j = 1, \dots, N-1, \end{cases} \quad (3.7)$$

and as solutions of mixed problems in the two side regions $R_0^{i+1} = \{(x, t) : 0 \leq x < x_{1/2}, t_i \leq t < t_{i+1}\}$ and $R_N^{i+1} = \{(x, t) : x_{N-1/2} \leq x < 1, t_i \leq t < t_{i+1}\}$:

$$\begin{cases} (2.2), & t > t_i, \quad x > 0, \\ \underline{v}_h|_{t=t_i} = v_1^i, & x > 0, \\ \underline{m}_h|_{x=0} = 0, \end{cases} \quad (3.8)$$

and

$$\begin{cases} (2.2), & t > t_i, \quad x < 1, \\ \underline{v}_h|_{t=t_i} = v_N^i, & x < 1, \\ \underline{m}_h|_{x=1} = 0. \end{cases} \quad (3.9)$$

And we set

$$v_j^{i+1} = \frac{1}{h} \int_{x_{j-1}}^{x_j} v_h(x, t_{i+1} - 0) dx, \quad 1 \leq j \leq N. \quad (3.10)$$

Therefore the approximate solutions $v_h = (\rho_h, m_h)$ are well defined in the region $\{0 \leq x \leq 1, 0 \leq t \leq T\}$ for any $T > 0$ since $\underline{\rho}_h \geq 0$.

For $t_i \leq t < t_{i+1}$, due to (2.5), we can obtain the expression of $(w_h(x, t), z_h(x, t))$ as follows:

$$\begin{aligned} w_h(x, t) &= \underline{w}_h(x, t) - \frac{\underline{w}_h + \underline{z}_h}{2}(t - t_i) + \frac{Q(\underline{\rho}_h)}{\underline{\rho}_h}(t - t_i), \\ z_h(x, t) &= \underline{z}_h(x, t) - \frac{\underline{w}_h + \underline{z}_h}{2}(t - t_i) + \frac{Q(\underline{\rho}_h)}{\underline{\rho}_h}(t - t_i), \end{aligned} \quad (3.11)$$

where \underline{w}_h and \underline{z}_h are Riemann invariants corresponding to the Riemann solution \underline{v}_h .

We prove the following uniform bound for the approximate solutions.

Lemma 3.1. *Suppose that the initial data (ρ_0, m_0) satisfy the following conditions:*

$$0 \leq \rho_0(x) \leq M_1, \rho_0 \not\equiv 0, |m_0(x)| \leq M_2 \rho_0(x). \quad (3.12)$$

And assume $Q(\rho)$ satisfies the assumption (A1). Then the approximate solutions (ρ_h, m_h) derived by the Godunov scheme are uniformly bounded in the region $\bar{I}_T \equiv \{(x, t) : 0 \leq x \leq 1, 0 \leq t \leq T\}$ for any $T > 0$; that is, there is a constant C independent of t such that

$$0 \leq \rho_h(x, t) \leq C, \quad |m_h(x, t)| \leq C \rho_h(x, t). \quad (3.13)$$

Proof. Assume that $0 < k < 1$. Firstly, for $0 \leq t < t_1$, the Riemann invariant properties imply that

$$\underline{w}_h(x, t) \leq \alpha_0, \quad \underline{z}_h(x, t) \geq -\alpha_0, \quad \text{and} \quad \underline{w}_h(x, t) - \underline{z}_h(x, t) \geq 0,$$

where

$$\alpha_0 = \max\left\{\sup_x w_0(x), -\inf_x z_0(x)\right\}. \quad (3.14)$$

Then it holds that

$$0 \leq \underline{\rho}_h(x, t) \leq (\theta \alpha_0)^{1/\theta}, \quad |\underline{m}_h(x, t)| \leq \alpha_0 \underline{\rho}_h(x, t).$$

From (A1), we have

$$\begin{aligned} w_h(x, t) &= \underline{w}_h(x, t)(1-t) + \left[\frac{\underline{w}_h - \underline{z}_h}{2} + \frac{Q(\underline{\rho}_h)}{\underline{\rho}_h} \right] t \\ &\leq \underline{w}_h(x, t)(1-t) + \left(\frac{\underline{\rho}_h^\theta}{\theta} + \frac{Q(\underline{\rho}_h)}{\underline{\rho}_h} \right) t \\ &\leq \underline{w}_h(x, t)(1-t) + \alpha_0 t \\ &\leq \alpha_0, \end{aligned} \quad (3.15)$$

and

$$\begin{aligned} z_h(x, t) &= \underline{z}_h(x, t)(1-t) + \left[\frac{Q(\underline{\rho}_h)}{\underline{\rho}_h} - \frac{\underline{w}_h + \underline{z}_h}{2} \right] t \\ &= \underline{z}_h(x, t)(1-t) + \left(\frac{Q(\underline{\rho}_h)}{\underline{\rho}_h} - \frac{\underline{\rho}_h^\theta}{\theta} \right) t \\ &\geq \underline{z}_h(x, t)(1-t) - \alpha_0 t \\ &\geq -\alpha_0. \end{aligned} \quad (3.16)$$

Inductively, we can prove that for $t_i \leq t < t_{i+1}$,

$$w_h(x, t) \leq \alpha_0, \quad z_h(x, t) \geq -\alpha_0, \quad w_h(x, t) - z_h(x, t) = \underline{w}_h(x, t) - \underline{z}_h(x, t) \geq 0.$$

Then there is a constant $C > 0$ independent of h, k and t such that

$$0 \leq \rho_h(x, t) \leq C, \quad |m_h(x, t)| \leq C \rho_h(x, t).$$

This completes the proof of Lemma 3.1. □

Finally, we can choose the time mesh length $k = k(h)$. Let

$$\lambda = \max_{i=1,2} \left\{ \sup_{0 \leq \rho \leq C, |m| \leq C\rho} |\lambda_i(\rho, m)| \right\},$$

then we take

$$k = \frac{T}{n}, \quad \text{where } n = \max \left\{ \left[\frac{4\lambda T}{h} \right] + 1, \left[\frac{T}{2} \right] + 1 \right\}.$$

For this k , both the CFL condition and $0 < k < 1$ hold.

Set $g(v) = (0, Q(\rho) - m)^T$. Then (1.2)-(1.3) can be rewritten as

$$\begin{cases} v_t + f(v)_x = g(v), \\ v(x, 0) = v_0(x), \quad x \in (0, 1), \\ m(0, t) = m(1, t) = 0. \end{cases} \quad (3.17)$$

In the following, we will show that the approximate solutions constructed above admit a convergent subsequence whose limit is a weak entropy solution of problem (1.2)-(1.3). The convergence is achieved by the compensated compactness, the boundary conditions are verified in the sense of trace.

In view of the uniform L^∞ estimates given in Lemma 3.1, and the specific structure of system (1.2), then it is standard to apply the compensated compactness framework to the approximate solution $\{v_h\}$, to conclude that there exists a convergent subsequence, still labeled $\{v_h\}$, such that

$$(\rho_h(x, t), m_h(x, t)) \rightarrow (\rho(x, t), m(x, t)) \quad \text{a.e.} \quad (3.18)$$

Clearly, there is a positive constant C such that

$$0 \leq \rho(x, t) \leq C, \quad |m(x, t)| \leq C\rho(x, t) \quad \text{a.e.} \quad (3.19)$$

For any $\phi \in C^\infty(\bar{I}_T)$ satisfying $\phi(x, T) = 0$, $\phi(0, t) = \phi(1, t) = 0$, we consider the following integral identity

$$\int_0^T \int_0^1 (\rho_h \phi_t + m_h \phi_x) dx dt + \int_{t=0}^T \rho_h \phi dx = A(\phi) + R(\phi), \quad (3.20)$$

where

$$\begin{aligned} A(\phi) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} (\rho_h^i - \rho_j^i) \phi^i dx, \\ R(\phi) &= \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} (m_h - \underline{m}_h) \phi_x dx dt, \end{aligned}$$

with $\rho_h^i = \rho(x, t_i - 0)$. Similar to [31], with the help of Hölder inequality and the uniform bound of v_h , we have

$$A(\phi) \leq Ch^{1/2} \|\phi\|_{C^1} \rightarrow 0, \quad \text{as } h \rightarrow 0. \quad (3.21)$$

$$\begin{aligned} R(\phi) &\leq \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} |(Q(\underline{\rho}_h) - \underline{m}_h)(t - t_i)| |\phi_x| dx dt \\ &\leq Ch \|\phi\|_{C^1} \rightarrow 0, \quad \text{as } h \rightarrow 0. \end{aligned} \quad (3.22)$$

Then from (3.21)-(3.22), it gives

$$\lim_{h \rightarrow 0} \int_0^T \int_0^1 (\rho_h \phi_t + m_h \phi_x) dx dt + \lim_{h \rightarrow 0} \int_{t=0} \rho_h \phi dx = 0. \quad (3.23)$$

By virtue of the dominated convergence theorem to (3.23), we get

$$\int_0^T \int_0^1 (\rho \phi_t + m \phi_x) dx dt + \int_{t=0} \rho_0(x) \phi dx = 0. \quad (3.24)$$

For every function $\phi \in C^1(\bar{I}_T)$ satisfying $\phi(x, T) = 0$ for $0 \leq x \leq 1$ and $\phi(0, t) = \phi(1, t) = 0$ for $t \geq 0$, we consider the integral identity

$$\int_0^T \int_0^1 (m_h \phi_t + f_1(v_h) \phi_x + V(v_h) \phi) dx dt + \int_{t=0} m_h \phi dx = B(\phi) + S(\phi), \quad (3.25)$$

with $f_1(v) = m^2/\rho + \rho^\gamma/\gamma$, $V(v) = Q(\rho) - m$, and

$$\begin{aligned} B(\phi) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} (\underline{m}_h^i - m_j^i) \phi^i dx + \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} V(\underline{v}_h) \phi dx dt, \\ S(\phi) &= \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} [(m_h - \underline{m}_h) \phi_t \\ &\quad + (f_1(v_h) - f_1(\underline{v}_h)) \phi_x + (V(v_h) - V(\underline{v}_h)) \phi] dx dt. \end{aligned}$$

Using the uniform bound of v_h and $|m_h - \underline{m}_h| \leq k(|\underline{m}_h| + C'|\underline{\rho}_h|)$, we have

$$S(\phi) \leq Ch \|\phi\|_{C^1} \rightarrow 0, \quad \text{as } h \rightarrow 0. \quad (3.26)$$

Due to $m_h^i = \underline{m}_h^i + \int_{t_i}^{t_{i+1}} V(\underline{v}_h^i) dt$, then $B(\phi)$ can be bounded by

$$\begin{aligned} B(\phi) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} (\underline{m}_h^i - m_j^i) (\phi^i - \phi_j^i) dx \\ &\quad + \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} V(\underline{v}_h) (\phi - \phi_j^i) dx dt \\ &\quad + \sum_{i,j} \int_{t_i}^{t_{i+1}} \int_{x_{j-1}}^{x_j} [V(\underline{v}_h) - V(\underline{v}_h^i)] \phi_j^i dx dt \\ &\leq Ch^{1/2} \|\phi\|_{C^1} + Ch \|\phi\|_{C^1} + C\varepsilon \|\phi\|_{\infty}, \end{aligned}$$

where ε is an arbitrarily small constant. Then, it implies

$$B(\phi) \rightarrow 0, \quad \text{as } h \rightarrow 0, \quad \varepsilon \rightarrow 0. \quad (3.27)$$

Then due to (3.26)-(3.27) and the dominated convergence theorem, we obtain

$$\begin{aligned} \int_0^T \int_0^1 \left(m \phi_t + \left(\frac{m^2}{\rho} + P(\rho) \right) \phi_x \right) dx dt \\ + \int_0^T \int_0^1 (Q(\rho) - m) \phi dx dt + \int_{t=0} m_0 \phi dx = 0. \end{aligned} \quad (3.28)$$

For every weak and convex entropy pair (η, q) and every nonnegative smooth function ψ which has a compact support in I_T , we study the integral identity

$$\int_0^T \int_0^1 (\eta(v_h)\psi_t + q(v_h)\psi_x) dxdt = A(\psi_h) + R(\psi_h) + \Sigma(\psi_h) + S(\psi_h), \quad (3.29)$$

where

$$\begin{aligned} A(\psi_h) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} [\eta(v_h^i) - \eta(v_h^j)] \psi(x, t_i) dx, \\ R(\psi_h) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} [\eta(v_h^i) - \eta(v_h^j)] \psi(x, t_i) dx, \\ \Sigma(\psi_h) &= \int_0^T \Sigma\{\sigma[\eta] - [q]\} \psi(x(t), t) dt, \\ S(\psi_h) &= \int_0^T \int_0^1 [\eta(v_h) - \eta(v_h)] \psi_t + [q(v_h) - q(v_h)] \psi_x dxdt. \end{aligned}$$

Since (η, q) is a convex entropy pair and $\psi \geq 0$, similar to [31], we have

$$\begin{aligned} A(\psi_h) &\geq \sum_{i,j} \int_{x_{j-1}}^{x_j} [\eta(v_h^i) - \eta(v_h^j)] (\psi^i - \psi^j) dx \\ &\geq -Ch^{\alpha-1/2} \|\psi\|_{C_0^\alpha}, \quad 1/2 \leq \alpha \leq 1, \end{aligned} \quad (3.30)$$

$$\Sigma(\psi_h) \geq 0, \quad (3.31)$$

$$S(\psi_h) \geq -Ch \|\psi\|_{H_0^1}, \quad (3.32)$$

$$\begin{aligned} R(\psi_h) &= \sum_{i,j} \int_{x_{j-1}}^{x_j} \int_0^1 \nabla \eta(v_h^i + \theta(v_h^i - v_h^j)) (v_h^i - v_h^j) d\theta \psi^i dx \\ &\geq -\sum_i \int_0^1 \left(\int_0^1 \eta_m(v_h^i + \theta(v_h^i - v_h^i)) d\theta \cdot [Q(\underline{\rho}_h^i) - m_h^i](t - t_i) \psi^i \right) dx \\ &\quad -Ch. \end{aligned} \quad (3.33)$$

With the help of these above inequalities and the fact that $v_h \rightarrow v$ a.e., letting $h \rightarrow 0$, then we have the following entropy condition

$$\int_0^T \int_0^1 (\eta(v)\psi_t + q(v)\psi_x) dxdt + \int_0^T \int_0^1 \eta_m(v)V(v)\psi_t dxdt \geq 0. \quad (3.34)$$

Now we turn to the boundary conditions of weak solutions. The exact meaning of traces for weak solutions is given below. Let $v(x, t) = (\rho(x, t), m(x, t))$ be a weak solution of (1.2) obtain in (3.18). We introduce the generalized function $\mathcal{A} : C_0^1(\mathbb{R}^2) \rightarrow \mathbb{R}^2$ as follows: for $\phi \in C_0^1(\mathbb{R}^2)$,

$$\mathcal{A}(\phi) = -\int_0^T \int_0^1 [v\phi_t + f(v)\phi_x + g(v)\phi] dxdt, \quad (3.35)$$

with $f(v) = (m, m^2/\rho + \rho^\gamma/\gamma)^T$, $g(v) = (0, Q(\rho - m))^T$.

We take smooth $\zeta_0(t), \zeta_T(t), \xi_0(x), \xi_1(x)$ with

$$\begin{aligned} \zeta_0(0) &= 1, & \zeta_0(T) &= 1; & \zeta_T(0) &= 0, & \zeta_T(T) &= 1; \\ \xi_0(0) &= 1, & \xi_0(T) &= 1; & \xi_T(0) &= 0, & \xi_T(T) &= 1. \end{aligned} \quad (3.36)$$

For any $\chi(x)$, we define the generalized functions:

$$\begin{aligned} v^*(\cdot, 0)(\chi) &= \mathcal{A}(\chi \cdot \zeta_0) - \chi(0)\mathcal{A}(\xi_0 \cdot \zeta_0) - \chi(1)\mathcal{A}(\xi_0 \cdot \zeta_0); \\ v^*(\cdot, T)(\chi) &= -\mathcal{A}(\chi \cdot \zeta_T) + \chi(0)\mathcal{A}(\xi_0 \cdot \zeta_T) + \chi(1)\mathcal{A}(\xi_0 \cdot \zeta_T); \\ f^*(v)(0, \cdot)(\chi) &= \mathcal{A}(\xi_0 \cdot \chi); \\ f^*(v)(1, \cdot)(\chi) &= -\mathcal{A}(\xi_1 \cdot \chi), \end{aligned}$$

where $\chi \cdot \zeta_0(x, t) = \chi(x)\zeta_0(t)$ and so mean the tensor product.

Then we define the trace of v along the segments $(0, 1) \times \{0\}$ and $(0, 1) \times \{T\}$, and the trace of $f(v)$ along the segments $\{0\} \times (0, T)$ and $\{1\} \times (0, T)$ respectively as $v^*(\cdot, 0), v^*(\cdot, T), f^*(v)(0, \cdot)$ and $f^*(v)(1, \cdot)$. Similarly, for any $t \in (0, T)$, we can also define $v^*(\cdot, t)$ as the trace of v along the segment $(0, 1) \times \{t\}$. For any $x \in (0, 1)$, we can also define $f^*(v)(x, \cdot)$ as the trace of $f(v)$ along the segment $\{x\} \times (0, 1)$.

Similar to [7], we have the following lemma.

Lemma 3.2. *Let v satisfy (1.2) in distributional sense, then*

$$\begin{aligned} v^*(\cdot, 0)|_{(0,1)}, \quad v^*(\cdot, T)|_{(0,1)} &\in L_{loc}^\infty(0, 1) \\ f^*(v)(0, \cdot)|_{(0,T)}, \quad f^*(v)(1, \cdot)|_{(0,T)} &\in L_{loc}^\infty(0, T), \end{aligned}$$

and $\forall \phi \in C_0^1(\mathbb{R}^2)$,

$$\begin{aligned} &\int_0^T \int_0^1 [v\phi_t + f(v)\phi_x + g(v)\phi] dx dt \\ &= \int_0^1 v^*(x, T)\phi(x, T) dx - \int_0^1 v^*(x, 0)\phi(x, 0) dx \\ &\quad + \int_0^T f^*(v)(1, t)\phi(1, t) dt - \int_0^T f^*(v)(0, t)\phi(0, t) dt. \end{aligned} \quad (3.37)$$

Theorem 3.3. *Let $v_h(x, t) = (\rho_h(x, t), m_h(x, t))$ be the approximate solutions of (1.2)-(1.3) constructed in this section and $v(x, t) = (\rho(x, t), m(x, t))$ be the limit function obtained in (3.18). Then $v = (\rho, m)$ satisfy the initial-boundary conditions*

$$m^*(0, t) = m^*(1, t) = 0, \quad t \in (0, T) \quad (3.38)$$

$$v^*(x, 0) = v_0(x), \quad x \in (0, 1). \quad (3.39)$$

Proof. From (1.7)- (1.8), it gives, for any $\phi \in C_0^1(\mathbb{R}^2)$, that

$$\lim_{h \rightarrow 0} \left[\int_0^T \int_0^1 [v_h\phi_t + f(v_h)\phi_x + g(v_h)\phi] dx dt + \int_{t=0} v_h\phi dx - \int_{t=T} v_h\phi dx \right] = 0,$$

which implies

$$\int_0^T \int_0^1 [v\phi_t + f(v)\phi_x + g(v)\phi] dx dt + \lim_{h \rightarrow 0} \left[\int_{t=0} v_h\phi dx - \int_{t=T} v_h\phi dx \right] = 0, \quad (3.40)$$

Inserting (3.37) into (3.40), we get

$$\begin{aligned} & \lim_{h \rightarrow 0} \left[\int_{t=T}^1 v_h \phi \, dx - \int_{t=0}^1 v_h \phi \, dx \right] \\ &= \int_0^1 v^*(x, T) \phi(x, T) \, dx - \int_0^1 v^*(x, 0) \phi(x, 0) \, dx \\ & \quad + \int_0^T f^*(v)(1, t) \phi(1, t) \, dt - \int_0^T f^*(v)(0, t) \phi(0, t) \, dt. \end{aligned} \quad (3.41)$$

So the first component of above equality reads

$$\begin{aligned} & \int_0^1 \rho^*(x, T) \phi(x, T) \, dx - \int_0^1 \rho^*(x, 0) \phi(x, 0) \, dx + \int_0^T m^*(v)(1, t) \phi(1, t) \, dt \\ & \quad - \int_0^T m^*(v)(0, t) \phi(0, t) \, dt + \int_{t=0}^1 \rho_0(x) \phi \, dx - \int_{t=T}^1 \rho \phi \, dx = 0. \end{aligned} \quad (3.42)$$

Taking $\phi(x, t) = \zeta(x)\chi(t) \in C_0^1(\mathbb{R}^2)$ with $\zeta, \chi \in C_0^1(\mathbb{R})$, and $\chi(0) = 1, \chi(T) = 0, \zeta(0) = \zeta(1) = 0$ in (3.41), we have

$$\int_0^1 \rho^*(x, 0) \zeta(x) \, dx = \int_0^1 \rho_0(x) \zeta(x) \, dx,$$

which implies $\rho^*(x, 0) = \rho_0(x)$ on $(0, 1)$.

Similarly, using the second component of (3.41), it is easy to show that $m^*(x, 0) = m_0(x)$ on $(0, 1)$. Taking $\phi(x, t) = \zeta(x)\chi(t) \in C_0^1(\mathbb{R}^2)$ with $\zeta, \chi \in C_0^1(\mathbb{R})$, and $\chi(0) = \chi(T) = 0, \zeta(0) = 1, \zeta(1) = 0$ in (3.41), one can get

$$\int_0^1 m^*(0, t) \chi(t) \, dt = 0.$$

Therefore $m^*(0, t) = 0$ on $(0, T)$. It is similar to obtain that $m^*(1, t) = 0$ on $(0, T)$. This completes the proof Theorem 3.3 \square

4 Large time behavior of weak solution

In this section, we will prove the asymptotic behavior of the weak solution, namely, Theorem 2.7. For this purpose, we assume that (ρ, m) is an entropy weak solution in L^∞ such that

$$0 \leq \rho \leq M, \quad |m(x, t)| \leq C_0 \rho(x, t)$$

for some positive constants M and C_0 .

Due to conservation law of total mass, we have

$$\int_0^1 \rho(x, t) \, dx = \int_0^1 \rho_0(x) \, dx = \rho^* > 0. \quad (4.1)$$

Without the loss of generality, we assume $\rho^* < M$, otherwise one has $\rho \equiv M$ and $m = Q(M) = 0$, the trivial constant solution, or inconsistency in the case $Q(M) \neq 0$. For the same reason, we require in Theorem 2.7 the condition

$$Q(\rho^*) = 0,$$

to ensure $(\rho^*, 0)$ a equilibrium solution to (2)-(3).

In order to control the singularity near vacuum, the following lemma proved in [30] plays an important role.

Lemma 4.1. *Let $0 \leq \rho \leq M < \infty$. There are positive constants C_1, C_2 and C_3 such that*

$$C_1 \leq f_2(\rho, \rho^*), f_4(\rho, \rho^*) \leq C_2, f_1(\rho, \rho^*) \leq C_3 f_3(\rho, \rho^*). \quad (4.2)$$

This lemma is a direct consequence of the mean value theorem for $P(\rho)$.

Remark 4.2. We remark that, due to the convexity of $P(\rho) = \frac{1}{\gamma} \rho^\gamma$, it is clear that

$$f_4(\rho, \rho^*) \geq f_4(0, \rho^*) = \frac{1}{\gamma} (\rho^*)^{\gamma-1} \equiv C^*.$$

Set

$$y = - \int_0^x (\rho - \rho^*) dr, \quad (4.3)$$

then

$$y_x = -(\rho - \rho^*), \quad y_t = m. \quad (4.4)$$

Due to the conservation of mass we have

$$y(0) = y(1) = 0. \quad (4.5)$$

The momentum equation becomes

$$y_{tt} + y_t + \left(\frac{m^2}{\rho} \right)_x + (P(\rho) - P(\rho^*))_x = Q(\rho). \quad (4.6)$$

Multiplying y with (4.6) and integrating the resulting equation over $[0, 1]$, we have

$$\begin{aligned} & \frac{d}{dt} \int_0^1 (y_t y + \frac{1}{2} y^2) dx + \int_0^1 (f_3(\rho, \rho^*) - y_t^2 - \frac{m^2}{\rho} y_x) dx \\ &= \int_0^1 (Q(\rho) - Q(\rho^*)) y dx. \end{aligned} \quad (4.7)$$

Let

$$\eta_e = \frac{m^2}{2\rho} + \frac{P(\rho)}{\gamma-1}, \quad q_e = \frac{m^3}{2\rho^2} + \frac{\rho^{\gamma-1} m}{\gamma-1}$$

be the mechanical energy and related flux, respectively. We define

$$\eta_* = \eta_e - \frac{1}{\gamma-1} P'(\rho^*)(\rho - \rho^*) - \frac{1}{\gamma-1} P(\rho^*). \quad (4.8)$$

Thus, by the definition of weak entropy solution, the following entropy inequality holds in the sense of distribution:

$$\eta_{*t} + \frac{1}{\gamma-1} [P'(\rho^*)(\rho - \rho^*)]_t + q_{ex} + \frac{m^2}{\rho} - \frac{m}{\rho} Q(\rho) \leq 0. \quad (4.9)$$

By the conservation of mass and theory of divergence-measure fields [1], we have

$$\frac{d}{dt} \int_0^1 \eta_* dx + \int_0^1 \left(\frac{y_t^2}{\rho} - \frac{m}{\rho} Q(\rho) \right) dx \leq 0. \quad (4.10)$$

Now we add (4.7) to (4.10) $\times \lambda_0$, for some $\lambda_0 > 2M$ chosen later,

$$\frac{d}{dt} E(t) + D(t) \leq 0, \quad (4.11)$$

where

$$\begin{aligned} E(t) &= \int_0^1 \left(\lambda_0 \eta_* + y y_t + \frac{1}{2} y^2 \right) dx, \\ D(t) &= \int_0^1 \left(\frac{\lambda_0 - \rho^*}{\rho} y_t^2 - \lambda_0 \frac{m}{\rho} Q(\rho) + f_3(\rho, \rho^*) - (Q(\rho) - Q(\rho^*)) y \right) dx. \end{aligned} \quad (4.12)$$

We note, from (4.4), that

$$\begin{aligned} D(t) &= \int_0^1 \left(\frac{\lambda_0 - \rho^*}{\rho} y_t^2 + \lambda_0 \frac{1}{\rho} f_5(\rho, \rho^*) y_t y_x + f_4(\rho, \rho^*) y_x^2 + f_5(\rho, \rho^*) y_x y \right) dx \\ &\equiv \int_0^t I_1 + I_2 dx, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \frac{\lambda_0 - \rho^*}{\rho} y_t^2 + \lambda_0 \frac{1}{\rho} f_5(\rho, \rho^*) y_t y_x + \frac{1}{a_1} f_4(\rho, \rho^*) y_x^2 \\ I_2 &= \left(\frac{1}{a_2} + \frac{1}{a_3} \right) f_4(\rho, \rho^*) y_x^2 + f_5(\rho, \rho^*) y_x y. \end{aligned} \quad (4.13)$$

Due to $\lambda_0 > 2M$, it holds that

$$\frac{\lambda_0 - \rho^*}{\rho} \geq 1. \quad (4.14)$$

Now we claim that there is a positive constant $\delta_0 > 0$, such that

$$\lambda_0 = 2M + \delta_0, \quad (4.15)$$

satisfying

$$2 \sqrt{\frac{\lambda_0 - \rho^*}{\rho}} \sqrt{\frac{f_4(\rho, \rho^*)}{a_1}} > \frac{\lambda_0}{\rho} f_5(\rho, \rho^*), \quad (4.16)$$

or equivalently

$$\frac{(f_5(\rho, \rho^*))^2}{\rho^2} \lambda_0^2 - \frac{4}{a_1 \rho} f_4(\rho, \rho^*) \lambda_0 + \frac{4\rho^*}{a_1 \rho} f_4(\rho, \rho^*) < 0,$$

for all $\rho \in [0, M]$. In fact, one can define the polynomial in λ ,

$$F(\lambda) = \frac{(f_5(\rho, \rho^*))^2}{\rho^2} \lambda^2 - \frac{4}{a_1 \rho} f_4(\rho, \rho^*) \lambda + \frac{4\rho^*}{a_1 \rho} f_4(\rho, \rho^*).$$

It is easy to verify that under the assumption of (1.14),

$$\begin{aligned} F(2M) &< -\frac{4(M-\rho^*)}{a_1\rho} f_4(\rho, \rho^*) \\ &\leq -\frac{4(M-\rho^*)}{a_1M} C^* < 0, \end{aligned}$$

Therefore, by the continuity of $F(\lambda)$, there is a $\delta_0 > 0$ such that

$$F(2M + \delta_0) < -\frac{2(M-\rho^*)}{a_1M} C^*. \quad (4.17)$$

This verifies our claim (4.15). We now fix this λ_0 . With the help of the estimate (4.17), we conclude that there is a $\delta_1 > 0$ and $C_4 > 0$ such that

$$\delta_1 \left(\frac{y_t^2}{\rho} + y_x^2 \right) \leq I_1 \leq C_4 \left(\frac{y_t^2}{\rho} + y_x^2 \right). \quad (4.18)$$

We now take care of I_2 . Using the Poincaré's inequality for y ,

$$\int_0^1 y^2 dx \leq \frac{1}{2} \int_0^1 y_x^2 dx,$$

one has

$$\frac{1}{a_3} \int_0^1 f_4(\rho, \rho^*) y_x^2 dx \geq \frac{C^*}{a_3} \int_0^1 y_x^2 dx \geq \frac{2C^*}{a_3} \int_0^1 y^2 dx. \quad (4.19)$$

Thus we have

$$\int_0^1 I_2 dx \geq \int_0^1 \left\{ \frac{1}{a_2} f_4(\rho - \rho^*) y_x^2 + f_5(\rho, \rho^*) y_x y + \frac{2C^*}{a_3} y^2 \right\} dx.$$

The assumption (1.15) implies

$$0 \leq \int_0^1 I_2 dx \leq C_5 \int_0^1 y_x^2 dx, \quad (4.20)$$

for some positive constant C_5 . This estimate, together with (4.18) leads to

$$\delta_1 \int_0^1 \left(\frac{y_t^2}{\rho} + y_x^2 \right) dx \leq D(t) \leq (C_4 + C_5) \int_0^1 \left(\frac{y_t^2}{\rho} + y_x^2 \right) dx. \quad (4.21)$$

We now turn to $E(t)$. Using the expression of η_* , we have

$$E(t) = \int_0^1 \left(\frac{\lambda_0}{2\rho} y_t^2 + y y_t + \frac{1}{2} y^2 + \frac{\lambda_0}{\gamma-1} f_1(\rho, \rho^*) \right) dx. \quad (4.22)$$

In view of Lemma 4.1, one has

$$C_6 y_x^2 \leq \frac{\lambda_0}{\gamma-1} f_1(\rho, \rho^*) \leq C_7 y_x^2, \quad (4.23)$$

for two positive constants C_6 and C_7 .

By the fact that $\lambda_0 = 2M + \delta_0$ and $0 \leq \rho \leq M < \infty$, we have

$$\begin{aligned} \int_0^1 \left(\frac{\lambda_0}{2\rho} y_t^2 + yy_t + \frac{1}{2} y^2 \right) dx &\geq C_8 \left(\int_0^1 y_t^2 dx + \int_0^1 y^2 dx \right) \\ &\geq C_8 \int_0^1 y_t^2 dx, \end{aligned} \tag{4.24}$$

for some positive constant C_8 . On the other hand, by Poincaré’s inequality and Cauchy-Schwartz inequality, there is a positive constant C_9 such that

$$\begin{aligned} \int_0^1 \left(\frac{\lambda_0}{2\rho} y_t^2 + yy_t + \frac{1}{2} y^2 \right) dx &\leq C \left(\int_0^1 \frac{y_t^2}{\rho} dx + \int_0^1 y^2 dx \right) \\ &\leq C_9 \int_0^1 \left(\frac{y_t^2}{\rho} + y_x^2 \right) dx. \end{aligned} \tag{4.25}$$

Therefore, we have the following estimate for $E(t)$,

$$\alpha \int_0^1 (y_t^2 + y_x^2) dx \leq E(t) \leq \beta D(t), \tag{4.26}$$

for some positive constant α and β . One thus concludes from (4.26) and (4.11) that

$$\frac{d}{dt} E(t) + \frac{1}{\beta} E(t) \leq 0,$$

which leads to

$$E(t) \leq E(0) e^{-\frac{1}{\beta} t}. \tag{4.27}$$

Finally, we have from (4.26) that

$$\int_0^1 y_t^2 + (\rho - \rho^*)^2 dx \leq \frac{1}{\alpha} E(0) e^{-\frac{1}{\beta} t}. \tag{4.28}$$

This completes the proof of Theorem 2.7.

Acknowledgments

Part of this research was carried out when R. Pan was visiting Beijing Institute of Applied Physics and Computational Mathematics as an Intellegent Scholar during the summer of 2009. R. Pan is grateful for the generous support of the Beijing IAPCM and the warm hospitality of his hosts, Prof. Jiang and Prof. Ju.

References

- [1] G. Q. Chen and H. Frid, Divergence-measure fields and hyperbolic conservation laws. *Arch. Rat. Mech. Anal.* **147** (1999), pp 89-118.
- [2] G. Q. Chen and T. P. Liu, Zero relaxation and dissipation limits for hyperbolic conservation laws. *Comm. Pure Appl. Math.* **46** (1993), pp 755–781.

-
- [3] G.Q. Chen, C. D. Levermore and T. P. Liu, Hyperbolic conservation laws with stiff relaxation terms and entropy. *Comm. Pure Appl. Math.* **47** (1994), pp 787–830.
- [4] C. M. Dafermos and R. H. Pan, Global BV solutions for the p-system with frictional damping. *SIAM J. Math. Anal.*, **41** (2009), pp 1190–1205.
- [5] X. X. Ding, G.Q. Chen and P.Z. Luo, Convergence of the Lax-Friedrichs scheme for the isentropic gas dynamics. *Acta. Math. Scientia* **5** (1985), pp 415-472; **6** (1986), pp 75-120; **9** (1989), pp 43-44.
- [6] R. J. Diperna, Convergence of the viscosity method for isentropic gas dynamics. *Comm. Math. Phys.* **91** (1983), pp 1-30.
- [7] A. Heidrich, Global weak solutions to initial boundary problems for the wave equation with large data. *Arch. Ration. Mech. Anal.* **126** (1994), pp 333-368.
- [8] L. Hsiao and T. Luo, Stability of travelling wave solutions for a rate-type viscoelastic system. *Advances in Nonlinear Partial Differential Equations and Related Areas*, World Sci. Publishing, River Edge, NJ, 1998.
- [9] L. Hsiao and R. H. Pan, Nonlinear stability of two-mode shock profiles for a rate-type viscoelastic system with relaxation. *Chinese Ann. Math., Ser. B*, **20** (1999), pp 479–488.
- [10] L. Hsiao and R. H. Pan, Zero relaxation limit to centered rarefaction waves for a rate-type viscoelastic system. *J. Differential Equations*, **157** (1999), pp 20–40.
- [11] L. Hsiao and R. H. Pan, Nonlinear stability of rarefaction waves for a rate-type viscoelastic system. *Chinese Ann. Math. Ser. B*, **20** (1999), pp 223–232.
- [12] L. Hsiao and R. H. Pan, The linear stability of traveling wave solutions for a reacting flow model with source term. *Quart. Appl. Math.*, **58** (2000), pp 219–238.
- [13] F. M. Huang, P. Marcati and R. H. Pan, Convergence to Barenblatt solution for the compressible Euler equations with damping and vacuum. *Arch. Ration. Mech. Anal.* **176** (2005), pp 1-24.
- [14] F. M. Huang and R. H. Pan, Convergence rate for compressible Euler equations with damping and vacuum. *Arch. Ration. Mech. Anal.* **166** (2003), pp 359-376.
- [15] F. M. Huang, R. H. Pan and Y. Wang, Stability of contact discontinuity for Jin-Xin relaxation system. *J. Differential Equations*. **244** (2008), pp 1114-1140.
- [16] S. Jin and Z. P. Xin, The relaxation schemes for systems of conservation laws in arbitrary space dimensions. *Comm. Pure Appl. Math.* **48** (1995), pp 235-277.
- [17] C. Lattanzio and P. Marcati, The zero relaxation limit for the hydrodynamic Whitham traffic flow model. *J. Diff. Eqns.* **141** (1997), pp 150-178.
- [18] T. Li and H. L. Liu, Stability of a traffic flow model with nonconvex relaxation. *Commun. Math. Sci.* **3** (2005), pp 101-118.

- [19] T. Li and H. L. Liu, Critical thresholds in hyperbolic relaxation systems. *J. Differential Equations* **247** (2009), pp 33-48.
- [20] H. L. Liu, Asymptotic stability of relaxation shock profiles for hyperbolic conservation laws. *J. Differential Equations*, **192** (2003), pp 285-307.
- [21] H. L. Liu, The L^p stability of relaxation rarefaction profiles. *J. Differential Equations*, **171** (2001), pp 397-411.
- [22] T. P. Liu, Hyperbolic conservation laws with relaxation. *Comm. Math. Phys.* **108** (1987), pp 153-175.
- [23] P. L. Lions, B. Perthame and P.E. Souganidis, Existence and stability of entropy solution for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates. *Comm. Pure Appl. Math.* **49** (1996), pp 599-638.
- [24] P. L. Lions, B. Perthame and E. Tadmor, Kinetic formulation of the isentropic gas dynamics and p-systems. *Comm. Math. Phys.* **163** (1994), pp 415-431.
- [25] C. Mascia and R. Natalini. L^1 nonlinear stability of traveling waves for a hyperbolic system with relaxation. *J. Differential Equations*, **132** (1996), pp 275-292.
- [26] C. Mascia and K. Zumbrun, Stability of large-amplitude shock profiles of general relaxation systems, *SIAM J. Math. Anal.*, **37** (2005), pp 889-913.
- [27] M. Mei and T. Yang Convergence rates to travelling waves for a nonconvex relaxation model, *Proc. Roy. Soc. Edinburgh Sect. A*, **128** (1998), pp 1053-1068.
- [28] R. Natalini, Recent results on hyperbolic relaxation problems, in *Analysis of System of Conservation Laws*, Aachen, 1997, Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math. Vol. 99, pp 128-198.
- [29] R. H. Pan, The nonlinear stability of travelling wave solutions for a reacting flow model with source term. *Acta Math. Sci. (English Ed.)*, **19** (1999), pp 26-36.
- [30] R. H. Pan and K. Zhao, Initial boundary value problem for compressible Euler equations with damping. *Indiana Univ. Math. J.* **57** (2008), pp 2257-2282.
- [31] Y. C. Qiu and K. J. Zhang, On the relaxation limits of the hydrodynamic model for semiconductor devices. *Math. Models and methods in Appl. Science* **12** (2002), pp 333-363.
- [32] D. Serre and L. Xiao, Asymptotic behavior of large weak entropy solutions of the damped p-system. *J. P. Diff. Eqns.* **10** (1997), pp 355-368.
- [33] G. B. Whitham, *Linear and Nonlinear Waves*, Wiley, New York, 1974.
- [34] T. Yang and C. J. Zhu, Existence and non-existence of global smooth solutions for p-system with relaxation. *J. Diff. Eqns* **161** (2000), pp 213-227.

- [35] W. A. Yong, Existence and asymptotic stability of traveling wave solutions of a model system for reacting flow. *Nonlinear Anal.*, **26** (1996), pp 1791–1809.
- [36] B. Zhang, Convergence of the Godunov scheme for a simplified one-dimensional hydrodynamic model for semiconductor devices. *Comm. Math. Phys.* **157** (1993), pp 1-22.
- [37] H. J. Zhao, Nonlinear stability of strong planar rarefaction waves for the relaxation approximation of conservation laws in several space dimensions. *J. Differential Equations*, **163** (2000), pp 198-220.
- [38] C. J. Zhu, Asymptotic behavior of solutions for p-system with relaxation. *J. Differential Equations*, **180** (2002), pp 273-306.