

On the Failure of the Gorenstein Property for Hecke Algebras of Prime Weight

L. J. P. Kilford and Gabor Wiese

CONTENTS

- 1. Introduction
 - 2. Relation between Multiplicity and Gorenstein Defect
 - 3. Modular Symbols and Hecke Algebras
 - 4. Computational Results
 - 5. Further Results and Questions
- Acknowledgments
References

In this article we report on extensive calculations concerning the Gorenstein defect for Hecke algebras of spaces of modular forms of prime weight p at maximal ideals of residue characteristic p such that the attached mod- p Galois representation is unramified at p and the Frobenius at p acts by scalars. The results lead us to ask the question whether the Gorenstein defect and the multiplicity of the attached Galois representation are always equal to 2. We review the literature on the failure of the Gorenstein property and multiplicity one, discuss in some detail a very important practical improvement of the modular-symbols algorithm over finite fields, and include precise statements on the relationship between the Gorenstein defect and the multiplicity of Galois representations.

1. INTRODUCTION

In Wiles's proof of Fermat's last theorem [Wiles 95], an essential step was to show that certain Hecke algebras are Gorenstein rings. Moreover, the Gorenstein property of Hecke algebras is equivalent to the fact that Galois representations appear on certain Jacobians of modular curves precisely with multiplicity one. This article is concerned with the Gorenstein property and with the multiplicity-one question. We report previous work and exhibit many new examples in which multiplicity one and the Gorenstein property fail. We compute the multiplicity in these cases. Moreover, we ask the question, suggested by our computations, whether it is always equal to two if it fails.

We have first to introduce some notation. For integers $N \geq 1$ and $k \geq 2$ and a Dirichlet character $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ we let $S_k(\Gamma_1(N))$ be the \mathbb{C} -vector space of holomorphic cusp forms on $\Gamma_1(N)$ of weight k , and $S_k(N, \chi)$ the subspace on which the diamond operators act through the character χ . We now introduce some extra notation for Hecke algebras over specified rings.

Notation. 1.1. (Notation for Hecke algebras.) Whenever $S \subseteq R$ are rings and M is an R -module on which

2000 AMS Subject Classification: Primary 11F80;
Secondary 11F33, 11F25

Keywords: Multiplicities of Galois representations,
Gorenstein property, Hecke algebras, mod- p modular forms

the Hecke and diamond operators act, we let $\mathbb{T}_S(M)$ be the S -subalgebra inside the R -endomorphism ring of M generated by the Hecke and the diamond operators. If $\phi : S \rightarrow S'$ is a ring homomorphism, we let $\mathbb{T}_\phi(M) := \mathbb{T}_S(M) \otimes_S S'$, or with ϕ understood, $\mathbb{T}_{S \rightarrow S'}(M)$.

We will mostly be dealing with the Hecke algebras $\mathbb{T}_{\mathbb{Z}}(S_k(\Gamma_1(N)))$ and $\mathbb{T}_{\mathbb{Z}[\chi]}(S_k(N, \chi))$, their completions $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{Z}_p}(S_k(\Gamma_1(N)))$ and $\mathbb{T}_{\mathcal{O} \rightarrow \hat{\mathcal{O}}}(S_k(N, \chi))$, as well as their reductions $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))$ and $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(S_k(N, \chi))$. Here, p is a prime and $\mathcal{O} = \mathbb{Z}[\chi]$ is the smallest subring of \mathbb{C} containing all values of χ , $\hat{\mathcal{O}}$ is the completion at a prime above p , and $\mathcal{O} \rightarrow \mathbb{F}$ is the reduction modulo that prime. In Section 3, the reductions of the Hecke algebras are identified with Hecke algebras of mod- p modular forms, which are closely related to Hecke algebras of Katz modular forms over finite fields (see Section 2).

We choose a holomorphic cuspidal Hecke eigenform as the starting point of our discussion and treatment. So let $f \in S_k(N, \chi) \subseteq S_k(\Gamma_1(N))$ be an eigenform for all Hecke and diamond operators. It (more precisely, its Galois conjugacy class) corresponds to minimal ideals, both denoted by \mathfrak{p}_f , in each of the two Hecke algebras $\mathbb{T}_{\mathbb{Z}}(S_k(\Gamma_1(N)))$ and $\mathbb{T}_{\mathbb{Z}[\chi]}(S_k(N, \chi))$. We also choose maximal ideals $\mathfrak{m} = \mathfrak{m}_f$ containing \mathfrak{p}_f of residue characteristic p again in each of the two. Note that the residue fields are the same in both cases.

By work of Shimura and Deligne, one can associate to f (more precisely, to \mathfrak{m}) a continuous odd semisimple Galois representation

$$\rho_f = \rho_{\mathfrak{m}_f} = \rho_{\mathfrak{m}} : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{T}_{\mathbb{Z}}(S_k(N, \chi))/\mathfrak{m})$$

unramified outside Np satisfying $\text{Tr}(\rho_{\mathfrak{m}}(\text{Frob}_l)) \equiv T_l \pmod{\mathfrak{m}}$ and $\text{Det}(\rho_{\mathfrak{m}}(\text{Frob}_l)) \equiv l^{k-1}\chi(l) \pmod{\mathfrak{m}}$ for all primes $l \nmid Np$. In the case of weight $k = 2$ and level N , the representation $\rho_{\mathfrak{m}}$ can be naturally realized on the p -torsion points of the Jacobian of the modular curve $X_1(N)_{\mathbb{Q}}$. The algebra $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(N)))$ acts naturally on $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]$, and we can form the Galois module $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}] = J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p][\tilde{\mathfrak{m}}]$ with $\tilde{\mathfrak{m}}$ the maximal ideal of $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(N)))$, which is the image of \mathfrak{m} under the natural projection. Supposing that $\rho_{\mathfrak{m}}$ is absolutely irreducible, the main result of [Boston et al. 91] shows that the Galois representation $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}]$ is isomorphic to a direct sum of r copies of $\rho_{\mathfrak{m}}$ for some integer $r \geq 1$, which one calls the *multiplicity of $\rho_{\mathfrak{m}}$ on $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}]$* (cf. [Ribet and Stein 01]). We shall for short speak only about the multiplicity of $\rho_{\mathfrak{m}}$. One says that $\rho_{\mathfrak{m}}$ is a *multiplicity-one representation* or *satisfies multiplicity one* if $r = 1$. See [Mazur 77] for a similar

definition for $J_0(N)$ and Proposition 2.6 for a comparison.

The notion of multiplicity can be naturally extended to Galois representations attached to eigenforms f of weights $3 \leq k \leq p + 1$ for $p \nmid N$. This is accomplished by a result of Serre's that implies the existence of a maximal ideal $\mathfrak{m}_2 \subset \mathbb{T}_{\mathbb{Z}}(S_2(\Gamma_1(Np)))$ such that $\rho_{\mathfrak{m}_f} \cong \rho_{\mathfrak{m}_2}$ (see Proposition 2.3). One hence obtains the notion of multiplicity (on $J_1(Np)$) for the representation $\rho_{\mathfrak{m}_f}$ by defining it as the multiplicity of $\rho_{\mathfrak{m}_2}$. Moreover, in allowing twists by the cyclotomic character, it is even possible to treat arbitrary weights. The following theorem summarizes results on when the multiplicity in the above sense is known to be one.

Theorem 1.2. (Mazur, Edixhoven, Tilouine, Gross, Buzzard.) *Let $\rho_{\mathfrak{m}}$ be a representation associated with a modular cuspidal eigenform $f \in S_k(N, \chi)$ and let p be the residue characteristic of \mathfrak{m} . Suppose that $\rho_{\mathfrak{m}}$ is absolutely irreducible and that p does not divide N . If*

1. $2 \leq k \leq p - 1$ or
2. $k = p$ and $\rho_{\mathfrak{m}}$ is ramified at p or
3. $k = p$ and $\rho_{\mathfrak{m}}$ is unramified at p and $\rho_{\mathfrak{m}}(\text{Frob}_p)$ is not scalar,

then the multiplicity of $\rho_{\mathfrak{m}}$ is one.

This theorem is composed of [Mazur 77, Lemma 15.1], [Tilouine 87, Proposition 5.6], [Edixhoven 92, Theorem 9.2], as well as [Gross 90, Proposition 12.10] and [Buzzard 01, Theorem 6.1]. The following theorem by the second author tells us when the multiplicity is not one.

Theorem 1.3. [Wiese 07a, Corollary 4.5] *Let $\rho_{\mathfrak{m}}$ be as in the previous theorem. Suppose $k = p$ and that $\rho_{\mathfrak{m}}$ is unramified at p . If $p = 2$, assume also that a Katz cusp form over \mathbb{F}_2 of weight 1 on $\Gamma_1(N)$ exists that gives rise to $\rho_{\mathfrak{m}}$.*

If $\rho_{\mathfrak{m}}(\text{Frob}_p)$ is a scalar matrix, then the multiplicity of $\rho_{\mathfrak{m}}$ is greater than 1.

In Section 2 we explain how the Galois representation $J_1(Np)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}]$ is related to the different Hecke algebras evoked above and see in many cases of interest a precise relationship between the geometrically defined term *multiplicity* and the *Gorenstein defect* of these algebras. The latter can be computed explicitly, which is the subject of the present article. We now give the relevant definitions.

Definition 1.4. (The Gorenstein property.) Let A be a local Noetherian ring with maximal ideal \mathfrak{m} . Suppose first that the Krull dimension of A is zero, i.e., that A is Artinian. We then define the *Gorenstein defect* of A to be the minimum number of A -module generators of the annihilator of \mathfrak{m} (i.e., $A[\mathfrak{m}]$) minus 1; equivalently, this is the A/\mathfrak{m} -dimension of the annihilator of \mathfrak{m} minus 1. We say that A is *Gorenstein* if its Gorenstein defect is 0, and *non-Gorenstein* otherwise.

If the Krull dimension of A is positive, we inductively call A *Gorenstein* if there exists $x \in \mathfrak{m}$ not a zero divisor such that $A/(x)$ is a Gorenstein ring of smaller Krull dimension (see [Eisenbud 95, p. 532]; note that our definition implies that A is Cohen–Macaulay).

A (not necessarily local) Noetherian ring is said to be *Gorenstein* if all of its localizations at its maximal ideals are Gorenstein.

We will, for example, be interested in the Gorenstein property of $\mathbb{T}_{\mathbb{Z}}(S_k(\Gamma_1(N)))_{\mathfrak{m}}$. Choosing $x = p$ in the definition with p the residue characteristic of \mathfrak{m} , we see that this is equivalent to the Gorenstein defect of the finite-dimensional \mathbb{F}_p -algebra $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))_{\mathfrak{m}}$ being zero. Whenever we refer to the Gorenstein defect of the former algebra (over \mathbb{Z}), we mean that of the latter. Our computations will concern the Gorenstein defect of $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(S_k(\Gamma_1(N), \chi))_{\mathfrak{m}}$. See Section 2 for a comparison with the one not involving a character. It is important to remark that the Gorenstein defect of a local Artin algebra over a field does not change after passing to a field extension and taking any of the conjugate local factors.

We illustrate the definition by an example. The algebra $k[x, y, z]/(x^2, y^2, z^2, xy, xz, yz)$ for a field k is Artinian and local with maximal ideal $\mathfrak{m} := (x, y, z)$ and the annihilator of \mathfrak{m} is \mathfrak{m} itself, so the Gorenstein defect is $3 - 1 = 2$. We note that this particular case does occur in nature; a localization $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_2}(S_2(\Gamma_0(431)))_{\mathfrak{m}}$ at one maximal ideal is isomorphic to this, with $k = \mathbb{F}_2$ (see [Emerton 02, the discussion just before Lemma 6.6]). This example can also be verified with the algorithm presented in this paper.

We now state a translation of Theorem 1.2 in terms of Gorenstein defects, which is immediate from the propositions in Section 2.

Theorem 1.5. *Assume the setup of Theorem 1.2 and that one of the three conditions of that theorem is satisfied. We use notation as in the discussion of multiplicities above.*

If $k = 2$, then $\mathbb{T}_{\mathbb{Z}}(S_2(\Gamma_1(N)))_{\mathfrak{m}}$ is a Gorenstein ring.

If $k \geq 3$, then $\mathbb{T}_{\mathbb{Z}}(S_2(\Gamma_1(Np)))_{\mathfrak{m}_2}$ is, too. Supposing in addition that \mathfrak{m} is ordinary (i.e., $T_p \notin \mathfrak{m}$), then also $\mathbb{T}_{\mathbb{Z}}(S_k(\Gamma_1(N)))_{\mathfrak{m}}$ is Gorenstein. If, moreover, $p \geq 5$ or $\rho_{\mathfrak{m}}$ is not induced from $\mathbb{Q}(\sqrt{-1})$ (if $p = 2$) or $\mathbb{Q}(\sqrt{-3})$ (if $p = 3$), then $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(S_k(N, \chi))_{\mathfrak{m}}$ is Gorenstein as well.

We now turn our attention to computing the Gorenstein defect and the multiplicity in the case in which it is known not to be one.

Corollary 1.6. *Let $\rho_{\mathfrak{m}}$ be a representation associated with a cuspidal eigenform $f \in S_p(N, \chi)$ with p the residue characteristic of \mathfrak{m} . Assume that $\rho_{\mathfrak{m}}$ is absolutely irreducible, unramified at p such that $\rho_{\mathfrak{m}}(\text{Frob}_p)$ is a scalar matrix. Let r be the multiplicity of $\rho_{\mathfrak{m}}$ and d the Gorenstein defect of any of $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(S_k(N, \chi))_{\mathfrak{m}}$, $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))_{\mathfrak{m}}$, $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(Np)))_{\mathfrak{m}_2}$.*

Then the relation $d = 2r - 2$ holds.

Proof: The equality of the Gorenstein defects and the relation with the multiplicity are proved in Section 2, where we note that \mathfrak{m} is ordinary, since $a_p(f)^2 = \chi(p) \neq 0$ (e.g., by [Gross 90, p. 487]). \square

1.1 Previous Results on the Failure of Multiplicity One or the Gorenstein Property

Prior to the present work and the article [Wiese 07a], there have been some investigations into when Hecke algebras fail to be Gorenstein. In [Kilford 02], the first author showed, using MAGMA [Bosma et al. 1997], that none of the three Hecke algebras $\mathbb{T}_{\mathbb{Z}}(S_2(431, \chi_{\text{triv}}))$, $\mathbb{T}_{\mathbb{Z}}(S_2(503, \chi_{\text{triv}}))$, and $\mathbb{T}_{\mathbb{Z}}(S_2(2089, \chi_{\text{triv}}))$ is Gorenstein by explicit computation of the localization of the Hecke algebra at a suitable maximal ideal above 2, and in [Ribet and Stein 01], it is shown in a similar fashion that $\mathbb{T}_{\mathbb{Z}}(S_2(2071, \chi_{\text{triv}}))$ is not Gorenstein. These examples were discovered by considering elliptic curves E/\mathbb{Q} such that in the ring of integers of $\mathbb{Q}(E[2])$ the prime ideal (2) splits completely, and then doing computations with MAGMA.

There are also some results in the literature on the failure of multiplicity one within the torsion of certain Jacobians. In [Agashe et al. 06, Proposition 5.1], the following theorem is proved:

Theorem 1.7. (Agashe, Ribet, Stein.) *Suppose that E is an optimal elliptic curve over \mathbb{Q} of conductor N , with congruence number r_E and modular degree m_E , and that p is a prime such that $p \mid r_E$ but $p \nmid m_E$. Let \mathfrak{m} be the*

annihilator in $\mathbb{T}_{\mathbb{Z}}(S_2(N, \chi_{\text{triv}}))$ of $E[p]$. Then multiplicity one fails for \mathfrak{m} .

The authors give a table of examples; for instance, $\mathbb{T}_{\mathbb{Z}}(S_2(54, \chi_{\text{triv}}))$ does not satisfy multiplicity one at some maximal ideal above 3. It is not clear whether this phenomenon occurs infinitely often.

The first examples of the failure of multiplicity one were given in [Mazur and Ribet 92], where it is proved in Theorem 2 that the mod- p multiplicity of a suitable representation $\rho_{\mathfrak{m}}$ of level prime to N in the torsion of the Jacobian $J_0(p^3N)$ is greater than 1. The explicit example given there is of a representation of level 11 having mod-11 multiplicity greater than one in $J_0(11^3)[11]$.

In [Ribet 90], it is shown that the mod- p multiplicity of a certain representation in the Jacobian of the Shimura curve derived from the rational quaternion algebra of discriminant $11 \cdot 193$ is 2; this result inspired the calculations in [Kilford 02].

Let us finally mention that for $p = 2$, the Galois representations ρ with image equal to the dihedral group D_3 come from an elliptic curve over \mathbb{Q} . We observe that $D_3 = \text{GL}_2(\mathbb{F}_2)$. Any S_3 -extension K of the rationals can be obtained as the splitting field of an irreducible integral polynomial $f = X^3 + aX + b$. The 2-torsion of the elliptic curve $E : Y^2 = f$ consists precisely of the three roots of f and the point at infinity. So the field generated over \mathbb{Q} by the 2-torsion of E is K .

1.2 New Results

Using the modular-symbols algorithm over finite fields with an improved stop criterion (see Section 3), we performed computations in MAGMA concerning the Gorenstein defect of Hecke algebras of cuspidal modular forms of prime weights p at maximal ideals of residue characteristic p in the case of Theorem 1.3. All of our 384 examples have Gorenstein defect equal to 2, and hence their multiplicity is 2.

We formulate part of our computational findings as a theorem.

Theorem 1.8. *For every prime $p < 100$ there exists a prime $N \neq p$ and a Dirichlet character χ such that the Hecke algebra $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_p(N, \chi))$ has Gorenstein defect 2 at some maximal ideal \mathfrak{m} of residue characteristic p . The corresponding Galois representation $\rho_{\mathfrak{m}}$ appears with multiplicity two on the \mathfrak{m} -torsion of the Jacobian $J_1(Np)_{\mathbb{Q}}(\mathbb{Q})$ if p is odd, and of the Jacobian $J_1(N)_{\mathbb{Q}}(\mathbb{Q})$ if $p = 2$.*

Our computational results are discussed in more detail in Section 4.

1.3 A Question

Question 1.9. Let p be a prime. Let f be a normalized cuspidal modular eigenform of weight p , prime level $N \neq p$ for some Dirichlet character χ . Let $\rho_f : G_{\mathbb{Q}} \rightarrow \text{GL}_2(\overline{\mathbb{F}}_p)$ be the modular Galois representation attached to f . We assume that ρ_f is irreducible and unramified at p and that $\rho_f(\text{Frob}_p)$ is a scalar matrix.

Write $\mathbb{T}_{\mathbb{F}}$ for $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_p(N, \chi))$. Recall that this notation stands for the tensor product over $\mathbb{Z}[\chi]$ of a residue field \mathbb{F}/\mathbb{F}_p of $\mathbb{Z}[\chi]$ by the $\mathbb{Z}[\chi]$ -algebra generated inside the endomorphism algebra of $S_p(N, \chi)$ by the Hecke operators and by the diamond operators. Let \mathfrak{m} be the maximal ideal of $\mathbb{T}_{\mathbb{F}}$ corresponding to f .

Is the Gorenstein defect of the Hecke algebra $\mathbb{T}_{\mathbb{F}}$ localized at \mathfrak{m} , denoted by $\mathbb{T}_{\mathfrak{m}}$, always equal to 2?

Equivalently, is the multiplicity of the Galois representation attached to f always equal to 2?

This question was also raised by both Kevin Buzzard and James Parson in communications to the authors.

2. RELATION BETWEEN MULTIPLICITY AND GORENSTEIN DEFECT

In this section we collect results, some of which are well known, on the multiplicity of Galois representations, the Gorenstein defect, and relations between the two. Whereas the mod- p modular-symbols algorithm naturally computes mod- p modular forms (see Section 3), this rather geometrical section uses (mostly in the references) the theory of Katz modular forms over finite fields (see, e.g., [Edixhoven 97]). If $N \geq 5$ and $k \geq 2$, the Hecke algebra $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))$ is both the Hecke algebra of mod- p cusp forms of weight k on $\Gamma_1(N)$ and the Hecke algebra of the corresponding Katz cusp forms over \mathbb{F}_p . However, in the presence of a Dirichlet character, $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))$ has an interpretation only as the Hecke algebra of the corresponding mod- p cusp forms, and there may be differences with the respective Hecke algebra for Katz forms (see Carayol's lemma, which is [Edixhoven 97, Proposition 1.10]).

We start with the well-known result in weight 2 (see, e.g., [Mazur 77, Lemma 15.1] or [Tilouine 97]) that multiplicity one implies that the corresponding local Hecke factor is a Gorenstein ring.

Proposition 2.1. *Let \mathfrak{m} be a maximal ideal of $\mathbb{T} := \mathbb{T}_{\mathbb{Z}}(S_2(\Gamma_1(N)))$ of residue characteristic p , which may divide N . Denote by $\tilde{\mathfrak{m}}$ the image of \mathfrak{m} in $\mathbb{T}_{\mathbb{F}_p} := \mathbb{T} \otimes_{\mathbb{Z}} \mathbb{F}_p = \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(N)))$. Suppose that the Galois representation $\rho_{\mathfrak{m}}$ is irreducible and satisfies multiplicity one (see Section 1).*

Then as $\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}$ -modules one has

$$J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]_{\tilde{\mathfrak{m}}} \cong \mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}} \oplus \mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}},$$

and the localizations $\mathbb{T}_{\mathfrak{m}}$ and $\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}$ are Gorenstein rings. Similar results hold if one replaces $\Gamma_1(N)$ and $J_1(N)$ by $\Gamma_0(N)$ and $J_0(N)$.

Proof: For the proof we have to pass to $\mathbb{T}_{\mathbb{Z}_p} = \mathbb{T}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Z}_p$. We also denote by \mathfrak{m} the maximal ideal in $\mathbb{T}_{\mathbb{Z}_p}$ that corresponds to \mathfrak{m} . Let V be the \mathfrak{m} -part of the p -Tate module of $J_1(N)_{\mathbb{Q}}$. Multiplicity one implies that $V/\mathfrak{m}V$ is a 2-dimensional $\mathbb{T}/\mathfrak{m} = \mathbb{T}_{\mathbb{Z}_p}/\mathfrak{m} = \mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}/\tilde{\mathfrak{m}}$ -vector space, since

$$\begin{aligned} V/\mathfrak{m}V &\cong (V/pV)/\tilde{\mathfrak{m}} \cong (J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p])/\tilde{\mathfrak{m}} \\ &\cong (J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p])^{\vee}/\tilde{\mathfrak{m}} \cong (J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}])^{\vee}, \end{aligned}$$

where the self-duality comes from the modified Weil pairing that respects the Hecke action (see, e.g., [Tilouine 87, Lemme 4.1] or [Gross 90, p. 485]). Nakayama's lemma hence implies that V is a $\mathbb{T}_{\mathbb{Z}_p, \mathfrak{m}}$ -module of rank 2. Since one knows that $V \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ is a $\mathbb{T}_{\mathfrak{m}} \otimes \mathbb{Q}_p$ -module of rank 2, it follows that V is a free $\mathbb{T}_{\mathbb{Z}_p, \mathfrak{m}}$ -module of rank 2, whence $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]_{\tilde{\mathfrak{m}}}$ is a free $\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}$ -module of rank 2. Taking the $\tilde{\mathfrak{m}}$ -kernel gives $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}] = (\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}[\tilde{\mathfrak{m}}])^2$, whence the Gorenstein defect is zero. In the Γ_0 -situation, the same proof holds. \square

In the so-called ordinary case, we have the following precise relationship between the multiplicity and the Gorenstein defect, which was suggested to us by Kevin Buzzard. A proof can be found in [Wiese 07a, Corollaries 2.3 and 4.2].

Proposition 2.2. *Suppose $p \nmid N$ and let $M = N$ or $M = Np$. Let \mathfrak{m} be a maximal ideal of $\mathbb{T}_{\mathbb{Z}}(S_2(\Gamma_1(M)))$ of residue characteristic p and assume that \mathfrak{m} is ordinary, i.e., that the p th Hecke operator T_p is not in \mathfrak{m} . Assume also that $\rho_{\mathfrak{m}}$ is irreducible. Denote by $\tilde{\mathfrak{m}}$ the image of \mathfrak{m} in $\mathbb{T}_{\mathbb{F}_p} := \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(M)))$. Then the following statements hold:*

(a) *There is the exact sequence*

$$0 \rightarrow \mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}} \rightarrow J_1(M)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]_{\tilde{\mathfrak{m}}} \rightarrow \mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}^{\vee} \rightarrow 0$$

of $\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}$ -modules, where the dual is the \mathbb{F}_p -linear dual.

(b) *If d is the Gorenstein defect of $\mathbb{T}_{\mathbb{F}_p, \tilde{\mathfrak{m}}}$ and r is the multiplicity of $\rho_{\mathfrak{m}}$, then the relation*

$$d = 2r - 2$$

holds.

We now establish a relation between mod- p Hecke algebras of weights $3 \leq k \leq p+1$ for levels N not divisible by p and Hecke algebras of weight 2 and level Np . It is needed in order to compare the Hecke algebras in higher weight to those acting on the p -torsion of Jacobians and thus to make a link to the multiplicity of the attached Galois representations.

Proposition 2.3. *Let $N \geq 5$, $p \nmid N$, and $3 \leq k \leq p+1$. Let \mathfrak{m} be a maximal ideal of the mod- p Hecke algebra $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))$. Then there exists a maximal ideal \mathfrak{m}_2 of $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(Np)))$ and a natural surjection*

$$\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(Np)))_{\mathfrak{m}_2} \twoheadrightarrow \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))_{\mathfrak{m}}.$$

If \mathfrak{m} is ordinary, i.e., $T_p \notin \mathfrak{m}$, this surjection is an isomorphism.

Proof: From [Wiese 07b, Sections 5 and 6], whose notation we adopt for this proof, one obtains without difficulty the commutative diagram of Hecke algebras shown in Figure 1.

The claimed surjection can be read off. The ideal \mathfrak{m}_2 can be explicitly defined as the preimage of \mathfrak{m} (before localization). Then it necessarily holds that $\langle a \rangle_p - a^{k-2}$ is in \mathfrak{m}_2 for all $a \in (\mathbb{Z}/p\mathbb{Z})^{\times}$. In the ordinary situation, Proposition 2.2 shows that the upper-left horizontal arrow is in fact an isomorphism. That also the upper-right horizontal arrow is an isomorphism is explained in [Wiese 07b]. The result follows. \square

As pointed out by one of the referees, the result in the ordinary case was first obtained in [Hida 81]. In the next proposition we compare Hecke algebras for spaces of modular forms on $\Gamma_1(N)$ to those of the same level and weight, but with a Dirichlet character.

Proposition 2.4. *Let $N \geq 5$, $k \geq 2$, and let $\chi : (\mathbb{Z}/N\mathbb{Z})^{\times} \rightarrow \mathbb{C}^{\times}$ be a Dirichlet character. Let $f \in S_k(N, \chi) \subset S_k(\Gamma_1(N))$ be a normalized Hecke eigenform. Let further $\mathfrak{m}_{\overline{\chi}}$ be the maximal ideal in*

$$\begin{array}{ccccc}
 \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(Np))_{\mathfrak{m}_2}) & \twoheadrightarrow & \mathbb{T}_{\mathbb{F}_p}(J_1(Np)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p])_{\mathfrak{m}_2} & \twoheadrightarrow & \mathbb{T}_{\mathbb{F}_p}(H^1(\Gamma_1(N), V_{k-2}(\mathbb{F}_p)))_{\mathfrak{m}} \\
 \parallel & & \downarrow & & \uparrow \\
 \mathbb{T}_{\mathbb{Z}_p \rightarrow \mathbb{F}_p}(L)_{\mathfrak{m}_2} & \twoheadrightarrow & \mathbb{T}_{\mathbb{F}_p}(\overline{L})_{\mathfrak{m}_2} & \twoheadrightarrow & \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))_{\mathfrak{m}}
 \end{array}$$

FIGURE 1. Commutative diagram for the proof of Proposition 2.3.

$\mathbb{T}_{\mathbb{F}}^{\overline{\chi}} := \mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))$, and \mathfrak{m} the one in $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))$ of residue characteristic p for $p \nmid N$ belonging to f . If $k = 2$, suppose additionally that $\rho_{\mathfrak{m}}$ is irreducible. If $p = 2$, suppose that $\rho_{\mathfrak{m}}$ is not induced from $\mathbb{Q}(\sqrt{-1})$, and if $p = 3$, suppose that $\rho_{\mathfrak{m}}$ is not induced from $\mathbb{Q}(\sqrt{-3})$.

Then the Gorenstein defects of $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))_{\mathfrak{m}_{\overline{\chi}}}$ and $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_k(\Gamma_1(N)))_{\mathfrak{m}}$ are equal.

Proof: Write $\Delta := (\mathbb{Z}/N\mathbb{Z})^\times$ and let Δ_p be its p -Sylow subgroup. Let $\overline{\chi} : \Delta \rightarrow \mathbb{F}^\times$ be the reduction of χ obtained by composing χ with $\mathbb{Z}[\chi] \rightarrow \mathbb{F}$. Since the Gorenstein defect is invariant under base extension, it is no loss to work with $\mathbb{T}_{\mathbb{F}} := \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}}(S_k(\Gamma_1(N)))$. We still write \mathfrak{m} for the maximal ideal in $\mathbb{T}_{\mathbb{F}}$ belonging to f . Note that $\langle \delta \rangle - \overline{\chi}(\delta) \in \mathfrak{m}$ for all $\delta \in \Delta$.

We let Δ act on $\mathbb{T}_{\mathbb{F}}$ via the diamond operators, and we let $\mathbb{F}^{\overline{\chi}}$ be a copy of \mathbb{F} with Δ -action through the inverse of $\overline{\chi}$. We have

$$(\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta} = (\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}}) / (1 - \delta | \delta \in \Delta) \cong \mathbb{T}_{\mathbb{F}, \mathfrak{m}_{\overline{\chi}}}^{\overline{\chi}},$$

which one obtains by considering the duals, identifying Katz cusp forms with mod- p ones on $\Gamma_1(N)$, and applying Carayol’s lemma [Edixhoven 97, Proposition 1.10]. For the case $k = 2$, we should point the reader to the correction at the end of the introduction to [Edixhoven 06]. However, the statement still holds after localization at maximal ideals corresponding to irreducible representations. Moreover, the equality $(\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta} = \mathbb{T}_{\mathbb{F}, \mathfrak{m}}[\langle \delta \rangle - \overline{\chi}(\delta) | \delta \in \Delta]$ holds by definition.

Now [Wiese 07b, Lemma 7.3] tells us that the localization at \mathfrak{m} of the \mathbb{F} -vector space of Katz cusp forms of weight k on $\Gamma_1(N)$ over \mathbb{F} is a free $\mathbb{F}[\Delta_p]$ -module. Note that the standing hypothesis $k \leq p + 1$ of [Wiese 07b, Section 7] is not used in the proof of that lemma and see also [Wiese 07b, Remark 7.5]. From an elementary calculation one now obtains that $N_{\Delta} = \sum_{\delta \in \Delta} \delta$ induces an isomorphism

$$(\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta} \xrightarrow{N_{\Delta}} (\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta}.$$

We now take the $\mathfrak{m}_{\overline{\chi}}$ -kernel on both sides and obtain

$$\begin{aligned}
 \mathbb{T}_{\mathbb{F}, \mathfrak{m}_{\overline{\chi}}}^{\overline{\chi}}[\mathfrak{m}_{\overline{\chi}}] &\cong (\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta}[\mathfrak{m}_{\overline{\chi}}] \cong (\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})_{\Delta}[\mathfrak{m}] \\
 &\cong (\mathbb{T}_{\mathbb{F}, \mathfrak{m}} \otimes_{\mathbb{F}} \mathbb{F}^{\overline{\chi}})^{\Delta}[\mathfrak{m}] = \mathbb{T}_{\mathbb{F}, \mathfrak{m}}[\mathfrak{m}].
 \end{aligned}$$

This proves that the two Gorenstein defects are indeed equal. \square

The Gorenstein defect that we calculate on the computer is the number d of the following corollary, which relates it to the multiplicity of a Galois representation.

Corollary 2.5. *Let p be a prime, $N \geq 5$ an integer such that $p \nmid N$, k an integer satisfying $2 \leq k \leq p$, and $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ a character. Let $f \in S_k(N, \chi)$ be a normalized Hecke eigenform. Let further \mathfrak{m} denote the maximal ideal in $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))$ belonging to f . Suppose that \mathfrak{m} is ordinary and that $\rho_{\mathfrak{m}}$ is irreducible and not induced from $\mathbb{Q}(\sqrt{-1})$ (if $p = 2$) and not induced from $\mathbb{Q}(\sqrt{-3})$ (if $p = 3$). We define d to be the Gorenstein defect of $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))_{\mathfrak{m}}$ and r to be the multiplicity of $\rho_{\mathfrak{m}}$.*

Then the equality $d = 2r - 2$ holds.

We include the following proposition because it establishes equality of the two different notions of multiplicities of Galois representations in the case of the trivial character.

Proposition 2.6. *Let $N \geq 1$ and $p \nmid N$ and $f \in S_2(\Gamma_0(N)) \subseteq S_2(\Gamma_1(N))$ be a normalized Hecke eigenform belonging to maximal ideals $\mathfrak{m}_0 \subseteq \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_0(N)))$ and $\mathfrak{m}_1 \subseteq \mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_p}(S_2(\Gamma_1(N)))$ of residue characteristic p . Suppose that $\rho_{\mathfrak{m}_0} \cong \rho_{\mathfrak{m}_1}$ is irreducible.*

Then the multiplicity of $\rho_{\mathfrak{m}_1}$ on $J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]$ is equal to the multiplicity of $\rho_{\mathfrak{m}_0}$ on $J_0(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]$. Thus, if $p > 2$, this multiplicity is equal to one by Theorem 1.2.

Proof: Let $\Delta := (\mathbb{Z}/N\mathbb{Z})^\times$. We first remark that one has the isomorphism

$$J_0(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p]_{\mathfrak{m}_0} \cong ((J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[p])_{\Delta})_{\mathfrak{m}_0},$$

which one can, for example, obtain by comparing with the parabolic cohomology with \mathbb{F}_p -coefficients of the modular curves $Y_0(N)$ and $Y_1(N)$. Taking the \mathfrak{m}_0 -kernel yields

$$J_0(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}_0] \cong J_1(N)_{\mathbb{Q}}(\overline{\mathbb{Q}})[\mathfrak{m}_1],$$

since \mathfrak{m}_1 contains $\langle \delta \rangle - 1$ for all $\delta \in \Delta$. \square

3. MODULAR SYMBOLS AND HECKE ALGEBRAS

The aim of this section is to present the algorithm that we use for the computations of local factors of Hecke algebras of mod- p modular forms. It is based on mod- p modular symbols, which have been implemented in MAGMA [Bosma et al. 1997] by William Stein.

The bulk of this section deals with proving the main advance, namely a stop criterion (Corollary 3.8), which in practice greatly speeds up the computations in comparison with “standard” implementations, since it allows us to work with many fewer Hecke operators than indicated by the theoretical Sturm bound (Proposition 3.10). We shall list results proving that the stop criterion is attained in many cases. However, the stop criterion does not depend on them, in the sense that it being attained is equivalent to a proof that the algebra it outputs is equal to a direct factor of a Hecke algebra of mod- p modular forms.

Whereas for Section 2 the notion of Katz modular forms seems the right one, the present section works entirely with mod- p modular forms, the definition of which is also recalled. This is very natural, since all results in this section are based on a comparison with the characteristic-zero theory.

3.1 Mod- p Modular Forms and Modular Symbols

3.1.1 Mod- p Modular Forms. Let us for the time being fix integers $N \geq 1$ and $k \geq 2$, as well as a character $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ such that $\chi(-1) = (-1)^k$. Let $M_k(N, \chi)$ be the space of holomorphic modular forms for $\Gamma_1(N)$, Dirichlet character χ , and weight k . It decomposes as a direct sum (orthogonal direct sum with respect to the Petersson inner product) of its cuspidal subspace $S_k(N, \chi)$ and its Eisenstein subspace $\text{Eis}_k(N, \chi)$. As before, we let $\mathcal{O} = \mathbb{Z}[\chi]$. Moreover, we let \mathfrak{P} be a maximal ideal of \mathcal{O} above p with residue field \mathbb{F} , and $\widehat{\mathcal{O}}$ the completion of \mathcal{O} at \mathfrak{P} . Furthermore, let $K = \mathbb{Q}_p(\chi)$ be the field of fractions of $\widehat{\mathcal{O}}$, and let $\bar{\chi}$ be χ followed by the natural projection $\mathcal{O} \rightarrow \mathbb{F}$.

Denote by $M_k(N, \chi; \mathcal{O})$ the sub- \mathcal{O} -module generated by those modular forms whose (standard) q -expansion

has coefficients in \mathcal{O} . It follows from the q -expansion principle that

$$M_k(N, \chi; \mathcal{O}) \cong \text{Hom}_{\mathcal{O}}(\mathbb{T}_{\mathcal{O}}(M_k(N, \chi)), \mathcal{O})$$

and hence that $M_k(N, \chi; \mathcal{O}) \otimes_{\mathcal{O}} \mathbb{C} \cong M_k(N, \chi)$. We put

$$\begin{aligned} M_k(N, \bar{\chi}; \mathbb{F}) &:= M_k(N, \chi; \mathcal{O}) \otimes_{\mathcal{O}} \mathbb{F} \\ &\cong \text{Hom}_{\mathcal{O}}(\mathbb{T}_{\mathcal{O}}(M_k(N, \chi)), \mathbb{F}) \end{aligned}$$

and call the elements of this space *mod- p modular forms*. The Hecke algebra $\mathbb{T}_{\mathcal{O}}(M_k(N, \chi))$ acts naturally, and it follows that $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(M_k(N, \chi)) \cong \mathbb{T}_{\mathbb{F}}(M_k(N, \chi; \mathbb{F}))$. Similar statements hold for the cuspidal and the Eisenstein subspaces, and we use similar notation.

We call a maximal ideal \mathfrak{m} of $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(M_k(N, \chi; \mathcal{O}))$ (respectively, the corresponding maximal ideal of $\mathbb{T}_{\mathcal{O} \rightarrow \widehat{\mathcal{O}}}(M_k(N, \chi; \mathcal{O}))$) *non-Eisenstein* if

$$S_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} \cong M_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}.$$

Otherwise, we call \mathfrak{m} *Eisenstein*.

We now include a short discussion of minimal and maximal primes, in view of Proposition 3.5. Write $\mathbb{T}_{\widehat{\mathcal{O}}}$ for $\mathbb{T}_{\mathcal{O} \rightarrow \widehat{\mathcal{O}}}(S_k(N, \chi))$. Let \mathfrak{m} be a maximal ideal of $\mathbb{T}_{\widehat{\mathcal{O}}}$. It corresponds to a $\text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F})$ -conjugacy class of normalized eigenforms in $S_k(N, \bar{\chi}; \mathbb{F})$. This means that for each $n \in \mathbb{N}$, the minimal polynomial of T_n acting on $S_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}$ is equal to a power of the minimal polynomial of the coefficient a_n of each member of the conjugacy class. Similarly, a minimal prime \mathfrak{p} of $\mathbb{T}_{\mathcal{O} \rightarrow \widehat{\mathcal{O}}}(S_k(N, \chi))$ corresponds to a $\text{Gal}(\overline{\mathbb{Q}}_p/K)$ -conjugacy class of normalized eigenforms in $S_k(N, \chi; \mathcal{O}) \otimes_{\mathcal{O}} K$.

Suppose that \mathfrak{m} contains minimal primes \mathfrak{p}_i for $i = 1, \dots, r$. Then the normalized eigenforms corresponding to the \mathfrak{p}_i are congruent to one another modulo a prime above p . Conversely, every congruence arises in this way. Thus, a maximal ideal \mathfrak{m} of $\mathbb{T}_{\widehat{\mathcal{O}}}$ is Eisenstein if and only if it contains a minimal prime corresponding to a conjugacy class of Eisenstein series. Since it is the reduction of a reducible representation, the mod- p Galois representation corresponding to an Eisenstein prime is reducible. It should be possible to show the converse, too.

3.1.2 Modular Symbols. We now recall the modular-symbols formalism and prove two useful results on base change and torsion. The main references for the definitions are [Stein 07] and [Wiese 08].

Let R be a ring, $\Gamma \leq \text{SL}_2(\mathbb{Z})$ a subgroup, and V a left $R[\Gamma]$ -module. Recall that $\mathbb{P}^1(\mathbb{Q}) = \mathbb{Q} \cup \{\infty\}$ is the set of cusps of $\text{SL}_2(\mathbb{Z})$, which carries a natural $\text{SL}_2(\mathbb{Z})$ -action via fractional linear transformations. We define

the R -modules

$$\begin{aligned} \mathcal{M}_R &:= R[\{\alpha, \beta\} \mid \alpha, \beta \in \mathbb{P}^1(\mathbb{Q})] \\ &\quad / \langle \{\alpha, \alpha\}\{\alpha, \beta\} + \{\beta, \gamma\} + \{\gamma, \alpha\} \\ &\quad \mid \alpha, \beta, \gamma \in \mathbb{P}^1(\mathbb{Q}) \rangle \end{aligned}$$

and $\mathcal{B}_R := R[\mathbb{P}^1(\mathbb{Q})]$. They are connected via the boundary map $\delta : \mathcal{M}_R \rightarrow \mathcal{B}_R$, which is given by $\{\alpha, \beta\} \mapsto \beta - \alpha$. Both are equipped with the natural left Γ -actions. Also let $\mathcal{M}_R(V) := \mathcal{M}_R \otimes_R V$ and $\mathcal{B}_R(V) := \mathcal{B}_R \otimes_R V$ with the left diagonal Γ -action. We call the Γ -coinvariants

$$\begin{aligned} \mathcal{M}_R(\Gamma, V) &:= \mathcal{M}_R(V)_\Gamma \\ &= \mathcal{M}_R(V) / \langle (x - gx) \mid g \in \Gamma, x \in \mathcal{M}_R(V) \rangle \end{aligned}$$

the space of (Γ, V) -modular symbols. Furthermore, the space of (Γ, V) -boundary symbols is defined as the Γ -coinvariants

$$\begin{aligned} \mathcal{B}_R(\Gamma, V) &:= \mathcal{B}_R(V)_\Gamma \\ &= \mathcal{B}_R(V) / \langle (x - gx) \mid g \in \Gamma, x \in \mathcal{B}_R(V) \rangle. \end{aligned}$$

The boundary map δ induces the boundary map $\mathcal{M}_R(\Gamma, V) \rightarrow \mathcal{B}_R(\Gamma, V)$. Its kernel is denoted by $\mathcal{CM}_R(\Gamma, V)$ and is called the space of cuspidal (Γ, V) -modular symbols.

Let now $N \geq 1$ and $k \geq 2$ be integers and $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow R^\times$ a character, i.e., a group homomorphism, such that $\chi(-1) = (-1)^k$ in R . Write $V_{k-2}(R)$ for the homogeneous polynomials of degree $k - 2$ over R in two variables, equipped with the natural $\Gamma_0(N)$ -action. Denote by $V_{k-2}^\chi(R)$ the tensor product $V_{k-2}(R) \otimes_R R^\chi$ for the diagonal $\Gamma_0(N)$ -action that on R^χ comes from the isomorphism $\Gamma_0(N)/\Gamma_1(N) \cong (\mathbb{Z}/N\mathbb{Z})^\times$ given by sending $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ to d followed by χ^{-1} .

We use the notation $\mathcal{M}_k(N, \chi; R)$ for $\mathcal{M}(\Gamma_0(N), V_{k-2}^\chi(R))$, as well as similarly for the boundary and the cuspidal spaces. The natural action of the matrix $\eta = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ gives an involution on all of these spaces. We will denote by the superscript $+$ the subspace invariant under this involution, and by the superscript $-$ the anti-invariant one. On all modules discussed so far, one has Hecke operators T_n for all $n \in \mathbb{N}$ and diamond operators. For a definition, see [Stein 07].

Lemma 3.1. *Let R, Γ , and V be as above and let $R \rightarrow S$ be a ring homomorphism. Then*

$$\mathcal{M}(\Gamma, V) \otimes_R S \cong \mathcal{M}(\Gamma, V \otimes_R S).$$

Proof: This follows immediately from the fact that tensoring and taking coinvariants are both right exact. \square

Proposition 3.2. *Let R be a local integral domain of characteristic zero with principal maximal ideal $\mathfrak{m} = (\pi)$ and residue field \mathbb{F} of characteristic p . Also let $N \geq 1, k \geq 2$, be integers and $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow R^\times$ a character such that $\chi(-1) = (-1)^k$. Suppose (i) that $p \geq 5$ or (ii) that $p = 2$ and N is divisible by a prime that is 3 modulo 4 or by 4 or (iii) that $p = 3$ and N is divisible by a prime that is 2 modulo 3 or by 9. Then the following statements hold:*

- (a) *If $k \geq 3$, then $\mathcal{M}_k(N, \chi; R)[\pi] = (V_{k-2}^\chi(\mathbb{F}))^{\Gamma_0(N)}$.*
- (b) *If $k = 2$ or if $3 \leq k \leq p + 2$ and $p \nmid N$, then $\mathcal{M}_k(N, \chi; R)[\pi] = 0$.*

Proof: The conditions ensure that the group $\Gamma_0(N)$ does not have any stabilizer of order $2p$ for its action on the upper half-plane. Hence, by [Wiese 08, Theorem 6.1], the modular-symbols space $\mathcal{M}_k(N, \chi; R)$ is isomorphic to $H^1(\Gamma_0(N), V_{k-2}^\chi(R))$. The arguments are now precisely those of the beginning of the proof of [Wiese 07b, Proposition 2.6]. \square

3.1.3 Hecke Algebras of Modular Symbols and the Eichler–Shimura Isomorphism. From Lemma 3.1 one deduces a natural surjection

$$\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(\mathcal{M}_k(N, \chi; \mathcal{O})) \rightarrow \mathbb{T}_{\mathbb{F}}(\mathcal{M}_k(N, \bar{\chi}; \mathbb{F})). \quad (3-1)$$

In the same way, one also obtains

$$\begin{aligned} \mathbb{T}_{\mathcal{O}}(\mathcal{M}_k(N, \chi; \mathcal{O})) &\rightarrow \mathbb{T}_{\mathcal{O}}(\mathcal{M}_k(N, \chi; \mathcal{O})/\text{torsion}) \\ &\cong \mathbb{T}_{\mathcal{O}}(\mathcal{M}_k(N, \chi; \mathbb{C})), \end{aligned} \quad (3-2)$$

where one uses for the isomorphism that the Hecke operators are already defined over \mathcal{O} . Similar statements hold for the cuspidal subspace.

We call a maximal prime \mathfrak{m} of $\mathbb{T}_{\mathcal{O} \rightarrow \hat{\mathcal{O}}}(\mathcal{M}_k(N, \chi; \mathcal{O}))$ (respectively the corresponding prime of $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(\mathcal{M}_k(N, \chi; \mathcal{O}))$) *nontorsion* if

$$\mathcal{M}_k(N, \chi; \hat{\mathcal{O}})_{\mathfrak{m}} \cong (\mathcal{M}_k(N, \chi; \hat{\mathcal{O}})/\text{torsion})_{\mathfrak{m}}.$$

This is equivalent to the height of \mathfrak{m} being 1. Proposition 3.2 tells us some cases in which all primes are nontorsion.

Theorem 3.3. (Eichler–Shimura.) *There are isomorphisms respecting the Hecke operators*

- (a) $M_k(N, \chi) \oplus S_k(N, \chi)^\vee \cong \mathcal{M}_k(N, \chi; \mathbb{C})$,
- (b) $S_k(N, \chi) \oplus S_k(N, \chi)^\vee \cong \mathcal{CM}_k(N, \chi; \mathbb{C})$,
- (c) $S_k(N, \chi) \cong \mathcal{CM}_k(N, \chi; \mathbb{C})^+$.

Proof: Parts (a) and (b) are [Diamond and Im 95, Theorem 12.2.2], together with the comparison of [Wiese 08, Theorem 6.1]. We use that the space of antiholomorphic cusp forms is dual to the space of holomorphic cusp forms. Part (c) is a direct consequence of (b). \square

Corollary 3.4. *There are isomorphisms*

$$\begin{aligned} \mathbb{T}_{\mathcal{O}}(S_k(N, \chi)) &\cong \mathbb{T}_{\mathcal{O}}(\mathcal{CM}_k(N, \chi; \mathbb{C})) \\ &\cong \mathbb{T}_{\mathcal{O}}(\mathcal{CM}_k(N, \chi; \mathbb{C})^+), \end{aligned}$$

given by sending T_n to T_n for all positive n .

3.2 The Stop Criterion

Although it is impossible to determine a priori the dimension of the local factor of the Hecke algebra associated with a given modular form mod p , Corollary 3.8 implies that the computation of Hecke operators can be stopped when the algebra generated has reached a certain dimension that is computed along the way. This criterion has turned out to be extremely useful and has made possible some of our computations that would not have been feasible using the Hecke bound naively. See Section 4 for a short discussion of this issue.

3.2.1 Some Commutative Algebra. We collect some useful statements from commutative algebra, which will be applied to Hecke algebras in the sequel.

Proposition 3.5. *Let R be an integral domain of characteristic zero that is a finitely generated \mathbb{Z} -module. Write \widehat{R} for the completion of R at a maximal ideal of R and denote by \mathbb{F} the residue field and by K the fraction field of \widehat{R} . Let furthermore A be a commutative R -algebra that is finitely generated as an R -module. For any ring homomorphism $R \rightarrow S$ write A_S for $A \otimes_R S$. Then the following statements hold:*

- (a) *The Krull dimension of $A_{\widehat{R}}$ is less than or equal to 1. The maximal ideals of $A_{\widehat{R}}$ correspond bijectively under taking preimages to the maximal ideals of $A_{\mathbb{F}}$. Primes \mathfrak{p} of height 0 that are contained in a prime of height 1 of $A_{\widehat{R}}$ are in bijection with primes of A_K under extension (i.e., $\mathfrak{p}A_K$), for which the notation \mathfrak{p}^e will be used.*

Under these correspondences, one has $A_{\mathbb{F}, \mathfrak{m}} \cong A_{\widehat{R}, \mathfrak{m}} \otimes_{\widehat{R}} \mathbb{F}$, and $A_{K, \mathfrak{p}^e} \cong A_{\widehat{R}, \mathfrak{p}}$.

- (b) *The algebra $A_{\widehat{R}}$ decomposes as*

$$A_{\widehat{R}} \cong \prod_{\mathfrak{m}} A_{\widehat{R}, \mathfrak{m}},$$

where the product runs over the maximal ideals \mathfrak{m} of $A_{\widehat{R}}$.

- (c) *The algebra $A_{\mathbb{F}}$ decomposes as*

$$A_{\mathbb{F}} \cong \prod_{\mathfrak{m}} A_{\mathbb{F}, \mathfrak{m}},$$

where the product runs over the maximal ideals \mathfrak{m} of $A_{\mathbb{F}}$.

- (d) *The algebra A_K decomposes as*

$$A_K \cong \prod_{\mathfrak{p}} A_{K, \mathfrak{p}^e} \cong \prod_{\mathfrak{p}} A_{\widehat{R}, \mathfrak{p}},$$

where the products run over the minimal prime ideals \mathfrak{p} of $A_{\widehat{R}}$ that are contained in a prime ideal of height 1.

Proof: Since $A_{\widehat{R}}$ is a finitely generated \widehat{R} -module, $A_{\widehat{R}}/\mathfrak{p}$ with a prime \mathfrak{p} is an integral domain that is a finitely generated \widehat{R} -module. Hence, it is either a finite field or a finite extension of \widehat{R} . This proves that the height of \mathfrak{p} is less than or equal to 1. The correspondences and the isomorphisms of part (a) are easily verified. The decompositions in parts (b) and (c) hold, since \widehat{R} is Henselian, and hence any finite algebra is the product of its localizations. Part (d) follows by tensoring (b) over \widehat{R} with K . \square

Similar decompositions for A -modules are derived by applying the idempotents of the decompositions of part (b).

Proposition 3.6. *Assume the setup of Proposition 3.5 and let M, N be A -modules that as R -modules are free of finite rank. Suppose that*

- (a) *$M \otimes_R \mathbb{C} \cong N \otimes_R \mathbb{C}$ as $A \otimes_R \mathbb{C}$ -modules, or*
 (b) *$M \otimes_R \bar{K} \cong N \otimes_R \bar{K}$ as $A \otimes_R \bar{K}$ -modules.*

Then for all prime ideals \mathfrak{m} of $A_{\mathbb{F}}$ corresponding to height-1 primes of $A_{\widehat{R}}$, the equality

$$\dim_{\mathbb{F}}(M \otimes_R \mathbb{F})_{\mathfrak{m}} = \dim_{\mathbb{F}}(N \otimes_R \mathbb{F})_{\mathfrak{m}}$$

holds.

Proof: As for A , we also write M_K for $M \otimes_R K$ and similarly for N and \widehat{R} , \mathbb{F} , etc. By choosing an isomorphism $\mathbb{C} \cong \bar{K}$, it suffices to prove part (b). Using Proposition 3.5, part (d), the isomorphism $M \otimes_R \bar{K} \cong N \otimes_R \bar{K}$ can be rewritten as

$$\bigoplus_{\mathfrak{p}} (M_{K, \mathfrak{p}^e} \otimes_K \bar{K}) \cong \bigoplus_{\mathfrak{p}} (N_{K, \mathfrak{p}^e} \otimes_K \bar{K}),$$

where the sums run over the minimal primes \mathfrak{p} of $A_{\widehat{R}}$ that are properly contained in a maximal prime. Hence, an isomorphism $M_{K,\mathfrak{p}^e} \otimes_K \bar{K} \cong N_{K,\mathfrak{p}^e} \otimes_K \bar{K}$ exists for each \mathfrak{p} . Since for each maximal ideal \mathfrak{m} of $A_{\widehat{R}}$ of height 1 we have by Proposition 3.5

$$M_{\widehat{R},\mathfrak{m}} \otimes_{\widehat{R}} K \cong \bigoplus_{\mathfrak{p} \subseteq \mathfrak{m} \text{ min.}} M_{K,\mathfrak{p}^e}$$

and similarly for N , we get

$$\begin{aligned} \dim_{\mathbb{F}} M_{\mathbb{F},\mathfrak{m}} &= \text{rk}_{\widehat{R}} M_{\widehat{R},\mathfrak{m}} = \sum_{\mathfrak{p} \subseteq \mathfrak{m} \text{ min.}} \dim_K M_{K,\mathfrak{p}^e} \\ &= \sum_{\mathfrak{p} \subseteq \mathfrak{m} \text{ min.}} \dim_K N_{K,\mathfrak{p}^e} = \text{rk}_{\widehat{R}} N_{\widehat{R},\mathfrak{m}} \\ &= \dim_{\mathbb{F}} N_{\mathbb{F},\mathfrak{m}}. \end{aligned}$$

This proves the proposition. □

3.2.2 The Stop Criterion. We begin with the following proposition.

Proposition 3.7. *Let \mathfrak{m} be a maximal ideal of $\mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(\mathcal{M}_k(N, \chi; \mathcal{O}))$ that is nontorsion and non-Eisenstein. Then the following statements hold:*

- (a) $\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} \cong \mathcal{M}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}$.
- (b) $2 \cdot \dim_{\mathbb{F}} S_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = \dim_{\mathbb{F}} \mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}$.
- (c) If $p \neq 2$, then

$$\dim_{\mathbb{F}} S_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = \dim_{\mathbb{F}} \mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}^+.$$

Proof: Part (c) follows directly from part (b) by decomposing $\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})$ into a direct sum of its plus and its minus parts. Statements (a) and (b) will be concluded from Proposition 3.6. More precisely, it allows us to derive from Theorem 3.3 that

$$\begin{aligned} \dim_{\mathbb{F}} ((\mathcal{M}_k(N, \chi; \mathcal{O})/\text{torsion}) \otimes_{\mathcal{O}} \mathbb{F})_{\mathfrak{m}} \\ = \dim_{\mathbb{F}} (\text{Eis}_k(N, \bar{\chi}; \mathbb{F}) \oplus S_k(N, \bar{\chi}; \mathbb{F}) \oplus S_k(N, \bar{\chi}; \mathbb{F})^{\vee})_{\mathfrak{m}} \end{aligned}$$

and

$$\begin{aligned} \dim_{\mathbb{F}} ((\mathcal{CM}_k(N, \chi; \mathcal{O})/\text{torsion}) \otimes_{\mathcal{O}} \mathbb{F})_{\mathfrak{m}} \\ = 2 \cdot \dim_{\mathbb{F}} (S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}. \end{aligned}$$

The latter proves part (b), since \mathfrak{m} is nontorsion. Since by the definition of a non-Eisenstein prime, $\text{Eis}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = 0$, and again since \mathfrak{m} is nontorsion, it follows that

$$\dim_{\mathbb{F}} \mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = \dim_{\mathbb{F}} \mathcal{M}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}},$$

which implies part (a). □

We will henceforth often regard non-Eisenstein nontorsion primes as in the proposition as maximal primes of $\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F})) = \mathbb{T}_{\mathcal{O} \rightarrow \mathbb{F}}(S_k(N, \chi))$.

Corollary 3.8. (Stop criterion.) *Let \mathfrak{m} be a maximal ideal of $\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))$ that is non-Eisenstein and nontorsion.*

- (a) One has

$$\dim_{\mathbb{F}} \mathcal{M}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = 2 \cdot \dim_{\mathbb{F}} \mathbb{T}_{\mathbb{F}}(\mathcal{M}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}$$

if and only if

$$\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} \cong \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}.$$

- (b) One has

$$\dim_{\mathbb{F}} \mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}} = 2 \cdot \dim_{\mathbb{F}} \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}$$

if and only if

$$\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} \cong \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}.$$

- (c) Assume $p \neq 2$. One has

$$\dim_{\mathbb{F}} \mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})_{\mathfrak{m}}^+ = \dim_{\mathbb{F}} \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}$$

if and only if

$$\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} \cong \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})^+)_{\mathfrak{m}}.$$

Proof: We prove only (a), since (b) and (c) are similar. From part (b) of Proposition 3.7 and the fact that the \mathbb{F} -dimension of the algebra $\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}$ is equal to that of $S_k(N, \bar{\chi}; \mathbb{F})$, since they are dual to each other, it follows that

$$2 \cdot \dim_{\mathbb{F}} \mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} = \dim_{\mathbb{F}} (\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}}.$$

The result is now a direct consequence of equations (3–1) and (3–2) and Corollary 3.4. □

Note that the first line of each statement uses only modular symbols and not modular forms, but it allows us to make statements involving modular forms. This is the aforementioned stop criterion; the computation of Hecke operators can be stopped if this equality is reached.

We now list some results concerning the validity of the equivalent statements of Corollary 3.8.

Proposition 3.9. *Let $p \geq 5$ be a prime, $k \geq 2$, and $N \geq 5$ with $p \nmid N$ integers, \mathbb{F} a finite extension of \mathbb{F}_p , $\bar{\chi} :$*

$(\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{F}^\times$ a character, and \mathfrak{m} a maximal ideal of $\mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))$ that is non-Eisenstein and nontorsion. Suppose (i) that $2 \leq k \leq p-1$ or (ii) that $k \in \{p, p+1\}$ and \mathfrak{m} is ordinary. Then

$$\begin{aligned} \mathbb{T}_{\mathbb{F}}(S_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} &\cong \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F}))_{\mathfrak{m}} \\ &\cong \mathbb{T}_{\mathbb{F}}(\mathcal{CM}_k(N, \bar{\chi}; \mathbb{F})^+)_{\mathfrak{m}}. \end{aligned}$$

Proof: Using the comparison with group cohomology of [Wiese 08, Theorem 6.1], the result follows under assumption (i) from [Edixhoven 06, Theorem 5.2] and is proved under assumption (ii) in [Wiese 07b, Corollary 6.9] for the case of the group $\Gamma_1(N)$ and no Dirichlet character. The passage to a character is established by [Wiese 07b, Theorem 7.4] and the remark following it. One identifies the mod- p modular forms appearing with corresponding Katz forms using Carayol’s lemma [Edixhoven 97, Proposition 1.10]. \square

We end this section by stating the so-called Sturm bound (also called the Hecke bound), which gives the best a priori upper bound for how many Hecke operators are needed to generate the entire Hecke algebra. We need it in our algorithm only in cases in which it is theoretically not known that the stop criterion will be reached. This will enable the algorithm to detect whether the Hecke algebra on modular symbols is not isomorphic to the corresponding one on cuspidal modular forms.

Proposition 3.10. (Sturm bound.) *The Hecke algebra $\mathbb{T}_{\mathbb{Z}[\chi] \rightarrow \mathbb{F}}(S_k(N, \chi))$ can be generated as an algebra by the Hecke operators T_l for all primes l smaller than or equal to*

$$\frac{kN}{12} \prod_{\substack{q|N \\ q \text{ prime}}} \left(1 + \frac{1}{q}\right).$$

Proof: This follows from [Stein 07, Theorem 9.18]. \square

3.3 Algorithm

In this section we present a sketch of the algorithm that we used for our computations. The MAGMA code, an example, and a manual are published as supplemental material to this article and are available online.¹

Algorithm 3.11.

INPUT: Integers $N \geq 1, k \geq 2$, a finite field \mathbb{F} , a character $\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{F}^\times$, and for each prime l less than or

equal to the Sturm bound an irreducible polynomial $f_l \in \mathbb{F}[X]$.

OUTPUT: An \mathbb{F} -algebra.

- $M \leftarrow \mathcal{CM}_k(N, \chi; \mathbb{F}), l \leftarrow 1, L \leftarrow$ empty list.
- repeat
 - $l \leftarrow$ next prime after l .
 - Compute T_l on M and append it to the list L .
 - $M \leftarrow$ the restriction of M to the f_l -primary subspace for T_l , i.e., to the largest subspace of M on which the minimal polynomial of T_l is a power of f_l .
 - $A \leftarrow$ the \mathbb{F} -algebra generated by the restrictions to M of T_2, T_3, \dots, T_l .
- until $2 \cdot \dim(A) = \dim(M)$ [the stop criterion] or $l >$ Sturm bound.
- return A .

The f_l should, of course, be chosen as the minimal polynomials of the coefficients $a_l(f)$ of the normalized eigenform $f \in S_k(N, \chi; \bar{\mathbb{F}})$ whose local Hecke algebra one wants to compute. Suppose the algorithm stops at the prime q . If q is greater than the Sturm bound, the equivalent conditions of Corollary 3.8 do not hold. In that case, the output should be disregarded. Otherwise, A is isomorphic to a direct product of the form $\prod_{\mathfrak{m}} \mathbb{T}(S_k(N, \chi; \mathbb{F}))_{\mathfrak{m}}$, where the \mathfrak{m} are those maximal ideals such that the minimal polynomials of T_2, T_3, \dots, T_q on $\mathbb{T}(S_k(N, \chi; \mathbb{F}))_{\mathfrak{m}}$ are equal to powers of f_2, f_3, \dots, f_q . It can happen that A consists of more than one factor. Hence, one should still decompose A into its local factors. Alternatively, one can also replace the last line but one in the algorithm by

- until $((2 \cdot \dim(A) = \dim(M))$ and A is local) or $l >$ Sturm bound,

which ensures that the output is a local algebra. In practice, one modifies the algorithm such that a polynomial f_l need not be given for every prime l , but that the algorithm takes each irreducible factor of the minimal polynomial of T_l if no f_l is known. It is also useful to choose the order in which l runs through the primes. For example, one might want to take $l = p$ at an early stage with p the characteristic of \mathbb{F} if one knows that this operator is needed, which is the case in all computations concerning Question 1.9.

¹<http://www.expmath.org/expmath/volumes/17/17.1/Wiese/supplement.zip>.

Level	Wt	ResD	Dim	EmbDim	NilO	GorDef	#Ops	#{ $p < \text{HB}$ }	Gp
5939	5	3	12	3	5	2	5	366	D_7

TABLE 1. Sample entry of important characteristics of a computed local Hecke algebra.

4. COMPUTATIONAL RESULTS

In view of Question 1.9, we produced 384 examples of odd irreducible continuous Galois representations $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\overline{\mathbb{F}}_p)$ that are completely split at p . The results are documented in tables that are published as supplemental material to this article. The complete data, which can be processed by our MAGMA package, are also available online.²

The Galois representations were created either by class field theory or from an irreducible integer polynomial whose Galois group embeds into $\text{GL}_2(\overline{\mathbb{F}}_p)$. All examples but one are dihedral; the remaining one is icosahedral. For each of these an eigenform was computed giving rise to it. The Gorenstein defect of the corresponding local Hecke algebra factor turned out always to be 2, supporting Question 1.9.

The authors preferred to proceed in this way instead of computing all Hecke algebras mod p in weight p for all “small” primes p and all “small” levels, since nondihedral examples in which the assumptions of Question 1.9 are satisfied are very rare.

4.1 Table Entries

For every computed local Hecke algebra, enough data are stored to re-create it as an abstract algebra, and important characteristics are listed in the tables available online. A sample table entry appears as Table 1.

Each entry corresponds to the Galois conjugacy class of an eigenform f mod p with associated local Hecke algebra A . The first and second columns indicate the level and the weight of f . The latter is in all examples equal to the characteristic of the base field k (a finite extension of \mathbb{F}_p) of the algebra. Let \mathfrak{m}_A denote the maximal ideal of A . Then ResD stands for the degree of $K = A/\mathfrak{m}_A$ over \mathbb{F}_p . Let us consider $A \otimes_k K$. It decomposes into a direct product of a local K -algebra B and its $\text{Gal}(K/k)$ -conjugates. The K -dimension of B (which is equal to the k -dimension of A) is recorded in the fourth column.

Let \mathfrak{m}_B be the maximal ideal of B . The *embedding dimension* EmbDim is the K -dimension of $\mathfrak{m}_B/\mathfrak{m}_B^2$. By Nakayama’s lemma, this is the minimal number of B -generators for \mathfrak{m}_B . The *nilpotency order* NilO is the

maximal integer n such that \mathfrak{m}_B^n is not the zero ideal. The column GorDef contains the Gorenstein defect of B (which is the same as the Gorenstein defect of A).

By #Ops it is indicated how many Hecke operators were used to generate the algebra A , applying the stop criterion (Corollary 3.8). This is contrasted with the number of primes smaller than the Sturm bound (Proposition 3.10; it is also called the Hecke bound), denoted by $\#(p < \text{HB})$. One immediately observes that the stop criterion is very efficient. Whereas the Sturm bound is roughly linear in the level, in 365 of the 384 calculated examples, fewer than ten Hecke operators sufficed, in 252 examples even five were enough.

The final column contains the image of the mod- p Galois representation attached to f as an abstract group.

4.2 Dihedral Examples

All Hecke algebras except one in our tables correspond to eigenforms whose Galois representations are dihedral, since these are by far the easiest to obtain explicitly, since one can use class field theory. This is explained now.

Let p be a prime and d a square-free integer that is $1 \pmod 4$ and not divisible by p . We denote by K the quadratic field $\mathbb{Q}(\sqrt{d})$. Further, we consider an unramified character $\chi : \text{Gal}(\overline{\mathbb{Q}}/K) \rightarrow \overline{\mathbb{F}}_p^\times$ of order $n \geq 3$. We assume that its inverse χ^{-1} is equal to χ conjugated by σ , denoted by χ^σ , for σ (a lift of) the nontrivial element of $\text{Gal}(K/\mathbb{Q})$. The induced representation

$$\rho_\chi := \text{Ind}_{\text{Gal}(\overline{\mathbb{Q}}/K)}^{\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})}(\chi) : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\overline{\mathbb{F}}_p)$$

is irreducible, and its image is the dihedral group D_n of order $2n$. If l is a prime not dividing $2d$, we have

$$\rho_\chi(\text{Frob}_l) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

if $\left(\frac{d}{l}\right) = -1$, and

$$\rho_\chi(\text{Frob}_l) = \begin{pmatrix} \chi(\text{Frob}_\Lambda) & 0 \\ 0 & \chi^\sigma(\text{Frob}_\Lambda) \end{pmatrix}$$

if $\left(\frac{d}{l}\right) = 1$ and $l\mathcal{O}_K = \Lambda\sigma(\Lambda)$. This explicit description makes it obvious that the determinant of ρ_χ is the Legendre symbol $l \mapsto \left(\frac{d}{l}\right)$.

Since the kernel of χ corresponds to a subfield of the Hilbert class field of K , simple computations in the class group of K allow one to determine which primes split

²<http://www.expmath.org/expmath/volumes/17/17.1/Wiese/supplement.zip>.

completely. These give examples satisfying the assumptions of Question 1.9 (the Frobenius at p is the identity) if ρ_χ is odd, i.e., if $p = 2$ or $d < 0$.

We remark that for characters χ of odd order n , the assumption $\chi^{-1} = \chi^\sigma$ is not a big restriction, since any character can be written as $\chi = \chi_1\chi_2$ with $\chi_1^\sigma = \chi_1^{-1}$ and $\chi_2^\sigma = \chi_2$; hence the latter descends to a character of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ and the representation ρ_χ is isomorphic to $\rho_{\chi_1} \otimes \chi_2$.

All dihedral representations are known to come from eigenforms in the minimal possible weight with level equal to the (outside of p) conductor of the representation [Wiese 04, Theorem 1].

In the tables, we computed the Hecke algebras of odd dihedral representations as above in the following ranges. For each prime p less than 100 and each prime l less than or equal to the largest level occurring in the table for p , we chose d as plus or minus l such that d is 1 mod 4, and we let H run through all nontrivial cyclic quotients of the class group of $\mathbb{Q}(\sqrt{d})$ of order coprime to p . For each H we chose (unramified) characters χ of the absolute Galois group of $\mathbb{Q}(\sqrt{d})$ corresponding to H , up to Galois conjugacy and up to replacing χ by its inverse. Then χ is not the restriction of a character of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. By genus theory, the order of χ is odd, since the class number is, so we necessarily have $\chi^{-1} = \chi^\sigma$. We computed the local factor of $\mathbb{T}_{\mathbb{F}_p}(S_p(l, \binom{d}{\cdot}; \mathbb{F}_p))$ corresponding to ρ_χ if ρ_χ is odd and p is completely split. For the prime $p = 2$ we also allowed square-free integers d that are 1 mod 4 and whose absolute value is less than 5000.

4.3 Icosahedral Example

With the help of a list of polynomials provided by Gunter Malle [Malle 06], a Galois representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ with values in $\text{GL}_2(\overline{\mathbb{F}}_2)$ that is of prime conductor, completely split at 2, and thus satisfies the assumptions of Question 1.9 and whose image is isomorphic to the icosahedral group A_5 could be described explicitly. The modular forms in weight 2 predicted by Serre's conjecture were found, and the corresponding Hecke algebra turned out to have Gorenstein defect equal to 2.

Let $f \in \mathbb{Z}[X]$ be an irreducible polynomial of degree 5 whose Galois group, i.e., the Galois group of the normal closure L of $K = \mathbb{Q}[X]/(f)$, is isomorphic to A_5 . We assume that K is unramified at 2, 3, and 5. We have the Galois representation

$$\rho_f : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \twoheadrightarrow \text{Gal}(L/\mathbb{Q}) \cong A_5 \cong \text{SL}_2(\mathbb{F}_4).$$

We now determine its conductor and its traces. Let p be a ramified prime. Since the ramification is tame, the image

of the inertia group $\rho_f(I_p)$ at p is cyclic of order 2, 3, or 5. In the first case, the image of a decomposition group $\rho_f(D_p)$ at p is either equal to $\rho_f(I_p)$ or equal to $\mathbb{Z}/2\mathbb{Z} \times \rho_f(I_p)$. If the order of $\rho_f(I_p)$ is odd and $\rho_f(I_p) = \rho_f(D_p)$, then any completion of L at the unique prime above p is totally ramified and cyclic of degree $\#\rho_f(I_p)$, hence contained in $\mathbb{Q}_p(\zeta_p)$ for ζ_p a primitive p th root of unity. It follows that p is congruent to 1 mod $\#\rho_f(I_p)$. If the order of $\rho_f(I_p)$ is odd, but $\rho_f(I_p)$ is not equal to $\rho_f(D_p)$, then $\rho_f(D_p)$ is a dihedral group and the completion of L at a prime above p has a unique unramified quadratic subfield S . Thus, we have the exact sequence

$$0 \rightarrow \rho_f(I_p) \rightarrow \rho_f(D_p) \rightarrow \text{Gal}(S/\mathbb{Q}_p) \rightarrow 0.$$

On the one hand, it is well known that the conjugation by a lift of the Frobenius element of $\text{Gal}(S/\mathbb{Q}_p)$ acts on $\rho_f(I_p)$ by raising to the p th power. On the other hand, since the action is nontrivial, it also corresponds to inversion on $\rho_f(I_p)$, since the only elements of order 2 in $(\mathbb{Z}/3\mathbb{Z})^\times$ and $(\mathbb{Z}/5\mathbb{Z})^\times$ are -1 . As a consequence, p is congruent to -1 mod $\#\rho_f(I_p)$ in this case.

We hence have the following cases:

- (1) Suppose $p\mathcal{O}_K = \mathfrak{P}^5$. Then $p \equiv \pm 1 \pmod{5}$.
 - (a) If $p \equiv 1 \pmod{5}$, then $\rho_f|_{I_p} \sim \begin{pmatrix} \chi & 0 \\ 0 & \chi^{-1} \end{pmatrix}$ with χ a totally ramified character of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ of order 5.
 - (b) If $p \equiv -1 \pmod{5}$, then $\rho_f(D_p)$ is the dihedral group with 10 elements.
- (2) Suppose $p\mathcal{O}_K = \mathfrak{P}^3\Omega\mathfrak{A}$ or $p\mathcal{O}_K = \mathfrak{P}^3\Omega$.
 - (a) If $p \equiv 1 \pmod{3}$, then $\rho_f|_{I_p} \sim \begin{pmatrix} \chi & 0 \\ 0 & \chi^{-1} \end{pmatrix}$ with χ a totally ramified character of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ of order 3.
 - (b) If $p \equiv -1 \pmod{3}$, then $\rho_f(D_p)$ is the dihedral group with 6 elements.
- (3) Suppose that p is ramified, but that we are neither in case (1) nor in case (2). Then $\rho_f|_{I_p} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

By the definition of the conductor at p it is clear that it is p^2 in cases (1) and (2) and p in case (3). However, in cases (1)(a) and (2)(a) one can choose a character ϵ of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ of the same order as χ whose restriction to D_p gives the character χ . If one twists the representation ρ_f by ϵ , one finds also in these cases that the conductor at p is p .

Level	Wt	ResD	Dim	EmbDim	NilO	GorDef	#Ops	#($p < \text{HB}$)	G _p
89491	2	2	12	4	3	2	4	1746	A ₅

TABLE 2. Table entry for the icosahedral example.

Dimension	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	36	40	46	56	60
Number of algebras	206	58	25	3	24	6	20	3	12	3	5	4	2	1	2	2	4	1	2	1

TABLE 3. The number of times that each dimension appears.

An inspection of the conjugacy classes of the group $\text{SL}_2(\mathbb{F}_4)$ shows that the traces of ρ_f twisted by some character ϵ of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ are as follows. Let l be an unramified prime.

- If the order of Frob_l is 5, then the trace at Frob_l is $\epsilon(\text{Frob}_l)w$, where w is a root of the polynomial $X^2 + X + 1$ in $\mathbb{F}_2[X]$.
- If the order of Frob_l is 3, then the trace at Frob_l is $\epsilon(\text{Frob}_l)$.
- If the order of Frob_l is 1 or 2, then the trace at Frob_l is 0.

These statements allow the easy identification of the modular form belonging to an icosahedral representation.

We end this section with some remarks on our icosahedral example. It was obtained using the polynomial $x^5 - x^4 - 79x^3 + 225x^2 + 998x - 3272$. The corresponding table entry is shown as Table 2.

Hence, in level 89,491 and weight 2 there is a single eigenform g mod 2 up to Galois conjugacy whose first couple of q -coefficients agree with the traces of a twist of the given icosahedral Galois representation. From this one can deduce that the Galois representation ρ_g of g has an icosahedral image and is ramified only at 89,491. Since weight-lowering is not known in our case, we cannot prove that ρ_g coincides with a twist of the given one. It might, however, be possible to exclude the existence of two distinct icosahedral extensions of the rationals inside \mathbb{C} that ramify only at 89,491 by consulting tables. According to Malle, the icosahedral extension used has smallest discriminant among all totally real A_5 -extensions of the rationals in which 2 splits completely.

5. FURTHER RESULTS AND QUESTIONS

In this section we present some more computational observations for Hecke algebras under the assumptions of Question 1.9, which lead us to ask some more questions.

5.1 On the Dimension of the Hecke Algebra

From the data, we see that many even integers appear as dimensions of the \mathbb{T}_m . We know that the dimension must be at least 4, since this is the dimension of the smallest non-Gorenstein algebra that can appear in our case. This extends the results of [Kilford 02], where the dimensions of the Hecke algebras $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_2}(S_2(\Gamma_0(431)))$ and $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_2}(S_2(\Gamma_0(503)))$ localized at the non-Gorenstein maximal ideals are shown to be 4.

In Table 3 we see exactly how many times each dimension appears in our data. We observe that every even integer between 4 and 32 appears, and that the largest dimension is 60. The most common dimension is 4, which appears about half the time. However, since the dimension of the Hecke algebra attached to $S_k(\Gamma_1(N))$ increases with N and with k , this may be an artifact of the data being collected for “small” levels N and primes p .

It seems reasonable that there should be infinitely many cases with dimension 4, and plausible that every even integer greater than or equal to 4 should appear as a dimension infinitely many times. From the tables, we see that algebras of dimension 4 appear at very high levels, so they do not appear to be becoming rare as the dimension increases, but this may, of course, be an artifact of our data.

We note that not every example that arises from an elliptic curve in characteristic $p = 2$ has Hecke algebra with dimension 4; for example, the algebra $\mathbb{T}_{\mathbb{Z} \rightarrow \mathbb{F}_2}(S_2(\Gamma_0(2089)))$ localized at its non-Gorenstein maximal ideal has dimension 18. In level 18,097 there is a dimension-36 example arising from an elliptic curve.

5.2 On the Residue Degree

We will now solve an easy aspect of the question of the possible structures of non-Gorenstein local algebras occurring as local Hecke algebras. We assume for the couple of lines to follow the generalized Riemann hypothesis (GRH).

We claim that then the residue degrees of \mathbb{T}_m (in the notation of Question 1.9) are unbounded if we let p and

N run through the primes such that $p \neq N$ and N is congruent to 3 modulo 4.

This is so because class groups of imaginary quadratic fields $\mathbb{Q}(\sqrt{-N})$ have arbitrarily large cyclic factors of odd order, since the exponent of these class groups is known to go to infinity as N does, by the main result of [Boyd and Kisilevsky 72], which assumes GRH. So the discussion on dihedral forms in Section 4 immediately implies the claim.

5.3 On the Embedding Dimension

One can ask whether the embedding dimension of the local Hecke algebras in the situation of Question 1.9 is bounded if we allow p and N to vary. This, however, seems to be a difficult problem. The embedding dimensions occurring in our tables are 3 (299 times), 4 (78 times) and 5 (7 times).

The embedding dimension d is related to the number of Hecke operators needed to generate the local Hecke algebra, in the sense that at least d Hecke operators are needed. Probably, d Hecke operators can be found that do generate, but they need not be the first d prime Hecke operators, of course. However, as our tables suggest, in most cases the actual computations were done using very few operators, and there are 99 of the 384 cases in which the computation finished after only d operators.

ACKNOWLEDGMENTS

The authors would like to thank Kevin Buzzard for informing them about the level-23 and weight-59 example, for many helpful conversations and communications, and for his comments on a first draft of this paper. They would like to thank Amod Agashe for comments on an early version and William Stein for the use of his computers MECCA and NERON for computations; without these, we would not have been able to pursue these computations as far as we did. The authors are grateful to Gunter Malle for providing them with a very useful list of A_5 -polynomials. The first author would also like to thank Edray Goins for helpful conversations. The second author is indebted to Bas Edixhoven for very enlightening explanations.

The authors would also like to thank the referees for suggesting improvements to the paper.

REFERENCES

[Agashe et al. 06] Amod Agashe, Kenneth A. Ribet, and William A. Stein. “The Modular Degree, Congruence Primes and Multiplicity One.” Available online (<http://modular.math.washington.edu/papers/ars-congruence/>), 2006.

[Bosma et al. 1997] W. Bosma, J. Cannon, and C. Playoust. “The Magma Algebra System I: The User Language.” *J. Symb. Comp.* 24:3–4 (1997) 235–265.

[Boston et al. 91] Nigel Boston, Hendrik W. Lenstra, Jr., and Kenneth A. Ribet. “Quotients of Group Rings Arising from Two-Dimensional Representations.” *C. R. Acad. Sci. Paris Sér. I Math.* 312:4 (1991), 323–328.

[Boyd and Kisilevsky 72] David W. Boyd and H. Kisilevsky. “On the Exponent of the Ideal Class Groups of Complex Quadratic Fields.” *Proc. Amer. Math. Soc.* 31 (1972), 433–436.

[Buzzard 01] K. Buzzard. “Appendix.” In *Arithmetic Algebraic Geometry (Park City, UT, 1999)*, IAS/Park City Math. Ser. 9. Providence, Amer. Math. Soc., 2001.

[Diamond and Im 95] Fred Diamond and John Im. “Modular Forms and Modular Curves.” In *Seminar on Fermat’s Last Theorem (Toronto, ON, 1993–1994)*, pp. 39–133, CMS Conf. Proc. 17. Providence, Amer. Math. Soc., 1995.

[Edixhoven 92] Bas Edixhoven. “The Weight in Serre’s Conjectures on Modular Forms.” *Invent. Math.* 109:3 (1992), 563–594.

[Edixhoven 97] Bas Edixhoven. “Serre’s Conjecture,” In *Modular Forms and Fermat’s Last Theorem (Boston, MA, 1995)*, pp. 209–242. New York: Springer, 1997.

[Edixhoven 06] Bas Edixhoven. “Comparison of Integral Structures on Spaces of Modular Forms of Weight Two, and Computation of Spaces of Forms mod 2 of Weight One.” *J. Inst. Math. Jussieu* 5:1 (2006), 1–34.

[Eisenbud 95] David Eisenbud. *Commutative Algebra*, Graduate Texts in Mathematics, 150. New York: Springer-Verlag, 1995.

[Emerton 02] Matthew Emerton. “Supersingular Elliptic Curves, Theta Series and Weight Two Modular Forms.” *J. Amer. Math. Soc.* 15:3 (2002), 671–714.

[Gross 90] B. Gross. “A Tameness Criterion for Galois Representations Associated to Modular Forms (mod p).” *Duke Math. Journal* 61:2 (1990), 445–517.

[Hida 81] H. Hida. “On Congruence Divisors of Cusp Forms as Factors of the Special Values of Their Zeta Functions.” *Invent. Math.* 64 (1981), 221–262.

[Kilford 02] L. J. P. Kilford. “Some Non-Gorenstein Hecke Algebras Attached to Spaces of Modular Forms,” *J. Number Theory* 97:1 (2002), 157–164.

[Malle 06] G. Malle. Personal communication, 2006.

[Mazur 77] B. Mazur. “Modular Curves and the Eisenstein Ideal.” *Inst. Hautes Études Sci. Publ. Math.* 47, (1977), 33–186.

[Mazur and Ribet 92] B. Mazur and K. A. Ribet. “Two-Dimensional Representations in the Arithmetic of Modular Curves.” *Astérisque* 196–197 (1992), 215–255.

[Ribet 90] Kenneth A. Ribet. “Multiplicities of Galois Representations in Jacobians of Shimura Curves.” In *Festschrift in Honor of I. I. Piatetski-Shapiro on the Occasion of his Sixtieth Birthday, Part II (Ramat Aviv, 1989)*, pp. 221–236, Israel Math. Conf. Proc. 3. Jerusalem: Weizmann.

- [Ribet and Stein 01] Kenneth A. Ribet and William A. Stein. “Lectures on Serre’s Conjectures.” In *Arithmetic Algebraic Geometry (Park City, UT, 1999)*, pp. 143–232, IAS/Park City Math. Ser. 9. Providence: Amer. Math. Soc., 2001.
- [Stein 07] William Stein. *Modular Forms: A Computational Approach*, Graduate Studies in Mathematics, 79. Providence, American Mathematical Society, 2007.
- [Tilouine 87] Jacques Tilouine. “Un sous-groupe p -divisible de la jacobienne de $X_1(Np^r)$ comme module sur l’algèbre de Hecke.” *Bull. Soc. Math. France* 115:3 (1987), 329–360.
- [Tilouine 97] Jacques Tilouine. “Hecke Algebras and the Gorenstein Property.” In *Modular Forms and Fermat’s Last Theorem (Boston, MA, 1995)*, pp. 327–342. New York: Springer, 1997.
- [Wiese 04] Gabor Wiese. “Dihedral Galois Representations and Katz Modular Forms.” *Doc. Math.* 9 (2004), 123–133.
- [Wiese 07a] Gabor Wiese. “Multiplicities of Galois Representations of Weight One.” *Algebra & Number Theory* 1:1 (2007), 67–85.
- [Wiese 07b] Gabor Wiese. “On the Faithfulness of Parabolic Cohomology as a Hecke Module over a Finite Field.” *J. Reine Angew. Math.* 606 (2007), 79–103.
- [Wiese 08] Gabor Wiese. “On Modular Symbols and the Cohomology of Hecke Triangle Surfaces.” To appear in *International J. of Number Theory*, 2008.
- [Wiles 95] Andrew Wiles. “Modular Elliptic Curves and Fermat’s Last Theorem.” *Ann. of Math. (2)* 141:3 (1995), 443–551.

L. J. P. Kilford, Department of Mathematics, University of Bristol, Bristol BS8 1TW, United Kingdom (l.kilford@gmail.com)

Gabor Wiese, Institut für Experimentelle Mathematik, Universität Duisburg-Essen, Ellernstraße 29, 45326 Essen, Germany (gabor.wiese@uni-due.de)

Received January 29, 2007; accepted October 2, 2007.