

BOUNDEDNESS AND CONVERGENCE OF SOLUTIONS OF DUFFING'S EQUATION

Dedicated to Professor Ryoji Shizuma on his 60-th birthday

KENICHI SHIRAIWA

In this paper, we shall discuss boundedness of solutions of the equation

$$(1) \quad x'' + f(x)x' + g(x) = e(t) \quad (' = d/dt)$$

under suitable conditions. And we shall discuss asymptotic stability of a periodic solution and convergence of solutions for the equation

$$(2) \quad x'' + cx' + g(x) = e(t)$$

for a positive constant c and a periodic function $e(t)$ under some restricted conditions.

This work is motivated by the work of H. Kawakami [3], which gives some numerical computations on the equation $x'' + kx' + x^3 = B \cos t$ for positive constants k and B . Kawakami's work contains many interesting informations on Duffing's equation of the above type, and this paper deals with only small part of his work. There is also very interesting experimental results by C. Hayashi, Y. Ueda and H. Kawakami [2]. Also, our paper heavily depends on the work of W. S. Loud [4].

§1. The boundedness theorem

The equation (2) is a special case of (1), and the equation (1) is equivalent to the following system of equations.

$$(3) \quad \begin{cases} x' = y \\ y' = -f(x)y - g(x) + e(t) \end{cases}$$

THEOREM 1. *In the equation (1) we assume the following conditions*

Received June 1, 1976.

(a), (b) and (c).

(a) *There exists a solution of (1) under any initial condition.*

(b) *There exist positive constants c and E such that*

$$f(x) \geq c \quad \text{and} \quad |e(t)| \leq E .$$

(c) *$g(x)$ is a differentiable function satisfying the following conditions (i), (ii) and (iii).*

(i) *$g'(x)$ is bounded on any finite interval.*

(ii) *$g'(x) \geq 0$.*

(iii) *$\lim_{x \rightarrow \infty} g(x) > E$ and $\lim_{x \rightarrow -\infty} g(x) < -E$.*

By the condition (c), $g(x)$ is a monotone increasing function, and there exist numbers x_1 and x_2 ($x_1 < x_2$) such that

$$g(x_1) = -E \quad \text{and} \quad g(x_2) = E .$$

Let $x(t)$ be any solution of (1). Then there exists a number t_0 such that

$$x_1 - 4E/c^2 \leq x(t) \leq x_2 + 4E/c^2$$

and

$$|x'(t)| \leq 4E/c \quad \text{for any } t \geq t_0 .$$

Our proof is similar to that of Theorem 1 of W. S. Loud [4]. He assumed that $g'(x) \geq b$ for some positive constant b in his paper and got an additional information. But the essential part of his argument is applicable in our case, too.

Proof of Theorem 1. From the equation (5) we obtain the following equation.

$$(4) \quad y \frac{dy}{dx} = -f(x)y - g(x) + e(t)$$

We consider the following equations.

$$(5) \quad y \frac{dy}{dx} = -cy - g(x) + E \quad (y \geq 0)$$

$$(6) \quad y \frac{dy}{dx} = -cy - g(x) - E \quad (y \leq 0)$$

By our assumption (c) (i) the existence and uniqueness of solutions of (5) and (6) are guaranteed.

Also, the following inequalities hold by our assumption (b).

$$(7) \quad \frac{dy}{dx} \text{ in (4)} \leq \frac{dy}{dx} \text{ in (5)} \quad \text{for } y > 0 .$$

$$(8) \quad \frac{dy}{dx} \text{ in (4)} \leq \frac{dy}{dx} \text{ in (6)} \quad \text{for } y < 0 .$$

LEMMA 1. *Any solution of (5) has the following properties in the upper half plane.*

(d) *It has a positive slope to the left of the curve $-cy - g(x) + E = 0$, a negative slope to the right of this curve, and reaches a maximum of y when crossing the curve.*

(e) *Any solution starting with $x < x_2$ must cross the line $x = x_2$ for positive value.*

(f) *To the right of the line $x = x_2$ the solution is concave downward. It leaves the upper half plane as t increases with $x > x_2$.*

A similar statement holds for the equation (6).

Our proof is similar to that of Loud [4] and is omitted.

Let y_1 be a positive number. Let C_1 be the solution of the equation (5) starting from (x_2, y_1) to its next intersection with the x -axis at $(x_3, 0)$, and let C_2 be the solution of the equation (6) starting from $(x_3, 0)$ to its next intersection with the line $x = x_1$ at $(x_1, -y)$. The existence of all the intersections mentioned above is guaranteed by Lemma 1.

LEMMA 2. *Let C be the arc consisting of C_1 and C_2 . If $y_1 \geq 4E/c$, then $|y| < y_1$ at any point of C other than (x_2, y_1) .*

Our proof is a slight modification of that of Loud [4] and is omitted.

Let B_1 be the solution of (5) starting from $(x_2, 4E/c)$ to its next intersection with the x -axis at $(a_1, 0)$. Let B_2 be the solution of (6) starting from $(a_1, 0)$ to its next intersection with the line $x = x_1$ at (x_1, b_1) . Then $b_1 > -4E/c$ by Lemma 2. Let B_3 be the line segment of $x = x_1$ from (x_1, b_1) to $(x_1, -4E/c)$.

Let B_4 be the solution of (6) starting from $(x_1, -4E/c)$ to its next intersection with the x -axis at $(a_2, 0)$. Let B_5 be the solution of (5) starting from $(a_2, 0)$ to its next intersection with the line $x = x_2$ at (x_2, b_2) .

Then $b_2 < 4E/c$ by the latter half of Lemma 2. Let B_6 be the line segment of $x = x_2$, from (x_2, b) to $(x_2, 4E/c)$.

Then the curve B consisting of these six curves B_i ($1 \leq i \leq 6$) surrounds the closed region D , which is homeomorphic to a disk. (Cf. Fig. 1)




Fig. 1

Using the inequalities (7), (8) and the similar argument in the proof of Theorem 1 of Loud [4], we can prove that any solution of (4) starting outside D enters into D and never leaves D afterward.

By Lemma 2 $|y| \leq 4E/c$ for any point of D . And at $(x_2, 4E/c)$, $\frac{dy}{dx} = -c$ by (5), and the boundary of D is concave downward by Lemma 1(f). Therefore, it is easy to see that D is contained in the rectangle $[x_1 - 4E/c^2, x_2 + 4E/c^2] \times [-4E/c, 4E/c]$. This completes the proof of Theorem 1.

COROLLARY. *In addition to the conditions (a), (b) and (c) of Theorem 1, we assume the following two conditions (g) and (h).*

(g) $f(x)$ and $e(t)$ are continuous, and $f(x)$ satisfies the local Lipschitz condition.

(h) $e(t)$ is periodic of period τ ($\tau > 0$).

Then the equation (1) has a periodic solution of period τ .

Proof. By the assumption (g), there exists a unique solution $x = \psi_1(t; x_0, y_0)$, $y = \psi_2(t; x_0, y_0)$ of (3) satisfying the condition

$$(\psi_1(0; x_0, y_0), \psi_2(0; x_0, y_0)) = (x_0, y_0)$$

for each $(x_0, y_0) \in \mathbb{R}^2$. Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a map defined by

$$(9) \quad T(x_0, y_0) = (\psi_1(\tau; x_0, y_0), \psi_2(\tau; x_0, y_0)).$$

We call this map T the *Poincaré transformation* associated with the periodic system (1) (or (3)).

By the assumption (h) and the uniqueness of the solution of (3), it is easy to verify the following equality for each t .

$$(10) \quad (\psi_1(t + \tau; x_0, y_0), \psi_2(t + \tau; x_0, y_0)) = (\psi_1(t; T(x_0, y_0)), \psi_2(t; T(x_0, y_0)))$$

Therefore, each fixed point of T defines a periodic solution of period τ of (1) and vice versa.

By the proof of Theorem 1 T maps the closed region D into D , and D is homeomorphic to a disk. Therefore, there exists a fixed point of T by the Brouwer's fixed point theorem. This completes the proof of this corollary.

§2. The convergence theorem

In this section we shall prove a convergence theorem for the equations of type (2) with periodic $e(t)$. The equation (2) is equivalent to the following equation.

$$(11) \quad \begin{cases} x' = y \\ y' = -cy - g(x) + e(t) \end{cases}$$

Throughout this section we assume the following conditions.

A (i) $e(t)$ is a continuous periodic function of period τ ($\tau > 0$), and E is a positive constant such that $|e(t)| \leq E$. The existence of such E is guaranteed by our assumption.

A (ii) $g(x)$ is a differentiable function of class C^1 such that $g'(x) \geq 0$,

$\lim_{x \rightarrow \infty} g(x) > E$, $\lim_{x \rightarrow -\infty} g(x) < -E$, and $g'(x) = 0$ only on a countable subset of the real numbers.

A (iii) c is a positive constant.

By A (ii) the following property is clear.

PROPERTY 1. $g(x)$ is strictly monotone increasing, and there exist numbers x_1 and x_2 ($x_1 < x_2$) such that

$$g(x_1) = -E \quad \text{and} \quad g(x_2) = E .$$

Under the assumptions A (i), A (ii) and A (iii) we get the following properties by Theorem 1 and its Corollary.

PROPERTY 2. There exists at least one periodic solution of period τ for the equation (11).

PROPERTY 3. Any periodic solution $x = \psi(t)$, $y = \psi_2(t)$ of the equation (11) satisfies the following inequalities for all t .

$$\begin{aligned} x_1 - 4E/c^2 &\leq \psi_1(t) \leq x_2 + 4E/c^2 , \\ |\psi_2(t)| &\leq 4E/c . \end{aligned}$$

Set $H(\alpha) = \sup \{g'(x); -\alpha \leq x \leq \alpha\}$ for any positive number α . By our assumption A (ii), $H(\alpha)$ is finite.

THEOREM 2. Assume the conditions A (i), A (ii) and A (iii). Let n be a positive integer and let $x = \psi_1(t)$, $y = \psi_2(t)$ be a non-constant periodic solution of period $n\tau$ for the equation (11).

Suppose that $|\psi_1(t)| \leq \beta$ for all t and $c^2 > H(\beta)$. Then the periodic solution $x = \psi_1(t)$, $y = \psi_2(t)$ is asymptotically stable.

This is a generalization of Theorem 2 of Loud [4].

Proof. The variation equation corresponding to the periodic solution $x = \psi_1(t)$, $y = \psi_2(t)$ is given by

$$(12) \quad \begin{cases} x' = y \\ y' = -cy - g'(\psi_1(t))x . \end{cases}$$

By our assumption, the variation equation (12) is a linear differential equation with continuous periodic coefficients. Therefore, to prove that $x = \psi_1(t)$, $y = \psi_2(t)$ is asymptotically stable, it is sufficient to show that all characteristic exponents of the equation (12) have negative real parts

(Cf. Chap. 13, Th. 1.4 Coddington-Levinson [1]).

Now, the following Lemma 3 is an easy consequence of the Floquet theory and the theory of linear differential equations with constant coefficients.

LEMMA 3. *Let $x' = P(t)x$ be a linear differential equation, where x is an n -dimensional vector and $P(t)$ is a continuous periodic $n \times n$ matrix function with period σ ($\sigma > 0$). Then the following three conditions are equivalent.*

- (a) *The identically zero solution is asymptotically stable.*
- (b) *All characteristic exponents have negative real parts.*
- (c) *Any solution approaches to the origin as t tends to infinity.*

By Lemma 3, it is sufficient to prove the following Lemma 4 for the completion of Theorem 2.

LEMMA 4. *Under the assumption of Theorem 2, any solution $(x(t), y(t))$ of (12) tends to the origin as t tends to infinity.*

Proof of Lemma 4. From the assumption $|\psi_1(t)| \leq \beta$ and the definition of $H(\beta)$, we have

$$(13) \quad 0 \leq g'(\psi_1(t)) \leq H(\beta).$$

Since $\psi_1(t)$ is assumed to be non-constant and $g'(x) = 0$ on a countable subset of \mathbf{R} , there exists some t_1 such that $g'(\psi_1(t_1)) \neq 0$. Then we have the following Property 4 by the periodicity and continuity of $g'(\psi_1(t))$.

PROPERTY 4. *There exist positive constants δ and γ satisfying the following condition (B).*

- (B) *For any number t_2 there exists a number t_3 ($t_3 > t_2$) such that*

$$g'(\psi_1(t)) \geq \delta \quad \text{for } t_3 \leq t \leq t_3 + \gamma.$$

Let y_1 be any positive number. Now, we shall construct a bounded closed region $D(y_1)$ as follows.

Let L_1 be the line segment of $y = -cx + y_1$ from $P_1(0, y_1)$ to $P_2(y_1/c, 0)$. Let L_2 be the line segment of $x = y_1/c$ from $P_2(y_1/c, 0)$ to $P_3(y_1/c, -H(\beta)y_1/c^2)$. Let L_3 be the line segment of $y = -H(\beta)y_1/c^2$ from $P_3(y_1/c, -H(\beta)y_1/c^2)$ to $P_4(0, -H(\beta)y_1/c^2)$.

Let L_4 be the line segment of $y = -cx - H(\beta)y_1/c^2$ from

$P_4(0, -H(\beta)y_1/c^2)$ to $P_5(-H(\beta)y_1/c^3, 0)$. Let L_5 be the line segment of $x = -H(\beta)y_1/c^3$ from $P_5(-H(\beta)y_1/c^3, 0)$ to $P_6(-H(\beta)y_1/c^3, H(\beta)^2y_1/c^4)$. Let L_6 be the line segment of $y = H(\beta)^2y_1/c^4$ from $P_6 = (-H(\beta)y_1/c^3, H(\beta)^2y_1/c^4)$ to $P_7(0, H(\beta)^2y_1/c^4)$.

Since $c^2 > H(\beta)$ by our assumption, P_7 is below P_1 on the y -axis. Let L_7 be the line segment of $x = 0$ from $P_7(0, H(\beta)^2y_1/c^4)$ to $P_1(0, y_1)$.

Then the polygonal curve L consisting of L_i ($1 \leq i \leq 7$) is a closed curve, and the closed region $D(y_1)$ is defined to be the region bounded by L . (Fig. 2)




Fig. 2

Using the inequality (13), it is easy to verify that the vector field defined by the equation (12) always points either inside of $D(y_1)$ or tangent to its boundary L on any point of L . Therefore, any solution (12)

starting from the point of $D(y_1)$ at any time t_0 stays on $D(y_1)$ for any $t \geq t_0$. Thus, the origin $O(0, 0)$ is a stable singularity of the equation (12).

Now, we shall prove the following properties.

PROPERTY 5. *Any solution of (12) starting from a point of the triangle OP_1P_2 except the line segment OP_2 reaches to the positive half of the x -axis in a finite time interval.*

PROPERTY 6. *Any solution of (12) starting from a point of the rectangle $OP_2P_3P_4$ either reaches to the negative half of the y -axis in a finite time interval or stays on the rectangle $OP_2P_3P_4$ and tends to the origin as t tends to infinity.*

From Properties 5, 6 and the similar properties for the solutions of (12) starting from the points of the triangle OP_4P_5 and the rectangle $OP_5P_6P_7$, we can easily prove Lemma 3, and this finishes the proof of Theorem 2.

Proof of Property 5. It is easy to see that any solution $x = x(t)$, $y = y(t)$ of (12) starting from a point of the triangle OP_1P_2 except the line segment OP_2 can leave the triangle OP_1P_2 only from the line segment OP_2 . Since $x'(t) = y(t) > 0$ as far as $(x(t), y(t))$ stays on the triangle OP_1P_2 except the line segment OP_2 , we can assume, without loss of generality, that there exists a positive number x_0 such that $x(t) \geq x_0 > 0$ as far as $(x(t), y(t))$ stays on the triangle OP_1P_2 .

Suppose that the solution $(x(t), y(t))$ stays on the triangle OP_1P_2 for any $t \geq t_1$. Then using the condition (B) and (12), we have $y'(t) \leq -\delta x_0 < 0$ for infinitely many time intervals of length at least γ . But this contradicts the boundedness of $y(t)$ which follows immediately from our assumption that $(x(t), y(t))$ stays on the triangle OP_1P_2 for $t \geq t_1$. This completes the proof of Property 5.

Proof of Property 6. It is easy to verify that any solution $(x(t), y(t))$ of (12) starting from a point of the rectangle $OP_2P_3P_4$ can leave the rectangle $OP_2P_3P_4$ only from the side OP_4 . Therefore, any solution of (12) starting from a point of the rectangle $OP_2P_3P_4$ leaves the fourth quadrant if and only if it leaves the rectangle $OP_2P_3P_4$.

Suppose that $(x(t_0), y(t_0))$ belongs to the rectangle $OP_2P_3P_4$ but not

the origin. Let K be the supremum of the numbers t_1 such that for $t_0 \leq t \leq t_1$ the solution $(x(t), y(t))$ stays on the rectangle $OP_2P_3P_4$.

Case 1. Let K be finite. Then it is clear that $(x(K), y(K))$ belongs to the rectangle $OP_2P_3O_4$. Now, we shall prove that $x(K) = 0$ and $y(K) < 0$. Thus, the solution $(x(t), y(t))$ reaches the negative half of the y -axis in a finite time interval and gets out of the rectangle $OP_2P_3P_4$ there.

By the definition of K and the equation (12), it is easy to see that $x(t)$ is monotone decreasing for $t_0 \leq t \leq K$, $x(K) \geq 0$, and $y(K) \leq 0$.

Suppose that $x(K) > 0$. If $y(K) < 0$, then $x'(K) = y(K) < 0$. And it is easy to see that the solution $(x(t), y(t))$ stays on the fourth quadrant for $K \leq t \leq K + k_0$ for a suitable positive k_0 . This contradicts the definition of K . Therefore, $y(K) = 0$.

If $y(K) = 0$, then $x'(K) = y(K) = 0$ and $y'(K) = -g'(\psi_1(t))x(K) \leq 0$ by our assumption. Using (13) and Property 4, we easily see that the solution $(x(t), y(t))$ stays on the fourth quadrant for $K \leq t \leq K + k_1$ for a suitable positive k_1 . But this contradicts the definition of K and establishes that $x(K) = 0$.

Next, if $y(K) = 0$, then $(x(K), y(K)) = (0, 0)$. Now, by the uniqueness of the solution of (12) under the same initial condition, $(x(t), y(t)) = (0, 0)$ for all t . This contradicts our assumption that $(x(t_0), y(t_0)) \neq (0, 0)$. Therefore, $y(K) < 0$.

Case 2. Let $K = \infty$. Then the solution $(x(t), y(t))$ stays on the rectangle $OP_2P_3P_4$ for all $t \geq t_0$, and $x(t)$ is monotone decreasing for $t \geq t_0$ since $x'(t) = y(t) \leq 0$.

In this case, we shall show that $\lim_{t \rightarrow \infty} x(t) = \lim_{t \rightarrow \infty} y(t) = 0$,

Since $x(t)$ is monotone decreasing and $x(t) \geq 0$ for $t \geq t_0$, $x(t)$ converges to $x_0 \geq 0$ as t tends to infinity.

Suppose that $x_0 > 0$. Since $(x(t), y(t))$ stays on the rectangle $OP_2P_3P_4$ for $t \geq t_0$, there exists a sequence $\{t_k\}$ such that

$$\lim_{k \rightarrow \infty} t_k = \infty \quad \text{and} \quad \lim_{k \rightarrow \infty} y(t_k) = y_0 \quad \text{for some } y_0 \leq 0.$$

Then $\lim_{k \rightarrow \infty} x(t_k) = x_0$ since $\lim_{t \rightarrow \infty} x(t) = x_0$. Thus, the following property is proved.

PROPERTY 7. $\lim_{k \rightarrow \infty} t_k = \infty, \lim_{k \rightarrow \infty} x(t_k) = x_0 \geq 0, x(t)$ is monotone decreasing for $t \geq t_0, x(t) \geq x_0$ for $t \geq t_0$ and $\lim_{k \rightarrow \infty} y(t_k) = y_0 \leq 0$.

And we are now assuming that $x_0 > 0$. Further, we assume that $y_0 < 0$. Then there exists a positive number ℓ such that $y_0/x_0 < -\ell$.

Let ε be a positive number, and we put $Q_1 = (x_0 + \varepsilon, y_0), Q_2 = (x_0, y_0 + c\varepsilon)$ and $Q_3 = (x_0, y_0 + (-c + H(\beta)/\ell)\varepsilon)$. Then the slope of Q_1Q_2 is $-c$, and the slope of Q_1Q_3 is $-c + H(\beta)/\ell$. Since the point (x_0, y_0) is below the line $y = -\ell x$ by our assumption, the triangle $Q_1Q_2Q_3$ is also below the line $y = -\ell x$ for a small ε . We choose such ε . (Fig. 3)




Fig. 3

If (x, y) is below the line $y = -\ell x$ and is in the rectangle $OP_2P_3P_4$, then by using (13) it is easy to see that any solution of (12) satisfies the following inequalities at the point (x, y) .

$$(14) \quad -c \leq \frac{dy}{dx} = -c - g(\psi_1(t)) \frac{x}{y} \leq -c + H(\beta)/\ell$$

By (14) and the definition of the triangle $Q_1Q_2Q_3$, we conclude that

any solution of (12) starting from a point in the triangle $Q_1Q_2Q_3$ can leave it only from its side Q_2Q_3 . By Property 7, there exists a t_k such that $(x(t_k), y(t_k))$ lies in the triangle $Q_1Q_2Q_3$. Therefore, $(x(t), y(t))$ must stay in the triangle $Q_1Q_2Q_3$ for all $t \geq t_k$.

By (12) and the construction of the triangle $Q_1Q_2Q_3$, $x'(t) = y(t) \leq y_0 + c\varepsilon < -\ell x_0 < 0$ for $t \geq t_k$. Therefore, $x(t)$ cannot be bounded below. This contradicts to $\lim_{t \rightarrow \infty} x(t) = x_0$.

By the above argument, $y_0 = 0$.

By the same argument as above, we can prove the following property.

PROPERTY 8. *Let $\{t_k\}$ be any sequence such that $\lim_{k \rightarrow \infty} t_k = \infty$ and $y(t_k)$ converges as k tends to infinity. Then $\lim_{k \rightarrow \infty} y(t_k) = 0$.*

Therefore, $\lim_{t \rightarrow \infty} y(t)$ exists and equal to 0. Thus, we have obtained the following equalities.

$$(15) \quad \lim_{t \rightarrow \infty} x(t) = x_0 > 0, \quad \lim_{t \rightarrow \infty} y(t) = 0$$

Let ε be a positive number such that

$$(16) \quad \varepsilon \leq \min \{ \delta x_0 / (2c), \delta x_0 \gamma / 2 \},$$

where δ and γ are given in Property 4.

By (15) there exists a number \bar{t} such that

$$(17) \quad |x(t) - x_0| < \varepsilon \quad \text{and} \quad |y(t)| < \varepsilon \quad \text{for } t \geq \bar{t}.$$

By Property 7 and (17), it follows that

$$(18) \quad 0 < x_0 \leq x(t) < x_0 + \varepsilon \quad \text{and} \quad -\varepsilon < y(t) \leq 0 \quad \text{for } t \geq \bar{t}.$$

By condition (B) of Property 4, there exists a number $\bar{t}_1 \geq \bar{t}$ such that $g'(\psi_1(t)) \geq \delta$ for $\bar{t}_1 \leq t \leq \bar{t}_1 + \gamma$. Therefore, by (16) and (18) we have the following inequalities for $\bar{t}_1 \leq t \leq \bar{t}_1 + \gamma$.

$$(19) \quad \begin{aligned} y'(t) &= -cy(t) - g'(\psi_1(t))x(t) \\ &< c\varepsilon - \delta x_0 \leq c\delta x_0 / (2c) - \delta x_0 \\ &= -\delta x_0 / 2 \end{aligned}$$

By the mean value theorem, (16), (18) and (19), we have the following inequalities.

$$y(\bar{t}_1 + \gamma) < y(\bar{t}_1) - (\delta x_0/2)\gamma \leq -\delta x_0\gamma/2 \leq -\varepsilon$$

But this contradicts to the latter half of (18).

By the above, we conclude that $x_0 = 0$, that is $\lim_{t \rightarrow \infty} x(t) = 0$.

Now, using a similar argument as above, we can prove that $\lim_{t \rightarrow \infty} y(t)$ exists and is equal to 0.

Thus, we finish the proof of Property 6, and this completes the proof of Theorem 2.

COROLLARY 1. *Assume the conditions A (i), A (ii) and A (iii). Let $A = \max \{|x_1 - 4E/c^2|, |x_2 + 4E/c^2|\}$, where x_1 and x_2 satisfy the equation $g(x_1) = -E$ and $g(x_2) = E$. Further, assume that $c^2 > H(A)$.*

Then every non-constant periodic solution of period $n\tau$ (n a positive integer) of the equation (11) is asymptotically stable.

Proof. This is an immediate consequence of Theorem 2 and Property 3.

COROLLARY 2. *In addition to the assumptions of Corollary 1, we assume that $e(t)$ is non-constant. Then there exists a non-constant periodic solution $x = \psi_1(t), y = \psi_2(t)$ of period τ for the equation (11) such that any periodic solution of period $n\tau$ (for a suitable positive integer n) for the equation (11) coincides with the solution $x = \psi_1(t), y = \psi_2(t)$.*

Proof. By Property 2 there exists a periodic solution $x = \psi_1(t), y = \psi_2(t)$ of period τ for the equation (11). Since $e(t)$ is non-constant by our assumption, there is no constant solution for the equation (11). Therefore, the periodic solution $x = \psi_1(t), y = \psi_2(t)$ is non-constant, and it may be considered as a periodic solution of period $n\tau$ for the equation (11) for an arbitrary positive integer n .

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be the Poincaré transformation associated with the equation (11). Then there is a one-to-one correspondence between the set of the periodic solution of period $n\tau$ for the equation (11) and the set of the fixed points of T^n (n times iterate of T).

By Corollary 1, every periodic solution of period $n\tau$ is asymptotically stable. Therefore, every fixed point of T^n is completely stable, and it is contained in a compact region D defined in the proof of Theorem 1. Since a completely stable fixed point is isolated, there are only a finite number of fixed points of T^n for a fixed n .

The index of a completely stable fixed point is $+1$, and the sum of all the indices of the fixed points of T^n is equal to 1 for every n . Therefore, there exists only one fixed point of T^n for each fixed n . This must correspond to the periodic solution $x = \psi_1(t)$, $y = \psi_2(t)$. This completes the proof of Corollary 2.

THEOREM 3. *Under the same assumption of Corollary 2 of Theorem 2, there exists a unique periodic solution $x = \psi_1(t)$, $y = \psi_2(t)$ of period τ for the equation (11) such that for any solution $x = x(t)$, $y = y(t)$ of (11) the following equalities hold.*

$$\lim_{t \rightarrow \infty} |x(t) - \psi_1(t)| = \lim_{t \rightarrow \infty} |y(t) - \psi_2(t)| = 0$$

That is any solution of (11) converges to a unique periodic solution of period τ of (11).

Proof. Let $x = \psi_1(t)$, $y = \psi_2(t)$ be the unique periodic solution of period τ for the equation (11), and let $x = x(t)$, $y = y(t)$ be any solution of (11). Set $z(t) = x(t) - \psi_1(t)$.

Since $x(t)$ and $\psi_1(t)$ are solutions of the equation (2), $z(t)$ satisfies the following equation.

$$(20) \quad z'' + cz' + g'(\delta(t))z = 0,$$

where $g'(\delta(t))z(t) = g(x(t)) - g(\psi_1(t))$ and $\delta(t)$ is a number between $x(t)$ and $\psi_1(t)$ for each t .

By our assumption $g'(\delta(t))$ satisfies the following inequality.

$$(21) \quad 0 \leq g'(\delta(t)) \leq H(A) \quad \text{for sufficiently large } t.$$

When Property 4 holds for $g'(\delta(t))$, then we can prove that $\lim_{t \rightarrow \infty} z(t) = \lim_{t \rightarrow \infty} z'(t) = 0$ as in the proof of Lemma 4. Therefore, it is sufficient to prove our theorem in case that Property 4 does not hold.

In this case, for any positive numbers δ and γ , there exists a number t_1 such that for any $t_2 > t_1$ there exists a number t_3 ($t_2 \leq t_3 \leq t_2 + \gamma$) such that $g'(\delta(t_3)) < \delta$. Therefore, there exists a sequence $\{t_k\}$ such that

$$\lim_{k \rightarrow \infty} t_k = \infty \quad \text{and} \quad \lim_{k \rightarrow \infty} g'(\delta(t_k)) = 0.$$

By the equality $g'(\delta(t))z(t) = g(x(t)) - g(\psi_1(t))$ and the boundedness

of $z(t) = x(t) - \psi_1(t)$ for sufficiently large t , we have $\lim_{k \rightarrow \infty} (g(x(t_k)) - g(\psi_1(t_k))) = 0$.

By our assumption, $|x(t)| \leq A, |\psi_1(t)| \leq A$ for sufficiently large t , and g is a homeomorphism between $[-A, A]$ and $[g(-A), g(A)]$. Therefore, $g^{-1}|[g(-A), g(A)]$ is uniformly continuous, and we can conclude that $\lim_{k \rightarrow \infty} (x(t_k) - \psi_1(t_k)) = 0$.

The equation (20) is equivalent to the following equation.

$$(22) \quad \begin{cases} z' = u \\ u' = -cu - g'(\delta(t))z \end{cases}$$

By a similar argument as in the first part of the proof of Lemma 4 using (21), we can prove that any solution of (22) is bounded for large t . Therefore, there exists a subsequence $\{t'_k\}$ of $\{t_k\}$ such that $\{u(t'_k)\}$ converges as k tends to infinity. Put $\lim_{k \rightarrow \infty} u(t'_k) = u_0$.

Since $\lim_{k \rightarrow \infty} z(t_k) = \lim_{k \rightarrow \infty} (x(t_k) - \psi_1(t_k)) = 0$, we have $\lim_{k \rightarrow \infty} z(t'_k) = 0$. Now, we shall prove that $u_0 = 0$, that is $\lim_{k \rightarrow \infty} u(t'_k) = 0$.

Suppose that $u_0 \neq 0$, say $u_0 > 0$. Then, using a similar construction as in the first part of the proof of Lemma 4, we can prove the following property.

PROPERTY 9. *There exists a positive number μ such that for any solution $z = z(t), u = u(t)$ of the equation (22) starting from a point sufficiently near the point $(0, u_0)$ the following inequality holds.*

$$u(t) \leq u_0 - \mu \quad \text{for sufficiently large } t.$$

But this contradicts our assumption $\lim_{k \rightarrow \infty} u(t'_k) = u_0$.

The case $u_0 < 0$ also leads to a contradiction. Thus, $u_0 = 0$. Therefore, we have proved that there exists a sequence $\{t'_k\}$ such that

$$(23) \quad \lim_{k \rightarrow \infty} z(t'_k) = \lim_{k \rightarrow \infty} u(t'_k) = 0.$$

By Corollary 2 of Theorem 2, $x = \psi_1(t), y = \psi_2(t)$ is an asymptotically stable periodic solution of (11). Therefore, any solution of (11) sufficiently near to the solution $(\psi_1(t), \psi_2(t))$ for some t must converge to the solution $(\psi_1(t), \psi_2(t))$ as t tends to infinity. And (23) is equivalent to the following (24).

$$(24) \quad \lim_{k \rightarrow \infty} (x(t'_k) - \psi_1(t'_k)) = \lim_{k \rightarrow \infty} (y(t'_k) - \psi_2(t'_k)) = 0$$

This completes the proof.

REFERENCES

- [1] Coddington-Levinson: Theory of ordinary differential equations, McGraw-Hill (1955).
- [2] C. Hayashi, Y. Ueda and H. Kawakami: Transformation theory as applied to the solution of non-linear differential equations of the second order, *Int. J. Non-Linear Mechanics*, **4** (1969), 235-255.
- [3] H. Kawakami: Qualitative study on the solutions of Duffing's equation, Thesis (1973), Kyoto University.
- [4] W. S. Loud: Boundedness and convergence of solutions of $x'' + cx' + g(x) = e(t)$, *Duke Math. J.*, **24** (1957), 63-72.

Department of Mathematics
College of General Education
Nagoya University