

A REMARK ON DIFFERENTIABLE STRUCTURES ON REAL PROJECTIVE $(2n-1)$ -SPACES

KENICHI SHIRAIWA

Dedicated to the memory of Professor TADASI NAKAYAMA

The main objective of this paper is to study the action of the group of differentiable structures Γ_{2n-1} on the $(2n-1)$ -sphere S^{2n-1} on the diffeomorphism classes on the real projective $(2n-1)$ -space P^{2n-1} by connected sum. This is done by considering universal covering spaces of the connected sum $P^{2n-1} \# \Sigma$, where Σ is an exotic $(2n-1)$ -sphere.

Throughout this paper all the manifolds considered are oriented, compact, and connected. Also the word *differentiable* is meant C^∞ -differentiable.

1. Let M, N be n -dimensional differentiable manifolds. If there exists an orientation preserving diffeomorphism of M onto N , then we shall denote it by $M \approx N$. The manifold M with orientation reversed is denoted by $-M$.

Let $M_1 \# M_2$ be the connected sum of two n -manifolds M_1 and M_2 . It is known that the connected sum operation is associative and commutative up to orientation preserving diffeomorphism. The sphere S^n serves as identity element (Cf. J. Milnor [2], M. Kervaire- J. Milnor [1]).

Let Σ be a smooth combinatorial n -sphere, which is called an exotic sphere. Then $\Sigma \# (-\Sigma) \approx S^n$. Thus the set of all the orientation preserving diffeomorphism classes of the exotic spheres forms a group under connected sum, which is denoted by Γ_n . We shall denote the class of Σ by $\{\Sigma\}$.

Let M be a differentiable n -manifold. Let Σ be an exotic n -sphere such that $M \# \Sigma \approx M$. Let $\Delta(M)$ be the subset of Γ_n consisting of the classes of such Σ .

PROPOSITION 1. $\Delta(M)$ is a subgroup of Γ_n , and $M \# \Sigma_1 \approx M \# \Sigma_2$, for exotic spheres Σ_1, Σ_2 , if and only if $\{\Sigma_1\} - \{\Sigma_2\} \in \Delta(M)$.

Proof. Let $\{\Sigma_1\}, \{\Sigma_2\} \in \Delta(M)$. Then

Received June 28, 1965.

$$M \# (\Sigma_1 \# \Sigma_2) \approx (M \# \Sigma_1) \# \Sigma_2 \approx M \# \Sigma_2 \approx M.$$

Also, $M \# (-\Sigma_1) \approx (M \# \Sigma_1) \# (-\Sigma_1) \approx M \# (\Sigma_1 \# (-\Sigma_1)) \approx M \# S^n \approx M$.

Thus $\mathcal{A}(M)$ is a subgroup of Γ_n .

Secondly, let $M \# \Sigma_1 \approx M \# \Sigma_2$. Then $M \# \Sigma_1 \# (-\Sigma_2) \approx M \# \Sigma_2 \# (-\Sigma_2) \approx M \# S^n \approx M$, that is $\{\Sigma_1 \# (-\Sigma_2)\} \in \mathcal{A}(M)$. Conversely, let $\{\Sigma_1 \# (-\Sigma_2)\} \in \mathcal{A}(M)$, then $M \approx M \# \Sigma_1 \# (-\Sigma_2)$. Adding Σ_2 from the right, we have $M \# \Sigma_2 \approx M \# \Sigma_1$.

Let $[G]$ be the order of a group G . Then we have the following.

Corollary. *The action of Γ_n on the orientation preserving diffeomorphism classes of a manifold M by connected sum is completely determined by $\mathcal{A}(M)$. In particular, the number of the orientation preserving diffeomorphism classes of M obtained by connected sum with exotic spheres is equal to $[\Gamma_n/\mathcal{A}(M)]$.*

2. Let M be a differentiable n -manifold such that the fundamental group $\pi_1(M)$ is a finite group of order $p = [\pi_1(M)]$. Let \tilde{M} be the universal covering space of M . Let N be a simply connected differentiable n -manifold.

PROPOSITION 2. *Under the above assumption and $n \geq 3$, the universal covering manifold of $M \# N$ is $\tilde{M} \# N \cdots \# N$ (p -factors of N).*

Proof. Let D^n be the unit disc in the euclidean n -space R^n . Let $f : D^n \rightarrow M$, $g : D^n \rightarrow N$ be differentiable imbeddings such that f preserves orientation and g reverses orientation. Then $M \# N$ is obtained from the disjoint sum

$$(M - f(0)) + (N - g(0))$$

by identifying $f(tx)$ with $g((1-t)x)$ for each $x \in S^{n-1} = \partial D^n$ and each $0 < t < 1$, where 0 is the origin of R^n .

Let $\pi : \tilde{M} \rightarrow M$ be the natural projection of the universal covering manifold \tilde{M} onto M . We can assume without loss of generality that $f(D^n)$ is contained in an open connected set U of M such that for each connected component U_i ($i = 1, 2, \dots, p$) of $\pi^{-1}(U)$, $\pi_i = \pi|_{U_i} : U_i \rightarrow U$ is a diffeomorphism onto. Let $f_i = \pi_i^{-1} \circ f : D^n \rightarrow U_i \subset \tilde{M}$. Then f_i is an orientation preserving diffeomorphism into.

Let N_1, \dots, N_p be p copies of N . We shall denote a point of N_i corresponding to $x \in N$ by x_i . Let $g_i : D^n \rightarrow N_i$ ($i = 1, 2, \dots, p$) be copies of g , that is $g_i(x) = (g(x))_i$. Then $\tilde{M} \# N_1 \cdots \# N_p$ is obtained from the disjoint sum

$$(\tilde{M} - \bigcup_{i=1}^p f_i(0)) + (N_1 - g_1(0)) + \cdots + (N_p - g_p(0))$$

by identifying $f_i(tx)$ with $g_i((1-t)x)$ for each $x \in S^{n-1}$, each $0 < t < 1$, and $i = 1, 2, \dots, p$.

Let $\pi' : M \# N_1 \# \dots \# N_p \rightarrow M \# N$ be the map defined by

$$\begin{aligned} \pi'(x) &= \pi(x) && \text{for } x \in (\tilde{M} - \bigcup_{i=1}^p f_i(0)) \text{ and} \\ \pi'(x_i) &= x && \text{for } x_i \in N_i - g_i(0), (i = 1, 2, \dots, p). \end{aligned}$$

Then π' is well defined and a local diffeomorphism onto $M \# N$. Thus, $\tilde{M} \# N_1 \# \dots \# N_p$ is a covering manifold of $M \# N$.

On the other hand, $\tilde{M} \# N_1 \# \dots \# N_p$ is simply connected as is seen by the following lemma. Therefore, the proposition is proved.

Lemma. *Let M, N be simply connected n -manifolds and $n \geq 3$. Then $M \# N$ is simply connected.*

Let $f : D^n \rightarrow M, g : D^n \rightarrow N$ be imbeddings as before. Then the connected sum $M \# N$ is obtained topologically from the disjoint sum $(M - f(\text{Int } D^n)) + (N - g(\text{Int } D^n))$ by identifying $f(x)$ with $g(x)$ for $x \in S^{n-1}$, where $\text{Int } D^n$ is the interior of D^n . Let K (resp. L) be the image of $(M - f(\text{Int } D^n))$ (resp. $N - g(\text{Int } D^n)$). Then $M \# N = K \cup L$ and $K \cap L$ is homeomorphic to S^{n-1} by f (or g). Thus $K \cap L$ is simply connected. Since K and $M - f(0)$ have the same homotopy type, K is simply connected by our assumption that M is simply connected and $n \geq 3$. By the same reason L is simply connected. It follows from the van Kampen's theorem that $M \# N = K \cup L$ is simply connected.

Let S^{2n-1} be the unit sphere in complex n -space C^n . That is a point of S^{2n-1} has a form (z_1, \dots, z_n) , where z_i is a complex number and $\sum_{i=1}^n z_i \bar{z}_i = 1$. Let $\lambda = \exp(2\pi i/p)$, where p is a natural number. Let $T : S^{2n-1} \rightarrow S^{2n-1}$ be transformation defined by $T(z_1, \dots, z_n) = (\lambda^{q_1} z_1, \dots, \lambda^{q_n} z_n)$, where q_1, \dots, q_n are integers prime to p . Then T is a fixed point free transformation of period p . The orbit space $S^{2n-1}/T = L_p(q_1, \dots, q_n) = L_p$ is an oriented differentiable manifold in a natural way. The natural projection $\eta : S^{2n-1} \rightarrow L_p$ is the universal covering. If $p = 2, L_2 = P^{2n-1}$, the real projective $(2n-1)$ -space. It is well known that $\pi_1(L_p) = Z_p$, the group of integers modulo p .

COROLLARY. *The universal covering space of $L_p \# \Sigma$, where Σ is an exotic*

sphere, is $S^{2n-1} \# \Sigma \# \cdots \# \Sigma \approx \Sigma \# \cdots \# \Sigma$ (p -factors of Σ) ($n \geq 2$).

In particular, the universal covering space of $P^{2n-1} \# \Sigma$ is $S^{2n-1} \# \Sigma \# \Sigma \approx \Sigma \# \Sigma$.

In conclusion, we have the following theorem.

THEOREM. *There exist at least $[2\Gamma_{2n-1}]$ distinct differentiable structures on P^{2n-1} up to orientation preserving diffeomorphism, where $[2\Gamma_{2n-1}]$ is the order of $2\Gamma_{2n-1}$.*

Also, there exist at least $[p\Gamma_{2n-1}]$ distinct differentiable structures on a lens space L_p .

Proof. Let Σ be an exotic sphere such that $P^{2n-1} \# \Sigma \approx P^{2n-1}$. Then the universal covering spaces must be diffeomorphic, that is $\Sigma \# \Sigma \approx S^{2n-1}$. Thus $\Delta(P^{2n-1}) \subset {}_2\Gamma_{2n-1}$, where ${}_2\Gamma_{2n-1}$ is the subgroup of Γ_{2n-1} consisting of the elements of order two. Thus

$$[\Gamma_{2n-1}/\Delta(P^{2n-1})] \geq [\Gamma_{2n-1}/{}_2\Gamma_{2n-1}] = [2\Gamma_{2n-1}].$$

Our theorem follows from the fact that $P^{2n-1} \# \Sigma$ is homeomorphic to P^{2n-1} and the Corollary of Proposition 1.

The same argument shows the latter half of the theorem.

According to the work of Kervaire-Milnor [1], Γ_{4m-1} ($m \geq 2$) contains a cyclic group of order $2^{2m-2} (2^{2m-1} - 1) \cdot \text{numerator}(4B_m/m)$, where B_m denotes the m -th Bernoulli number. Therefore, there exist at least $2^{2m-3} (2^{2m-1} - 1) \cdot \text{numerator}(4B_m/m)$ distinct differentiable structures on P^{4m-1} ($m \geq 2$). For example, 14 on P^7 , 496 on P^{11} , 4064 on P^{15} , and etc..

REFERENCES

- [1] M. Kervaire-J. Milnor, Groups of homotopy spheres, I, *Ann. of Math.*, **77** (1963), 504-537.
- [2] J. Milnor, Sommes de variétés différentiables et structures différentiables des sphères, *Bull. Soc. Math. France* **87** (1959), 439-444.
- [3] M. Hirsch-J. Milnor, Some curious involutions of spheres, *Bull. A.M.S.* **70** (1964), 372-377.

Nagoya University